

VOL 96 No. 05 MAY, 2019

Print ISSN 0019 - 6479

E - ISSN 0974 - 9365

₹. 80

THE INDIAN VETERINARY JOURNAL

SINCE - 1924

Journal of the
INDIAN VETERINARY ASSOCIATION

ESTD - 1922

Regd. No. Sl. No. 96/1967



No. 11, Pasumpon Muthuramalinga Thevar Salai (Chamiers Road),
Nandanam, Chennai - 600 035, Tamil Nadu, India

Tel. : +91 44 2435 1006

Email : ivj83@yahoo.com

ONLINE : www.ivj.org.in

THE INDIAN VETERINARY JOURNAL

(Official Organ of the Indian Veterinary Association)

EDITORIAL COMMITTEE

Dr A.V. KRISHNAN, Chief Editor
B.V.Sc., M.V.Sc (Path.)

Dr S. SUKUMAR, Managing Editor
B.V.Sc., M.V.Sc (Vet. Micro), Ph.D. (Biotech)

Dr V. Titus George, Editor
B.V.Sc., M.V.Sc., Ph.D. (Patho)

Dr I. Ponnu Pandian, Editor
B.V.Sc

Dr K. Venukopalan, Editor
B.V.Sc., M.V.Sc., Ph.D. (Poul.)

EDITORIAL BOARD

CHAIRMAN

Dr. Chirantan Kadian
B.V.Sc & A.H.
President, Indian Veterinary Association

MEMBERS

Prof. Dr C. Balachandran
M.V.Sc, Ph.D., PGDAJ, PGDEVP, DICVP
FAO Fellow, FIAVP, FNAVS, FASAW
Vice-Chancellor
Tamilnadu Veterinary and Animal Sciences University
Madhavaram Milk Colony, Chennai - 600 051.

Prof. Dr. Gurdial Singh
B.V.Sc & A.H, MVSc, Ph.D, FNAVS, FIAVA
Vice Chancellor, LUVAS, Hisar.

Dr. P.K. Panwar
BVSc & AH, MVSc,
Department of Animal Husbandry
Uttar Pradesh

Dr. Subhash Chander Ahlawat
BVSc & AH.
Department of Animal Husbandry
Haryana

MANAGING COMMITTEE OF INDIAN VETERINARY ASSOCIATION

President :
Dr. Chirantan Kadian

Secretary General :
Dr. R.S. Patel

Treasurer :
Dr. I. Ponnu Pandian

Joint Secretary :
Dr. M.K. Srikanth

Vice Presidents :
Dr. L. Chandrasekaran, HQ, Tamilnadu
Dr. Arun Sirkeck, Himachal Pradesh

Dr. S.G. Yalagod, Karnataka
Dr. Ilang, Nagaland
Dr. Ramji Lal Meena, Rajasthan
Dr. Sudhir Kumar, Uttar Pradesh

Zonal Secretaries :
Dr. Amit Nain, Punjab
Dr. R. Vasantha Rayalu, Andhra Pradesh
Dr. Bhuban Ch. Sarmah, Assam
Dr. Umesh Balaji Sontakke, Maharashtra
Dr. Pankaj Kumar, Jharkhand

Managing Editor, IVJ :
Dr. S. Sukumar, Tamil Nadu

**LIFE MEMBERS OF INDIAN VETERINARY ASSOCIATION
(2013 ONWARDS)**

274	Dr Dilip Kumar	Bihar	288	Dr Asish Ashokrao Pawar	Maharashtra
275	Dr Dharmvir Kumar	Bihar	289	Dr Nameeta Nitin Nadkarni	Maharashtra
276	Dr Bam Shankar Kumar	Bihar	290	Dr Loveleen Maria Vaz	Goa
277	Dr Kamlesh Kumar Singh	Bihar	291	Dr Dasudi Nagesh Ramarao	Karnataka
278	Dr Raghu Nath Kohli	Delhi	292	Dr Renu Gupta	Haryana
279	Dr Surendra Kishore Jain	Uttar Pradesh	293	Dr Jagmohan Singh	Chhattisgarh
280	Dr Shalini Parihar	Uttar Pradesh	294	Dr (Col.) Mudgal	Haryana
281	Dr Suman Biswas	West Bengal	295	Dr Sha Arun	Karnataka
282	Dr Kunal Batabyal	West Bengal	296	Prakasha Rao	Andhra Pradesh
283	Dr (Col.) Surinder Singh	Chandigarh	297	Surender Singh	Delhi
284	Dr Ninan Jacob	Puducherry	298	Suresh Kumar	Delhi
285	Dr Sudheer B. Arumbaka	Andhra Pradesh	299	Mitra Ranjan Panda	Odisha
286	Dr Jyoti Ranjan Mohanty	Odisha	300	Gunaram Saikia	Assam
287	Dr Pardeep Kumar Yadav	New Delhi	301	Bhabatosh Pal	Himachal Pradesh

Associate Member of IVA - 2001

M/s. Indovax Private Limited, Corporate Heights,
SCO - 24, Sector - 14, Gurgaon - 122 001.

Corporate Member of IVA - 2000

M/s. Indian Immunologicals Ltd.,
Road No.44, Jubilee Hills,
Hyderabad - 500 033.

Corporate Member of IVA - 2008

M/s. Vetnex RFCL Ltd.,
A-3, Okhla Industrial Area, Phase-1,
New Delhi - 110 020.

Corporate Member of IVA - 2013

BOVIAN HEALTHCARE PVT. LTD.
501, SSR Corporate Park,
13/6 Mathura Road, NH - 2.
FARIDABAD - 121003 (Haryana)

Nanobubble Aquaculture System: Its Effect Towards Immune Response and Infection of *Vibrio* sp. in Vannamei Shrimp (*Litopenaeus vannamei*)

Gunanti Mahasri¹, Laksmi Sulmatiwi, Sudarno, Prayogo,
Galang Damar Pamenang and Ade Irmalia Harifa

Fisheries and Marine Faculty, Universitas Airlangga, Indonesia

(Received : November, 2018 406/18 Accepted : January, 2019)

Abstract

The application of nanobubble had a significant effect on the Vannamei shrimp (*Litopenaeus vannamei*) immune response against *Vibrio* sp. with the best THC and DHC results (45.23 cells/ml and 30.23%). The highest average TPC of *Vibrio* sp. in shrimp hepatopancreas was on aerator treatment, which was 3.32×10^5 (CFU / g). The highest average TPC of *Vibrio* sp. was on aerator treatment, which was 3.43×10^6 (CFU / ml). The results of bacterial identification showed that the types of bacteria were *Vibrio vulnificus* and *Vibrio alginolitycus*.

Key words: Nanobubble, Immune Response, Pathogen Infection, Vannamei Shrimp

Water quality is one of the key factors of aquaculture success (Effendi, 2004). Bad water quality gives an opportunity to the *Vibrio* sp. infecting Vannamei shrimp health (Lorenzon *et al.*, 2001; Supito *et al.*, 2008). Nanobubble is technology can be used to improve water quality by decomposing the organic matter (Ebina *et al.*, 2013; Chiba and Takahashi, 2007; Mahasri, 2016). The purpose of this study was to determine the effect of nanobubble aquaculture systems against immune response and *Vibrio* sp. infection on Vannamei shrimp (*Litopenaeus vannamei*).

Materials and Methods

The materials used in this study were 240 Vannamei shrimp size 7-10 cm with a weight of 4-6 gr, pond water with salinity 18-20 ppt, aerator and nanobubble generator. This study consisted of 2 factors, A (nanobubble and aerator) and B (0 days, 10 days, 20 days, and 30 days). The optimization used in this study is

for 1 hour in the morning at 6 am and in the afternoon at 4 pm.

The Vannamei shrimp blood was charged on the hemocytometer and then counted using a hand counter and with its calculation formula for Total Haemocyte Count (THC) (Brock *et al.*, 1991). Making blood-thinning preparations was to take the blood of Vannamei shrimp with a 1 ml syringe that has been given EDTA as an anticoagulant (Mahasri, 2007; Darwantin *et al.*, 2016). The number of Differential Haemocyte Count (DHC) cells was calculated to be up to 100 cells and the percentage was sought (Martin *et al.*, 1985).

The TPC calculation method was carried out aseptically on the laminar flow by means of samples that had been dissolved in sterile physiological NaCl taken 100 μ l and diluted multi-level (10⁻¹ to 10⁻⁵ dilutions), then 100 μ l was taken from 10⁻³, 10⁻⁴ dilutions and 10⁻⁵ using the Duplo spread plate method on Nutrient agar media. The next step was incubating the incubator for 18-24 hours at room temperature (28-30°C) and after that, the colony was counted which grows based on SNI 01-2332.3-2006 (Kusdarwati *et al.*, 2016).

Bacterial isolation was carried out aseptically on laminar flow with the streak plate method, by taking one sample which had been dissolved in sterile physiological NaCl and scratched on TCBS media agar and incubated for 18-24 hours at room temperature (28-30°C). The colonies that grew later would be chosen based on differences in colour and size of the colonies that appeared purified. Purification was done by taking a culture selected and grown on NA saline (+ 2% NaCl) with the streak plate method. The purified culture incubated

¹Corresponding author : Email : mahasritot@gmail.com

Table I. THC and DHC of Vannamei shrimp treated by aerator and nanobubble

Technology	Day	THC	DHC
Nanobubble	0	30	18.89
	10	40.68	22.68
	20	41.5	27.65
	30	45.23	30.23
Aerator	0	30	17.98
	10	27.6	16.6
	20	33	17.61
	30	29.63	15.6

Table II. TPC from Vannamei shrimp hepatopancreas and water media treated by aerator and nanobubble.

Technology	Day	TPC	
		Hepatopancreas	Water
Nanobubble	0	30.5	3.05
	10	18.75	1.525
	20	12.875	1.0475
	30	3.3	0.33
Aerator	0	33.25	3.425
	10	27	2.55
	20	10.45	0.9575
	30	16	1.35

for 18-24 hours at room temperature (28-30°C) (Letchumanan *et al.*, 2015). The pure colonies are then used for bacterial identification (Lubis *et al.*, 2014). Identification of bacteria was done by morphological observations, Gram staining, and biochemical testing.

Results and Discussion

The results of the THC and DHC could be seen in Table I which showed a significant difference between nanobubble treatment with aerators ($p < 0.05$).

THC and DHC in Vannamei shrimp with nanobubble treatment increased because, in nanobubble systems, oxygen in the waters could last longer so that it is utilized for the biological activity of organisms. Nanobubble systems produced dissolved oxygen for a long time so that it met the needs of organisms and decomposition of organic matter (Chiba and Takahasi, *loc. cit.*). An increase in DHC indicates that the shrimp has improved health status.

TPC from shrimp hepatopancreas and in water media showed in Table II.

TPC on the aerator treatment showed a poor value of water quality than nanobubble since the ammonia and nitrite values were above the normal range for Vannamei shrimp. Poor water quality could cause stress on cultivated shrimp so that the growth of shrimp would be disrupted and reduced the body's resistance to infection and cause death (Diansari *et al.*, *loc. cit.*).

The results of the identification of *Vibrio* sp. on Vannamei shrimp and water media could be seen in Table III.

The biochemical test results showed that the bacterial samples were *Vibrio vulnificus* and *Vibrio alginolyticus*. Several species of *Vibrio* bacteria were reported to be disease agents that caused a decrease in shrimp production, namely *V. vulnificus*, *V. splendidus* and *V. damsela* infection in *Penaeus monodon* larvae and *Vibrio alginolyticus* infection in *Penaeus orientalis* larvae (Wang *et al.*, 2005).

Nitrite and ammonia levels in the aerator treatment tended to be high, 1.25 ppm and 1.5 - 1.7 ppm. It was considered above the safe limit for Vannamei shrimp cultivation. The quality

Table III. Presence of *Vibrio* sp. in Vannamei shrimp treated by aerator and nanobubble

No.	Technology	Sample	Bacteria
1	Aerator	Water	<i>Vibrio vulnificus</i> (99,64%)
		Hepatopancreas	<i>Vibrio alginolitycus</i> (98,77%)
2	Nanobubble	Water	<i>Vibrio alginolitycus</i> (52,04%)
		Hepatopancreas	<i>Vibrio vulnificus</i> (89,58%) <i>Vibrio alginolitycus</i> (82,41%)

Table IV. Water Quality of Vannamei shrimp culture treated by aerator and nanobubble

Parameter	Technology	
	Nanobubble	Aerator
DO (ppm)	3,9 – 10,8	2,81 – 4,65
Temperature (°C)	27 – 29	27 – 29
Salinity (ppt)	19 – 20	19 – 20
pH	8 – 8,6	7,2 – 7,8
Nitrite (ppm)	0,57	1,25
Ammonia (ppm)	0,006	1,5 – 1,7

of nanobubble water was more stable compared to ordinary bubbles (Ebina *et al.*, *loc. cit.*). If air bubbles were not easily broken, the dissolved oxygen content in the waters would also be more stable.

Summary

Nanobubble could reduce the *Vibrio* sp. infection in Vannamei shrimp culture and maintain the water quality.

References

Brock, T., M. Madigan, J. M. Martinko and J. Parker, (1991) *Biology of microorganisms*, Prentice Hall, New Jersey, 6th Edition, 1104 p.

Chiba, K. and M. Takahashi, (2007) *U.S. Patent Application No. 10/591,979*.

Darwanti, K. and R. Sidik, (2016) Efisiensi Penggunaan Imunostimulan dalam Pakan Terhadap Laju Pertumbuhan, Respon Imun dan Kelulushidupan Udang Vannamei (*Litopenaeus vannamei*), *Jurnal Biosains Pascasarjana*, **18**(2).

Diansari, R. R., E. Arini and T. Elfitasari, (2013) Pengaruh Kepadatan Yang Berbeda Terhadap Kelulushidupan Dan Pertumbuhan Ikan Nila (*Oreochromis niloticus*) Pada Sistem Resirkulasi Dengan Filter Zeolit, *JAMT*, **2**(3) : 37-45.

Ebina, K., Shi, K., Hirao, M., Hashimoto, J., Kawato, Y., Kaneshiro, S., Morimoto, T., Koizumi, K. and Yoshikawa, H. (2013). Oxygen and air nanobubble water solution promote the growth of plants, fishes, and mice. *PLoS One*, **8**(6) : e65339.

Ebina, K., K. Shi, M. Hirao, J. Hashimoto, Y. Kawato, S. Kaneshiro, T. Morimoto and I. Effendi, (2004) Pengantar Akuakultur. Penebar Swadaya, Jakarta, Vol 1, 184 p.

Kusdarwati, R, Sudarno, Rozi and D. D. Nindarwi, (2016) Petunjuk Praktikum Analisis Penyakit Ikan I, Fakultas Perikanan dan Kelautan Universitas Airlangga, Surabaya, 23-34 p.

Lorenzon, S., M. Francese, V. J. Smith and E. A. Ferrero, (2001) Heavy metals affect the circulating haemocyte number in the shrimp *Palaemon elegans*, *Fish & shellfish immunology*, **11**(6) : 459-472 p.

Leichumanan, V., W. F. Yin, L. H. Lee and K. G. Chan, (2015) Prevalence and antimicrobial susceptibility of *Vibrio parahaemolyticus* isolated from retail shrimps in Malaysia, *Frontiers in microbiology*, **6** : 33.

Lubis, D. A., H. Syawal and M. Riauaty, (2014) Identifikasi Bakteri Patogen pada Ikan Nila (*Oreochromis niloticus*) di Kecamatan Marpoyan Damai Kota Pekanbaru, *Jurnal Dinamika Pertanian*, **2** : 12-15.

Mahasri, G., (2007) Protein membrane imunogenik zoothamnium penaei sebagai bahan pengembangan imunostimulan pada udang windu (*Penaeus monodon fabricus*) terhadap zoothamniosis, Program Pasca Sarjana Universitas Airlangga Surabaya, Surabaya.

Mahasri, G. (2016) Pengembangan teknologi nanobubble dan sistem budidaya terpadu pada nila dan ikan lele sebagai upaya pengendalian penyakit dan peningkatan produksi perikanan. Laporan Akhir Intensif Riset Sinas RT 10.04. Lembaga Penelitian dan Inovasi Universitas Airlangga Surabaya, Surabaya.

Supito, D., Adiwijaya, J. Maskar and S. Damang, (2008) Teknik Budidaya Udang Windu Intensif dengan *Green Water System* Melalui Penggunaan Pupuk Nitrat dan Penambahan Sumber Karbon, *Jurnal Perikanan*, **7** : 38 – 53.

Wang, Y. B., Z. R. Xu and M. S. Xia, (2005) The effectiveness of commercial probiotics in northern white shrimp *Penaeus vannamei* ponds, *Fisheries science*, **71**(5) : 1036-1041.

Linko, K. and Paloheimo, M. (1989) Monitoring of the inspired and end-tidal oxygen, carbon dioxide and nitrous oxide concentrations: clinical applications during anaesthesia and recovery. *J. Clin. Monit. Comput.*, 5: 149-156.

Ramakrishnan, K.S. (2012) Evaluation of buprenorphine and butorphanol as preanaesthetic in midazolam-ketamine-isoflurane anaesthesia for ovariohysterectomy in cats. M.V.Sc. thesis submitted to The Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamil Nadu.

Senthilkumar, S. (2013) Cardiopulmonary and stress response to low flow isoflurane anaesthesia with opioid analgesics in cattle. Ph.D. thesis submitted to Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamilnadu.

Thangadurai, R. (2012) General anaesthesia with intermittent positive pressure ventilation for thorocotomy in cattle with

traumatic pericarditis. Ph.D. thesis submitted to The Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamil Nadu.

Tranquilli, W.J., Thurmon, J.C. and Grimm, K.A. (2007) Lumb and Jones' Veterinary Anesthesia and Analgesia. 4th edn., Blackwell Publishers, Iowa, USA. pp. 513.

Turner, P., Kerr, C.L., Healy, A. J. and Taylor, M. (2006) Effect of meloxicam and butorphanol on minimum alveolar concentration of isoflurane in rabbits. *Am. J. Vet. Res.*, 67: 770-774.

Yamashita, K., Okano, N., Yamshita, M., Umar, M.A., Kushiro, T. and Muir, W.W. (2008) Effects of carprofen and meloxicam with or without butorphanol on minimum alveolar concentration of sevoflurane in dogs. *J. Vet. Med. Sci.*, 70: 29-35.

Indian Vet. J., May 2019, 96 (05) : 37 - 39

Oxygen Dissolved Nanobubble Technology Improved the Quality of Pacific White Shrimp Cultivation

Gunanti Mahasri¹, Ade Irmalia Harifa and Sudarno

Department of Fish Health Management, Faculty of Fisheries and Marine, Universitas Airlangga, Surabaya 60115, Indonesia.

(Received : October, 2018 358/18 Accepted : December, 2018)

Abstract

This study aims to determine the effect of nanobubble on dissolved oxygen to reduce Presumptive Vibrio Count (PVC) in Pacific White Shrimp (*Litopenaeus vannamei*) aquaculture system. The results of the analysis showed that the use of nanobubble had a significant effect ($p < 0,05$) in dissolved oxygen and reduce PVC while it had no significant difference in PVC on aquaculture media. The use of nanobubble technology provided optimal dissolved oxygen for Pacific White Shrimp aquaculture and lower PVC value of *Vibrio* sp. in the hepatopancreas compared to aerators treatment.

Key words: Nanobubble, Dissolved Oxygen, Quality, Pacific White Shrimp

The high loss due to *Vibrio* sp. bacte-

rial infection is one of the challenges on Pacific White Shrimp. One of the technologies developed to improve the quality of aquaculture is nanobubble since it can be made available for a long time so as to maintain the dissolved oxygen in the waters remains stable and related to the respiratory and metabolic systems (Mahasri et al., 2018; Takahashi and Chiba, 2007). The purpose of this study was to determine its effect on dissolved oxygen (DO) and Presumptive Vibrio Count (PVC) on Pacific White Shrimp.

Materials and Methods

This study used 240 of Pacific White Shrimp (7-10 cm, 4-6 g) and 30 of each treatment. The treatments were A1 (nanobubble), A2 (aerator), B1 (0 days), B2 (10 days), B3 (20 days) and B4 (30 days). The dissolved oxygen was measured by DO meter tool. PVC was a calculation method to determine the number of *Vibrio* sp. bacteria

¹Corresponding author : Email : mahasritot@gmail.com