

EFFECT OF DIFFERENT CONCENTRATION OF ALPHA TOCOPHEROL SUPPLEMENTATION ON HUMAN BONE MARROW MESENCHYMAL STEM CELL VIABILITY (IN VITRO STUDY)

by Ni Putu Mira Sumarta

Submission date: 28-Apr-2021 09:55AM (UTC+0800)

Submission ID: 1571936235

File name: 4._AJM-26.pdf (156.29K)

Word count: 1518

Character count: 8637

EFFECT OF DIFFERENT CONCENTRATION OF ALPHA TOCOPHEROL SUPPLEMENTATION ON HUMAN BONE MARROW MESENCHYMAL STEM CELL VIABILITY (*IN VITRO* STUDY)

JEFRY WAHYUDI S¹, NI PUTU MIRA SUMARTA^{2*}, GATOT BAYDOWI¹ AND COEN PRAMONO D.²

²
(Received 25 September, 2018; accepted 15 November, 2018)

Key words : Autogenous bone grafts, Bone marrow mesenchymalstem cell, Alpha tocopherol, Viability

Abstract– Bone defect can be reconstructed with autogenous bone graft and considered as an ideal for bone graft material. The process of bone regeneration that occurs because of the proliferation and differentiation of human bone mesenchymal stem cell marrow. Alpha tocopherol is a co-factor that can help osteogenesis as micronutrient supplementation. Biocompatibility of this supplement still needs to be proven. This study is to determine the effect of alpha tocopherol supplementation on human bone marrow mesenchymal stem cell viability. Experimental studies performed in vitro on human bone marrow mesenchymal stem cell. This study was divided into 2 main groups: alpha tocopherol concentrations group (25 µM/mL, 50 µM/mL, 75 µM/mL) and control group. Each group is analyzed with MTT assay observed living cells. Alpha tocopherol with 25 µM/mL concentration obtained high results (107,86%) than other samples ($p > 0,05$ in all subjects observed). Optimum concentration of Alpha tocopherol is 25 µM/ml.

INTRODUCTION

Bone defects could be treated by reconstruction procedure using one's own body termed as autogenous bone graft (Randall M, 2004). Autogenous bone graft is considered as an ideal bone graft because of its osteogenic characteristic because it contains many living cells, osteoconductive meaning that it has a woven structure that functions as a scaffold for a growth media or host tissue vascular regeneration, and also has osteoinductive property which is the ability to secrete growth factors that could induce proliferation and differentiation processes of the bone marrow mesenchymal stem cell to turn into osteoprogenitor cells and has the ability to stimulate osteoprogenitor cells to differentiate into osteoblast and could start the process of new bone formation (Axelrad *et al.*, 2009; Choi, 2008).

The failure of bone graft could happen because of periosteum damage thus reconstruction with non-vascularized autogenous bone graft often fails

marked by graft resorption or incorporation between grafts and recipient bone do not occur (Pramono, 2011). The proliferation stage is the early stage to examine cell viability. Because of this condition the role of growth factor and co factor with its osteoinductive property holds an important role to increase proliferation process and bone marrow mesenchymal stem cell tissue differentiation. Alpha tocopherol is a cofactor that could help supplement micronutrients in the bone marrow mesenchymal stem cell culture media in the osteogenesis process (Granjeiro *et al.*, 2005).

Alpha tocopherol could increase bone marrow mesenchymal stem cell proliferation, inducing osteoblast differentiation, and increase collagen fiber formation (Pradel *et al.*, 2008). Biocompatibility of this supplement still needs to be proven. This study is an in vitro experimental study with alpha tocopherol concentrations starting 25µM/mL, 50 µM/mL, 75 µM/mL with the aim to determine the effect of alpha tocopherol supplementation on human bone marrow mesenchymal stem cell

*Corresponding author's email: niputu.mira@fkg.nair.ac.id

viability.

MATERIALS AND METHODS

This study was a true experimental laboratory study divided into 2 groups, the treatment group, examining the impact of alpha tocopherol supplementation with concentrations of 25 µM/mL, 50 µM/mL, and 75 µM/mL, and the control group, without alpha tocopherol towards human bone marrow mesenchymal stem cell (hBMSC) viability *in vitro*. The sample was taken from human bone marrow mesenchymal stem cell stock with 3 times passages at Stem Cell Research and Development Center in the Institute of Tropical Disease Center, Airlangga University. Every treatment group consisted of 7 samples.

Thawing process was conducted on human bone marrow mesenchymal stem cell until monolayer was formed on a petridish after 7 days. The monolayer was split, then moved into 96 wheel microplate with every wheel given about 5×10^4 cells with α -MEM 100 mL, incubated until the cell reached 80% confluence. After the cell achieved 80% confluence, it was administered with staged Alpha tocopherol (25 mM/mL, 50 mM/mL, 75 mM/mL) and α -MEM 50 iM/mL then incubated for 20 hours in 37°C. MTT Assay was done after 4 hours of incubation, optical density (OD) was evaluated using Elisa Reader, and viable cells were counted using formula below:

$$\% \text{ living cells} = \frac{\text{OD treatment} + \text{OD media} \times 100\%}{\text{OD cell control} + \text{OD media}}$$

Note: % living cells : The percentage of living cells after the examination.

OD treatment : OD value in every sample after Elisa Reader result examination.

OD media : Optical density stem cell value in control media.

OD cell control : Optical density stem cell value in control cell.

Toxicity was determined when cell viability is <60% (Freshney, 2000). The data was analysed statistically using One-way ANOVA, $p < 0.05$ was considered significantly different.

RESULT

The result of this study also showed that all treatment and control groups revealed no living cell percentage above 60% which indicated that none of

the groups has cytotoxicity (Figure 1). The decline of living cells percentage were documented in 50 µM/mL and 75 µM/mL groups, however they were not significantly different ($p=0.682$; $p > 0.05$).




Fig. 1. The amount of living cells percentages showed bone marrow mesenchymal stem cell viability in each concentration.

DISCUSSION

Mesenchymal stem cells are progenitor cells that have the ability of differentiation. Mesenchymal stem cells could originate from fibroblastic cells, muscle cells, bone cells, tendons, ligaments and fatty tissues (Kaveh, 2011; Banfi, 2009). Alpha tocopherol induced the differentiation of mesenchyme cells including fat cells, osteoblasts, myoblasts and chondrocytes. Alpha tocopherol deficiency can decrease the proliferation of chondrocytes, the number of cells to osteoblasts and disturbance towards the synthesis of matrix (Tecla M et al., 2010). This study was conducted to determine the effect of alpha tocopherol supplementation on human bone marrow mesenchymal stem cell viability. Alpha tocopherol has an important role in collagen synthesis and marks differentiation into osteoblast (Pradel, 2008).

The result of the study proved that various concentrations (25 µM/mL, 50 µM/mL, and 75 µM/mL) of Alpha tocopherol could influence Bone Marrow Mesenchymal Stem Cell viability. Upon the concentration of 25 µM/mL, the percentage of living cells was 107.86%; highest amongst the other concentrations. This is possibly a result of the characteristic of Alpha tocopherol as a lipid soluble antioxidant thus preventing free radicals in the cell membrane. In addition, alpha tocopherol could increase the proliferation of mesenchymal stem cells inducing osteoblast differentiation and increasing collagen fibres production. The administration of Alpha tocopherol 50 µM/mL caused decline to the percentage of living cells due to the stationary phase or decrease in cell count in compliance to the cell

cycle. Administration of a relatively high dose of alpha tocopherol would influence the presence of fatty acid in the mitochondria membrane. This is in accordance to the study by Lauridsen and Jansen (2011) which stated that administration of alpha tocopherol dose needs to be considered because of the variation in composition of membrane fatty acid and its relation with gene regulation.

CONCLUSION

Various concentrations of alpha tocopherol influenced hBMSC viability. Alpha tocopherol at 25 µM/ml concentration showed the highest percentage of living cells.

REFERENCES

- Axelrad, W., Kakar, S. and Einhorn, T. 2009. Biological and biophysical technologies for enhancement of fracture repair. In : Bucholz R.W., Heckman, J.D., Court-Brown, C., Tornetta, P., Koval, J.K., Wirth, M.A. *Rockwood and Green's Fractures in Adults*. (Lippincott william & wilkins). p 105.
- Banfi A, Muraglia A and Dozin B. 2000. Proliferation kinetics and differentiation potential of ex vivo expanded human bone marrow stromal cells: implication for their use in cells therapy, *Exp. Hematology*, 28: pp 707-715.
- Choi, K.M. 2008. Effect of ascorbic acid on bone marrow derived mscs proliferation and differentiation, department and engineering, dongguk university, Korea. pp 586-587.
- Freshney, I.R. 2000. *Culture of Animal Cells Manual of Basic Technique*. New York: A John Wiley and Sons Ltd.
- Granjeiro J, Oliver R, Bustos-Valenzuel J, Sogayar M, and Taga R. 2005. Bone morphogenetic proteins ; from structure to clinical use, *braz j med & biol res*. pp 1463-1473
- Kaveh, K. 2011. Mesenchymal stem cell, osteogenic lineage and bone tissue engineering, *journal of animal and veterinary advances*. 10 (17) : 2317-2330.
- Lauridsen, C. and Jansen, S.K. 2012. Alpha tocopherol incorporation in mitochondria and microsome upon supranutritional vitamin E supplementation. *Gene Nutrition*. 7 (4) : 475-482.
- Pradel, W., Mal, R., Gedrange, T. and Lauer, G. 2008. Cell passage and composition of culture medium effects proliferation and differentiation of human osteoblast-like cells from facial bone. *Journal of Physiology and Pharmacology*. 47-58.
- Pramono, C. 2011. Mandibular reconstruction using non-vascularized autogenous bone graft applied in decorticated cortical bone. *It J. maxillofac. Surg*. 22 (1) : 47-56.
- Randall, M. 2004. *Bony Reconstruction of the Jaws*. (London: Bc decker inc.) 783-790.
- Tecla, M., Temu, Ke-Ying, W., Philip, A., Gruppiso and Chanika, P. 2010. The Mechanism of Ascorbic acid-induce differentiation of ATDC5 Chondrogenic cells. *American Journal of Physiology-Endocrinology and Metabolism*. 299 (2) : 325-334.

EFFECT OF DIFFERENT CONCENTRATION OF ALPHA TOCOPHEROL SUPPLEMENTATION ON HUMAN BONE MARROW MESENCHYMAL STEM CELL VIABILITY (IN VITRO STUDY)

ORIGINALITY REPORT

8%

SIMILARITY INDEX

4%

INTERNET SOURCES

4%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Maria Luz Alonso-Alonso, Laura García-Posadas, Yolanda Diebold. "Extracellular Vesicles from Human Adipose-Derived Mesenchymal Stem Cells: A Review of Common Cargos", Stem Cell Reviews and Reports, 2021
Publication 2%
- 2 repository.unair.ac.id
Internet Source 1%
- 3 Longgui Zhang, Ting Su, Bin He, Zhongwei Gu. "Self-assembly Polyrotaxanes Nanoparticles as Carriers for Anticancer Drug Methotrexate Delivery", Nano-Micro Letters, 2014
Publication 1%
- 4 silo.pub
Internet Source 1%
- 5 Submitted to Universitas Airlangga
Student Paper 1%

6 Waya Rahmaning Gusti Agrippina, Prihartini Widiyanti, Helmy Yusuf. "Synthesis and Characterization of Bacterial Cellulose - *Garcinia mangostana* Extract as Anti Breast Cancer Biofilm Candidate", Journal of Biomimetics, Biomaterials and Biomedical Engineering, 2017
Publication 1 %

7 onlinelibrary.wiley.com
Internet Source 1 %

8 peerj.com
Internet Source 1 %

9 www.ejbiotechnology.info
Internet Source 1 %

Exclude quotes Off

Exclude matches < 7 words

Exclude bibliography On

EFFECT OF DIFFERENT CONCENTRATION OF ALPHA TOCOPHEROL SUPPLEMENTATION ON HUMAN BONE MARROW MESENCHYMAL STEM CELL VIABILITY (IN VITRO STUDY)

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
