

Flavonoid from Stem Bark of *Aquilaria microcarpa*

by Alfinda Novi Kristanti

Submission date: 09-Mar-2020 05:29PM (UTC+0800)

Submission ID: 1272167519

File name: 2018-EJCHEM_Volume_61_Issue_2_Pages_313-316.pdf (532.51K)

Word count: 2034

Character count: 10267

Flavonoid from Stem Bark of *Aquilaria microcarpa*

A. N. Kristanti^{1*}, M. Tanjung¹ and O. P. Rahayu¹

¹Laboratory of Organic Chemistry, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115, Indonesia

THIS study reported the isolation of 7-hydroxy-5,3',4'-trimethoxyflavone (**1**) from stem bark of *Aquilaria microcarpa*, a species belonging to Thymelaeaceae family. This report is our continuation study about this plant. Previously, we reported the presence of 5,3',4'-trihydroxy-7-methoxyflavon and 6-hydroxy-2-(2-phenylethyl) chromon. The chemical structure of compound (**1**) has been elucidated based on spectroscopy method including FTIR, NMR and HRESI-MS analysis.

Keywords: 7-hydroxy-5,3',4'-trimethoxyflavone; flavonoid, *Aquilaria microcarpa*; Thymelaeaceae

Introduction

The Thymelaeaceae is a family of dicotyledonous plants mainly found in the tropics and subtropics. They are mostly trees and shrubs, but include also a few vines and herbaceous plants. It is recognized that Thymelaeaceae composed by 45 genus [1]. Among these 45 genera, *Aquilaria*, *Daphne*, *Gnidia* dan *Phaleria* are the most widely studied and reported genus. The genus of *Aquilaria* is distributed widely in Asia. *A. sinensis*, *A. malaccensis*, *A. hirta*, *A. crassna*, and *A. agallocha* are the widely studied species of *Aquilaria*, but some species have not been investigated, such as *A. acuminata*, *A. baillonii*, *A. citrinicarpa*, *A. filaria*, *A. grandiflora*, *A. microcarpa* etc. Certain trees of *Aquilaria* produce the fragrant resinous wood, called agarwood, eaglewood, gaharu, kanankoh, jinkoh, chen xiang or tram in different region [2,3]. Other people also call it aloeswood or agalloch [3]. This agarwood could be used as incense which has been widely present in the religious ceremony of traditional culture [4,5] and used also as traditional medicine [2,6-10]. Due to the odor produced, agarwood now is a material for perfume and aromatherapy, in addition to all the other benefits since antiquity. The *Aquilaria* is of prime importance owing to its richness in a variety of different classes of natural product, especially sesquiterpenes and chromones. Other classes of compounds are flavonoid, benzophenone, diterpenoid, triterpenoid and lignan.

In continuation of our research about the secondary metabolites contained in *A. microcarpa*, we report here the isolation and structural elucidation of a flavonoid, 7-hydroxy-5,3',4'-trimethoxyflavone (**1**) (Figure 1) from the stem bark of this plant. Previously, we reported the presence of two phenolic compounds, namely 5,3',4'-trihydroxy-7-methoxyflavon or known as 3'-hydroxy genkwainin and 6-hydroxy-2-(2-phenylethyl) chromon from the same plant [11]. Compound (**1**) has only ever been reported once in the plant by some Chinese researchers. It was first isolated and identified from the leaf of *Murraya paniculata*, but the spectroscopy data was not available [12]. There is not any publication reported about the presence of this compound in *Aquilaria*. even in Thymelaeaceae.




Fig. 1. Structure of 7-hydroxy-5,3',4'-trimethoxyflavone (1)

Experimental

General experimental

The UV spectrum was measured with Shimadzu series 1800 spectrophotometer (Kyoto, Japan). The FTIR spectrum was recorded in KBr

*Corresponding author e-mail: alfinda-n-k@fst.unair.ac.id.
DOI : 10.21608/ejchem.2018.2173.1174

powder with the Diffuse Reflectance Method on spectrophotometer IRTracer 100 (Shimadzu, Kyoto, Japan). NMR Spectra (^1H , APT, HMBC and HMQC) were recorded on JEOL 400 ECA spectrometer (JEOL, Tokyo, Japan) using CDCl_3 as a solvent and internal standard. The mass spectrum was recorded by HRESI-MS (Waters LCT Premier XE, Waters Corp. Milford, MA, USA).

Plant Material

Samples of stem bark of *A. microcarpa* were collected in October 2015 from Conservation forest Bukit Bangkirai, Samboja, Samarinda, East Kalimantan, Indonesia and the voucher specimen (UA-TAm011015) was deposited at the Herbarium of Universitas Airlangga. The stem bark was cleaned, air dried under the shade, cut into small pieces and milled.

Extraction and Isolation


The dried stem barks of *A. microcarpa* (1.69 kg) were macerated with n-hexane at room temperature for 2x24 hours followed by maceration using methanol for 3x24 hours. The methanol extract was evaporated under reduced pressure to give a crude methanol extract (120 g). Into this extract was then added water and partitioned with ethyl acetate. The ethyl acetate extract obtained was evaporated under reduced pressure to give a crude ethyl acetate extract (10.8 g). Ethyl acetate extract was separated by flash chromatography on silica gel using the mixture of n-hexane-ethyl acetate as eluent. Elution was done by increasing the polarity gradually by increasing the amount of ethyl acetate to give five major fractions A-E. TLC on fraction E (250 mg) using eluent chloroform showed a potential spot. Purification of this fraction using flash chromatography twice with eluent n-hexane-ethyl acetate and n-hexane-chloroform respectively, yielded compound (1) (20 mg).

Results and Discussion

Compound (1) was obtained as a yellow solid (mp 172 – 174°C) after passing through a series of isolation steps, including extraction followed by separation and purification using chromatographic techniques. The molecular formula is $\text{C}_{18}\text{H}_{17}\text{O}_6$, whereas that of the protonated molecule $[\text{M}+\text{H}]^+$ is $\text{C}_{18}\text{H}_{18}\text{O}_6$ at m/z 329.1024 (calcd. 329.1025) by the HR-ESI-MS. The UV spectrum showed two absorption bands, benzoyl and cinamoyl moiety, characteristic of flavonoid skeleton at

λ_{max} 269 (4.19) and 340 (4.24) nm. The IR spectrum showed absorption bands at ν_{max} 3280, 3093, 2924, 1618, 1587, 1517, 1261, 831 cm^{-1} indicating the presence of a hydroxyl, aromatic group, conjugated carbonyl and methoxy group. ^1H -NMR spectrum showed a singlet signal at δ_{H} 6.60 ppm which was a characteristic of flavon group. There were also five signals of aromatic protons which appeared at δ_{H} 7.53 ppm (H_6 , dd, $J = 8.6$ and 2.1 Hz); 7.34 ppm (H_2 , d, $J = 2.1$ Hz); 6.98 ppm (H_8 , d, $J = 8.6$ Hz). These three signals described the existence of ABX system. Two other signals appeared as doublet at δ_{H} 6.50 and 6.38 ppm. Both of signals indicated that these two protons were in meta position. In addition, it was also observed the presence of three singlet signals of methoxy group at δ_{H} 3.99; 3.97 and 3.89 ppm (Table 1). ^{13}C -NMR ATP spectrum showed 18 completely separated signals, which is consistent with the structure of substituted flavone. Through the spectrum it can be seen that this compound was composed by six methins (CH) (δ_{C} 120.2; 111.2; 108.8; 104.8; 98.2 and 92.8 ppm), three methoxy (OCH_3) (δ_{C} 56.2; 56.1; and 55.9 ppm), and 9 quaternary carbon (δ_{C} 182.4; 165.5; 164.7; 164.1; 162.3; 152.2; 149.3; 123.8 dan 105.6 ppm) (Table 1).

The HMQC spectrum gave the information about the correlation between directly-bonded H on C, so through this spectrum we obtained the information about which proton is attached to which carbon. In the HMBC spectrum, it was observed the presence of long range correlations between the proton signal aromatic H-8 (δ_{H} 6.98) with three quaternary carbon signals at δ_{C} 164.7 (C-7), 162.3 (C-8a), 105.6 (C-4a) and one methin at δ_{C} 98.2 (C-6). The appearance another long range correlations between the proton signal aromatic H-6 (δ_{H} 6.38) with three quaternary carbon signals at δ_{C} 165.5 (C-5), 164.7 (C-7), 105.6 (C-4a) and one methin at δ_{C} 92.8 (C-8) placed certainly the methoxy at the C-5 and not at C-7 in the ring A. This argument was supported by the existence of the correlation of three methoxys with three oxyaryl carbons, which are $-\text{OCH}_3$ (δ_{H} 3.99) with an oxyaryl carbon (δ_{C} 152.2; C-4'); $-\text{OCH}_3$ (δ_{H} 3.97) with an oxyaryl carbon (δ_{C} 149.3; C-3') and $-\text{OCH}_3$ (δ_{H} 3.89) with an oxyaryl carbon (at δ_{C} 165.5; C-5). All of data obtained from NMR are presented in Table 1. Correlations observed in HMBC spectrum are described in Fig. 2. Therefore, compound (1) was elucidated as 7-hydroxy-5, 3', 4'-trimethoxyflavone.



4
Fig. 2. HMBC correlations of compound (1)

TABLE 1. ¹H, ¹³C and HMBC NMR data of compound (1) in CDCl₃ (δ in ppm)

No. Atom	δ _H (mult, J Hz)	δ _C	HMBC
2	-	164.1	
3	6.60 (s)	104.8	C-2, C-4, C-1', C-4a
4	-	182.4	
4a	-	105.6	
5	-	165.5	
6	6.38 (d, J = 2.2 Hz)	98.2	C-4a, C-5, C-7, C-8
7	-	164.7	
8	6.50 (d, J = 2.2 Hz)	92.8	C-4a, C-6, C-7, C-8a
8a	-	162.3	
1'	-	123.8	
2'	7.34 (d, J = 2.1 Hz)	108.8	C-4', C-6'
3'	-	149.3	
4'	-	152.2	
5'	6.98 (d, J = 8.6 Hz)	111.2	C-1', C-3'
6'	7.53 (dd, J = 8.6 & 2.1 Hz)	120.2	C-2', C-4'
5-OCH ₃	3.89 (s)	55.9	C-5
3'-OCH ₃	3.97 (s)	56.1	C-3'
4'-OCH ₃	3.99 (s)	56.2	C-4'

The spectrum data was compared with similar compound, namely 5,7,3',4'-tetramethoxyflavone which was isolated and identified from *Kaempferia parviflora*, reported by Sutthanut [13].

Conclusion

A flavonoid namely 7-hydroxy-5,3',4'-trimethoxyflavone was successfully isolated and identified from stem bark of *Aquilaria microcarpa*,

a species belonging to Thymelaeaceae. This is the first time that this compound reported found in *Aquilaria*, even in Thymelaeaceae.

Acknowledgements

This research was supported by Ministry of Research, Technology and Higher Education, Republik Indonesia (Penelitian Unggulan Perguruan Tinggi, 2016).

References

1. Herber, B.E., "The Families and Genera of Vascular Plants", ed. K. Kubitzki and C. Bayer, Vol. 5, Springer-Verlag, Berlin (2003).
2. Li, W., Cai, C.H., Dong, W.H., Guo, Z.K., Wang, H., Mei, W.L. and Dai, H.F., 2-(2-phenylethyl) chromone derivatives from Chinese agarwood induced by artificial holing. *Fitoterapia*, **98**, 117 (2014).
3. Ueda, J., Imamura, L., Tezuka, Y., Tran, Q.L., Tsuda, M. and Kadota, S., New sesquiterpene from Vietnamese agarwood and its induction effect on brain-derived neurotrophic factor mRNA expression in vitro. *Bioorg. Med. Chem*, **14**, 3571 (2006).
4. Alkhatlan, H.Z., Al-Hazimi, H.M., Al-Dhalaan, F.S. and Mousa, A.A., Three 2-(2-phenylethyl) chromones and two terpenes from agarwood. *Nat Prod Res*, **19** (4), 367 (2005).
5. Yang, D.L., Wang, H., Guo, Z.K., Dong, W.H., Mei, W.L. and Dai, H.F., A new 2-(2-phenylethyl) chromone derivative in Chinese agarwood "Qi-Nan" from *Aquilaria sinensis*. *J Asian Nat Prod Res*, **16** (7), 770 (2014).
6. Bhuiyan, M.N.I., Begum, J. and Bhuiyan, M.N.H., Analysis of essential oil of eaglewood tree (*Aquilaria agallocha* Roxb.) by gas chromatography mass spectrometry. *Bangladesh J. Pharmacol.*, **4**, 24 (2009).
7. Kim, Y.C., Lee, E.H., Lee, Y.M., Kim, H.K., Song, B.K., Lee, E.J. and Kim, H.M., Effect of the aqueous extract of *Aquilaria agallocha* stems on the immediate hypersensitivity reactions. *J Ethnopharmacol*, **58**, 31 (1997).
8. Nakanishi, T., Inada, A., Nishi, M., Yamagata, E. and Yoneda, K., A new and a known derivatives of 2-(2-phenylethyl) chromone from a kind of agarwood ("kanankoh" in Japanese) originating from *Aquilaria agallocha*. *J. Nat. Prod.*, **49**(6), 1106 (1986).
9. Yagura, T., Iro, M., Kiuchi, F., Honda G. and Shimada, Y., Four new 2-(2-phenylethyl) chromone derivatives from withered wood of *Aquilaria sinensis*. *Chem Pharm Bull*, **51**(5), 560 (2003).
10. Zhou, M., Wang, H., Suolangjiba, Kou, J. and Yu, B., Antinociceptive and anti-inflammatory activities of *Aquilaria sinensis* (Lour.) Gilg. leaves extract. *J Ethnopharmacol*, **117**, 345 (2008).
11. Kristanti, A.N., Tanjung, M. and Rahayu, O.P., Phenolic compounds from *Aquilaria microcarpa* stem bark. *J. Chem. Technol. Metall.*, **52**, 1111 (2017).
12. Shan, J., Wang, X.Z., Ma, Y.D., Yang, R.J., Li, X.W. and Jin, Y.R., Studies on flavonoids from leaves of *Murraya paniculata* L. *Zhongguo Yaoxue Zazhi*, **45**, 1910 (2010).
13. Sutthanut, K., Sripanidkulchai, B., Yenjai, C. and Jay, M., Simultaneous identification and quantitation of 11 flavonoid constituents in *Kaempferia parviflora* by gas chromatography. *J. Chromatogr. A*, **1143**, 227 (2007).

(Received 28/11/2017;
accepted 12/2/2018)

Flavonoid from Stem Bark of Aquilaria microcarpa

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Mahabusarakam, W.. "Xanthone derivatives from *Cratoxylum cochinchinense* roots", *Phytochemistry*, 200603
Publication 2%
- 2 Tjahjandarie, Tjitjik, Ratih Saputri, and Mulyadi Tanjung. "Methyl 2,5-Dihydroxy-4-(3'-methyl-2'-butenyl)benzoate", *Molbank*, 2016.
Publication 2%
- 3 doaj.org
Internet Source 2%
- 4 www.thieme-connect.com
Internet Source 1%
- 5 P. Z. Xian Li. "A new perylenequinone from the fruit bodies of *Bulgaria inquinans*", *Journal of Asian Natural Products Research*, 12/1/2006
Publication 1%
- 6 doczz.net
Internet Source 1%

7	Jeevan Kumar Prasain, Masashi Ueki, Piotr Stefanowicz, Hiroyuki Osada. "Rapid screening and identification of cytochalasins by electrospray tandem mass spectrometry", <i>Journal of Mass Spectrometry</i> , 2002	1 %
Publication		
8	Hinna Hamid, S. Tarique Abdullah, Mohammed Ali, M. Sarwar Alam. " New Phytoconstituents from the Aerial Parts of . ", <i>Pharmaceutical Biology</i> , 2008	1 %
Publication		
9	ptti.or.id Internet Source	1 %
10	mdpi.com Internet Source	1 %
11	Wu, Wen-Bin, Gao-Chao Yue, Qi-Lin Huang, Ling-Ling Sun, and Wen Zhang. "A new compound from an endophytic fungus <i>Alternaria tenuissima</i> ", <i>Journal of Asian Natural Products Research</i> , 2014.	1 %
Publication		
12	publikationen.bibliothek.kit.edu Internet Source	1 %
13	Yeon Choi, Gyu Yon, Kyung Hong, Dae Yoo, Chun Choi, Woo-Kyu Park, Jae Kong, Young Kim, Shi Ryu. " BACE-1 Inhibitory Phenolic	1 %

Components from the Seeds of ", Planta Medica, 2008

Publication

14	bdt.d.biblioteca.ufpb.br Internet Source	1 %
15	russianpatents.com Internet Source	1 %
16	Hyoung-Geun Kim, Hyun-Ji Oh, Jung-Hwan Ko, Hae Seong Song, Yeong-Geun Lee, Se Chan Kang, Dae Young Lee, Nam-In Baek. "Lanceoleins A–G, hydroxychalcones, from the flowers of <i>Coreopsis lanceolata</i> and their chemopreventive effects against human colon cancer cells", <i>Bioorganic Chemistry</i> , 2019 Publication	1 %
17	www.journal.csj.jp Internet Source	1 %
18	Arora, Arti, Susanne Valcic, Santiago Cornejo, Muralee G. Nair, Barbara N. Timmermann, and Daniel C. Liebler. "Reactions of Genistein with Alkylperoxyl Radicals", <i>Chemical Research in Toxicology</i> , 2000. Publication	<1 %
19	V. Kontrimavičiūtė, O. Mathieu, L. Balas, R. Escale, Jean Pierre Blayac, F. M. M. Bressolle. "Ibogaine and Noribogaine: Structural Analysis and Stability Studies. Use of LC - MS to	<1 %

Determine Alkaloid Contents of the Root Bark of ", Journal of Liquid Chromatography & Related Technologies, 2007

Publication

20

Jayasinghe, L.. "A new ecdysteroid, 2-deoxy-5@b,20-dihydroxyecdysone from the fruits of *Diploclisia glaucescens*", *Steroids*, 200305

Publication

<1 %

21

Nabila Brahmi, Monica Scognamiglio, Severina Pacifico, Aida Mekhoukhe, Khodir Madani, Antonio Fiorentino, Pietro Monaco. "1 H NMR based metabolic profiling of eleven Algerian aromatic plants and evaluation of their antioxidant and cytotoxic properties", *Food Research International*, 2015

Publication

<1 %

22

kb.psu.ac.th

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Flavonoid from Stem Bark of Aquilaria microcarpa

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
