



Asian Pacific Journal of Tropical Biomedicine

[Latest issue](#) [All issues](#)

[Search in this journal](#)

Volume 5, Issue 12

Pages 977-1064 (December 2015)

[Download full issue](#)

[< Previous vol/issue](#)

[Next vol/issue >](#)

Research article [Open access](#)

Cocktail of *Theileria equi* antigens for detecting infection in equines

Shimaa Abd El-Salam El-Sayed, Mohamed Abdo Rizk, Mohamed Alaa Terkawi, Ahmed Mousa, ... Ikuo Igarashi

Pages 977-981

[Download PDF](#) [Article preview](#)

Research article [Open access](#)

Antiplasmodial activity of traditional polyherbal remedy from Odisha, India: Their potential for prophylactic use

Prakash Bangalore Nagendrappa, Jean-François Franetich, Frederick Gay, Audrey Lorthiois, ... Dominique Mazier

Pages 982-986

[Download PDF](#) [Article preview](#)

Research article [Open access](#)

Deletion of *Salmonella enterica* serovar typhimurium *sipC* gene

Maryam Safarpour Dehkordi, Abbas Doosti, Asghar Arshi

Pages 987-991

[Download PDF](#) [Article preview](#)

Research article [Open access](#)

Anticancer activity of *Cyanotheca* sp. strain extracts from Egypt: First record

Nermin Adel El Semary, Manar Fouada

Pages 992-995

[Download PDF](#) Article preview 

Research article [Open access](#)

Influence of CD133⁺ expression on patients' survival and resistance of CD133⁺ cells to anti-tumor reagents in gastric cancer

De-Hu Chen, Rui-Qi Lu, Xiao-Chun Ni, Ju-Gang Wu, ... Ji-Wei Yu

Pages 996-1004

[Download PDF](#) Article preview 

Research article [Open access](#)

Vascular endothelial growth factor before and after locoregional treatment and its relation to treatment response in hepatocellular carcinoma patients

Heba Sedrak, Noaman El-Garem, Mervat Naguib, Heba El-Zawahry, ... Lila Rashed

Pages 1005-1009

[Download PDF](#) Article preview 

Research article [Open access](#)

Combined treatment of 3-hydroxypyridine-4-one derivatives and green tea extract to induce hepcidin expression in iron-overloaded β -thalassemic mice

Supraanee Upanan, Kanjana Pangjit, Chairat Uthaipibull, Suthat Fucharoen, ... Somdet Srichairatanakool

Pages 1010-1017

[Download PDF](#) Article preview 

Research article [Open access](#)

Anti-breast cancer properties and toxicity of *Dillenia suffruticosa* root aqueous extract in BALB/c mice

Latifah Saiful Yazan, Yong Sze Ong, Nur Elena Zaaba, Razana Mohd Ali, ... Yin Sim Tor

Pages 1018-1026

[Download PDF](#) Article preview 

Research article [Open access](#)

Effect of sucrose and potassium nitrate on biomass and saponin content of *Talinum paniculatum* Gaertn. hairy root in balloon-type bubble bioreactor

Yosephine Sri Wulan Manuhara, Alfinda Novi Kristanti, Edy Setiti Wida Utami, Arif Yachya

Pages 1027-1032

[Download PDF](#) Article preview 

Research article [Open access](#)

Gas chromatography mass spectrometry analysis and *in vitro* antibacterial activity of essential oil from *Trigonella foenum-graecum*

Moniruzzaman, Shahinuzzaman, Ahsanul Haque, Rahima Khatun, Zahira Yaakob

Pages 1033-1036

[Download PDF](#) Article preview 

Research article [Open access](#)

Cytotoxic activity of crude extracts and fractions from *Premna odorata* (Blanco), *Artocarpus camansi* (Blanco) and *Gliricidia sepium* (Jacq.) against selected human cancer cell lines

Ourlad Alzeus G. Tantengco, Sonia D. Jacinto

Pages 1037-1041

[Download PDF](#) [Article preview](#) 

Research article [Open access](#)

Screening for anti-pancreatic lipase properties of 28 traditional Thai medicinal herbs

Ananya Dechakhamphu, Nattapong Wongchum

Pages 1042-1045

[Download PDF](#) [Article preview](#) 

Research article [Open access](#)

Qualitative analysis of catechins from green tea GMB-4 clone using HPLC and LC-MS/MS

Erna Susanti, Ciptati, Retty Ratnawati, Aulanni'am, Achmad Rudijanto

Pages 1046-1050

[Download PDF](#) [Article preview](#) 

Research article [Open access](#)

Evaluation of febrile neutropenic patients hospitalized in a hematology clinic

Mücahit Görük, Mehmet Sinan Dal, Tuba Dal, Abdullah Karakus, ... Orhan Ayyildiz

Pages 1051-1054

[Download PDF](#) [Article preview](#) 

Research article [Open access](#)

The frequency and antimicrobial resistance patterns of nosocomial pathogens recovered from cancer patients and hospital environments

Aymen Mudawe Nurain, Naser Eldin Bilal, Mutasim Elhadi Ibrahim

Pages 1055-1059

[Download PDF](#) [Article preview](#) 

Research article [Open access](#)

Seroepidemiology of *Toxoplasma* infection in blood donors in Jahrom District, Southern Iran

Mohammad Hassan Davami, Morteza Pourahmad, Rasoul Baharlou, Abdolreza Sotoodeh Jahromi, ... Mohsen Kalantari

Pages 1060-1064

[Download PDF](#) [Article preview](#) 

ISSN: 2221-1691

Copyright © 2019 Hainan Medical University. Production and hosting by Elsevier B.V.

ELSEVIER

[About ScienceDirect](#) [Remote access](#) [Shopping cart](#) [Advertise](#) [Contact and support](#) [Terms and conditions](#) [Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the [use of cookies](#).

Copyright © 2019 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

 RELX™

Editorial Board

[About the journal](#)

Editors-in-Chief

Prof. Dr. Jeffrey M. Bethony

Department of Microbiology, Immunology & Tropical Medicine, School of Medicine and Health Sciences, The George Washington University, Washington DC, USA

E-mail: jbethony@gwu.edu

Prof. Dr. Santiago Mas-Coma

Departamento de Parasitología, Facultad de Farmacia, Universidad de Valencia, Valencia, Spain

E-mail: S.Mas.Coma@uv.es

Prof. Dr. Malcolm K. Jones

School of Veterinary Science, Faculty of Science, The University of Queensland, Queensland, Australia

E-mail: m.jones@uq.edu.au

Prof. Dr. Vanessa Steenkamp

Department of Pharmacology, University of Pretoria, Pretoria, South Africa

E-mail: vanessa.steenkamp@up.ac.za

Nazni Bte Hj. Wasi Ahmad

Senior Research Officer, Medical Entomology Unit, Institute for Medical Research, Kuala Lumpur, Malaysia

E-mail: nazni@imr.gov.my

President Executive-Editor-in-Chief

Prof. Shunhai Qu

Hainan Medical University Journal Publisher, Hainan Medical University, Haikou, China

E-mail: qu.sh@163.com

Deputy Editors-in-Chief

Prof. Dr. Jong-Yil Chai

Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, Seoul, Korea

Dr. Pierre Roques

ImmunoVirology Division, Institute for Emerging Diseases and Innovative Therapies, Fontenay-aux-Roses, France

Dr. Alcides Troncoso

Department of Infectious Diseases, School of Medicine, Buenos Aires University, Buenos Aires, Argentina

Dr. Stephen Munga

Centre for Global Health Research, Kenya Medical Research Institute, Nairobi, Kenya

Associate Editors-in-Chief

Prof. Giuseppe La Torre

Department of Public Health and Infectious Diseases, Sapienza University of Rome, Rome, Italy

Prof. Odir Antônio Dellagostin

Biotechnology Unit, Center for Technology Development, Federal University of Pelotas, Pelotas, Brazil

Prof. Sung-Jong Hong

Department of Medical Environmental Biology, Chung-Ang University College of Medicine, Seoul, Korea

Prof. Hassan Vatandoost

Department of Entomology and Vector Control, School of Public Health and National Institute of Health Research, Teheran University of Medical Sciences, Teheran, Iran

Prof. Maria José Ferreira

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Lisbon, Lisbon, Portugal

Dr. Leonard E. G. Mboera

National Institute for Medical Research, Dar es Salaam, Tanzania

Prof. Dr. Iván Dario Vélez

Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia

Editorial Directors

Prof. Dr. Viroj Wiwanitkit

Department of Laboratory Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Assoc. Prof. Dr. Saravanan Thangamani

Department of Pathology, University of Texas Medical Branch (UTMB), Galveston, USA

[SUBMIT ARTICLE](#)

[SUBSCRIBE](#)

[POPULAR ARTICLES](#)

[JOIN AS REVIEWER](#)

[GET EMAIL ALERTS](#)

[RECOMMEND](#)

Prof. Dr. Laura Bongiovanni
Faculty of Veterinary Medicine, University of Teramo, Teramo, Italy

Statistic Editors

Assoc. Prof. Khaled Khatab
Health Economics Modelling and Medical Statistics, Faculty of Health and Wellbeing, Sheffield Hallam University, Sheffield, UK

Assist. Prof. Sarath Chandra Janga
Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, USA

Executive Editors

Prof. Dr. Salah Akkal
Department of Chemistry, University of Constantine, Constantine, Algeria

Dr. Hervé Hoste
French National Institute for Agricultural Research, Université de Toulouse, Toulouse, France

Prof. Thomas N. Jr. Tully
Department Veterinary Clinical Sciences, Louisiana State University School of Veterinary Medicine, Baton Rouge, USA

Prof. Dr. Barbara R Conway
Department of Pharmacy, University of Huddersfield, Huddersfield, UK

Executive Members of Editorial Board

Assoc. Prof. Vittorio Sambri
Unit of Microbiology, Department of Specialistic, Diagnostic and Experimental Medicine, University of Bologna, Bologna, Italy

Prof. Dr. Qiushui He
Department of Microbiology and Immunology, University of Turku, Turku, Finland

Assist. Prof. Dr. Bindu Sukumaran
Emerging Infectious Diseases Programme, Duke-NUS Medical School, Singapore

Prof. Dr. Suhail Ahmad
Department of Microbiology, Kuwait University, Kuwait City, Kuwait

Prof. Indra Vythilingam
Department of Parasitology, University of Malaya, Kuala Lumpur, Malaysia

Prof. Dr. Wael Mohamed Abou EL-Makarem EL-Deeb
Department of Clinical Studies, College of Veterinary Medicine and Animal Resources, King Faisal University, Al-Ahsa, Saudi Arabia

Assoc. Prof. Dr. Ben Slama Karim
Département de Biotechnologie, Université de Tunis El Manar, Tunis, Tunisia

Prof. Bing Huang
Department of Animal Parasitology, Shanghai Veterinary Research Institute, Chinese Academy of Agricultural Science, Shanghai, China

Prof. Xiao-Nong Zhou
National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention, Shanghai, China

Dr. Nahid Einollahi
Department of Lab Medical Sciences, Faculty of Allied Medical Sciences, Tehran University of Medical Sciences, Tehran, Iran

Prof. Dr. Munir Aktas
Department of Parasitology, College of Veterinary Medicine, Firat University, Elazig, Turkey

Dr. José Manuel Lorenzo Rodríguez
Researcher in Meat Technology Centre of Galicia, Meat Technology Centre of Galicia, Galicia, Spain

Prof. Dr. Jeong Hwan Shin
Department of Laboratory Medicine, Busan Paik Hospital, Inje University College of Medicine, Busan, Korea

Luis Ignacio González Granada
Teaching assistant, Immunodeficiencies Unit, Hospital 12 de Octubre, Madrid, Spain

Dr. Abhishek Mathur
Executive Director (R&D), NCS Group, Nagpur, Maharashtra, India

Dr. Ahmed Ismail
Consultant Clinical Pathologist & Chief Lab Services, Supreme Council of Health, Doha, Qatar

Dr. Américo David Rodríguez Ramírez
Centro Regional de Investigación en Salud Pública, Chiapas, Mexico

Prof. Xue-Jie Yu
Department of Pathology, University of Texas Medical Branch, Texas, USA

Prof. Arun Kumar
Department of Biochemistry, Shri Shankaracharya Institute of Medical Sciences, Bilai, India

Prof. Dr. Chia-Kwung Fan
Department of Parasitology, Taipei Medical University College of Medicine, Taipei

Assist. Prof. Dr. Cyrille Bisseye
Centre de Recherche Biomoléculaire Pietro Annigoni (CERBA)/LABIOGENE, UFR/SVT, Ouagadougou, Burkina Faso

Prof. Dr. Dongmi Kwak
College of Veterinary Medicine, Kyungpook National University, Daegu, Korea

Dr. Donovan Anthony McGrowder
Chemical Pathology Department, Faculty of Medical Sciences, University Hospital of The West Indies, Kingston, Jamaica

Dr. Farouk El Allaki
Epidemiology and Surveillance Section, Canadian Food Inspection Agency, Quebec, Canada

Prof. Dr. Gabriel Trueba
Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito, Ecuador

Assoc. Prof. Gordana J. Dragovic Lukic
Department of Pharmacology, Clinical Pharmacology and Toxicology, School of Medicine, University of Belgrade, Belgrade, Serbia

Dr. Hanspeter Marti
Swiss Tropical and Public Health Institute, Basel, Switzerland

Prof. Dr. Herbert Auer
Department of Medical Parasitology, Medical University of Vienna, Vienna, Austria

Hetron Mweemba Munang'andu
Department of Basic Sciences and Aquatic Medicine, Norwegian School of Veterinary Science, Oslo, Norway

Dr. Irma Khachidze
Department of Behavior, Cognition Functions and Human Psychophysiology, Beritashvili Institute of Physiology, Tbilisi, Georgia

Dr. Jean-François Faucher
Praticien hospitalier en maladies infectieuses et tropicales, CHU de Besançon, Service des maladies infectieuses et tropicales, Hôpital Minjoz, Besançon cedex, France

Dr. José A. Oteo
Infectious Diseases Department, Center of Rickettsioses and Vector-borne Diseases, Hospital San Pedro-CIBIR, Logroño, Spain

Dr. Juraj Majtán
Section of Molecular and Applied Zoology, Institute of Zoology, Slovak Academy of Sciences, Bratislava, Slovakia

Dr. Kalman Imre
Department of Animal Production and Veterinary Public Health, Faculty of Veterinary Medicine, Banat University of Agricultural Sciences and Veterinary Medicine Timisoara, Timisoara, Romania

Dr. Kingsley Badu
Noguchi Memorial Institute for Medical Research, University of Ghana, College of Health Science, Accra, Ghana

Lala Ravaomanarivo
Département d'Entomologie, de la Faculté des Sciences d'Antananarivo, University of Antananarivo, Antananarivo, Madagascar

Prof. Dr. Liwang Cui
Department of Entomology, The Pennsylvania State University (PSU), Pennsylvania, USA

Assist. Prof. Dr. Luís R. Silva
Laboratory for Process Engineering, Environment, Biotechnology and Energy (LEPABE), Faculty of Engineering (FEUP)
University of Porto, Porto, Portugal

Dr. M. Fawzi Mahomoodally
Department of Health Sciences, Faculty of Science, University of Mauritius, Reduit, Mauritius

Mamadou C Baldé
Institut Pasteur de Guinée, Cheikh Anta Diop University, Kindia, Guinée

Prof. Dr. Milkyas Endale
Department of Chemistry, Hawassa University, Hawassa, Ethiopia

Dr. Mochammad Hatta
Molecular Biology and Immunology Laboratory for Infectious Diseases, Department Microbiology, Faculty Medicine
Hasanuddin University, Makassar, Indonesia

Moses Samje
Department of Medicine, Faculty of Health Sciences, University of Bamenda, Bambili, Cameroon

Assist. Prof. Muftah A. M. Shushni
Department of Pharmacognosy, University of Tripoli, Tripoli, Libya

Dr. Muhammad Akram
Department of Eastern Medicine and Surgery, Faculty of Medical and Health Sciences, The University of Poonch, Azad Jammu and Kashmir, Pakistan

Dr. Musso Munyeme
Department of Disease Control, School of Veterinary Medicine, University of Zambia, Lusaka, Zambia

Prof. Dr. Nguyen Hoang Loc
Department of Biotechnology, College of Sciences, Hue University, Hue, Vietnam

Dr. Prof. Obembe Olawole
Department of Biological Sciences, Covenant University, Ota, Nigeria

Prof. Dr. Pius Tshimankinda Mpiana
Department of Chemistry, Faculty of science, University of Kinshasa, Kinshasa, Democratic Republic of the Congo

Prof. Dr. Polrat Wilairatana
Department of Clinical Tropical Medicine, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

Dr. Pravin Malla Shrestha
Energy Biosciences Institute, University of California, Berkeley, USA

Dr. Rebuma Firdessa
Institute for Molecular Infection Biology, University of Wuerzburg, Wuerzburg, Germany

Prof. Rinaldo Poncio Mendes
Faculdade de Medicina de Botucatu, Universidade Estadual Paulista Júlio de Mesquita Filho, Botucatu, Brazil

Dr. Robin A J Nicholas
Mycoplasma Group, Department of Statutory and Exotic Bacteria, Veterinary Laboratories Agency (Weybridge), Surrey, United Kingdom

Dr. Assoc. Prof. Samath D Dharmaratne
Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka

Dr. Assist. Prof. Song Gao
Department of Pharmacological and Pharmaceutical Sciences, The University of Houston, Houston, USA

Dr. Assist. Prof. Tarig Mohamed Saad Alnour
Department of Microbiology, Alzaiem Alazhari University, Khartoum, Sudan

Dr. Tashi Tobgay
Vector-Borne Disease Control Programme, Ministry of Health Royal Government of Bhutan, Thimphu, Bhutan

Dr. Thomas Dorlo
Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden

Prof. Tohru Gono
Medical Mycology Research Center, Chiba University, Chiba, Japan

Dr. Vania Giacomet
Department of Paediatrics, University of Milan, Luigi Sacco Hospital, Milano, Italy

Dr. Veasna Duong
Virology Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

Assist. Prof. Dr. Weaam N. E. Ebrahim
Department of Pharmacognosy and Medicinal Plants, Faculty of Pharmacy, Mansoura University, Mansoura, Egypt.

Dr. Xuanming Shi
Department of Physiology, University of Texas Southwestern Medical Center, Dallas, TX, USA

Assist. Prof. Dr. Yong Song
Faculty of Medicine, Dentistry and Health Sciences, The University of Western Australia, Perth, Australia


Dr. Yu Pang
National Center for Tuberculosis Control and Prevention,
Chinese Center for Disease Control and Prevention, Beijing, China

Assist. Prof. Yuki Eshita
Department of Infectious Disease Control, Faculty of Medicine, Oita University, Oita, Japan



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbOriginal article <http://dx.doi.org/10.1016/j.apjtb.2015.09.009>Effect of sucrose and potassium nitrate on biomass and saponin content of *Talinum paniculatum* Gaertn. hairy root in balloon-type bubble bioreactorYosephine Sri Wulan Manuhara^{1*}, Alfinda Novi Kristanti², Edy Setiti Wida Utami¹, Arif Yachya³¹Laboratory of Plant Tissue Culture, Biology Department, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia²Laboratory of Organic Chemistry, Chemistry Department, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia³Laboratory of Plant Tissue Culture, Biology Department, Faculty of Mathematics and Natural Sciences, PGRI Adi Buana University, Surabaya, Indonesia

ARTICLE INFO

Article history:

Received 15 Jun 2015

Received in revised form 10 Jul,

2nd revised form 29 Jul 2015

Accepted 2 Aug 2015

Available online 20 Oct 2015

Keywords:

Talinum paniculatum Gaertn.

Saponin

Hairy root

Liquid culture

Balloon-type bubble bioreactor

ABSTRACT

Objective: To increase biomass and saponin production in hairy root culture of *Talinum paniculatum* Gaertn. (*T. paniculatum*) in balloon-type bubble bioreactor (BTBB).**Methods:** Hairy roots which were collected from leaf explants of *T. paniculatum* were infected by *Agrobacterium rhizogenes* strain LB510. The hairy roots were cultivated at 400 mL Murashige and Skoog liquid medium without growth regulator (MS0) in 1000 mL BTBB. Each BTBB had 2 g hairy roots as initial inoculum and these cultures were treated with various concentrations of sucrose (3%, 4%, 5%, 6% w/v) and potassium nitrate (0.5, 1.0, 1.5 and 2.0 strength of MS medium). Cultures were maintained for 14 days. Fresh and dry weights of hairy roots at the end of culture were investigated.**Results:** Various concentrations of sucrose influenced the biomass accumulation of hairy roots. Maximum biomass was reached by MS medium supplemented with 6% sucrose and it was approximately threefold higher than control. Culture supplemented with potassium nitrate at 2.0 strength of MS0 could increase biomass accumulation of hairy roots until 0.14 g dry weight and it was almost threefold higher than control. However, the maximum saponin content was obtained by MS medium supplemented with 5% sucrose and 2.0 strength potassium nitrate of MS.**Conclusions:** Based on this research, those conditions can be used to produce biomass and saponin of hairy root of *T. paniculatum* in the large scale.

1. Introduction

Java ginseng [*Talinum paniculatum* Gaertn. (*T. paniculatum*)] has been used in pharmaceutical industries for source of saponins, flavonoids, tannins, triterpenes or sterols, and polyphenols. Saponins of *T. paniculatum* are accumulated in roots. Ability and effectiveness of saponins on many medicinal treatments have been scientifically proven. Saponins were

reported to be able to enhance viability, motility and number of spermatozoa. Saponins also act as a anti-inflammatory agent, have androgenic potency, are able to induce cell differentiation through receptor cells [1], and could increase body resistance to disease [2]. *T. paniculatum* needs 3–4 years to produce the maximum saponins in the root. Root culture technology could be a solution to fill saponins demand in the market. This technology is important to be developed for plant preservation and increasing saponin content in roots.

Transformation by using *Agrobacterium rhizogenes* (*A. rhizogenes*) as a mediator to transfer transfer-DNA (T-DNA) into plant DNA is shortly alternative to produce roots. The T-DNA contains genes encoding enzymes for the synthesis of the phytohormones cytokinin and auxin, and of specific opine. The expression of oncogenes in Ri plasmid is indicated by adventive roots formation in infected area of explants. These adventive

*Corresponding author: Yosephine Sri Wulan Manuhara, Laboratory of Plant Tissue Culture, Biology Department, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia.

Tel: +62 (0)817584872

E-mail: wulanmanuhara@gmail.com

Peer review under responsibility of Hainan Medical University.

Foundation Project: Supported by Grant from Universitas Airlangga, Surabaya, Indonesia with Grant No. 8714/UN3/KR/2013.

roots are called hairy roots. In recent decades, hairy root culture has been widely used to produce various types of secondary metabolites that are naturally present in the roots. It was previously reported that *rolC* from *A. rhizogenes* T-DNA was shown to stimulate the production of secondary metabolites in the transformed plant cells of different plants. It was revealed that hairy roots enhanced the amount of glycyrrhizin in *Glycyrrhiza glabra* [3], plumbagine in *Plumbago rosea* [4], saponin in *Bacopa monnieri* [5], anthraquinones in *Polygonum multiflorum* [6], and polyphenols in *Momordica charantia* [7]. Organic nutrients in plants play a role in growth, development and accumulation of secondary metabolites. Growth and synthesis of secondary metabolite in hairy roots are also influenced by the nutritional quality. The effect of sucrose and nitrate concentrations in culture medium had been investigated. Biomass growth and withanolide A, production of *Withania somnifera* hairy roots were affected by different carbon sources in the Murashige and Skoog liquid medium (MS medium) [8]. Biomass and metabolite accumulation were also affected by NH_4 : NO_3 ratio in balloon-type bubble bioreactor (BTBB) culture of *Eurycoma longifolia* adventitious roots [9].

In previous studies, hairy roots of *T. paniculatum* that were cultivated in solid MS medium without growth regulator (MSO medium) grew slowly. Hairy root growth was accelerated after subculture in semi-solid MSO medium, but limited by oxygen supply and space growth. Therefore, hairy roots must be subcultured to liquid medium. Liquid culture has some advantages, for example, oxygen demand is filled by agitation or aeration, culture space limitation is solved by widening bioreactor chamber, pH medium is under control, nutrients are more homogenized and available for all parts of explants. Culture of *T. paniculatum* hairy roots in BTBB has mainly focused on aeration rates and inoculum densities. The best aeration rate and inoculum density for biomass and saponin content were reached at 2 g/400 mL and 0.25 vvm respectively [10]. In this research, various concentrations of sucrose and nitrogen are determined to increase biomass and saponin content.

2. Materials and methods

2.1. Materials

T. paniculatum was obtained from Botanical Garden of Purwodadi, Indonesia. *T. paniculatum* leaves were used as explants to initiate hairy roots. *A. rhizogenes* LB510 was infected to *T. paniculatum* leaves to induce hairy roots. The bacteria were obtained from Research Center of Biotechnology, Indonesian Institute of Sciences, Indonesia.

2.2. Explant sterilization and induction of hairy roots

The leaf explants of *T. paniculatum* were washed briefly with detergent and rinsed with running tap water. After rinsing, the explants were sterilized with 10% (v/v) clorox for 5 min, and then rinsed 3 times with sterile water. After sterilized and rinsed, the explants were shaken gently. The sterile leaf explants were cut $\pm 1 \text{ cm}^2$ and ready to be infected by *A. rhizogenes* LB510. The bacteria were grown in liquid Luria-Bertani medium at 110 r/min, and $(28 \pm 2)^\circ\text{C}$ for 2 days. The suspension of *A. rhizogenes* LB510 was diluted with liquid

MSO medium, then 100 $\mu\text{mol/L}$ acetosyringone was added. The explants were submerged in the solution for 5 min and were shaken gently. After that, the explants were drained on sterile filter paper and then transferred to MSO agar. The explants were incubated at $(28 \pm 2)^\circ\text{C}$ under dark condition for 2 days. In the next step, the explants were transferred to MSO solid medium supplemented with 500 mg/L cefotaxime. The explants were incubated at $(28 \pm 2)^\circ\text{C}$ under dark condition for a week, and then were moved into MSO semi-solid medium (5 g/L agar) for a week. Successful transformation was known with hairy root formation from the edge of the leaf explants. In the end of incubation time, hairy roots were excised from explants and then transferred to 250 mL Erlenmeyer flask. The flask contained 50 mL liquid MSO supplemented with 500 mg/L cefotaxime. Hairy roots cultures were agitated at 90 r/min, $(28 \pm 2)^\circ\text{C}$, under dark condition for a week. At the end of culture, hairy roots were ready to use as inoculum on liquid culture in BTBB.

2.3. Liquid culture condition in BBTB

Volumes of BBTB vessel were 1000 mL with working volumes between 200 and 500 mL. About 400 mL of MSO medium was placed in 1000 mL Erlenmeyer flask and sterilized on autoclave at 121°C for 20 min. Liquid MSO medium was transferred into BTBB aseptically. Initial inoculum was 2 g hairy roots and all BTBB cultures were aerated at flow rate of 0.25 vvm.

2.4. Treatment of sucrose and potassium nitrate at various concentrations

These experiments had eight BTBB cultures. Each culture was in the same conditions, such as 2 g hairy roots as inoculum, 0.25 vvm of air flow rate and 400 mL volume of MSO medium. In this test, eight BTBB were treated with various concentrations of sucrose [3%, 4%, 5%, 6% (w/v)] and various concentrations of potassium nitrate (0.5, 1.0, 1.5 and 2.0 strength of MS medium). Cultures were incubated under dark condition at 25°C for 14 days. The changes of pH, conductivity and total sugar in culture medium were checked every two days. Conductivity and total sugar content in medium culture were checked with hand conductometer (Ezdo) and hand refractometer (Atago). At the end of cultivation, biomass and saponin content were measured.

2.5. Saponin analyses

Saponin content of hairy roots was analyzed qualitatively by using thin layer chromatography and quantitatively by using high performance liquid chromatography (HPLC). Hairy roots were dried at 50°C for 5 days and then were ground with mortar. About 100 mg powders of hairy roots were immersed in 10 mL ethanol, and then heated at 80°C in water bath for 30 min. The hairy root extract was concentrated at 80°C for 3 h until a volume of 0.2 mL. Extract and standard saponin (Calbiochem) were spotted on silica gel GF254 and were eluted by using eluent propanol: water (14:3). Spot was detected by spraying with anisaldehyde 0.5 mL, acetic acid glacial 10.0 mL, ethanol 85.0 mL, and sulfuric acid 5.0 mL and then heated at 110°C for 6–10 min. Standard saponin will be dark green color.

Quantitative measurement of saponin was analyzed by using HPLC system (Agilent Q-TOF 6530 L) equipped with C18 columns. The mobile phase was mixtures of 0.1% formic acid in water grade and acetonitrile (40:60, v/v), and the injection volume was 0.2 μ L. The wavelength of detection was set at 299 nm.

3. Results

3.1. Induction of hairy roots

The hairy roots of *T. paniculatum* were transformed successfully by *A. rhizogenes* LB510 as shown in Figure 1A–C. Hairy roots as inoculum on liquid culture in BTBB were shown in Figure 1D. In the preliminary study, we did the transformation with two strains of *Agrobacterium*, *A. rhizogenes* LB510 and YMB072001, with different durations of infection and different explants (leaf and stem). The highest transformation efficiency was obtained by *A. rhizogenes* strain LB510 with 5-min infection at bacterial concentration and OD600 = 0.1 (78.43%), whereas for *A. rhizogenes* strain YMB072001, the highest transformation efficiency was obtained in 10-min infection (71.47%). Meanwhile, transformation by two strains of *A. rhizogenes* leaf explants obtained higher transformation efficiency than by stem explants. In *A. rhizogenes* strain LB510, transformation efficiency was 73.33% (leaf explants) and 66.67% (stem explants), whereas in *A. rhizogenes* strain YMB072001, transformation efficiency was 60.00% (leaf explants) and 56.67% (stem explants).

3.2. Effects of sucrose at various concentrations on biomass and saponin content

Sucrose at different concentrations was tested as carbon source for hairy root growth of *T. paniculatum* in BTBB. The results showed that different concentrations of sucrose provided different growth and saponin content in each hairy root culture (Figure 2). Hairy root culture with supplemented sucrose 6% had the maximum dry weight. The saponin content of *T. paniculatum* hairy roots which were cultured in medium supplemented with sucrose at range 3%–5% was higher than in medium supplemented with 6% sucrose. In this research, the maximum saponin content reached 71 365 mg/L/g dry weight in the medium supplemented with 5% sucrose (Figure 2 and Table 1).

Detection of hairy root growth of *T. paniculatum* in various concentrations of sucrose was done by measurement of total

sugar, pH and electric conductivity (EC) medium during 14-day cultivation (Figure 3). Total sugar of all treatments at different sucrose concentrations in BTBB for 14 days showed the decrease trend (Figure 3A).

In this research, the uptake of macronutrients and micronutrients by hairy roots was shown by changes of EC medium. During the first week, the EC value in all culture medium decreased gradually (Figure 3C). The biggest changes of total sugar and EC values were obtained in medium culture supplemented with 5% sucrose and the changes indicated the ability of hairy roots to absorb sucrose, macronutrients and micronutrients. During the cultivation, the pH value of all culture medium decreased (Figure 3B).




Figure 2. Biomass and saponin content of *T. paniculatum* hairy roots in various concentrations of sucrose after 14-day culture.

Table 1

Saponin content of *T. paniculatum* hairy roots in various concentrations of sucrose and potassium nitrate for 14 days.

Treatment	Sucrose concentration (%)	Potassium nitrate strength (\times MS)	Saponin (mg/L/g dry weight)
Sucrose concentration (%)	3.0		70717
	4.0		70717
	5.0		71365
	6.0		44030
Potassium nitrate strength (\times MS)		0.5	63299
		1.0	70717
		1.5	70726
		2.0	70730




Figure 1. Induction of *T. paniculatum* hairy roots in MS0 medium.

A: 9 days old in solid medium containing 500 mg/L cefotaxime; B: 16 days old in semi-solid medium containing 500 mg/L cefotaxime; C: 23 days old in liquid medium containing 500 mg/L cefotaxime; D: 30 days old as inoculum in liquid culture in BTBB.




Figure 3. Total sugar content, pH, and EC of culture medium of *T. paniculatum* hairy roots during 14 days cultivation in various concentration of sucrose.




Figure 4. Biomass and saponin content of *T. paniculatum* hairy roots at various concentrations of potassium nitrate after 14-day culture.

3.3. Effect of potassium nitrate at various concentrations on biomass and saponin content

The hairy root biomass of *T. paniculatum* at various potassium nitrate concentrations is shown in Figure 4. Supplement with potassium nitrate at 1–2 strength of MS medium could increase hairy root biomass after 14-day cultivation. The biomass and saponin content were increased at potassium nitrate 1–2 strength.

Hairy root growth of *T. paniculatum* was shown by profile of total sugar, pH, and EC medium during 14-day cultivation (Figure 5). Total sugar and EC values from the start till the end of culture showed down-trend line (Figure 5A and 5C). The saponin content of *T. paniculatum* hairy roots was analyzed by two methods. The results of qualitative and quantitative test by




Figure 5. Profile of total sugar content, pH and EC medium of *T. paniculatum* hairy roots during 14-day cultivation in various concentrations of potassium nitrate.




Figure 6. Chromatogram of ethanol extract of *T. paniculatum* hairy roots after cultured in 1000 mL BTBB containing 400 mL MS medium with various concentrations of sucrose and potassium nitrate for 14 days.

1: Saponin standard; 2: At 3% of sucrose w/v; 3: At 4% of sucrose w/v; 4: At 5% of sucrose w/v; 5: At 6% of sucrose w/v; 6: At 1 potassium nitrate strength of MS medium; 7: At 0.5 potassium nitrate strength of MS medium; 8: At 1.5 potassium nitrate strength of MS medium; 9: At 2 potassium nitrate strength of MS medium.

using thin layer chromatography and HPLC are shown in Figure 6 and Table 1 respectively.

4. Discussion

Sucrose is the source of carbon that is widely used in cell culture of plants, animals, fungi and bacteria. Cells use sucrose for energy and biosynthesis, including the biosynthesis of secondary metabolites. The maximum absorption of sucrose and the other nutrients increased dry weight and saponin content of hairy roots that were cultivated in medium supplemented with 5% sucrose (Figure 2).

Ability of *T. paniculatum* hairy roots to consume sucrose, macronutrients and micronutrients had correlation with pH medium. The initial pH medium was around 5.3 after autoclaving. The cultures supplemented with 3%, 4% and 6% sucrose underwent a little changes of total sugar content and EC value during cultivation. These changes correlated with pH value of culture medium. Decrease of pH in culture medium supplemented with 3%, 4% and 6% sucrose was drastical and it begun on the second day of cultivation. The pH medium was changed from 5.3 to 4.0, so the culture medium became too acid for hairy root growth. Acidity of medium did not cause the maximization of absorption of sucrose, macronutrients and micronutrients. Contrary with the other cultures, the change of

total sugar content and EC value in medium supplemented with 5% sucrose was bigger than medium supplemented with 3%, 4%, and 6% sucrose. It was also correlated with pH changes in medium during cultivation. The pH changes of medium supplemented with 5% sucrose were not drastical like the others. The pH value around 4.5–5.0 in medium was begun from the second day till the end of culture, so the absorption of sucrose, macronutrients and micronutrients was maximum. Therefore, growth of hairy roots was enhanced until 2 weeks. It is shown that hairy roots cultured were at exponential phase. This phenomenon was supported by the earlier study which reported that adventitious root growth typically exhibited a lag phase from 0 to 1 week, an exponential phase from 1 to 5 weeks, a stationary phase from 5 to 6 weeks, and a declining phase thereafter [11]. Concentrations of sucrose also effected on growth and saponin content of *T. paniculatum* hairy roots. Sucrose at concentrations of 3%–6% in culture medium was able to increase dry weight and saponin content of *T. paniculatum* hairy roots, but 5% sucrose was known as the best concentration of sucrose to enhance dry weight and saponin content. The effect of sucrose concentrations on growth and production of withanolide A was also reported on *Withania somnifera* hairy root culture. Among the various concentrations of sucrose (1%–8% w/v) tested, 3% sucrose favored the highest accumulation of biomass, whereas 4% sucrose concentration favored the highest production of withanolide A [9]. MS liquid medium containing sucrose 3% with an initial pH 5.8 favored the maximum biomass accumulation of *Solanum trilobatum* L. hairy roots [12].

Furthermore, hairy root culture of *Glycyrrhiza inflata* in MS medium with various concentrations of sucrose obtains the maximum biomass and glycyrrhizin in MS medium containing 12% and 6% sucrose, respectively [13]. Concentration of sucrose in medium also influenced the biomass production of *Artemisia vulgaris* hairy roots. Biomass accumulation was the highest when the medium was supplemented with 40 g/L sucrose [14].

Nitrogen is an essential element that composes most plant body, such as 1.5%–2.0% of dry weight of plant body and around 16.0% of total protein of plant body. In the cytoplasmic roots, nitrate as a nitrogen source is saved in vacuole and transported by xylem to all around the plant body. Nitrate is also as a key enzyme in amylum biosynthesis pathway, so that the carbon bone for amylum is shifted to carbon bone for nitrogen assimilation. The changes of amylum biosynthesis pathway to nitrogen assimilation pathway will increase proteins and enhance the plant growth rate. Generally, biomass growth rate has a direct correlation to secondary metabolite contents. All treatments of various concentrations of potassium nitrate enhanced dry weight and saponin content of *T. paniculatum* hairy roots. It indicated that the hairy roots were still in growth during cultivation of 14 days. In this research, the activity of hairy roots to absorb potassium nitrate for biomass growth and saponin formation was proved by decreasing trend line of EC curve. Enhancement of potassium nitrate from 950 mg/L (0.5 strength) to 3800 mg/L (2.0 strength) influenced biomass and saponin content of hairy roots of *T. paniculatum*. Dry weight and saponin content tended to increase with increasing concentrations of potassium nitrate (Figure 4 and Table 1).

Various concentrations of potassium nitrate in this research also affected the ratio of ammonium to nitrate and differentiation of total nitrogen concentration in each culture medium.

However, the results of dry weight and saponin content of *T. paniculatum* hairy roots proved that the ratio of ammonium to nitrate and different concentrations of total nitrogen influenced the production of secondary metabolites. This phenomenon was also reported in another production of secondary metabolites. The accumulation of secondary metabolites in the adventitious roots of *Eurycoma longifolia* was affected by $\text{NH}_4^+ : \text{NO}_3^-$ ratio. The secondary metabolites increased with NH_4^+ , and the highest levels were obtained when NH_4^+ was the sole nitrogen source, which was contrasted with the root growth results [9]. Productivity of biomass and bioactive compounds through bioreactor culture was also affected by medium salt strength, such as accumulation of bioactive compounds in the adventitious roots of *Eleutherococcus koreanum* Nakai, which indicated that $1/2$ MS was the optimal salt strength for the production of both biomass and bioactive compounds [15].

Various concentrations of sucrose in liquid culture by using MS0 medium could influence growth of hairy roots of *T. paniculatum*. The maximum biomass was reached in culture supplemented with 6% sucrose and approximately three fold higher than control. Culture supplemented with potassium nitrate at 2.0 strength of normal MS0 medium could increase dry weight of hairy roots until almost threefold higher than control. However, the maximum saponin content was obtained in culture supplemented with 5% sucrose and 2 strength potassium nitrate of normal MS0. Based on this research, those conditions can be used to produce biomass and saponin of hairy root of *T. paniculatum* in the large scale.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

We thank Dr. Ermawaty for providing *A. rhizogenes* strains LB510 and YMB072001. This research was supported by grant from Universitas Airlangga, Surabaya, Indonesia with Grant No. 8714/UN3/KR/2013.

References

- [1] Zuo G, Guan T, Chen D, Li C, Jiang R, Luo C, et al. Total saponins of *Panax ginseng* induces K562 cell differentiation by promoting internalization of the erythropoietin receptor. *Am J Chin Med* 2009; **37**(4): 747-57.
- [2] Hu XY, Neill SJ, Cai WM, Tang ZC. Activation of plasma membrane NADPH oxidase and generation of H_2O_2 mediate the induction of PAL activity and saponin synthesis by endogenous elicitor in suspension-cultured cells of *Panax ginseng*. *J Integr Plant Biol* 2003; **45**(12): 1434-41.
- [3] Mehrotra S, Kukreja AK, Khanuja SPS, Mishra BN. Genetic transformation studies and scale up of hairy root culture of *Glycyrrhiza glabra* in bioreactor. *Elect J Biotechnol* 2008; **11**(2): 1-7.
- [4] Yogananth N, Jothi Basu M. TLC method for determination of plumbagin in hairy root culture of *Plumbago rosea* L. *Glob J Biotechnol Biochem* 2009; **4**(1): 66-9.
- [5] Majumdar S, Garai S, Jha S. Genetic transformation of *Bacopa monnieri* by wild type strains of *Agrobacterium rhizogenes* stimulates production of bacopa saponins in transformed calli and plants. *Plant Cell Rep* 2011; **30**(5): 941-54.
- [6] Thiruvengadam M, Praveen N, Kim EH, Kim SH, Chung IM. Production of anthraquinones, phenolic compounds and biological

- activities from hairy root cultures of *Polygonum multiflorum* Thunb. *Protoplasma* 2014; **251**(3): 555-66.
- [7] Thiruvengadam M, Praveen N, Maria John KM, Yang YS, Kim SH, Chung IM. Establishment of *Momordica charantia* hairy root cultures for the production of phenolic compounds and determination of their biological activities. *Plant Cell Tissue Organ Cult* 2014; **118**(3): 545-57.
- [8] Praveen N, Murthy HN. Synthesis of withanolide A depends on carbon source and medium pH in hairy root culture of *Withania somnifera*. *Ind Crops Prod* 2012; **35**(1): 241-3.
- [9] Lulu T, Park SY, Ibrahim R, Paek KY. Production of biomass and bioactive compounds from adventitious roots by optimization of culturing conditions of *Eurycoma longifolia* in balloon-type bubble bioreactor system. *J Biosci Bioeng* 2015; **119**(6): 712-7.
- [10] Manuhara YSW, Yachya A, Kristanti AN. Effect of aeration and inoculum density on biomass and saponin content of *Talinum paniculatum* Gaertn. hairy roots in balloon-type bubble bioreactor. *J Pharm Biomed Sci* 2012; **2**(4): 47-52.
- [11] Cui XH, Murthy HN, Jin YX, Yim YH, Kim JY, Paek KY. Production of adventitious root biomass and secondary metabolites of *Hypericum perforatum* L. in a balloon type airlift reactor. *Bioresour Technol* 2011; **102**(21): 10072-9.
- [12] Shilpha J, Satish L, Kavikkul M, Joe Virgin Lorgia M, Ramesh M. Methyl jasmonate elicits the solasodine production and antioxidant activity in hairy root cultures of *Solanum trilobatum* L. *Ind Crops Prod* 2015; **71**: 54-64.
- [13] Wongwicha W, Tanaka H, Shoyama Y, Putalun W. Methyl jasmonate elicitation enhances glycyrrhizin production in *Glycyrrhiza inflata* hairy roots cultures. *Z Naturforsch C* 2011; **66**: 423-8.
- [14] Sujatha G, Zdravković-Korać S, Čalić D, Flamini G, Ranjitha Kumari BD. High-efficiency *Agrobacterium rhizogenes*-mediated genetic transformation in *Artemisia vulgaris*: hairy root production and essential oil analysis. *Ind Crops Prod* 2013; **44**: 643-52.
- [15] Lee EJ, Paek KY. Enhanced productivity of biomass and bioactive compounds through bioreactor culture of *Eleutherococcus koreanum* Nakai adventitious roots affected by medium salt strength. *Ind Crops Prod* 2012; **36**(1): 460-5.



Asian Pacific Journal of Tropical Biomedicine

48

H Index

Country [Netherlands](#) -  [SJR Ranking of Netherlands](#)

Subject Area and Category [Biochemistry, Genetics and Molecular Biology](#)
[Biochemistry, Genetics and Molecular Biology \(miscellaneous\)](#)

Publisher [Medknow Publications](#)

Publication type Journals


ISSN 22211691

Coverage 2011-ongoing

 [Join the conversation about this journal](#)


Quartiles

Biochemistry, Genetics and Molecular Biology (miscellaneous)









2012 2013 2014 2015 2016 2017 2018

SJR



Citations per document

Total Cites Self-Cites 



Asian Pacific Journal of Tropical Biomedicine

Q2 Biochemistry, Genetics and Molecular Biology...
best quartile

SJR 2018
0.49

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scima
```

Leave a comment

Name

Email

(will not be published)



I'm not a robot

reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2019. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)
