


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

Natural Product Communications

Table of Contents

Volume 15 Issue 5, May 2020

Special Collection in honor of Professor Dr. Gerald Blunden's 80th Birthday Review

Aspects in the Total Syntheses of Higher Terpenoids Starting From Wieland–Miescher Ketone and Its Derivative: A Review

Hisahiro Hagiwara 

First Published May 29, 2020

[Abstract](#)
> [Preview](#)



Review

Phytoconstituents of Genus *Micromelum* and Their Bioactivity—a Review

Tin Myo Thant, Nanik Siti Aminah , Alfinda Novi Kristanti, Rico Ramadhan, Hnin Thanda Aung, Yoshiaki Takaya

First Published May 29, 2020

[Abstract](#)
> [Preview](#)



 **Quantitative Analysis of 7 Compounds in *Diospyros lotus* Leaf Extract and Its Biological Effects on Neuroprotection and Antineuroinflammation**


Yu Jin Kim , Hye-Sun Lim, Joo-Hwan Kim, MinKyun Na , Soo-Jin Jeong

First Published May 28, 2020

[Abstract](#)
> [Preview](#)



Special Collection in Honor of Professor Dr. Gerald Blunden's 80th Birthday
Original Article

 **23-O-Substituted-2,3-Dehydrosilybins Selectively Suppress Androgen Receptor-Positive LNCaP Prostate Cancer Cell Proliferation**

Ziran Jiang, Arman Sekhon, Yogeshwari Oka, Guanglin Chen, Nagat Alrubati, Jasleen Kaur, Alexia Orozco, Qiang Zhang, Guangdi Wang, Qiao-Hong Chen 

First Published May 27, 2020

[Abstract](#)
> [Preview](#)



Natural Products as Sources of Cytotoxic Compounds with Potential for Development into Anti-cancer Agents
Original Article

 **In Vitro Antitumor Activities of Biologically Active Extracts and Isolated Phytochemicals From *Rabdosia latifolia* C. Y. Wu Et H. W. Li in Addenda**

Guang-Ping Liang , Ji-Xin Li, Jue Zhang, Ming Zhang, Jun Yang, Guang Ping Liang, Jie Wu

First Published May 27, 2020

[Abstract](#)
> [Preview](#)



Short Communication

 **Oleanane-Type Saponins From the Aerial Parts of *Panax bipinnatifidus***

Nguyen Huu Tung , Nguyen Thi Hoang Anh, Chu Van Tan, Pham Trung Hieu, Pham Quang Duong, Le Dang Quang, Vu Dinh Hoang

First Published May 20, 2020

[Abstract](#)
> [Preview](#)



Natural Products as Sources of Cytotoxic Compounds with Potential for Development into Anti-cancer Agents
Original Article

 **Mechanism Analysis of Coix Seed in Gastric Cancer Treatment Based on Biological Network Modules**


Fengbin Zhang, Xiaoyan Liu, Bingjie Huo, Bing Li , Ruixing Zhang

First Published May 20, 2020

[Abstract](#)
> [Preview](#)



Bioactive Natural Products from Terrestrial and Marine Resources, Especially Terpenes, but Not Limited to
Original Article

 **Chemical Composition, Antioxidant, and Antimicrobial Activities of the Essential Oils From *Artemisia annua* L. Growing Wild in Tajikistan**

Farukh S Sharopov , Aminjon Salimov, Sodik Numonov, Abduahad Safomuddin, Mahinur Bakri, Tojiddin Salimov, William N Setzer , Maidina Habasi

First Published May 20, 2020

[Abstract](#)
> [Preview](#)



Bioactive Natural Products from Terrestrial and Marine Resources, Especially Terpenes, but not limited to
Original Article

Chemical Composition of Volatile Oils of Fresh and Air-Dried Buds of Cannabis chemovars, Their Insecticidal and Repellent Activities

Amira S. Wanas, Mohamed M. Radwan, Suman Chandra, Hemant Lata, Zlatko Mehmedic, Abbas Ali, KHC Baser, Betul Demirci, Mahmoud A. ElSohly 

First Published May 19, 2020

[Abstract](#)
> [Preview](#)



Natural Products as Sources of Cytotoxic Compounds With Potential for Development Into Anti-Cancer Agents
Original Article

***Pinus maritima* Extract Induces Apoptosis in Human Malignant Melanoma Cells via ROS/Caspase-3 Signaling**

Sunisa Thaichinda, Salunya Tanchaen , Takuro Kanekura, Yuko Higashi, Pornpen Dararat, Kiyoshi Kikuchi, Thamthiwat Nararatwanchai

First Published May 19, 2020

[Abstract](#)
> [Preview](#)



 **Antimycobacterial Activity of Milemarinol, a New Squalene-Type Triterpene, and Other Isolate?**


Samyra I. S. Boeno, Michel de S. Passos, Mariana Félix, Sanderson D. Calixto, Almir R. C. Júnior, Lara F. Barbosa Siqueira, Michelle F. Muzitano, Raimundo Braz-Filho, Ivo J. Curcino Vieira

First Published May 18, 2020

[Abstract](#)
> [Preview](#)



Original Article

 **Synthesis of 3,15-Disuccinate-12-Coumarin Substituted Andrographolide Derivatives and Their Antiplatelet Aggregation Activities In Vitro**

Xue Li , Jiang-Wei Wang, Bin Huang, Zhi-Xiang Peng, Yuan-Yuan Zhang, Shi-Yun Zhao

First Published May 12, 2020

[Abstract](#)
> [Preview](#)



 **Nonacosan-10-ol and *n*-Alkanes in Needles of *Pinus halepensis***

Biljana Nikolić , Marina Todosijević, Iris Đorđević, Jovana Stanković, Zorica S. Mitić, Vele Tešević, Petar D. Marin

First Published May 12, 2020

[Abstract](#)
> [Preview](#)





A Survey of Antifungal Activity of Selected South African Plant Species Used for the Treatment of Skin Infections

Ndzalama Shikwambana, Salome Mamokone Mahlo

First Published May 12, 2020

[Abstract](#)
> [Preview](#)



Special Collection in Honor of Professor Dr. Gerald Blunden's 80th Birthday
Original Article



Chemical Profiling of *Astragalus membranaceus* Roots (Fish.) Bunge Herbal Preparation and Evaluation of Its Bioactivity

Valentina Santoro, Valentina Parisi, Massimiliano D'Ambola, Chiara Sinisgalli, Magnus Monné , Luigi Milella, Rosario Russo, Lorella Severino, Alessandra Braca , Nunziatina De Tommasi

First Published May 12, 2020

[Abstract](#)
> [Preview](#)



Short Communication



Piceid Nanoparticles Stabilized by Anionic Phospholipids for Transdermal Delivery

Noriyuki Uchida , Masayoshi Yanagi, Hiroki Hamada

First Published May 12, 2020

[Abstract](#)
> [Preview](#)



Original Article



Nonacosan-10-ol and *n*-Alkanes in Leaves of *Pinus pinaster*

Biljana Nikolić , Marina Todosijević, Iris Đorđević, Jovana Stanković, Zorica S. Mitić, Vele Tešević, Petar D. Marin

First Published May 12, 2020

[Abstract](#)
> [Preview](#)



Network Pharmacology Research and Preliminary Verification of Gegen Qinlian Decoction for the Treatment of Non-Alcoholic Fatty Liver Disease

Pengfei Hao , Yiyi Xiong, Hezhen Wu, Yanfang Yang

First Published May 11, 2020

[Abstract](#)
> [Preview](#)



Natural Products as Sources of Cytotoxic Compounds with Potential for Development into Anti-Cancer Agents
Original Article



Cytotoxic Triterpenes From *Trametes orientalis*

Kun-Min Xiao, Jun-Ming Wang, Ling-Yun Zhou, Lei Guo, Da-Cong Tan, Qi Yan, Yan Hua 

First Published May 11, 2020

[Abstract](#)
> [Preview](#)



Flavonoids and Related Compounds as Natural Products
Original Article



The Role of C-8 OH on the Antioxidant Activity of Norwogonin and Isowogonin

Qiushan Zhang, Jin Shao, Tong Zhao, Lei He , Huiping Ma, Linlin Jing 

First Published May 8, 2020

[Abstract](#)
> [Preview](#)



Terpenoids and Related Compounds – Structure, Synthesis, and Biological Activity
Preface

Preface

Motoo Tori

First Published May 7, 2020

[Abstract](#)



Original Article

Novel 1,3-Benzodioxole From Marine-Derived Actinomycete in East Vietnam Sea

Dao Phi Thi, Huong Doan Thi Mai, Duc Danh Cao, Quyen Vu Thi, Mai Anh Nguyen, Hong Minh Le Thi, Dang Thach Tran, Van Minh Chau, Van Cuong Pham 

First Published May 7, 2020

[Abstract](#)
> [Preview](#)



Flavonoids and Related Compounds as Natural Products
Original Article

Anthocyanin Profile and Antioxidant Property of Anti-asthma Flowers of *Cordyline terminalis* (L.) Kunth (Agavaceae)

Christopher Adaku , Irene Skaar, Robert Byamukama, Monica Jordheim , Øyvind M. Andersen

First Published May 7, 2020

[Abstract](#)
> [Preview](#)



Original Article



Justicidin B Inhibits PDGF-BB-Induced Proliferation and ECM Accumulation in Mesangial Cells via Nrf2/HO-1 and Akt/mTOR Signaling Pathway

Zhong-zhu Ai , Shan-shan Zhou, Song-tao Wu, Zong-chao Hong , Yu Zhang, Meng-fan Wang, Yan-fang Yang, He-zhen Wu

First Published May 7, 2020

[Abstract](#)
> [Preview](#)



Natural Product Communications

Editorial Board

[Hide All](#)

Editor-in-Chief

Dr. Pawan K. Agrawal

Natural Product, Inc

Honorary Editor

Professor Gerald Blunden

University of Portsmouth, UK

Editors

**Professor Vladimir I.
Kalinin**

G.B. Elyakov Pacific Institute of Bioorganic
Chemistry, Russia

Professor Phan Van Kiem

Vietnam Academy of Science and Technology
(VAST), Vietnam

**Professor Carmen Martin-
Cordero**

University of Seville, Spain

**Professor Manfred G.
Reinecke**

Texas Christian University, USA

Professor Sergio Rosselli

Università Degli Studi di Palermo, Italy

Professor Tadaaki Satou	International University of Health and Welfare, Japan
Professor William N. Setzer	University of Alabama in Huntsville, USA
Professor Ping-Jyun Sung	National Museum of Marine Biology and Aquarium
Professor Yasuhiro Tezuka	Hokuriku University, Japan
Professor David E. Thurston	King's College London, UK

Coordinating Editor

Chandan Agrawal	Natural Product, Inc
------------------------	----------------------

Advisory Editorial Board

Professor Giovanni Appendino	Universita del Pimonte Orientale, Italy
Professor Yoshinori Asakawa	Tokushima Bunri University, Japan
Professor Vassya Bankova	Bulgarian Academy of Science, Bulgaria
Professor Anna R. Bilia	University of Florence, Italy
Professor Maurizio Bruno	University of Palermo, Italy
Professor Geoffrey Cordell	Natural Products Inc.
Professor Fatih Demirci	Anadolu University, Turkey
Professor Francesco Epifano	D'Annunzio University of Chieti-Pescara, Rovigo, Italy
Professor Antonio Evidente	University of Naples "Federico II", Napoli, Italy
Professor Ana Cristina Figueiredo	Universidade de Lisboa, Portugal
Professor Mary J. Garon	The University of Queensland, Australia

Professor Cristina Gracia-Viguera	CEBAS-CSIC, Spain
Dr. Christopher Gray	University of New Brunswick, Canada
Professor Dominique Guillaume	Université de Reims-Champagne-Ardenne, France
Professor Hisahiro Hagiwara	Niigata University, Japan
Professor Judith Hohmann	University of Szeged, Hungary
Professor Tsukasa Iwashina	National Museum of Nature and Science, Japan
Professor Niel A. Koorbanally	University of KwaZulu-Natal, South Africa
Professor George A. Kraus	Iowa State University, USA
Professor Chiaki Kuroda	Rikkyo University, Japan
Professor M. A. Lacaille-Dubois	Université de Bourgogne, France
Professor Shoei-Sheng Lee	National Taiwan University
Professor Yoshihiro Mimaki	Tokyo University of Pharmacy and Life Sciences (TUPL), Japan
Professor Robert W. Owen	German Cancer Research Centre, Germany
Professor M. Soledade C. Pedras	University of Saskatchewan, Canada
Professor Luc Pieters	University of Antwerp, Belgium
Professor Phila Raharivelomanana	Université de la Polynésie Française, French Polynesia
Professor Jesús M. Rodilla	Universidade da Beira Interior, Covilhã, Portugal
Professor Stefano Serra	C.N.R.-Istituto di Chimica del Riconoscimento Molecolare, Italy
Dr. Bikram Singh	CSIR-Institute of Himalayan Bioresource, India
Professor Alexios-Leandros Skaltsounis	University of Athens, Greece
Professor John I. Sorensen	University of Manitoba, Canada

Professor John E. Sorenson

University of Manitoba, Canada

Professor Marina Stefova

Saints Cyril and Methodius University of Skopje,
Macedonia

Professor Valentin Stonik

G.B. Elyakov Pacific Institute of Bioorganic
Chemistry, Russia

Professor Félix Tomi

Université de Corse-CNRS, France

Professor Karin Valant-
Vetschera

University of Vienna, Austria

Professor Johannes van
Staden

University of KwaZulu-Natal, South Africa

Dr. Alessandro Venditti

Università di Roma "La Sapienza", Roma, Italy

Assoc. Professor Yasunori
Yaoita

Tohoku Medical and Pharmaceutical University,
Japan

More about this journal



Description



Aims and Scope



Abstracting/Indexing



Submit Paper

Phytoconstituents of Genus *Micromelum* and Their Bioactivity—a Review

Natural Product Communications
Volume 15(5): 1–15
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1934578X20927124
journals.sagepub.com/home/npx



Tin Myo Thant^{1,2}, Nanik Siti Aminah¹ , Alfinda Novi Kristanti¹, Rico Ramadhan¹, Hnin Thanda Aung³, and Yoshiaki Takaya⁴

Abstract

The genus *Micromelum* belongs to the Rutaceae family. As its rich bioactive constituents its stems, flowers, leaves, and roots have been used in traditional medicine, for the treatment of various diseases from ancient time. Phytochemically, many bioactive compounds, including coumarins, polyoxygenated flavonoids, phenylpropanoic acid derivatives, quinolone alkaloids, and also carbazole alkaloids, have been reported as secondary metabolites of the *Micromelum* spp. including many new compounds. Therefore, *Micromelum* spp. are considered potential for drug leads. In this article, we present an overview of secondary metabolites isolated from genus *Micromelum* and their bioactivities that have been reported between 1982 and 2019.

Keywords

Micromelum, Rutaceae, coumarins, flavonoids, quinolone and carbazole alkaloids

Received: November 27th, 2019; Accepted: April 22nd, 2020.

Introduction to Chemical Constituents of Genus *Micromelum* and Their Bioactivity

Micromelum spp. (Rutaceae) consist of 9 species found mostly in Samoa, Fiji, and Tonga islands, Australia, Southeast Asia, southern China, Ceylon, northeastern India, and West Pakistan.^{1,2} *Micromelum* spp. are known for being rich in coumarins such as the 6- and 8-prenylated coumarins. In addition, the carbazole alkaloids, dihydrocinnamic acid derivatives, and flavonoids have been isolated from it.^{3,4} The leaves and stems of *Micromelum* species have been found to contain coumarins, phenylpropanoic acid derivatives, polyoxygenated flavonoids, and also alkaloids.^{5,6} The following *Micromelum* species have been well investigated due to their pharmaceutical properties.

M. integerrimum is a tree that grows up to 8 m high. Its young parts have rust-colored pubescent. It usually grows in moist mountain forests, maritime thickets in sandy soil; near sea level to 2000 m and is widely distributed in China, Bhutan, Cambodia, India, Laos, Myanmar, Nepal, Thailand, and Vietnam.⁷ *M. integerrimum* is highly diversified for its secondary metabolites, and apart from acridone and carbazole alkaloids, a variety of coumarin derivatives, especially 6,7-di and 7,8-disubstituted coumarin core structures, having at least a prenyl unit have been isolated. Naturally occurring coumarin derivatives from *Micromelum* spp. exhibit a variety of biological activities including anti-corpulence, cytotoxicity, anti-platelet, and antimutagenicity.³

M. minutum is a small spineless tree that can reach up to 3 m in height. It is widely distributed in Southeast Asia and in Pacific islands. The stems, flowers, leaves, and roots are used pharmaceutically for a variety of indications.^{8,9} The plant is known to contain coumarins, some of which are active compounds showing strong cytotoxic activities. The leaves are traditionally used to treat fever and giddiness. The poultice of the boiled roots is used for ague.¹⁰ In Indochina, the roasted and crushed leaves are rubbed to skin irritated by scabies and are considered to be emmenagogues.¹¹ Previous phytochemical investigations on the different parts of *M. minutum* have showed the presence of coumarins, triterpenes, alkaloids, and phenylpropanoids.¹²

M. falcatum is a tree of 1-3 m high and its leaflet blades are alternate, ovate to lanceolate, in equilateral. Flowers are ellipsoid or broad in bud. Petals are white, oblong, outside glabrous, or

¹Dept of Chemistry, Fac of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia

²Department of Chemistry, Mandalay Degree College, Mandalay, Myanmar

³Department of Chemistry, Mandalay University, Myanmar

⁴Faculty of Pharmacy, Meijo University, Yagotoyama, Tempaku, Nagoya, Japan

Corresponding Author:

Nanik Siti Aminah, Dept of Chemistry, Fac of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia.

Email: nanik-s-a@fst.unair.ac.id






Figure 1. Coumarins from *Micromelum minutum*, *M. integerrimum*, *M. falcatum*, and *M. zeylanicum*.

pubescent. Fruits are ellipsoid to ovoid, 1- or 2-seeded. Its flowering period is from January to April. It is usually found in the mountain area at heights below 1200 m in China, Cambodia, Laos, Myanmar, Thailand, and Vietnam.¹³ Its leaves and roots are used for treating infected wounds, odynolysis, rheumatism, muscular atrophy, and insect bites.^{4,14} *M. falcatum* has been found to contain several coumarins, such as 7-oxygenated coumarins, dihydrocinnamic acid derivatives, the dimeric indole alkaloid (yuehchukene), the carbazole alkaloid, 5,6-pyranoglycozoline, the

6-prenylated coumarin, micromelin, and the 8-prenylated coumarins, phebalosin and murpanidin.^{6,15,16} The dimeric indole alkaloid yuehchukene was found to have potent anti-implantation activity.¹⁷

M. compressum is a tree described as unarmed and glabrous. Its leaves are odd-pinnate, the leaflets are alternate, and the fruits are ovoid. Phytochemical efforts on the CH_2Cl_2 -methanol extract of *M. compressum* have resulted in the isolation and identification of 3 polymethoxylated flavones.¹




Figure 2. Coumarins isolated from *Micromelum falcatum*.

M. hirsutum is a tree of up to 10 m height with greenish-yellow flowers and aromatic leaves. It is widely distributed in peninsular Malaysia (Melaka and Pahang northward), Andaman Islands, Myanmar, Laos, Thailand, and Vietnam.¹⁸ In 2010, Rodphukdeekul research group investigated antitumor activity of the dichloromethane extract from branches of *M. hirsutum* on human B lymphoma cells, Ramos. The result demonstrated that the extract has antitumor potential against B lymphoma.¹⁹ In Myanmar its leaves are widely used as spices in curries. Phytochemical analysis of this species has reported the presence of many carbazole alkaloids, as well as the γ -lactone compounds.¹⁸

M. glanduliferum is a tropical or subtropical small tree found in Thailand. Coumarins and some carbazole alkaloids compounds, such as osthol, phebalosin, minumicrolin, murrangatin, murralongin, micormarin A, microminutinin, micromelin, and 2,7-dihydroxy-3-formyl-1-(30-methyl-20-butenyl)carbazole and 7-methoxy heptaphylline, were reported from the roots of *M. glanduliferum*.²⁰

The chemical analysis of leaves and stems of *M. zeylanicum* has reported an oxazole alkaloid, O-methylhalfordinol, a coumarin, micromelin, and a flavone, 5-hydroxy-3,3',4',7-pentamethoxyflavone. In addition, the stems contained 6-formyl-7-methoxycoumarin, and the leaves contained the carbazole alkaloid, koenigine, and *p*-sitosterol.²¹

In this review the articles were collected from 1982 to 2019. The main purpose of this article was to evaluate the diversity of secondary metabolites isolated from *Micromelum* spp. in order to provide a complete reference of the diverse secondary metabolites from *Micromelum* spp. To the best of our knowledge this is the first review about secondary metabolites isolated from genus *Micromelum*.

Phytochemical Aspects of *Micromelum* Spp. and Their Tested Bioactivities

Coumarins

Coumarins are highly active biological substances and major secondary metabolites isolated from *Micromelum* spp. They are widely used in medicine, perfumes, cosmetics, and laser dyes.⁶ Coumarins have antioxidant activity and they preserve the




Figure 3. Some coumarins isolated from *Micromelum integerrimum* and *M. minutum*.

levels of other antioxidants in human plasma. Coumarins also inhibited lipid peroxidation and increased the activity of antioxidants, superoxide dismutase, and catalase.²² Many bioactive coumarins have been reported from genus *Micromelum*, and the basic core coumarin compound, umbelliferone (1), was isolated from the roots of *M. minutum*.⁸ Scopoletin compound 7-hydroxy-6-methoxy-2H-chromen-2-one (2) was isolated from the fruits of *M. minutum*.²³ and 6-hydroxy-7-methoxy-2H-chromen-2-one (3) was isolated from the stems and leaves of *M. integerrimum*.²⁴ Another coumarin, 6-formyl-7-methoxycoumarin (4), was isolated from the leaves and stems of *M. zeylanicum*.²¹




Figure 4. Coumarins from *Micromelum minutum* and *M. falcatum*.




Figure 5. Coumarins from the *Micromelum minutum*.

7-Methoxyprenylated coumarin, osthonon (**5**), and osthol (**6**) were isolated from the roots of *M. minutum*.⁸ Microfalcrin (**7**) was isolated from the leaves of *M. falcatum*.¹⁶ This compound (**7**) was also found in *M. minutum*⁸ (Figure 1).

One-prenylated coumarins named micromarin-F (**8**), -G (**9**), and -H (**10**) were isolated from the stems of *M. minutum*.²⁵ The compound murralongin-1 (**11**) was isolated from the roots of *M. minutum*.⁸ Murralongin-2 (**12**) and murralonginol (**13**) were isolated from the fruits of *M. minutum*.^{23,26} 7-Oxygenated coumarins, phebalosin (**14**), murrangatin (**15**), murrangatin acetate (**16**), murracarpin (**17**), and minumicrolin (**18**) were isolated from the roots of *M. minutum*⁸ (Figure 1). Microminutin (**19**)

was isolated from the acetone extract of the stems of *M. minutum*.²⁵ Some coumarins, such as 7-methoxy-8-(4'-methyl-3'-furyl)coumarin (**20**), microminutin B (**21**), microminutin C (**22**), and micromeloside A (**23**), were isolated from the finely powdered fruits of *M. falcatum*.⁴ In 2014, Kouloura et al reported microcoumaririn (**24**) from the leaves of *M. falcatum*. The anti-inflammatory properties of compounds (**24**, **32**) were evaluated by measuring the inhibition of the proinflammatory mediator nuclear factor kappa B induction and nitric oxide (NO) production but these 2 compounds did not show any inhibition.¹⁶




Figure 6. Coumarins from the *Micromelum minutum*, *M. integerrimum*, and *M. falcatum*.

Micromelin or 7-methoxy-6-((1R,2R,5R)-5-methyl-4-oxo-3,6-dioxabicyclo[3.1.0]hexan-2-yl)-2H-chromen-2-one (**25**) was isolated from the fruits of *M. minutum*.²³ This compound is common to all species of *Micromelum* so far examined and can perhaps be considered as a chemotaxonomic marker for the genus.²⁷ The compound acetyldihydromicromelin A (**26**) was isolated from the aerial parts of *M. minutum*.²⁶ One-prenylated coumarin, namely hydramicromelin D (**27**), was isolated from the twigs of *M. integerrimum*.³ Hopeyhopol (**28**) and marmesin (**29**) were isolated from the whole plants of *M. minutum*.²⁸ (-) Decursinol (**30**) was isolated from the finely powdered fruits of *M. falcatum*.⁴ One-prenylated coumarins named micromarin-A (**31**), -B (**32**), and -C (**33**) were isolated from the seeds of *M. minutum*²² and fruits of *M. falcatum*¹⁶ (Figure 3).

Ester coumarinoids, (3*S*,4*S*)-3,4-dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-ethylenebutyl 3-methylbutanoate (**34**), (3*R*,4*S*)-3,4-dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl-3-methylbutanoate (**35**), and 3-hydroxy-4-methoxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl-3-ethylbutanoate (**36**), were isolated from the finely powdered fruits of *M. falcatum*.^{4,16} (Figure 4). Another ester coumarin, 7-methoxy-8-(2-hydroxymethyl-1-*O*-isovaleryl-4-butenyl) coumarin (**37**), was isolated from the stem bark of *M. falcatum*.²⁹

Murralonginol isovalerate-1 (**38**) was isolated from the stems of *M. minutum*.²³ Toxicity of compound (**37**) isolated

from *M. falcatum* was tested against brine shrimp larvae using a 96-well plate assay and the compound (**37**) exhibited an LC₅₀ value of 6.8 μ M, indicating that it was a potent toxic natural product and its in vitro antiproliferative activities against lung cancer (HvEvc) cell lines by the MTT method, and (**37**) displayed moderate activity against the HvEvc cell line with IC₅₀ value of 35.7.²⁹ Two-prenylated coumarins, murralonginol isovalerate-2 (**39**), minutuminolate (**40**), and 7-demethylmurralonginol isovalerate (**41**), were isolated from the fruits and roots of *M. minutum*^{8,23} (Figure 4). Micromarininate (**42**) was isolated from the EtOAc extract fruits of *M. falcatum*⁴ (Figure 4).

Five other geranylcoumarins, such as 8-hydroxyisocapnolactone-2',3'-diol (**43**), isolated from *M. minutum* have in vitro antiplasmodial activity against *Plasmodium falciparum* FCR-3 and D-10 strains with IC₅₀ values of 6.39 μ g/mL (16.99 μ M) and 24.23 mg/mL (64.45 μ M), respectively.¹⁰ 8-methoxycapnolactone (**44**), 3'',4''-dihydrocapnolactone (**45**), 2',3' epoxyisocapnolactone (**46**), 8-hydroxy-3'',4''-dihydrocapnolactone-2',3'-diol (**47**), and 8,4-dihydroxy-3,4-dihydrocapnolactone-2,3-diol (**48**) were isolated from the leaves of *M. minutum*.^{10,11,30} (Figure 5). These coumarins contained a lactone ring in their structure. In 2009, Ratna et al reported the cytotoxicity of these lactone coumarins on T-lymphoblastic leukemia (CEM-SS), promyelocytic leukemia (HL60), cervical cancer (HeLa), and liver cancer (HepG2) cell lines. Among the tested compounds, (**43**) showed the most active with IC₅₀ values of 2.9, 2.5, 6.9, and 5.9 μ g/mL, respectively.¹¹ In 2014, the same group investigated the cytotoxic activity of compound (**43**) on MCF-7 and T47D cells. The results showed IC₅₀ values of 8 μ g/mL (21.2 μ M) and 4 μ g/mL (10.6 μ M) respectively.³¹

Two monoterpene coumarins, minutin A (**49**) and minutin B (**50**), were isolated from the leaves of *M. minutum* (Figure 5) and their cytotoxic activity against *Leishmania major* and cancer cells were already studied. It has been reported that minutin B (**50**) has strong cytotoxic activity against the cancer cell lines such as SBC3, A549, K562, and K562/ADM with IC₅₀ values of 8.8, 10.1, 16.9, and 10.1 μ M, respectively.³²

The 7-methoxylated coumarin, micromelosidester (**51**), was isolated from the leaves of *M. falcatum*.¹⁶ Two compounds microminutin (**52**) and 6-methoxymicrominutin (**53**) were isolated from the leaves of *M. falcatum*.⁴ These 2 compounds are different from other compounds shown previously, because the prenyl attached on coumarin core forms a bicyclic structure. Other 2-prenylated coumarins named integerrimelin (**54**), (1*R*,3*R*,4*R*,6*S*)-4-(7-Methoxy-2-oxo-2H-chromen-6-yl)-1-methyl-3,6-dioxabicyclo[3.1.0]hexan-2-yl acetate (**55**), and scopolin (**56**) were isolated from the twigs, stems, and leaves of *M. integerrimum*.^{3,24,33} A glycoside coumarin 7-methoxy-8-(1-hydroxy-2-*O*- β -glucopyranosyl-3-methyl-4-butene-1-yl) coumarin (**57**) was isolated from the stem bark of *M. falcatum*.²⁹ Another coumarin, micromelumside D (**58**), was isolated from the aerial portions of *M. minutum*⁶ (Figure 6).

In 2018, Kassim et al reported a lignan sesamin (**118**) (Figure 7) together with 3 coumarins, hydramicromelinin (**59**), micromelinin (**60**), and 1-*O*- β -D-glucopyranosylmarmesin (**61**)




Figure 7. Miscellaneous compounds from the *Micromelum integerrimum*, *M. hirsutum*, *M. minutum*, and *M. falcatum*.

(Figure 8) from *M. minutum* methanol bark extract. Of the reported compounds, the ORAC measurements of compounds **59** and **118** showed potent antioxidant activity with the values of 5539 and 4031 $\mu\text{mol TE/g}$, respectively.³⁴

Alkaloids

Alkaloids have inspired the development of several antibacterial drugs through the synthesis of quinine serendipitously yielding the quinolones, structural alteration of azomycin yielding metronidazole, and working with the quinoline scaffold yielding bedaquiline. In other drugs, alkaloids are present as scaffold substructures, eg linezolid and trimethoprim. Alkaloids remain the focus of much research. Their development as antibacterial drugs pursued within academia, industry, and joint-ventures.³⁵ A large number of carbazole alkaloids have been

isolated from higher plants of the genera *Murraya*, *Glycosmis*, *Micromelum*, and *Clausena*, all belonging to the family rutaceae.^{36,37}

The quinoldione alkaloids 2-(3-hydroxy-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinolin-3-yl)acetate (**62**) and 3-hydroxy-1-methyl-3-(2-oxopropyl)quinoline-2,4(1H,3H)-dione (**63**) along with 2 quinolinone alkaloids *N*-methylflindersine (**64**), 4-hydroxy-3-methoxy-1-methyl-2(1H)-quinolinone (**65**), and the *N*-methylswietenidine-B (**66**) were reported from the stem of *M. falcatum* (Figure 9). The toxicity of all compounds (**62-66**) was tested in the brine shrimp larvae assay, and their LD_{50} values were 143, 355.0, 1.39, 2020.0, and 70.5 mg/mL, respectively, indicating that compound (**64**) was a potent toxic substance.¹⁷

The alkaloids myosmine (**67**), halfordinol (**68**), and *O*-methylhalfordinol (**69**) were reported from the leaves and




Figure 8. Coumarins from the *Micromelum minutum*.

young stems of *M. zeylanicum*.²¹ The methyl carbazole-3-carboxylate (**70**) was isolated from the leaves of *M. integerrimum*³⁸ (Figure 9).

Increased death rates due to lung cancer have necessitated the search for potential novel anticancer compounds such as carbazole derivatives. Carbazoles are aromatic heterocyclic




Figure 9. Alkaloids from *Micromelum falcatum*, *M. zeylanicum*, and *M. integerrimum*.

organic compounds, with a tricyclic structure composed of 2 benzene rings each fused onto a 5-membered nitrogen-containing ring with anticancer, antibacterial, and anti-inflammatory activity.

Anti-TB bioassay-directed fractionation led to the isolation of 6 carbazole alkaloids from the CH_2Cl_2 extract of the stem bark of *M. birsutum*. The carbazoles including 3-methylcarbazole (**71**), 3-formylcarbazole (**72**), lansine (**73**), 3-formyl-6-methoxycarbazole (**74**), and methyl carbazole-3-carboxylate (**75**) were reported from the stem barks of *M. birsutum*.¹⁸ A acridone alkaloid, 1,3-dihydroxy-4-methoxy-10-methylacridone (**76**), was isolated from the leaves of *M. integerrimum*³⁸ (Figure 10).

Two carbazole alkaloids, 2,7-dihydroxy-3-formyl-1-(30-methyl-20-butenyl)carbazole (**77**) and 7-methoxy heptaphylline (**78**), were isolated from the roots of *M. glanduliferum* by Siridechakorn et al in 1985. They might be a useful chemotaxonomic marker of the species *M. glanduliferum* and could be used to differentiate *M. glanduliferum* from other species of *Micromelum*.²⁰ Micromeline (**79**) was isolated from the stem barks of *M. birsutum*.¹⁸ The alkaloid compounds koenimbine (**80**), koenine (**81**), koenigine (**82**), and koenidine (**83**) were isolated from the leaves and young stems of *M. zeylanicum*²¹ (Figures 10 and 11).

In addition, a carbazole alkaloid, mahanine (**84**), was isolated from the leaves of *M. minutum* (Figure 11). It has been shown to exhibit a wide range of pharmacological effects including antimutagenicity against heterocyclic amines, antimicrobial activity against Gram positive bacteria, an anti-inflammatory effect, and it appeared to be cytotoxic against several cancer cell lines.³⁵

Phenethyl cinnamides, micrometam A (**85**), micrometam D (**86**), micrometam E (**87**), and micrometam B (**88**), and a core of novel marine compound, micrometam C (**89**), were reported from the whole plant of *M. falcatum*^{15,39} (Figure 11).

Flavones

A number of phytochemical studies dealing with polymethoxylated flavones from *Micromelum* spp. have been reported and a large number of flavonol-type metabolites were identified with the occurrence of partially methylated derivatives.¹

Polymethoxylated flavones, 5-hydroxy-3,4',7,8-tetramethoxyflavone (**90**), 5,7-dihydroxy-3,4',8-trimethoxyflavone (**91**), 5-hydroxy-3,4',7,8-tetramethoxyflavone (**92**), and 5-hydroxy-3,4',6,7,8-pentamethoxyflavone (**93**), were reported from *M. minutum*.⁴⁰ Other polymethoxylated flavones such as 3,5,7,4'-tetramethoxyflavone (**94**), 3,5,7,8,4'-pentamethoxyflavone (**95**), and 3,5,6,7,4'-pentamethoxyflavone (**96**) were also reported from the leaves of *M. compressum*¹ (Figure 12).

Three compounds, 5,7-dihydroxy-4',6,8-trimethoxyflavone (**97**), 7-hydroxy-4',5,6,8-tetramethoxyflavone (**98**), and 7-hydroxy-4',6,8-trimethoxyflavone-5-ylacetate (**99**), were isolated from the aerial parts of *M. minutum*.²⁶ 5-hydroxy-3,3',4',7,8-pen-




Figure 10. Alkaloids from *Micromelum minutum*, *M. birsutum*, *M. zeylanicum*, *M. glanduliferum*, *M. falcatum*, and *M. integerrimum*.

tamethoxyflavone (**100**) was also isolated from the leaves and stems of *M. zeylanicum*²¹ (Figure 12).

Coumarin and Flavone Condensed Binary Compounds

The compounds 8-((1*S*,2*R*)-1-hydroxy-2-(5-hydroxy-6, 8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4*H*-chromen-7-yl)oxy)-3-methylbut-3-enyl)-7-methoxy-2*H*-chromen-2-one (**101**),

8-((1*S*,2*R*)-1-hydroxy-3-methyl-2-(5,6,8-trimethoxy-2-(4-methoxyphenyl)-4-oxo-4*H*-chromen-7-yl)oxy)but-3-enyl)-7-methoxy-2*H*-chromen-2-one (**102**), and 7-((1*S*,2*R*)-1-hydroxy-1-(7-methoxy-2-oxo-2*H*-chromen-8-yl)-3-methylbut-3-en-2-yl)oxy)-6,8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4*H*-chromen-5-yl acetate (**103**) were reported from the above ground parts of *M. minutum*²⁶ (Figure 13).




Figure 11. Alkaloids from *Micromelum falcatum*.

Miscellaneous Compounds

The stigmasterol (**104**) was isolated from the leaves of *M. minutum*.¹⁰ Two triterpenes, 5(6)-gluten-3 α -ol (**105**) and 5(6)-gluten-3-on (**106**), were isolated from the leaves' and barks' crude petroleum ether and chloroform extracts of *M. minutum*.¹¹ The derivative of oleic acid with trivial name micro-molide (**107**) was isolated from the stem barks of *M. hirsutum*.¹⁸ (Figure 14).

In 2014, Zhi-yao et al have reported 2 phenylpropanoids, microintegerrin A (**108**) and microintegerrin B (**109**), from the air-dried stems and leaves of *M. integerrimum*. These 2 compounds were tested for their cytotoxicity against 3 cancer cell lines (HeLa, A549, and BGC-823) but did not show any activity.²⁴ Dihydrocinnamic methyl ester derivatives, 3-(5-((2R,3R,4R)-3,4-dihydroxy-4-methyl-5-oxotetrahydrofuran-2-yl)-2-hydroxy-4-methoxyphenyl)




Figure 12. Flavones from *Micromelum minutum*, *M. zeylanicum*, and *M. compressum*.

propanoic acid (**110**) and secomicromelin (**111**), were isolated from the fruits of *M. falcatum*⁴ (Figures 7 and 14). Three dihydrocinnamic acid derivatives, 3,4-dihydro-1,2-secomicrominin (**112**), 3,4-dihydro-1,2-secomicrominin methylester (**113**), and 3,4-dihydro-1,2-secomicrominin-9-*O*-glucoside (**114**), were isolated from the leaves of *M. falcatum*.¹⁴ Other 3 glucosides of phenylpropanoic acid derivatives,




Figure 14. Miscellaneous compounds from *Micromelum minutum* and *M. birsutum*.

micromelumosides A (**115**), micromelumosides B (**116**), and micromelumosides C (**117**), and sesamin (**118**) were reported from the aerial parts of *M. minutum*⁶ (Figure 7).




Figure 13. Coumarin and flavone condensed compounds from *Micromelum minutum*.

Conclusion

Micromelum spp. possess very rich sources of bioactive secondary metabolites. As a member of rutaceae family it contains mainly coumarins (from simple to prenyl, and geranyl), alkaloids (quinolidine and carbazole), polymethoxyflavones, and a few glycosides and phenylpropanoic acid derivatives (Table 1). The coumarins are the most abundant compounds from *Micromelum* spp., and some compounds, such as 8-hydroxyisocapnolactone-2',3'-diol, have positive effects on the anticancer activity. Currently, at least 118 compounds have been reported from different plant parts of *M. integerrimum*, *M. minutum*, *M. falcatum*, *M. compressum*, *M. glanduliferum*, *M. birsutum*, and *M. zeylanicum* (Table 1). Among them, *M. integerrimum*, *M. minutum*, and *M. falcatum* were well studied chemically and pharmacologically and others, *M. compressum*, *M. glanduliferum*, *M. birsutum*, and *M. zeylanicum*, have been studied less. So, the review suggests that there is a great potential for researchers to search drug lead secondary metabolites from the genus *Micromelum*.

Acknowledgments

Appreciation is conveyed to Prof. M. Iqbal Choudhary from H.E.J. Research Institute of Chemistry, International Center of Chemical and Biological Sciences, University of Karachi, Karachi-75270, Pakistan, for his help in improving this review article.

Table 1. Isolated Compounds Reported From *Micromelum* Spp.

No. of compounds	Plant parts	Name of species	References
Simple coumarins			
Umbelliferone (1)	Roots	<i>M. minutum</i>	8,13
7-Hydroxy-6-methoxy-2H-chromen-2-one (2)	Fruits	<i>M. minutum</i>	23
6-Hydroxy-7-methoxy-2H-chromen-2-one (3)	Leaves and stems	<i>M. integerrimum</i>	24
6-Formyl-7-methoxycoumarin (4)	Leaves and stems	<i>M. zeylanicum</i>	21
Osthenon (5)	Roots	<i>M. minutum</i>	8
Prenylated coumarins			
Osthol (6)	Leaves	<i>M. minutum</i>	8
Microfalcrin (7)	Roots	<i>M. falcatum</i>	16
Micromarin-F (8)	Stems	<i>M. minutum</i>	25
Micromarin-G (9)	Stems	<i>M. minutum</i>	25
Micromarin-H (10)	Stems	<i>M. minutum</i>	25
Murralongin-1 (11)	Fruits	<i>M. minutum</i>	8,23
Murralongin-2 (12)	Fruits	<i>M. minutum</i>	8,23
Murralonginol (13)	Fruits	<i>M. minutum</i>	8,23
Phebalosin (14)	Roots	<i>M. minutum</i>	8
Murrangatin (15)	Roots	<i>M. minutum</i>	8
Murrangatin acetate (16)	Roots	<i>M. minutum</i>	8
Murracarpin (17)	Roots	<i>M. minutum</i>	8
Minumicrolin (18)	Roots	<i>M. minutum</i>	8
Microminutin (19)	Stem	<i>M. minutum</i>	25
7-Methoxy-8-(4'-methyl-3'-furyl)coumarin (20)	Fruits	<i>M. falcatum</i>	4
Microminutin B (21)	Fruits	<i>M. falcatum</i>	4
Microminutin C (22)	Fruits	<i>M. falcatum</i>	4
Micromeloside A (23)	Fruits	<i>M. falcatum</i>	4
Microcoumaririn (24)	Leaves	<i>M. falcatum</i>	16
Micromelin (25)	Fruits	<i>M. minutum</i>	23
Acetyldihydromicromelin A (26)	Aerial parts	<i>M. minutum</i>	26
Hydramicromelin D (27)	Twigs	<i>M. integerrimum</i>	3
Hopeyhopol (28)	Whole plants	<i>M. minutum</i>	28
Marmesin (29)	Whole plants	<i>M. minutum</i>	28
(-)-Decursinol (30)	Fruits	<i>M. falcatum</i>	4
Micromarin-A (31)	Stems	<i>M. falcatum</i>	16,22
Micromarin-B (32)	Stems	<i>M. minutum</i>	16,22
Micromarin-C (33)	Stems	<i>M. minutum</i>	22
(3 <i>S</i> ,4 <i>S</i>)-3,4-Dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-ethylenebutyl 3-methylbutanoate (34)	Fruits	<i>M. falcatum</i>	4
(3 <i>R</i> ,4 <i>S</i>)-3,4-Dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl 3-methylbutanoate (35)	Fruits	<i>M. falcatum</i>	4
3-Hydroxy-4-methoxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl 3-ethylbutanoate (36)	Fruits	<i>M. falcatum</i>	4
7-Methoxy-8-(2-hydroxymethyl-1-O-isovaleryl-4-butenyl) coumarin (37)	Stem bark	<i>M. falcatum</i>	29
Murralonginol isovalerate-1 (38)	Stems	<i>M. minutum</i>	23
Murralonginol isovalerate-2 (39)	Roots	<i>M. minutum</i>	8
Minutuminolate (40)	Roots	<i>M. minutum</i>	8
7-Demethylmurralonginol isovalerate (41)	Roots	<i>M. minutum</i>	23
Micromarinate (42)	Fruits	<i>M. falcatum</i>	4
8-Hydroxyisocapnolactone-2',3'-diol (43)	Leaves	<i>M. minutum</i>	10,11,28,32
8-Methoxycapnolactone (44)	Leaves	<i>M. minutum</i>	10,11,28,32
3'',4''-Dihydrocapnolactone (45)	Leaves	<i>M. minutum</i>	10,11,28,32
2',3'-Epoxyisocapnolactone (46)	Leaves	<i>M. minutum</i>	10,11,28,32
8-Hydroxy-3'',4''-dihydrocapno lactone-2',3'-diol (47)	Leaves	<i>M. minutum</i>	10,11,32
8,4-Dihydroxy-3,4-dihydro capnolactone-2,3-diol (48)	Leaves	<i>M. minutum</i>	10,11,32
Minutin A (49)	Leaves	<i>M. minutum</i>	10,11,32
Minutin B (50)	Leaves	<i>M. minutum</i>	10,11,32
Micromelosidester (51)	Leaves	<i>M. falcatum</i>	4

(Continued)

Table 1. Continued

No. of compounds	Plant parts	Name of species	References
Microminutinin (52)	Leaves	<i>M. falcatum</i>	4
6-Methoxymicrominutinin (53)	Leaves	<i>M. falcatum</i>	16,25
Integerrimelin (54)	Twigs	<i>M. integerrimum</i>	3
(1R,3R,4R,6S)-4-(7-Methoxy-2-oxo-2H-chromen-6-yl)-1-methyl-3,6-dioxabicyclo[3.1.0]hexan-2-yl acetate (55)	Leaves	<i>M. integerrimum</i>	33
scopolin (56)	Stems and leaves	<i>M. integerrimum</i>	24
7-Methoxy-8-(1-hydroxy-2-O- β -glucopyranosyl-3-Methyl-4-butene-1-yl) coumarin (57)	Stem bark	<i>M. falcatum</i>	29
Micromelumside D (58)	Aerial parts	<i>M. minutum</i>	6
Hydramicromelinin (59)	Bark	<i>M. minutum</i>	34
Micromelinin (60)	Bark	<i>M. minutum</i>	34
1-O- β -D-Glucopyranosylmarmesin (61)	Bark	<i>M. minutum</i>	34
Alkaloids			
2-(3-Hydroxy-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinolin-3-yl)acetate (62)	Stems	<i>M. falcatum</i>	17
3-Hydroxy-1-methyl-3-(2-oxopropyl)quinoline-2,4(1H,3H)-dione (63)	Stems	<i>M. falcatum</i>	17
N-Methylflindersine (64)	Stems	<i>M. falcatum</i>	17
4-Hydroxy-3-methoxy-1-methyl-2(1 H)-quinolinone (65)	Stems	<i>M. falcatum</i>	17
N-Methylswietenidine-B (66)	Stems	<i>M. falcatum</i>	17
Myosmine (67)	Leaves and stem	<i>M. zeylanicum</i>	21
Halfordinol (68)	Leaves and stem	<i>M. zeylanicum</i>	21
O-Methylhalfordinol (69)	Leaves and stem	<i>M. zeylanicum</i>	21
Methyl carbazole-3-carboxylate (70)	Leaves	<i>M. integerrimum</i>	38
3-Methylcarbazole (71)	Stem barks	<i>M. hirsutum</i>	18
3-Formylcarbazole (72)	Stem barks	<i>M. hirsutum</i>	18
Lansine (73)	Stem barks	<i>M. hirsutum</i>	18
3-Formyl-6-methoxycarbazole (74)	Stem barks	<i>M. hirsutum</i>	18
Methyl carbazole-3-carboxylate (75)	Leaves	<i>M. integerrimum</i>	38
1,3-Dihydroxy-4-methoxy-10-methylacridone (76)	Stem bark	<i>M. hirsutum</i>	18
2,7-Dihydroxy-3-formyl-1-(30-methyl-20-butenyl)carbazole (77)	Roots	<i>M. glanduliferum</i>	20
7-Methoxy heptaphylline (78)	Roots	<i>M. glanduliferum</i>	20
Micromeline (79)	Stem bark	<i>M. hirsutum</i>	18
Koenimbine (80)	Leaves and stem	<i>M. zeylanicum</i>	21
Koenine (81)	Leaves and stem	<i>M. zeylanicum</i>	21
Koenigine (82)	Leaves and stem	<i>M. zeylanicum</i>	21
Koenidine (83)	Leaves and stem	<i>M. zeylanicum</i>	21
Mahanine (84)	Roots	<i>M. minutum</i>	35
Miscellaneous compounds			
Micrometam A (85)	Whole plants	<i>M. falcatum</i>	15
Micrometam D (86)	Whole plants	<i>M. falcatum</i>	15
Micrometam E (87)	Whole plants	<i>M. falcatum</i>	15
Micrometam B (88)	Whole plants	<i>M. falcatum</i>	15
Micrometam C (89)	Whole plants	<i>M. falcatum</i>	15
Polymethoxylated flavones			
5-Hydroxy-3,4',7,8-tetramethoxyflavone (90)	Aerial parts	<i>M. minutum</i>	40
5,7-Dihydroxy-3,4',8-trimethoxyflavone (91)	Aerial parts	<i>M. minutum</i>	40
5-Hydroxy-3,4',7,8-tetramethoxyflavone (92)	Aerial parts	<i>M. minutum</i>	40
5-Hydroxy-3,4',6,7,8-pentamethoxyflavone (93)	Leaves	<i>M. minutum</i>	40
3,5,7,4'-Tetramethoxyflavone (94)	Leaves	<i>M. compressum</i>	1
3,5,7,8,4'-Pentamethoxyflavone (95)	Leaves	<i>M. compressum</i>	1

(Continued)

Table 1. Continued

No. of compounds	Plant parts	Name of species	References
3,5,6,7,4'-Pentamethoxyflavone (96)	Leaves	<i>M. compressum</i>	1
5,7-Dihydroxy-4',6,8-trimethoxyflavone (97)	Aerial parts	<i>M. minutum</i>	26
7-Hydroxy-4',5,6,8-tetramethoxyflavone (98)	Aerial parts	<i>M. minutum</i>	26
7-Hydroxy-4',6,8-trimethoxyflavone -5-ylacetate (99)	Aerial parts	<i>M. minutum</i>	26
5-Hydroxy-3,3',4',7,8-pentamethoxyflavone (100)	Leaves and stems	<i>M. zeylanicum</i>	21
Coumarin and flavone condensed compounds			
8-((1 <i>S</i> ,2 <i>R</i>)-1-Hydroxy-2-(5-hydroxy-6,8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4H-chromen-7-yloxy)-3-methylbut-3-enyl)-7-methoxy-2H-chromen-2-one (101)	Aerial parts	<i>M. minutum</i>	26
8-((1 <i>S</i> ,2 <i>R</i>)-1-Hydroxy-3-methyl-2-(5,6,8-trimethoxy-2-(4-methoxyphenyl)-4-oxo-4H-chromen-7-yloxy)but-3-enyl)-7-methoxy-2H-chromen-2-one (102)	Aerial parts	<i>M. minutum</i>	26
7-((1 <i>S</i> ,2 <i>R</i>)-1-Hydroxy-1-(7-methoxy-2-oxo-2H-chromen-8-yl)-3-methylbut-3-en-2-yloxy)-6,8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4H-chromen-5-yl acetate (103)	Aerial parts	<i>M. minutum</i>	26
Stigmasterol and triterpenes			
Stigmasterol (104)	Leaves	<i>M. minutum</i>	10
5(6)-Gluten-3 α -ol (105)	Leaves and barks	<i>M. minutum</i>	23
5(6)-Gluten-3-one (106)	Leaves and barks	<i>M. minutum</i>	23
Derivative of oleic acid and phenylpropanoic acid			
Micromolide (107)	Stem barks	<i>M. birsutum</i>	18
Microintegerrin A (108)	stems and leaves	<i>M. integerrimum</i>	24
Microintegerrin B (109)	stems and leaves	<i>M. integerrimum</i>	24
3-(5-((2 <i>R</i> ,3 <i>R</i> ,4 <i>R</i>)-3,4-Dihydroxy-4-methyl-5-oxotetrahydrofuran-2-yl)-2-hydroxy-4-methoxyphenyl)propanoic acid (110)	fruits	<i>M. falcatum</i>	4
Secomicromelin (111)	fruits	<i>M. falcatum</i>	4
3,4-Dihydro-1,2-secomicromininin (112)	leaves	<i>M. falcatum</i>	14
3,4-Dihydro-1,2-secomicromininin methylester (113)	leaves	<i>M. falcatum</i>	14
3,4-Dihydro-1,2-secomicromininin-9- <i>O</i> -glucoside (114)	leaves	<i>M. falcatum</i>	14
Micromelumosides A (115)	aerial parts	<i>M. minutum</i>	6
Micromelumosides B (116)	aerial parts	<i>M. minutum</i>	6
Micromelumosides C (117)	aerial parts	<i>M. minutum</i>	6
Sesamin (118)	bark	<i>M. minutum</i>	34


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID ID

Nanik Siti Aminah  <https://orcid.org/0000-0002-2767-6006>

References

- Vidar WS, Macabeo APG, Knorn M, Kohls P, Aguinaldo AM. Polymethoxylated flavones from *Micromelum compressum*. *Biochem Syst Ecol.* 2013;50:48-50. doi:10.1016/j.bse.2013.03.010
- Singh RK. Lectotypification of the name *Micromelum integerrimum* (Rutaceae). *Phytotaxa.* 2016;255(2):172-174. doi:10.11646/phytotaxa.255.2.7
- Phakhodee W, Pattarawarapan M, Pongparn P, Laphookhieo S. Naturally occurring prenylated coumarins from *Micromelum integerrimum* twigs. *Phytochem Lett.* 2014;7:165-168. doi:10.1016/j.phytol.2013.11.009
- Suthiwong J, Sriphana U, Thongsri Y, Promsuwan P, Prariyachaitigul C, Yenjai C. Coumarinoids from the fruits of *Micromelum falcatum*. *Fitoterapia.* 2014;94:134-141. doi:10.1016/j.fitote.2014.02.004
- Susidarti RA. Chemical Constituents and Biological Activities of *Micromelum Minutum* (Rutaceae) and Two *Eugenia* Species (Myrtaceae). [PhD thesis]. Universiti Putra Malaysia; 2003.
- Sahakitpichan P, Tanpatanan W, Chimnoi N, Ruchirawat S, Kanchanapoom T. Glucosides of phenylpropanoic acid derivatives and coumarins from *Micromelum minutum*. *Phytochem Lett.* 2015;14:12-16. doi:10.1016/j.phytol.2015.08.009

7. Zhang D, Hartley TG, Mabberley DJ. *Flora of China, FOC.* ; 1998;11; 51.
8. Lekphrom R, Kanokmedhakul K, Sangsopha W, Kanokmedhakul S. A new coumarin from the roots of *Micromelum minutum*. *Nat Prod Res.* 2016;30(21):2383-2388. doi:10.1080/14786419.2016.1185720
9. Tantishaiyakul V, Pummangura S, Chaichantipwath C, McLaughlin JL, Division LS. phebalsin from the stem bark of *Micromelum minutum*. *J Nat Prod*;49(1):49-50.
10. Susidarti R, Rahmani M, Ali A. 8-Methoxycapnolactone and stigmaterol From *Micromelum minutum*. *Maj Farm.* 2007;18(2):105-109.
11. Asmah Susidarti R, Rahmani M, Ismail HBM, et al. Cytotoxic activity of coumarins from *Micromelum minutum*. *Pharm Biol.* 2009;47(2):182-185. doi:10.1080/13880200802436513
12. Hasan CM, Ahsan M. Chemical constituents, cytotoxic activities and traditional uses of *Micromelum minutum* (Rutaceae): a review. *Pharm Pharmacol Int J.* 2019;7(5):229-236.
13. Ramesh B, Pugalendi KV. Antioxidant role of umbelliferone in STZ-diabetic rats. *Life Sci.* 2006;79(3):306-310. doi:10.1016/j.lfs.2006.01.005
14. Kamperdick C, Phuong NM, Sung TV, Schmidt J, Adam G. Coumarins and dihydrocinnamic acid derivatives from *Micromelum falcatum*. *Phytochemistry.* 1999;52(8):1671-1676. doi:10.1016/S0031-9422(99)00243-5
15. Luo X, Huang Y, Zhang S, Yin H, Li C, Li Q. Five new phenethyl cinnamides from the mangrove associates *Micromelum falcatum*. *Biochem Syst Ecol.* 2014;56:191-195. doi:10.1016/j.bse.2014.05.018
16. Kouloura E, Danika E, Kim S, et al. Rapid identification of coumarins from *Micromelum falcatum* by UPLC-HRMS/MS and targeted isolation of three new derivatives. *Molecules.* 2014;19(9):15042-15057. doi:10.3390/molecules190915042
17. Luo XM, Qi SH, Yin H, Gao CH, Zhang S. Alkaloids from the stem bark of *Micromelum falcatum*. *Chem Pharm Bull.* 2009;57(6):600-602. doi:10.1248/cpb.57.600
18. Ma C, Case RJ, Wang Y, et al. Anti-Tuberculosis constituents from the stem bark of *Micromelum hirsutum*. *Planta Med.* 2005;71(3):261-267. doi:10.1055/s-2005-837826
19. Rodphukdeekul S, Sueblinvong T, Reungrungsi N, Chansrinoyom C. Anti-tumor activity of *Micromelum hirsutum* extract. *Thai J Pharmacol.* 2010;32(1):106-110.
20. Siridechakorn I, Ritthiwigrom T, Laphookhieo S. Coumarins and carbazole alkaloids from the roots of *Micromelum glanduliferum*. *Biochem Syst Ecol.* 2012;40:69-70. doi:10.1016/j.bse.2011.10.002
21. Bowen IH, Perera KPWC. Alkaloids, coumarins and flavonoids of *Micromelum zeylanicum*. *Phytochemistry.* 1982;21(2):433-437. doi:10.1016/S0031-9422(00)95283-X
22. Koriem KMM, Aminuddin ME, Kader AS. Antihyperglycemic SNR. Antihyperlipidemic and Antiapoptotic Activities of *Micromelum minutum* Seeds in Diabetic Rats. *J Mol Genet Med.* 2013
23. Lekphrom R, Kanokmedhakul S, Kukongviriyapan V, Kanokmedhakul K. C-7 oxygenated coumarins from the fruits of *Micromelum minutum*. *Arch Pharm Res.* 2011;34(4):527-531. doi:10.1007/s12272-011-0402-y
24. Wang Z-Y, He W-J, Zhou W-B, et al. Two new phenylpropanoids from *Micromelum integerrimum*. *Chin J Nat Med.* 2014;12(8):619-622. doi:10.1016/S1875-5364(14)60094-7
25. Ito C, Otsuka T, Ruangrunsi N, Furukawa H. Chemical constituents of *Micromelum minutum*. Isolation and structural elucidation of new coumarins. *Chem Pharm Bull.* 2000;48(3):334-338. doi:10.1248/cpb.48.334
26. Das S, Baruah RH, Sharma RP, Barua JN, Kulanthaivel P, Herz W. 7-methoxycoumarins from *Micromelum minutum*. *Phytochemistry.* 1984;23(10):2317-2321. doi:10.1016/S0031-9422(00)80543-9
27. Yun-Cheung Ket al. *Micromelum*: a key genus in the chemosystematics of the Clausenea. *Biochem Syst Ecol.* 1988;16(5):485-489.
28. Rahman MS, Rahman MZ, Hasan CM, Rashid MA, Kaiser MA. A review on phytochemicals from some medicinal plants of Bangladesh. *J. Pbar. Nutri. Sci.* 2011;1(1):87-95. doi:10.6000/1927-5951.2011.01.01.14
29. Luo X, He W, Yin H, et al. Two new coumarins from *Micromelum falcatum* with cytotoxicity and brine shrimp larvae toxicity. *Molecules.* 2012;17(6):6944-6952. doi:10.3390/molecules17066944
30. Rahmani M, Susidarti RA, Ismail HBM, et al. Coumarins from Malaysian *Micromelum minutum*. *Phytochemistry.* 2003;64(4):873-877. doi:10.1016/j.phytochem.2003.08.007
31. Susidarti RA, Jenie RI, Ikawati M, Putri DDP, Meiyanto E. Cytotoxic activity and apoptosis induction of 8-hydroxyisocapnolactone-2',3'-diol and its combination with Doxorubicin on MCF-7 and T47D cells. *J Appl Pharm Sci.* 2014;4(6):089-097.
32. Sakunpak A, Matsunami K, Otsuka H, Panichayupakaranant P. Isolation of new monoterpene coumarins from *Micromelum minutum* leaves and their cytotoxic activity against *Leishmania major* and cancer cells. *Food Chem.* 2013;139(1-4):458-463. doi:10.1016/j.foodchem.2013.01.031
33. Phakhodee W, Laphookhieo S, Prior TJ, Rujiwatra A. 1R,3R,4R,6S)-4-(7-Methoxy-2-oxo-2H-chromen-6-yl)-1-methyl-3,6-dioxabicyclo-[3.1.0]hexan-2-yl acetate. *Acta Crystallogr Sect E Struct Rep Online.* 2012;68(Pt 12):3421-3422. doi:10.1107/S160536812047617
34. Kassim NK, Lim PC, Ismail A, Awang K. Isolation of antioxidative compounds from *Micromelum minutum* guided by preparative thin layer chromatography-2,2-diphenyl-1-picrylhydrazyl (PTLC-DPPH) bioautography method. *Food Chem.* 2019;272:185-191. doi:10.1016/j.foodchem.2018.08.045
35. Roy MK, Thalang VN, Trakoontivakorn G, Nakahara K. Mechanism of mahanine-induced apoptosis in human leukemia cells (HL-60). *Biochem Pharmacol.* 2004;67(1):41-51. doi:10.1016/j.bcp.2003.07.021
36. Schmidt AW, Reddy KR, Knölker H-J, Occurrence KH-J. Occurrence, biogenesis, and synthesis of biologically active carbazole alkaloids. *Chem Rev.* 2012;112(6):3193-3328. doi:10.1021/cr200447s
37. Chakraborty DP, Roy S. Carbazole alkaloids. III. *Fortschr Chem Org Naturst.* 1991;57:71-152. doi:10.1007/978-3-7091-9119-4_2
38. Yang X-L, Xie Z-H, Jiang X-J, Huang Y-B, Liu J-K, Yan-Bing Huang J-KL. A new acridone alkaloid from *Micromelum integerrimum*. *Chem Pharm Bull.* 2009;57(7):734-735. doi:10.1248/cpb.57.734

39. Tang H, Ge H, Chen Z-B, et al. Micrometam C protects against oxidative stress in inflammation models in zebrafish and RAW264.7 macrophages. *Mar Drugs*. 2015;13(9):5593-5605. doi: 10.3390/md13095593
40. Sohrab MH, Chowdhury R, Hasan CM, Rashid MA. Chemotaxonomic significance of polyoxygenated flavonoids from the leaves of *Micromelum minutum*. *Biochem Syst Ecol*. 2004;32(9):829-831. doi: 10.1016/j.bse.2003.11.009



x ⓘ

Free English Writing Tool

Grammarly makes sure everything you type is effective and mistake-free. Try now

DOWNLOAD Grammarly

Natural product communications

35

H Index

Country	United States - IIII SIR Ranking of United States
Subject Area and Category	Agricultural and Biological Sciences Plant Science Medicine Complementary and Alternative Medicine Medicine (miscellaneous) Pharmacology, Toxicology and Pharmaceutics Drug Discovery Pharmacology
Publisher	Natural Product Communications
Publication type	Journals
ISSN	15559475, 1934578X
Coverage	2008-ongoing
Scope	Natural Product Communications (NPC) is an open access, peer reviewed journal bringing studies on all aspects of natural products, including isolation, characterization, spectroscopic properties, biological activities, synthesis, structure-activity, biotransformation, biosynthesis, tissue culture and fermentation into one journal. It publishes communications, full papers, accounts and reviews covering the full breadth of chemistry, biochemistry, biotechnology, pharmacology, and chemical ecology of natural products. Readership includes: Academic and Industrial Scientists working in all aspects of Organic, Pharmaceutical, Medicinal, Bioorganic and Analytical Chemistry, Biotechnology, Biodiversity and Chemotaxonomy.
?	Homepage How to publish in this journal Contact Join the conversation about this journal









x ⓘ

Free English Writing Tool

Grammarly makes sure everything you type is effective and mistake-free. Try now

DOWNLOAD Grammarly

Quartiles	+
-----------	---



Natural product communications

Q2 Complementary and Alternative Medicine
best quartile

SJR 2018
0.28

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

Academic paper editing

Get Professional academic paper editing by Subject Area Native English Editors Today!

authorassists.com

OPEN

U **Unang Supratman** 5 months ago

I try to submit but can not entrance due to incorrect username and password.
Can you give me a new username and password?

I waiting for your further information.

reply



Melanie Ortiz 5 months ago

ScImago Team

Dear Unang,
thank you for contacting us.
Sorry to tell you that ScImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.
Unfortunately, we cannot help you with your request, we suggest you to contact the journal's editorial staff , so they could inform you more deeply. Best Regards, ScImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot

reCAPTCHA
[Privacy](#) · [Terms](#)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.