


J. Indian Chem. Soc. Vol. 96, No. 6
June 2019

JOURNAL OF THE INDIAN CHEMICAL SOCIETY



Published by Indian Chemical Society
92, Acharya Prafulla Chandra Road
Kolkata-700 009, India


In this issue (13 articles)

•  *Original Paper*


Theoretical study of structural, electronic and vibrational properties of a Ni-Salen Complex
Alireza Akbari

•  *Original Paper*

Prediction of charge transfer transition energies of the molecular complexes of PMDA with a series of methylbenzenes by TDDFT
Amit S. Tiwary

•  *Original Paper*


Chromium doping in Titania for making Dye Sensitized Solar cell with improved efficiency and stability
Narottam Prasad 1, K.R. Patel 2 and M. S. Roy 1

•  *Original Paper*


A Time-dependent Multi-reference Unified Cluster Cumulant Formulation to Study the Subdynamics of Quantum System Coupled both to Thermal and Stochastic Bath
Seikh Hannan Mandal

•  *Original Paper*


QSAR modeling and in silico designing of Tumor-Associated Carbonic Anhydrases XII inhibitors
Shalini Singh

•  *Original Paper*

Approach to zero band gap graphene sheets: A graph theoretical analysis
Bankim Chandra Ghosh

•  *Original Paper*

Synthesis of 1-aminomethyl-3-[4'-(4"-fluorobenzyloxy)-benzohydrazono]-5-ethylisatins as antifungal agents
Nisheeth Rastogi

•  *Original Paper*

Isolation, Characterization and improvement on a Wild strain of *Corynebacterium glutamicum* for L-glutamic acid production
*Subhadeep Ganguly

•  *Original Paper*


Simultaneous Estimation of Metoprolol Tartrate and Amlodipine Besylate by Absorbance Ratio Method
Kamal Shah *1 and Pradeep Mishra 2

•  *Original Paper*

Investigation on the rate of horizontal spread of Anthracene in a Sandy Clay Soil
Mizanur Rahman, Jyoti Prakash Sarania, and Prahash Chandra Sarma*

•  *Original Paper*

Mechanistic studies of oxidation of substituted oxoacids by IFC in the presence of 1,10-phenanthroline
M.Vellaisamy* and M.Sharmila

•  *Original Paper*

Influence of a drying phase on K-transformation in a continuously flooded limed and the corresponding unlimed soil
Tapa Debnath, Sayan Majumder* and Dipankar Saha

-
-  *Original Paper*

COUMARINS FROM MYANMAR EDIBLE FRUIT TREE (*CASIMIROA EDULIS*)

Khun Nay Win Tun a,b, Nanik Siti Aminah a*, Alfinda Novi Kristanti a, Rico Ramadhan a, Yoshiaki Takaya c, HninThanda Aung d

Honorary Editors

Inorganic and Analytical Chemistry Section

Professor Samaresh Bhattacharya
Department of Chemistry
Inorganic Chemistry Section
Jadavpur University
Kolkata - 700 032, India
E-mail: samaresh_b@hotmail.com
<http://www.jaduniv.edu.in/>

Organic and Biochemistry Section

Professor Akhilesh Kumar Verma,
Department of Chemistry,
University of Delhi, Delhi-110 007, India
E - mail: averma@acbr.du.ac.in,
akhilesh682000@gmail.com
<http://www.du.ac.in/du/>

Physical and Industrial Chemistry Section

Professor Pradeep Kumar Sharma
Department of Chemistry
Jai Narayan Vyas University
Jodhpur, India
E-mail: drpkvs27@yahoo.com
<http://www.jnvu.co.in/>

International Executive Editor

Bimal Krishna Banik, PhD. C. Chem. F. R. S. C.
Vice President of Research and Education Development
Community Health Systems of South Texas
Edinburg, Texas 78539, USA
E-mail: bimal.banik@chsst.org; bimalbanik10@gmail.com
Phone Numbers: 956-627-4094 and 281-813-2104

International Editors

Professor Takashiro Akitsu
Department of Chemistry, Faculty of Science, Tokyo University of
Science 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan Tel. +81-3-
5228-8271, Fax. +81-3-5261-4631
E-mail: akitsu@rs.kagu.tus.ac.jp
<http://www.rs.kagu.tus.ac.jp/akitsu/index.htm>

Professor Ebbe Nordlander
Department of Chemical Physica
Lund University
Box 117, 221 00 Lund, Sweden
E-mail: ebbe.nordlander@chemphys.lu.se
<https://www.lunduniversity.lu.se/>

Dr. Anil Kumar *FNA FASc FNASc*
J C Bose National Fellow
Physical & Materials Chemistry Division
C SIR-National Chemical Laboratory (NCL)
Pune 411008, INDIA
Email: a.kumar@ncl.res.in
<http://academic.ncl.res.in/a.kumar>

Editor (Guest)

Dr. Shivendu Ranjan, B.Tech, Ph.D, PDF, MICS, FBSS (India), FLS (London)
Scientist
DST-Centre for Policy Research
Lucknow, Uttar Pradesh, India
And
Senior Research Associate (Adjunct)
Faculty of Engineering & Built Environment
University of Johannesburg, Johannesburg
(Auckland Park Kingsway Campus)
South Africa

Email: shivenduranjan@gmail.com
<https://sites.google.com/view/shivenduranjan>



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Journal of the Indian Chemical Society

Country [India](#) - [IIII](#) [SJR Ranking of India](#)**Subject Area and Category**[Chemistry](#)[Electrochemistry](#)[Inorganic Chemistry](#)[Organic Chemistry](#)[Physical and Theoretical Chemistry](#)[Pharmacology, Toxicology and Pharmaceutics](#)[Drug Discovery](#)**Publisher**[Scientific Publishers](#)**Publication type**[Journals](#)**ISSN**

00194522

Coverage

1973-1985, 1987-1991, 1996-ongoing

Scope


The Journal of the Indian Chemical Society publishes original, fundamental, theoretical, experimental research work of highest quality in all areas of chemistry, biochemistry, medicinal chemistry, electrochemistry, agrochemistry, chemical engineering and technology, food chemistry, environmental chemistry, etc.

[Homepage](#)[How to publish in this journal](#)[Contact](#)[Join the conversation about this journal](#)

31

H Index

Quartiles









SJR



Citations per document





Journal of the Indian Chemical Society

Q4 Drug Discovery
best quartile

SJR 2018
0.11

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```


J. Indian Chem. Soc.,
Vol. 96, June 2019, pp. 1-4

Coumarins from Myanmar edible fruit tree (*Casimiroa edulis*)

Khun Nay Win Tun^{a,b}, Nanik Siti Aminah^{*a}, Alfinda Novi Kristanti^a, Rico Ramadhan^a, Yoshiaki Takaya^c and HninThanda Aung^d

^aDepartment of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia

E-mail: nanik-s-a@fst.unair.ac.id

^bDepartment of Chemistry, Taunggyi University, Shan State (South), Myanmar

^cFaculty of Pharmacy, Meijo University, Tempaku, Nagoya, Japan

^dDepartment of Chemistry, Mandalay University, Mandalay, Myanmar

Manuscript received online 03 April 2019, revised 22 May 2019, accepted 31 May 2019

This research expresses the phytochemical study from the Myanmar edible fruit tree, *Casimiroa edulis* (Rutaceae). The result revealed that the isolation and identification of two furanocoumarins (bergapten **1** and isopimpinellin **2**) from the stem bark of this plant. Their molecular structures were elucidated and identified by using NMR spectroscopy in combination with IR, UV and HRMS spectra data, respectively. Furthermore, these two compounds were investigated for their anti-diabetic activity. According to the result, bergapten **1** and isopimpinellin **2** are not essentially good for anti-diabetic activity. This is the first report of two furanocoumarins from the Myanmar edible fruit tree.

Keywords: *Casimiroa edulis*, furanocoumarins, NMR spectroscopy, anti-diabetic activity.

Introduction

Casimiroa edulis La Llave (Rutaceae) is a genus of comprising only 10 species which is distributed in tropical and subtropical regions, including Myanmar. Among then, *Casimiroa edulis* is the well-known species^{1,2}. The traditional used and the pharmacological studies of the leaves and seeds of *Casimiroa edulis* displayed various biological activities. In Myanmar, this plant is named Thar-kyar-thee.

Phytochemical studied from genus *Casimiroa* assist to the separation of various chemical constituents such as 18 flavonoids (isolated from leaves, fruits, and seeds), 24 alkaloids (isolated from leaves, fruits, bark, and seeds), and 16 coumarins (isolated from leaves, fruits, and seeds). Pharmacological investigations of isolated flavonoids are antioxidant, anti-mutagenic, solid tumor selective cytotoxicity, vasodilation and radical-scavenging activities, the investigations of isolated alkaloids are anti-mutagenic, solid tumor selective cytotoxicity, anti-hypertensive, cardiovascular activities, and the investigations of isolated coumarins are anti-coagulant, vasodilation and radical-scavenging, adipogenesis, solid tumor selective cytotoxicity activities, respectively³⁻¹⁴. This

paper deals with the structure elucidation of two furanocoumarins from the stem bark of *Casimiroa edulis*.

Experimental

General:

The infrared spectra (IR) were obtained on FTIR-8400S (Shimadzu) using KBr. UV spectra were examined in MeOH by using UV-Vis Shimadzu spectrometer. Column chromatography (CC) was achieved on silica gel (BW-820H). Analytical thin-layer-chromatography (TLC) was completed at room temperature on pre-coated Kieselgel silica gel 60 F₂₅₄ (Merck) aluminium plates. Melting points were determined by fisher john melting point apparatus. All NMR spectra were checked in CDCl₃ by using a Bruker Avance 3 (¹H: 600 MHz and ¹³C: 151 MHz) with TMS as an internal standard. Chemical shifts are described in part per million (δ , ppm) based on their solvent signal (chloroform-*d*: δ_{H} 7.26, δ_{C} 77.16) coupling constants *J* value in Hertz (Hz). HR-FAB-MS was attained with a JEOL JMS HX-110 mass spectrometer.

Plant material:

Casimiroa edulis La Llave was collected from Namp-see


Village, Taunggyi (Shan State), Myanmar, in August 2016, and was identified by the Professor of Botany Department, University of Taunggyi.

Extraction and isolation:

The dry sample of *Casimiroa edulis* (1 kg) was exhaustively extracted with MeOH (3 L) at room temperature for four weeks. The methanol extract was concentrated under pressure using rotator evaporator for dryness. Then, the methanolic extract was partition between *n*-hexane and MeOH (300 ml×3, v/v). The concentrated MeOH extract (50 g) was subjected to vacuum liquid chromatography (VLC) and eluted with *n*-hexane-EtOAc in a stepwise gradient solvent system (10:0, 6:4, 0:10, v/v) to afford several fractions. Column eluate was collected in 150 ml glass bottles to yield 28 fractions and examined under UV. Based on analytical TLC, the fractions were grouped and evaporated on a rotary evaporator to afford six fractions (J-1 to J-6). Fractions J-5 (2 g) were additionally purified by silica gel column chromatography with a gradient solvent system of *n*-hexane in EtOAc (10:0, 0:10) to give colourless needles of compound **1**, and pale-yellow crystalline solids of compound **2**.

Results and discussion

Compound **1** was obtained as colorless needle crystals. Its melting point was 186–206°C. Based on the molecular ion peak at m/z 217.0501 $[M + H]^+$ (Calcd. 216.0423) in the HR-FAB-MS, the molecular formula was determined as $C_{12}H_8O_4$. IR spectrum informed the characteristic of sharp bands at 3143, 3115, 3088, 3010, 2924, 2852, 1737, 1720, 1625 and 1581 cm^{-1} which could be assigned to $-OCH_3$, sp^2-CH , sp^3-CH , lactone $C=O$, $C=C$ ring skeletal stretching vibration, respectively. The UV spectrum showed the bands



Compds.	R ₁	R ₂
1	OCH ₃	H
2	OCH ₃	OCH ₃

Fig. 1. Chemical structure of compound **1** and **2**.

at λ_{max} 221, 249, 259, 268, and 311 nm, which were typical of a furanocoumarin skeleton. The 1H NMR spectrum was exhibited as an easily distinguishable pair of doublets at δ_H 6.28 (d, 1H, 9.8 Hz, H-3) and δ_H 8.16 (d, 1H, 9.8 Hz, H-4). It also displayed two furan doublets at δ_H 7.60 and 7.02 (each, d, 1H, 2.4 Hz, H-2' and H-3'), and an aromatic singlet at δ_H 7.14 (s, 1H, H-8). Combining analysis of ^{13}C NMR and DEPT spectra data, compound **1** displayed 12 carbons resonance, including five methine, one oxygenated and six quaternary carbons, respectively. In the 1H - 1H COSY experiment, the cross-peak correlations of H-3 with H-4, and the cross-peak correlations of H-2' through H-3' were observed. The long-range HMBC attachment from H-3 to C-2 and C-10, H-4 to C-2 and C-9, H-8 to C-6, C-7, C-9 and C-10, H-2'' to C-3'', C-6 and C-7, H-3'' to C-2', C-6 and C-7 were also observed. The remaining one singlet proton at δ 4.27 (integrating for three protons) showed the presence of methoxy group on the aromatic ring. This confirmation was deduced by HMBC spectrum, which confirms that methoxy proton showed the long-range correlation with the C-5 carbon at δ_C 149.6. Analysis of UV, IR, 1D, 2D NMR and HRMS spectra as well as on comparison with Literature, were allowed to determine the structure of compound **1** as bergapten^{15–17}.

Compound **2** was obtained as pale-yellow crystalline solids with a melting point at 178–180°C. Its molecular formula was assigned as $C_{13}H_{10}O_5$ based on HR-FAB-MS (m/z 247.0606 $[M + H]^+$, Calcd. 246.0528). The UV spectrum ex-




Fig. 2. 1H - 1H COSY correlations (a) and major 1H - ^{13}C long-range HMBC (b) correlation in compound **1**.




Fig. 3. 1H - 1H COSY correlations (a) and major 1H - ^{13}C long-range HMBC (b) correlation in compound **2**.

hibited the absorption maxima at 241, 248, 268 and 311 nm, which is consistent with the presence of a furocoumarin. The IR spectrum revealed the presence of an α,β -unsaturated lactone (ν_{\max} 1751 and 1718 cm^{-1}), methoxy group (3132 and 3159 cm^{-1}), C-H stretching vibration of sp^2 hydrocarbon (3084 and 3005 cm^{-1}), asymmetrical and symmetrical C-H stretching vibration of sp^3 hydrocarbon (2843 and 2953 cm^{-1}), and C=C ring skeleton stretching vibration (ν_{\max} 1604 and 1593 cm^{-1}) groups. ^{13}C NMR of compound **2** included 13 carbon signals, comprising 2 methoxy carbon signals, 4 methine carbon signals, and 7 quaternary carbon signals, which were classified by DEPT and HSQC experiment. Its ^1H NMR spectrum was exhibited a distinct pair of doublets at δ_{H} 6.27 and 8.11 (each, 1H, 9.8 Hz) assignable to C-3, C-4 protons, a furan doublet at δ_{H} 7.62 and 6.99 (each, 1H, 2.2 Hz) assignable to C-2', C-3' protons. The ^1H - ^1H COSY spectrum showed a correlation between H-3 and H-4 protons while H-2' proton had a correlation to the H-4' proton. In HMBC spectrum, the following correlations were observed; the proton signal at δ_{H} 6.27 was connected with C-2 and C-10, the proton signal at δ_{H} 8.11 was connected with C-2 and C-9, the proton signal at δ_{H} 7.62 was correlated with C-2', C-6 and C-7, the proton signal at δ_{H} 6.99 was correlated with C-3', C-6 and C-7. The remaining methoxy protons at δ 4.16 (integrating for 6 protons) were placed at C-5 and C-8 on the

Table 1. ^1H NMR and ^{13}C NMR data of compound **1** in CDCl_3 and bergapten ^1H in CDCl_3

Position	Compd. 1		Lit. [Ref. 16]	
	^1H NMR ^a	^{13}C NMR ^b	^1H NMR ^c	^{13}C NMR ^d
2	–	161.2	–	161.4
3	6.28 (d, 9.8 Hz)	112.6	6.27 (d, 9.8 Hz)	112.6
4	8.16 (d, 9.8 Hz)	139.2	8.15 (d, 9.8 Hz)	139.4
5	–	149.6	–	149.7
6	–	112.7	–	112.9
7	–	158.4	–	158.5
8	7.14 (s)	93.9	7.13 (s)	94.0
9	–	152.7	–	151.9
10	–	106.5	–	106.6
2'	7.60 (d, 2.4 Hz)	144.8	7.59 (d, 2.4 Hz)	144.9
3'	7.02 (d, 2.4 Hz)	105.0	7.02 (d, 2.4 Hz)	105.2
5-OMe	4.27 (s)	60.1	4.27 (s)	60.3
8-OMe	–	–	–	–

^aSpectra recorded in 600 MHz, ^bspectra recorded in 151 MHz, ^cspectra recorded in 400 MHz and ^dspectra recorded in 100 MHz.

Table 2. ^1H NMR and ^{13}C NMR data of compound **2** in CDCl_3 and isopimpinellin in CDCl_3

Position	Compd. 2		Lit. [Ref. 16]	
	^1H NMR ^a	^{13}C NMR ^b	^1H NMR ^c	^{13}C NMR ^d
2	–	160.4	–	160.4
3	6.27 (d, 9.8 Hz)	112.8	6.26 (d, 9.8 Hz)	112.8
4	8.11 (d, 9.8 Hz)	139.4	8.10 (d, 9.8 Hz)	139.3
5	–	144.3	–	144.3
6	–	114.8	–	114.8
7	–	150.0	–	150.0
8	–	128.2	–	128.2
9	–	143.7	–	143.7
10	–	107.6	–	107.6
2'	7.62 (d, 2.2 Hz)	145.1	7.61 (d, 2.3 Hz)	145.1
3'	6.99 (d, 2.2 Hz)	105.1	6.99 (d, 2.3 Hz)	105.1
5-OMe	4.16 (s)	61.7	4.16 (s)	61.7
8-OMe	4.16 (s)	60.8	4.14 (s)	60.8

^aSpectra recorded in 600 MHz, ^bspectra recorded in 151 MHz, ^cspectra recorded in 400 MHz and ^dspectra recorded in 100 MHz.

basic of long range HMBC correlations. With the aid of above data and previous report, compound **2** was identified as isopimpinellin^{16–18}.

Conclusions

In this study, two known compounds were obtained from the stem bark of *Casimiroa edulis*. The structure of isolated compounds identified as bergapten **1** and isopimpinellin **2** based on their spectroscopic data, such as UV-Vis, FTIR, 1D NMR, 2D NMR, and HRMS. The isolated compounds are not important for their anti-diabetic activity. A lot of researchers were reported these two compounds from many sources of natural product. This is the first report on the occurrence of bergapten and isopimpinellin from Myanmar edible fruit tree *Casimiroa edulis*.

Acknowledgements

This work was supported in part by Ailangga Development Scholarship (ADS) and "RISET MANDAT GRANT" of Universitas Airlangga, Surabaya, Indonesia. KNWT gracefully acknowledges Rectors, Vice-Rectors, and Professors from Universitas Airlangga.

Supporting Information

1D and 2D NMR spectrum are described.

References

1. M. Yamamoto, T. Tomita, M. Onjo, K. Ishihata, T. Kubo and S. Tominaga, *HortScience*, 2007, **42(6)**, 1329.
2. N. Satheesh, *American-Eurasian J. Agric. & Environ. Sci.*, 2015, **15(8)**, 1574.
3. A. Ito, L. A. Shamon, B. Yu, E. Mata-Greenwood, S. K. Lee, R. B. Breemen, R. G. Mehta, N. R. Farnsworth, H. H.S. Fong, J. M. Pezzuto and A. Douglas, *J. Agric. Food Chem.*, 1998, **46**, 3509.
4. A. S. Awaad, N. H. El-Sayed, D. J. Maitland and T. J. Mabry, *Pharm. Biol.*, 2006, **44(4)**, 258.
5. A. S. Awaad, N. A. Al-Jaber, G. A. Soliman, M. R. Al-Outhman, M. E. Zain, J. E. Moses and R. M. El-Meligy, *Phytother. Res.*, 2012, **26**, 452.
6. A. N. García-Argáeza, N. M. González-Lugo, H. Parra-Delgado, and M. Martínez-Vázquez, *Biochem. Syst. Ecol.*, 2005, **33**, 441.
7. A. S. Awaad, D. J. Maitland and S. M. Moneir, *Chem. Nat Compd.*, 2007, **43(5)**, 576.
8. A. E. M. Khaleel, *Monatshefte Fur Chemie*, 2002, **133**, 183.
9. B. Heneka, H. Rimpler, A. Ankli, O. Sticher, S. Gibbons and M. Heinrich, *Phytochemistry*, 2005, **66**, 649.
10. D. L. Dreyer, *J. Org. Chem.*, 1968, **33(9)**, 3577.
11. H. Nagai, T. Tanaka, T. Goto, T. Kusudo, N. Takahashi, and T. Kawada, *Biosci. Biotechnol. Biochem.*, 2014, **78(2)**, 296.
12. A. N. García-Argáeza, N. M. González-Lugob, H. Parra-Delgadob and M. Marínez-Vázquezb, *Z. Naturforsch.*, 2004, **59(b)**, 245.
13. R. Bertin, Z. Chen, M. Martínez-vázquez, A. García-argaéz and G. Froldi, *Phytomedicine*, 2014, **21**, 586.
14. X. U. Ya-ming, M. C. Ramirez-ahumada, F. A. Valeriote and A. A. L. Gunatilaka, *Chin. J. Nat. Med.*, 2011, **9(5)**, 334.
15. C. Chunyan, S. Bo, L. Ping and L. Jingmei, *J. Liq. Chrom. Relat. Tech.*, 2009, **32**, 136.
16. T. O'Neill, J. A. Johnson, D. Webster and C. A. Gray, *J. Ethnopharmacol.*, 2013, **147**, 232.
17. J. N. Ngwendson, E. Bedir, S. M. N. Efangé, C. O. Okunji, M. M. Iwu, B. G. Schuster and I. A. Khan, *Pharmazie*, 2003, **58**, 587.
18. G. Wang, Z. Zhou, C. Cheng, J. Yao and Z. Yang, *Vet. Parasitol.*, 2008, **158**, 144.