

Ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate

*by* Alfinda Novi Kristanti

---

**Submission date:** 09-Mar-2020 05:28PM (UTC+0800)

**Submission ID:** 1272167199

**File name:** 2017-molbank-M954-1.pdf (1.46M)

**Word count:** 2134

**Character count:** 18862

Short Note

# Ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate

Hery Suwito<sup>1,\*</sup>, Salma Zulqaida<sup>1</sup>, Kautsar Ul Haq<sup>1</sup>, Alfinda Novi Kristanti<sup>1</sup> and Indriani Indriani<sup>2</sup>

<sup>1</sup> Department of Chemistry, Faculty of Science and Technology, Airlangga University, Surabaya 60115, Indonesia; salmazulqaida@gmail.com (S.Z.); kautsar.ul.haq-2016@fst.unair.ac.id (K.U.H.); krisnosuwono@yahoo.com (A.N.K.)

<sup>2</sup> Department of Chemistry, Tadulako University, Palu 94118, Indonesia; indri.2707@gmail.com

\* Correspondence: herys08032002@yahoo.com; Tel.: +62-31-5922427

Received: 23 July 2017; Accepted: 9 August 2017; Published: 11 August 2017

**Abstract:** A one-pot multicomponent reaction has been used to synthesize the title compound, ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate by PTSA catalyzed Biginelli reaction. The chemical structure of the product was confirmed by spectroscopic evidence, FTIR, HRESI-MS, 1D-, and 2D NMR.

**Keywords:** one-pot multicomponent reaction; Biginelli reaction; furanyl DHPM


## 1. Introduction

Nitrogen containing heterocycles are widespread in many naturally occurring compounds. They build the core structures of many biologically active scaffolds as well as some industrial compounds. Among these, dihydropyrimidinone (DHPM) derivatives attract many researchers due to their simple synthesis protocol. The synthesis of DHPM was firstly reported by Italian chemist Pietro Biginelli in 1893 through simple one-pot multicomponent reaction of benzaldehyde, urea, and ethyl acetoacetate employing HCl as catalyst [1]. Furthermore, the attractiveness of DHPM derivatives are associated with their unique pharmacological activities such as antioxidant [2], anti-inflammation, antibacterial, antifungal [3], anti HIV [4], anticancer [5], and antihypertensive activities [6]. In continuation of our research for finding an anticancer agent acting as inhibitor of the mitotic kinesin Eg5, we have successfully synthesized a series of DHPM derivatives employing Biginelli reaction and herein we report a new DHPM derivative namely ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate (4).

## 2. Results and Discussion

The synthesis of the title compound (4) was conducted in one step reaction as presented in Scheme 1.

The synthesis of the title compound was performed by combining usual Biginelli procedure with methylation at hydroxyl methyl group of the furan ring as reported by Yoshida et al. [7]. The reaction was carried out in methanol in the presence of 0.025 mol % of PTSA as catalyst at room temperature. The usage of PTSA as a catalyst for Biginelli reaction was previously reported by Jin et al. [8] and Matache et al. [9]. Besides solvent, methanol was also acting as a methylation agent. The target molecule 4 was obtained as white solid in 40% yield with m.p of 178–180 °C. The structure of the compound 4 was established by FTIR, HRESI-MS, <sup>1</sup>H-, APT-, and 2D NMR spectral data.




**Scheme 1.** Synthesis of ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate (4).


The HRESIMS displayed a negative molecular ion at  $m/z$  293.1145 indicating a molecular formula of  $C_{14}H_{18}N_2O_5$  and 8 degrees of unsaturation (see Supplementary Material Figure S1). The FTIR spectrum exhibited absorption bands at 3238, 1710, 1656  $\text{cm}^{-1}$  indicating the presence of an  $\text{-NH}$  amide group, carbonyl group of an ester, and an amide respectively [10] (see Supplementary Material, Figure S2).

To facilitate the discussion, we provide the numbering of the title compound as displayed in Figure 1, while the NMR chemical shifts is displayed in Figure 2. The existence of tetrahydropyrimidine ring is proven by the following data. Two broad single signals at 5.94 and 8.53 ppm of  $^1\text{H-NMR}$  spectrum indicated two amide protons (see Supplementary Material, Figure S3), while the carbonyl group is assigned by the  $^{13}\text{C}$  signal at 154.0 ppm (see Supplementary Material, Figure S4). The  $\text{sp}^2$  two carbon atoms were confirmed by two  $^{13}\text{C}$  signals at 98.2 and 148.1 ppm, whereas a signal at 48.9 ppm indicated a  $\text{sp}^3$  carbon atom which attaches a proton that appeared as a doublet signal at 5.46 ppm. A methyl group attached in a  $\text{sp}^2$  carbon atom was observed as a singlet signal at 2.34 ppm. This information indicated that the title compound possessing 6-methyl-4-aryl-3,4-dihydropyrimidin-2-on scaffold, typical skeleton of Biginelli product. The existence of the ethyl ester group was confirmed by a  $^{13}\text{C}$  signal at 165.5 ppm indicated a carbonyl ester, whereas the ethyl group was observed by  $^1\text{H}$  signal as a quartet at 4.12 ppm indicated an oxy-methylene ( $^{13}\text{C}$  signal was observed at 60.1 ppm), and the methyl group appeared as a triplet at 1.19 ppm ( $^{13}\text{C}$  signal was observed at 14.3 ppm) [11,12].

Furthermore, in the aromatic area, two signals— $\delta_{\text{H}}$  (ppm) 6.05 (d,  $J = 3.1$  Hz, 1H) and 6.20 (d,  $J = 3.1$  Hz, 1H)—were observed, indicating the presence of two protons in an ortho position of H-9 and H-10 in a furan ring. The oxy-methylene fragment appeared as a singlet signal at  $\delta_{\text{H}}$  4.33 ppm or at 66.5 ppm in the carbon signal. A methoxy fragment attached in an aliphatic carbon atom was demonstrated by a singlet signal at  $\delta_{\text{H}}$  3.34 ppm or at 58.0 ppm in the carbon signal. The APT spectra showed 14 signals representing all carbon atoms of the title compound.



**Figure 1.** Numbering of the title compound.



**Figure 2.**  $^1\text{H}$ - (A) and  $^{13}\text{C}$ - (B) NMR chemical shifts ( $\text{CDCl}_3$ ) of the title compound.

The existence of two NH protons was also confirmed by HMQC experiment which showed the existence of two proton signals exhibiting no correlation with carbon atoms (see Supplementary Material, Figure S5). Based on HMBC experiment (see Supplementary Material, Figure S6), the presence of 6-methyl-4-aryl-3,4-dihydropyrimidin-2-one scaffold possessing ester substituent at C-5 was convinced by a correlation of proton H-4 ( $\delta_{\text{H}}$  5.46 ppm) with olefinic carbon and  $\alpha,\beta$ -unsaturated carbonyl ester  $\delta_{\text{C}}$  (ppm) 98.2 (C-5), 148.1 (C-6), 165.5 (C-14) respectively, carbonyl urea-type ( $\delta_{\text{C}}$  154.0 ppm, C-2), and two carbon from the furan ring— $\delta_{\text{C}}$  (ppm) 155.1 (C-8) and 106.7 (C-9). The presence of this skeleton was also convinced by a long range correlation of NH proton  $\delta_{\text{H}}$  8.53 ppm (H-1) with two  $\alpha,\beta$ -unsaturated olefinic carbon atoms (C-5 and C-6) and methyl olefinic ( $\delta_{\text{C}}$  18.6 ppm, C-7). Furthermore, the presence of furan ring was convinced by two aromatic protons— $\delta_{\text{H}}$  (ppm) 6.05 (H-9) and 6.20 (H-10)—which correlated with its three neighboring carbon atoms— $\delta_{\text{C}}$  (ppm) 155.1 (C-8), 151.6 (C-11), and 110.2 (C-10) or 106.7 (C-9). In addition, the presence of methoxymethylene substituent at C-11 was convinced by a long range correlation between methylene proton  $\delta_{\text{H}}$  4.33 with two carbon of furan ring (C-10 and C-11) and methoxy carbon ( $\delta_{\text{C}}$  58 ppm, C-13). The selected long range correlation is suitable with the structure of the title compound is displayed in Figure 3. The structure refinement parameters is displayed in Table 1. Based on the structure elucidation of the obtained spectrum, the title compound is a new compound.

**Table 1.** NMR data of Ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate in  $\text{CDCl}_3$

No. Atom	$\delta_{\text{H}}$ (mult, J Hz)	$\delta_{\text{C}}$ (ppm)	HMBC
1	8.53 (s, 1H)	—	C-5, C-6, C-7, C-14
2	—	154.0	—
3	5.94 (s, 1H)	—	C-4, C-5
4	5.46 (d, J = 3.0 Hz, 1H)	48.9	C-2, C-5, C-6, C-8, C-9, C-14
5	—	98.2	—
6	—	148.1	—
7	2.35 (s, 3H)	18.6	C-6, C-4, C-5, C-14
8	—	155.1	—
9	6.05 (d, J = 3.2 Hz, 1H)	106.7	C-8, C-10, C-11
10	6.20 (d, J = 3.2 Hz, 1H)	110.2	C-9, C-8, C-11
11	—	151.6	—
12	4.33 (s, 2H)	66.5	C-10, C-11, C-13
13	3.34 (s, 3H)	58.0	C-12
14	—	165.5	—
15	4.12 (q, 2H)	60.1	C-16
16	1.19 (t, J = 7.1 Hz, 3H)	14.3	C-15




Figure 3. Selected HMBC correlation for compound 4.

### 3. Materials and Methods

The chemicals used in the research were provided from the commercial sources and pro analysis or pro synthesis grade. The melting point was measured with a Fisher–Johns melting point apparatus 220 VAC (Fisher Scientific, Waltham, MA, USA) and uncorrected. The reaction progress and purity of the product were monitored by thin layer chromatography on silica gel GF<sub>254</sub> plates (E Merck, Darmstadt, Germany) and the spots were identified by UV lamp ( $\lambda$  254 nm). The mass spectra was recorded by HRESI-MS (waters LCT premier XE, Waters Corp, Milford, MA, USA). FTIR spectra was recorded in KBr powder with the Diffuse Reflectance Method on spectrophotometer IRTracer100 (Shimadzu, Kyoto, Japan). NMR spectra (<sup>1</sup>H, APT, HMBC, and HMQC) were recorded on JEOL 400 ECA spectrometer (JEOL, Tokyo, Japan) using CDCl<sub>3</sub> as a solvent and internal standard.

#### Synthesis Procedure of the Title Compound

The mixture of 2 mmol 5-hydroxymethyl furfural, 3 mmol urea, 3 mmol ethyl acetoacetate, 0.5 mmol PTSA, and 5 mL methanol was stirred for 24 h at room temperature, the reaction progress was monitored by TLC on a silica gel plate employing chloroform:ethyl acetate (1:2) as a mobile phase. After the reaction was completed, the reaction mixture was then subjected to silica gel column chromatography using chloroform:ethyl acetate (1:2) as eluent to yield 40% of title compound.

### 4. Conclusion

One-pot multicomponent Biginelli reaction was employed to conduct the synthesis of a new compound ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate.

**Supplementary Materials:** The following are available on line at [www.mdpi.com/1422-8599/2017/3/M954](http://www.mdpi.com/1422-8599/2017/3/M954), HRESIMS, FTIR, <sup>1</sup>H-, NMR, APT, HMQC, and HMBC spectra of the title compound are reported in the supplementary materials as Figures S1–S6 and structure refinement parameters as Table 1.

**Acknowledgments:** The authors gratefully acknowledge to Lembaga Penelitian dan Inovasi, Airlangga University for research funding through the Hibah Riset Mandat Grant 2017.

**Author Contributions:** H.S. brought out the idea, managed the research, and wrote the article. S.Z. carried out the synthesis, K.U.H. analyzed the spectra. I.I. elucidated the spectral data. A.N.K. elucidated the spectral data and corrected the draft. All the authors have read the draft.

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

1. Biginelli, P. Derivati Aldeiduredici Degli Eteri Acetile Dossal-Acetico. *Gazz. Chim. Ital.* **1893**, *23*, 360–413.

2. De Vasconcelos, A.; Oliveira, P.S.; Ritter, M.; Freitag, A.; Romano, L.; Quina, F.H.; Pizzuti, L.; Pereira, C.M.P.; Stefanello, F.M.; Barschak, A.G. Antioxidant capacity and environmentally friendly synthesis of dihydropyrimidin-(2H)-ones promoted by naturally occurring organic acids. *J. Biochem. Mol. Toxicol.* **2012**, *26*, 155–161. [[CrossRef](#)] [[PubMed](#)]
3. Bhatewara, A.; Jetti, S.R.; Kadre, T.; Paliwal, P.; dan Jain, S. Microwave-assisted synthesis and biological evaluation of dihydropyrimidinone derivatives as anti-inflammatory, antibacterial, and antifungal agents. *Int. J. Med. Chem.* **2013**, *2013*, 1–5. [[CrossRef](#)] [[PubMed](#)]
4. Kim, J.; Park, C.; Ok, T.; So, W.; Jo, M.; Seo, M.; Kim, Y.; Sohn, J.H.; Park, Y.; Ju, M.K.; et al. Discovery of 3,4-dihydropyrimidin-2(1H)-ones with inhibitory activity against HIV-1 replication. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 2119–2124. [[CrossRef](#)] [[PubMed](#)]
5. Sośnicki, J.G.; Struk, Ł.; Kurzawski, M.; Perużyńska, M.; Maciejewska, G.; dan Drożdżik, M. Regioselective synthesis of novel 4,5-diaryl functionalized 3,4-dihydropyrimidine-2(1H)-thiones via a non-biginielli-type approach and evaluation of their in vitro anticancer activity. *Org. Biomol. Chem.* **2014**, *12*, 3427–3440. [[CrossRef](#)] [[PubMed](#)]
6. Nagarajaiah, H.; Mukhopadhyay, A.; dan Moorthy, J.N. Biginelli reaction: An overview. *Tetrahedron Lett.* **2016**, *57*, 5135–5149. [[CrossRef](#)]
7. Yoshida, N.; Kasuya, N.; Haga, N.; Fukuda, K. Brand-new biomass-based vinyl polymers from 5-hydroxymethylfurfural. *Polym. J.* **2008**, *40*, 1164–1169. [[CrossRef](#)]
8. Jin, T.; Zhang, S.; Li, T. p-Toluenesulfonic Acid-catalyzed efficient synthesis of dihydropyrimidines: Improved high yielding protocol for the biginelli reaction. *Synth. Commun.* **2002**, *32*, 1847–1851. [[CrossRef](#)]
9. Matache, M.; Dobrota, C.; Bogdan, N.D.; Dumitru, I.; Ruta, L.L.; Paraschivescu, C.C.; Farcasanu, I.C.; Baciuc, I.; Funeriu, D.P. Synthesis of fused dihydro-pyrimido [4, 3-d] coumarins using Biginelli multicomponent reaction as key step. *Tetrahedron* **2009**, *65*, 5949–5957. [[CrossRef](#)]
10. Pretsch, E.; Bühlmann, P.; Badertscher, M. *Structure Determination of Organic Compounds, Tables of Spectral Data*, 4th Completely Revised ed.; Springer: Berlin, Germany, 2009; pp. 70, 134, 138.
11. Pavia, D.L.; Lampman, G.M.; Kriz, G.S.; Vyvyan, J.R. *Introduction to Spectroscopy*, 4th ed.; Brooks/Cole, Cengage Learning: Belmont, CA, USA, 2009; pp. 151, 178, 293.
12. Breitmaier, E. *Structure Elucidation by NMR in Organic Chemistry: A Practical Guide*, 3th Revised ed.; John Wiley & Sons, Ltd.: West Sussex, UK, 2002; pp. 11–12.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

# Ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 **Ângelo de Fátima, Taniris C. Braga, Leonardo da S. Neto, Bruna S. Terra et al. "A mini-review on Biginelli adducts with notable pharmacological properties", Journal of Advanced Research, 2015** 2%

Publication
- 2 **Tjitjik Tjahjandarie, Ratih Saputri, Mulyadi Tanjung. "5,9,11-Trihydroxy-2,2-dimethyl-3-(2-methylbut-3-en-2-yl)pyrano[2,3-a]xanthen-12(2H)-one from the Stem Bark of Calophyllum tetrapterum Miq.", Molbank, 2017** 2%

Publication
- 3 **derpharmachemica.com** 2%

Internet Source
- 4 **S. Nanjunda Swamy, Basappa, S. Naveen, B. Prabhuswamy, M. A. Sridhar, Javaregowda S. Prasad, Kanchugarakoppal S. Rangappa. "Synthesis and crystal structure analysis of 2-** 1%



# (4-methyl-2'-biphenyl)-4-amino-1,2,4-triazole-3-thiol", Structural Chemistry, 2006

Publication

---

5	<a href="https://patents.google.com">patents.google.com</a> Internet Source	1 %
6	<a href="http://www.oalib.com">www.oalib.com</a> Internet Source	1 %
7	J. G. Contreras, C. A. López, G. V. Seguel. " The Raman and IR Spectra of (C H ) N ZnBr ", Spectroscopy Letters, 2006 Publication	1 %
8	<a href="http://krishikosh.egranth.ac.in">krishikosh.egranth.ac.in</a> Internet Source	1 %
9	<a href="http://www.arkat-usa.org">www.arkat-usa.org</a> Internet Source	1 %
10	"Crystal Structure of Bis(2,4,6-trimethylphenyl)-phosphine Oxide", Molbank, 2017 Publication	1 %
11	"Readily Available Chiral Benzimidazoles-Derived Guanidines as Organocatalysts in the Asymmetric $\alpha$ -Amination of 1,3-Dicarbonyl Compounds", Molecules, 2017 Publication	1 %
12	<a href="http://mdpi.com">mdpi.com</a> Internet Source	1 %

---



13

Brown, Elvie E, Dehong Hu, Nehal I Abu-Lail, and Xiao Zhang. "Potential of nanocrystalline cellulose-fibrin nanocomposites for artificial vascular graft applications", *Biomacromolecules*, 2013.

Publication

---

<1 %

14

Nemanja Trišović, Nataša Valentić, Marko Erović, Tatjana Đaković-Sekulić, Gordana Ušćumlić, Ivan Juranić. "Synthesis, structure, and solvatochromic properties of pharmacologically active 5-substituted 5-phenylhydantoins", *Monatshefte für Chemie - Chemical Monthly*, 2011

Publication

---

<1 %

15

Hashem Sharghi, Mahboubeh Jokar. " Al O /MeSO H: A Novel and Recyclable Catalyst for One-Pot Synthesis of 3,4-Dihydropyrimidinones or Their Sulfur Derivatives in Biginelli Condensation ", *Synthetic Communications*, 2009

Publication

---

<1 %

16

Mulyadi Tanjung, Tjitjik Srie Tjahjandarie, Ratih Dewi Saputri, Baharrani Dwi Kurnia et al. " Calotetrapterins A-C, three new pyranoxanthenes and their cytotoxicity from the stem bark of Miq ", *Natural Product Research*, 2019

Publication

---

<1 %

17 [tel.archives-ouvertes.fr](http://tel.archives-ouvertes.fr) Internet Source <1 %

---

18 [www.beilstein-journals.org](http://www.beilstein-journals.org) Internet Source <1 %

---

19 [www.sumobrain.com](http://www.sumobrain.com) Internet Source <1 %

---

20 [www.thieme-connect.com](http://www.thieme-connect.com) Internet Source <1 %

---

21 Mihaela Matache, Cristian Dobrota, Niculina D. Bogdan, Daniel P. Funeriu. "Recent Developments in the Reactivity of the Biginelli Compounds", Current Organic Synthesis, 2011  
Publication <1 %

---

22 Robert-Jan van Putten, Jan C. van der Waal, Ed de Jong, Carolus B. Rasrendra, Hero J. Heeres, Johannes G. de Vries. "Hydroxymethylfurfural, A Versatile Platform Chemical Made from Renewable Resources", Chemical Reviews, 2013  
Publication <1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Ethyl 4-[5-(methoxymethyl)furan-2-yl]-6-methyl-2oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---