

Antioxidant activity of flavonoid compounds from the leaves of *Macaranga gigantea*

by Alfinda Novi Kristanti

Submission date: 09-Mar-2020 05:15PM (UTC+0800)

Submission ID: 1272164047

File name: 2014-JCPR-6-6-688-692.pdf (134.25K)

Word count: 2977

Character count: 14029



Antioxidant activity of flavonoid compounds from the leaves of *Macaranga gigantea*

Nanik Siti Aminah*, Alfinda Novi Kristanti and Mulyadi Tanjung

Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry,
Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia

ABSTRACT

Three flavonoid compounds have been isolated from the leaves of *Macaranga gigantea* (Euphorbiaceae) namely as glyasperin A (1), broussonflavonol F (2), apigenin (3). Their structures were elucidated by spectroscopic methods including UV, IR, HRESIMS, 1D and 2D NMR analysis. Compounds 1–3 were evaluated for their radical scavenging against 2,2-diphenyl-1-picrylhydrazyl (DPPH), showing their IC_{50} were 125.10, 708.54, and 518.01 μ M, respectively. The results indicate that as glyasperin A (1) more active than ascorbic acid (329.01 μ M).

Keywords: Glyasperin A, Broussonflavonol F, Apigenin, Flavonoid, *Macaranga gigantea*, Antioxidant


INTRODUCTION

Macaranga is one of the largest genus of the family Euphorbiaceae, comprising of about 300 species. In addition to in Indonesia, found in parts of Africa, Madagascar, Asia, the east coast of Australia, and the Pacific islands. The *Macaranga* plants are generally in the form of shrubs or trees, and grow in a place that gets a lot of sunlight in secondary forests or forests that have been damaged [1]. The phytochemical studies, this plants producing phenolic compounds, particularly flavonoid and stilbene derivatives. Structural variation of these derivatives occurs as a result of terpenoid substituents on various positions of aromatic rings. The terpenoid substituents identified include isoprenyl (C₅), geranyl (C₁₀), farnesyl (C₁₅) and geranyl-geranyl (C₂₀) [2,3,4,5]. The compound of flavonoid and stilbenoid from *Macaranga* plants exhibit various of bioactivity as antitumor, anticancer, antimalarial, antimicrobial, cyclooxygenase, and antioxidant [6,7,8,9,10,11]. In continuation of our phytochemical work of Indonesian *Macaranga* plants aiming to find new antioxidant compounds from *Macaranga gigantea*. In this paper, we report the isolation of flavonoid compounds, glyasperin A (1), broussonflavonol F (2), apigenin (3) from the methanol extract of the leaves of *Macaranga gigantea*. The antioxidant activity of compounds 1–3 against DPPH is also briefly described.

EXPERIMENTAL SECTION

The leaves of *Macaranga gigantea* were collected in July 2012 from Cinta Damai Village, District Banyuasin, South Sumatera, Indonesia. The plant was identified by Mr Ismail Rachman, Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia, and the voucher specimen was deposited in the herbarium. The dried and powdered leaves of *Macaranga gigantea* (1.5 kg) were macerated in methanol at room temperature three times, and the methanol extract was evaporated under reduced pressure to give a

dark brown residue (70 g). Furthermore, the methanol extract were partition with n-heksana and ethyl acetate. The ethyl acetate extract (40 g) was separated by vacuum liquid chromatography on silica gel eluted with *n* hexane-ethyl acetate mixture containing increasing amount of ethyl acetate (90:10, 80:20; 50:50 and 30:70) to give five major fraction A-E. On TLC analysis, fraction B (450 mg) showed two major spots on purification of this fraction using planar radial chromatography, and using *n* hexane-ethyl acetate (from 9:1 and 8:2) to yielded compound **1** (160 mg) and **2** (20 mg). The separation of fraction C (1.2 g) by flash chromatography with *n* hexane-ethyl acetate (from 8:1 and 7:3) to give three subfractions C₁-C₃. Further purification of subfraction C₃ (280 mg) by radial chromatography with *n* hexane- chloroform 3:7, and chloroform to give compound **3** (18 mg).



Glyasperin A (**1**), pale yellow solid, UV (MeOH) λ_{maks} nm (log ϵ): 205 (4.32); 230 sh (4.11); 254 (4.05); 271 (4.06); 347 (3.98); and 368 (3.98) nm; (MeOH + NaOH) 205 (4.80); 230 sh (4.18); 278 (4.05); 321 (3.91); and 413 (4.07); (MeOH + AlCl₃) 205 (4.30); 234 sh (4.04); 265 (4.10); 305 sh (3.68); 368 (3.75); and 434 (4.12) nm; (AlCl₃+ HCl) 206 (4.25); 234 sh (4.03); 264 (4.08); 306 sh (3.65); 348 (3.67); 368 (3.73); and 435 (4.11) nm; (NaOAc): 205 (4.89); 268 (4.08); 347 (3.73); and 435 (4.09) nm. IR (KBr) ν_{max} (cm⁻¹): 3321 (OH); 2964, 2912 (CH alkyl); 1645 (conj. C=O); and 1606-1448 (C=C aromatic). HRESIMS m/z [M-H]⁻ 421.1647 (calcd for C₂₅H₂₅O₆, 421.1651). ¹H NMR (500 MHz, acetone-d₆): see Table 1. ¹³C NMR (125 MHz, acetone-d₆): see Table 1.

Brousoflavonol F (**2**), pale yellow solid, UV (MeOH) λ_{maks} nm (log ϵ): 204 (4.36); 230 sh (4.19); 256 (4.20); 340 (3.58); 273 (4.10); 346 (4.02); and 364 (3.12) nm; (MeOH + AlCl₃) 204 (4.80); 232 sh (4.20); 270 (4.10); and 432 (4.27) nm; (AlCl₃+ HCl) 204 (4.25); 232 sh (4.09); 270 (4.12); 308 sh (3.69); 350 (3.70); 368 (3.80); and 432 (4.15) nm; (NaOAc): 204 (4.90); 266 (4.10); 346 (3.75); and 435 (4.11) nm. HRESIMS m/z [M-H]⁻ 421.1657 (calcd for C₂₅H₂₅O₆, 421.1651). ¹H NMR (400 MHz, acetone-d₆): see Table 2. ¹³C NMR (100 MHz, acetone-d₆): see Table 2. Apigenin (**3**), yellow solid, UV (MeOH) λ_{maks} : 205 (4.49); 228 sh (4.29); 275 (4.18); 300 (4.10); 341 (4.12), and 383 (4.00) nm, (MeOH + AlCl₃) 205 (4.50); 230 sh (4.25); 276 (4.17); 300 (4.08); 341 (4.11); and 382 (3.99) nm; (AlCl₃ + HCl) 205 (4.49); 228 sh (3.86); 276 (4.18); 300 (4.10); 343 (4.12); and 385 (4.00) nm; (NaOAc) 202 (4.76); 272 (4.18); 300 (4.05); 344 (4.01); and 356 (3.59) nm. EIMS: m/z (% relatif): 270 (M⁺, 100); 242 (16); 153 (17); and 121 (11). ¹H-NMR (500 MHz, acetone-d₆) δ_{H} (ppm): 13.00 (1H, s, 5-OH); 9.71 (1H, s, 7-OH); 9.28 (1H, s, 4'-OH); 7.92 (2H, d, *J* = 9.1 Hz, H-2/H-6); 7.01 (2H, d, *J* = 9.1 Hz, H-3/H-5); 6.62 (1H, s, H-3); 6.53 (1H, d, *J* = 2.5 Hz, H-8); and 6.24 (1H, d, *J* = 2.5 Hz, H-6).

DPPH scavenging activity test: Determination of the antioxidant activity of the isolated performed using reagent DPPH (2,2-diphenyl-1-pikrihidrazil) using methods of reduction of free radicals as measured by UV spectrometer at λ 517 nm [12,13,14]. Determination of antioxidant activity done by the dissolving a compounds assay with

methanol, then added solution of 0.1 M buffer acetate (pH 5.5) and added DPPH radical solution of 5.10^{-4} M. Determination of the inhibition of isolated compounds against DPPH radical was observed using a spectrometer at λ 517 nm after incubation for 30 min at 20°C.

RESULTS AND DISCUSSION

Three flavonoids, glyasperin A (1), broussonin F (2), apigenin (3) have been isolated from the leaves of *Macaranga gigantea*. Their structures were elucidated by spectroscopic methods including UV, IR, HRESIMS, 1D and 2D NMR spectrum.

Table 1. NMR spectroscopic data of glyasperin A (1)

No.C	δ_H (mult, J Hz)	δ_C	HMBC
2	-	147.0	-
3	-	136.4	-
4	-	176.4	-
4a	-	103.8	-
5	-	158.7	-
6	-	111.6	-
7	-	162.9	-
8	6.56 (s)	93.7	C-4a, C-6, C-7, C-8a
8a	-	155.5	-
1'	-	128.9	-
2'	8.00 (d, 2.5)	130.1	C-2, C-4', C-6'
3'	-	132.9	-
4'	-	157.9	-
5'	6.97 (d, 8.5)	115.6	C-1', C-3', C-4'
6'	7.91 (dd, 8.5; 2.5)	127.8	C-2', C-2'', C-4'
1''	3.30 (d, 7.4)	21.8	C-2, C-3, C-4', C-2''
2''	5.23 (m, 7.4)	123.2	C-2, C-3, C-4', C-2''
3''	-	131.5	-
4''	1.77 (s)	25.8	-
5''	1.64 (s)	18.0	C-1', C-3', C-4'
1'''	3.34 (d, 7.3)	29.9	C-3', C-4', C-2'', C-3''', C-4'''
2'''	5.33 (m, 7.3)	123.1	C-1', C-4', C-5''
3'''	-	131.5	-
4'''	1.74 (s)	25.8	C-2', C-3''', C-5''
5'''	1.74 (s)	17.8	C-2', C-3'', C-4''
3-OH	7.85 (br, s)	-	C-3
5-OH	12.41 (br, s)	-	C-4a, C-5, C-6
7-OH	9.78 (br, s)	-	C-7
4'-OH	8.89 (br, s)	-	C-4

Glyasperin A (1) was obtained as pale yellow solid, showed a quasimolecular ion $[M-H]^-$ at m/z 421.1647 consistent to the molecular formula $C_{25}H_{25}O_6$. The UV spectrum of 1 exhibited absorption maxima for a flavonol structure at λ_{max} 205, 230 sh, 254, 271, 347, and 368 nm, and showed a bathochromic shift on addition of $AlCl_3$, $AlCl_3 + HCl$, and $NaOAc$ [15]. The IR spectrum indicated absorptions for hydroxyl (3321 cm^{-1}), conjugated carbonyl (1645 cm^{-1}), and aromatic ($1606\text{--}1448\text{ cm}^{-1}$) groups. The 1H -NMR spectrum of compound 1 showed three aromatic proton signals for ABX system at δ_H 8.00 (1H, *d*, $J = 2.5$ Hz); 7.91 (1H, *dd*, $J = 8.5, 2.5$ Hz); and 6.97 ppm (1H, *d*, $J = 8.5$ Hz) corresponding to the group substituent at C-3' and C-4' in ring B flavonols which also showed an isoprenyl group at C-3'. A singlet signal at 6:56 ppm in the aromatic region of the 1H -NMR spectrum showed one isoprenyl group at A ring of flavonol structure. The existence of two isoprenyl chain of compound 1 showed the presence of four methyl groups (δ_H 1.74, 1.71, 1.70, and 1.60 ppm), two methylene groups (3:34 and 3:30 ppm), two vinyl groups (δ_H 5:33, and 5:23 ppm). In the ^{13}C NMR (APT experiment, Table 1) showed 22 carbon signals representing 25 carbon atoms were observed. Two of signals at δ_C 136.4 and 176.5 are characteristic for C-3 and C-4 of a flavonol structure [x]. Five carbon signals (δ_C 147.0; 155.5; 157.9; 158.7; and 162.9 ppm) characteristic for the region oksiaril signals which indicate that the structure is a derivative of kaempferol. The correlation of the one bond and the two/three bond 1H - ^{13}C compound 1 can be seen in the HMQC and HMBC spectra (Table-1). The presence of an isoprenyl group at C-6 shows in the HMBC spectrum, the long-range correlation between a proton signal chelate-OH group at δ_H 12.41 ppm with three quaternary atoms at δ_C 103.8 (C-4a), 111.6 (C-6), and 158.7 (C-5). This is supported by the correlation between a methylene proton signal at δ_H 3.30 ppm with three quaternary atoms at δ_C 111.6 (C-6), 158.7 (C-5); and 162.9 (C-7). Based on data from 1D and 2D NMR of compound 1 is 6,3'-

diisoprenylkaempferol or known as glyasperin A [16]. Other HMBC correlations consistent with the structure **1** are shown in Table 1.

Brousoflavonol F (**2**) was also obtained as pale yellow solid. The ion peak at m/z 421.1657 $[M-H]^-$ in the HRESIMS spectrum gave the molecular $C_{25}H_{25}O_6$, which was isomeric with **1**. Based on UV, 1H NMR, ^{13}C NMR and HMQC spectrum, the compound **2** identical with compound **1**. The placement of isoprenyl at C-8 shown in HMBC spectrum (Table 2). The presence of long-range correlations in the HMBC spectrum of **2** between the singlet proton of a chelated $-OH$ group at δ_H 12.09 with two quaternary at δ_C 104.1 (C-4a), 159.8 (C-5), and one methine carbon at δ_C 98.9 (C-6) showed isoprenyl at C-8. The presence of isoprenyl group at C-8, the correlation between a methylene proton signal at δ_H 3.55 with four quaternary atoms at δ_C 103.8 (C-4a), 133.3 (C-3''), 154.9 (C-8a), and 161.9 (C-7). Based on data from 1D and 2D NMR, compound **2** is 8,3'-diisoprenylkaempferol or known as brousoflavonol F [16]. Other HMBC correlations consistent with the structure **2** are shown in Table 2.

Table 2. NMR spectroscopic data of brousoflavonol F (**2**)

No.C	δ_H (mult, J Hz)	δ_C	HMBC
2	-	147.1	-
3	-	136.4	-
4	-	176.7	-
4a	-	104.1	-
5	-	159.8	-
6	6.34 (s)	98.9	C-4a, C-5, C-7, C-8
7	-	161.9	-
8	-	107.1	-
8a	-	154.9	-
1'	-	123.7	-
2'	8.04 (d, 2.4)	129.8	C-2, C-3', C-4', C-2''
3'	-	129.1	-
4'	-	157.7	-
5'	7.01 (d, 7.2)	128.2	C-2, C-2', C-4'
6'	8.05 (dd, 7.2; 2.4)	115.7	C-1', C-2', C-4'
1''	3.55 (d, 7.4)	22.2	C-5, C-6, C-7, C-2'', C-3''
2''	5.39 (d, 1.6)	123.3	C-1', C-4', C-5'
3''	-	133.3	-
4''	1.65 (s)	18.1	C-2'', C-3'', C-5''
5''	1.80 (s)	25.9	C-2'', C-3'', C-4''
1'''	3.40 (d, 7.6)	29.0	C-2'', C-3'', C-4'', C-2''', C-3'''
2'''	5.31 (t, 1.6)	123.0	C-1'', C-4'', C-5''
3'''	-	132.0	-
4'''	1.74 (s)	25.9	C-2''', C-3''', C-5'''
5'''	1.74 (s)	17.8	C-2'', C-3'', C-4''
3-OH	7.89 (br, s)	-	C-3
5-OH	12.09 (br, s)	-	C-4a, C-5, C-6
7-OH	6.58 (br, s)	-	C-7
4'-OH	8.96 (br, s)	-	C-4'




Figure 1. The difference of HMBC spectrum for compound **1** and **2**

Apigenin (**3**) was obtained as yellow solid, showed ion peak at m/z 270 $[M]^+$ in the EIMS spectrum. The UV spectrum of **3** exhibited absorption maxima for a flavon structure at λ_{max} 205, 228 sh, 275, 300, 341, and 383, and showed a bathochromic shift on addition of $AlCl_3$, $AlCl_3 + HCl$, and $NaOAc$. The 1H -NMR spectrum of compound **3** showed a pair of doublets ($J = 9.1$ Hz) in the aromatic region at δ_H 7.92 and 7.01 (each 2H), suggested the signal of a *p*-hydroxyphenyl at B ring. The presence of a pair of doublets ($J = 2.5$ Hz) at δ_H 6.53 and 6.24 in A ring corresponding to the proton at H-8 and H-6. The presence of a singlet signal at δ_H 6.62 suggested that the placement H-3 for a flavon structure. Based on data from UV, EIMS and 1H NMR, compound **3** is 5,7,4'-trihydroxyflavone or known as apigenin [17].

The antioxidant activity of glyasperin A (**1**), brousoflavonol F (**2**), apigenin (**3**) at different concentration (1000, 500, 250, 100, and 50 μM) were evaluated against the DPPH radical scavenging. The IC_{50} values of glyasperin A (**1**), brousoflavonol F (**2**), apigenin (**3**) showed radical scavenging activity with IC_{50} values 125.10, 708.54, and 518.01 μM , respectively. Ascorbic acid as positive control have IC_{50} 329.01 μM . The results of antioxidant activity showed glyasperin A (**1**) more active than ascorbic acid. Glyasperin A (**1**), and brousoflavonol F (**2**) are isomeric structure. The placement of isoprenyl at C-6 to increasing antioxidant activity than at C-8.

CONCLUSION

Two isomeric isoprenyl flavonols, glyasperin A (**1**) and brousoflavonol F (**2**) together apigenin (**3**) have been isolated from the leaves of *Macaranga gigantea*. Apigenin is a flavone derivative and first time found in the genera of *Macaranga*. The antioxidant activity of compounds **1-3** were evaluated by measuring their ability to scavenge the DPPH radical showed glyasperin A > apigenin > brousoflavonol F. The result indicate that compound **1** to give very high activity than ascorbic acid as positive control. The structure-activity relationship of compounds **1-2** against DPPH radical scavenging suggested that the presence of isoprenyl group at C-6 on glyasperin A more active than the isoprenyl group at C-8 on brousoflavonol F.

Acknowledgements

This research was supported by Directorate of Higher Education, Ministry of National Education, Republic of Indonesia (PUPT Airlangga University 2013). We would like to thank to Mr. Ismail Rachman from the Herbarium Bogoriense, Botanical Garden, Bogor, Indonesia for identifying the species.

REFERENCES

- [1] FR Blattner; K Weising; G Banfer; U Maschwitz; B Fiala, *Mol. Phyl. Evol.*, **2001**, 19, 331-344.
- [2] SC Fang; BC Shieh; RR Wu; CN Lin, *Phytochem.*, **1995**, 38(2), 535-537.
- [3] S Suttivaiyakit; S Unganont; P Suttivaiyakit; A Suksamrarn, A., *Tetrahedron*, **2002**, 58, 3619-3622.
- [4] T Mulyadi; EH Hakim; D Mujahidin; A Damawan; YM Syah, *J. Nat. Prod Comm.*, **2010**, 5, 1209-1211.
- [5] YM Syah; EL Ghisalberty, *J. Nat. Prod Comm.*, **2010**, 5, 219-222.
- [6] JA Beutler; RH Shoemaker; T Johnson; MR Boyd, *J. Nat. Prod.*, **1998**, 61, 1509-1512.
- [7] I Zakaria; N Ahmad; FM Jaafar; A Widyawaruyanti, *Fitoterapia*. **2012**, 83, 968-972.
- [8] TY Lim; YY Lim; CM Cule, *Food Chem.*, **2009**, 37, 304-317.
- [9] DS Jang; M Cuendet; AD Pawlus; LBS Kardono; K Kawanishi; R Farnsworth; HHS Fong; JM Pezzuto; AD Kinghorn, *Phytochem.*, **2004**, 65, 345-350.
- [10] DS Jang; M Cuendet; ME Hawthorne; LBS Kardono; K Kawanishi; HHS Fong; RG Mehta; JM Pezzuto; AD Kinghorn, *Phytochem.*, **2002**, 61, 867-873.
- [11] K Matsunami; I Takamori; T Shinzato; M Aramoto; K Kondo; H Otsuka; Y Takeda, *Chem. Pharm. Bull.*, **2006**, 54, 1403-1407.
- [12] T Mulyadi; DS Saputri; ST Tjitjik, *J. Chem. Pharm. Res.*, **2014**, 6(1):705-708.
- [13] ST Tjitjik; DS Saputri; T Mulyadi, *J. Chem. Pharm. Res.*, **2014**, 6(4):786-790.
- [14] N Madhusudhanan; I Vidya; T Laksmi, *J. Chem Pharm Res.*, **2013**, 5(10): 194-200.
- [15] E Hnawia; O Thoison; FG Voegelien; D Bourret; T Sevenet, *Phytochem.*, **1992**, 29, 2367-2368.
- [16] SC Fang; B Shieh; R Wu; C Lin, *Phytochem.*, **1995**, 38, 535-537.
- [17] J Nazaruk; J., *Acta Poloniae Pharmachem.*, **2006**, 63, 317-319.

Antioxidant activity of flavonoid compounds from the leaves of *Macaranga gigantea*

ORIGINALITY REPORT

20%
SIMILARITY INDEX

13%
INTERNET SOURCES

17%
PUBLICATIONS

0%
STUDENT PAPERS

PRIMARY SOURCES

1 www.science.gov 1%
Internet Source

2 Lawrence O. Manguro Arot, Lawrence A.D. Williams. "A flavonol glycoside from *Embelia schimperi* leaves", *Phytochemistry*, 1997 1%
Publication

3 210.75.237.14 1%
Internet Source

4 Ming-Bao Luan, Zi-Zheng Zou, Juan-Juan Zhu, Xiao-Fei Wang, Ying Xu, Zhi-Min Sun, Jian-Hua Chen. "Genetic diversity assessment using simple sequence repeats (SSR) and sequence-related amplified polymorphism (SRAP) markers in ramie", *Biotechnology & Biotechnological Equipment*, 2015 1%
Publication

5 jurnal.kimia.fmipa.unmul.ac.id 1%
Internet Source

6 es.scribd.com 1%
Internet Source

7	doaj.org Internet Source	1 %
8	hdl.handle.net Internet Source	1 %
9	Z. P. Xiao, H. K. Wu, T. Wu, H. Shi, B. Hang, H. A. Aisa. "Kaempferol and quercetin flavonoids from <i>Rosa rugosa</i> ", <i>Chemistry of Natural Compounds</i> , 2006 Publication	1 %
10	Toshio Fukai, Yue-Hu Pei, Taro Nomura, Chun-Quan Xu, Li-Jun Wu, Ying-Jie Chen. "Isoprenylated flavanones from <i>Morus cathayana</i> ", <i>Phytochemistry</i> , 1998 Publication	1 %
11	lume.ufrgs.br Internet Source	1 %
12	Chen, L.. "Flavonoids from <i>Disporum cantoniense</i> (Liliaceae)", <i>Biochemical Systematics and Ecology</i> , 200911 Publication	1 %
13	www.freshpatents.com Internet Source	1 %
14	A.H. Meriçli, G.V. Seyhan. " Constituents of . Leaves ", <i>Pharmaceutical Biology</i> , 2008 Publication	<1 %

15 Yi Lu, Tatsuya Umeda, Akihito Yagi, Kanzo Sakata, Tirthankar Chaudhuri, D.K. Ganguly, Scion Sarma. "Triterpenoid saponins from the roots of tea plant (*Camellia sinensis* var. *assamica*)", *Phytochemistry*, 2000
Publication

16 studylib.net
Internet Source

17 epublications.unilim.fr
Internet Source

18 www.j3.jstage.jst.go.jp
Internet Source

19 Achara Raksat, Wisanu Maneerat, Raymond J. Andersen, Stephen G. Pyne, Surat Laphookhieo. " Antibacterial Prenylated Isoflavonoids from the Stems of ", *Journal of Natural Products*, 2018
Publication

20 pubmedcentralcanada.ca
Internet Source

21 Augustin E Nkengfack, Anatole G.B Azebaze, Alain K Waffo, Zacharias T Fomum, Michelle Meyer, Fanie R van Heerden. "Cytotoxic isoflavones from *Erythrina indica*", *Phytochemistry*, 2001
Publication

22 Marwan Shabana, Mariam Gonaid, Maha Mahmoud Salama, Essam Abdel-Sattar. "Phenylalkylamine alkaloids from ", Natural Product Research, 2006
Publication <1 %

23 umexpert.um.edu.my
Internet Source <1 %

24 Brent J. Yoder, Shugeng Cao, Andrew Norris, James S. Miller et al. " Antiproliferative Prenylated Stilbenes and Flavonoids from from the Madagascar Rainforest", Journal of Natural Products, 2007
Publication <1 %

25 C. Georgopoulou. "Acretoside, a new sucrose ester from Aristolochia cretica *", Journal of Asian Natural Products Research, 12/1/2005
Publication <1 %

26 F. Ameer, Ivan R. Green, K. Krohn, M. Sitoza. " Strategies towards the Synthesis of 6 - , - Diethylcarbamyloxy - 1,4 - dimethoxy - 7 - naphthylboronic Acid ", Synthetic Communications, 2007
Publication <1 %

27 Jin-Bin Wu, Yih-Dih Cheng, Li-Ling Su, Chi-Wen Kuo, Shieng-Chu Kuo. "A flavonol C-glycoside from Moghania macrophylla", Phytochemistry, 1997
Publication <1 %

28

Maria Kotsos, Nektarios Aligiannis, Sofia Mitaku, Alexios-Leandros Skaltsounis, Catherine Charvala. " Chemistry of Plants from Crete: Stachyspinoside, a New Flavonoid Glycoside And iridoids from ", Natural Product Letters, 2001

Publication

<1 %

29

s3.amazonaws.com

Internet Source

<1 %

30

F. Peuzzdni, L. Verotta, C. B. Rogers, R. Colombo, B. Pedrotti, G. Balconi, E. Erba, M. D'incalci. "Cell Growth Inhibitor Constituents From Combretum Kraussii", Natural Product Letters, 1993

Publication

<1 %

31

Hasan Şahin, Aynur Sarı, Nurten Özsoy, Berna Özbek Çelik, Onur Koyuncu. " Two new phenolic compounds and some biological activities of Sibth. & Sm. subaerial parts ", Natural Product Research, 2018

Publication

<1 %

32

Jenkins, T.. "Flavonoids from the root-bark of Dioclea grandiflora", Phytochemistry, 199910

Publication

<1 %

33

Shimokawa, Yoko, Yusuke Akao, Yusuke Hirasawa, Khalijah Awang, A. Hamid A. Hadi, Seizo Sato, Chihiro Aoyama, Jiro Takeo, Motoo

<1 %

Shiro, and Hiroshi Morita. "Gneyulins A and B, Stilbene Trimers, and Noidesols A and B, Dihydroflavonol-C-Glucosides, from the Bark of *Gnetum gnemonoides*", Journal of Natural Products, 2010.

Publication

34

Shu-Geng Cao, Keng Yeow Sim, S. H. Goh. "Minor Methylated Pyranoamentoflavones from ", Natural Product Letters, 2006

Publication

<1 %

35

Xu, T.H.. "A new glycoside from *Polygala tenuifolia* Willd", Chinese Chemical Letters, 200807

Publication

<1 %

36

academicjournals.org

Internet Source

<1 %

37

kanazawa-u.repo.nii.ac.jp

Internet Source

<1 %

38

onlinelibrary.wiley.com

Internet Source

<1 %

39

www.hindawi.com

Internet Source

<1 %

40

www.pubmedcentral.nih.gov

Internet Source

<1 %

41

www.revistadechimie.ro

Internet Source

<1 %

42

Iliana Krasteva, Stefan Platikanov, Stefan Nikolov, Maki Kaloga. " Flavonoids from ",
Natural Product Research, 2007

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Antioxidant activity of flavonoid compounds from the leaves of *Macaranga gigantea*

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
