

# Copro-parasitological examinations and molecular determination of *Eimeria* species in Madura cattle reared on Madura Island, Indonesia

*by* Endang Suprihati

---

**Submission date:** 29-Mar-2022 10:18AM (UTC+0800)

**Submission ID:** 1795646541

**File name:** ological\_examinations\_and\_molecular\_determination\_of\_Eimeria.pdf (1.41M)

**Word count:** 3820

**Character count:** 19819



9

Contents lists available at ScienceDirect

Parasitology International

journal homepage: [www.elsevier.com/locate/parint](http://www.elsevier.com/locate/parint)

## Coproparasitological examinations and molecular determination of *Eimeria* species in Madura cattle reared on Madura Island, Indonesia<sup>☆</sup>

Poedji Hastutiek<sup>a</sup>, Nunuk Dyah Retno Lastuti<sup>a</sup>, Lucia Tri Suwanti<sup>a</sup>, Agus Sunarso<sup>a</sup>,  
Endang Suprihati<sup>a</sup>, Dyah Ayu Kurniawati<sup>a</sup>, Makoto Matsubayashi<sup>a,b,\*</sup>

<sup>a</sup> Department of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia

<sup>b</sup> Department of Veterinary Science, Graduate School of Life and Environmental Sciences, Osaka Prefecture University, Osaka, Japan

### ARTICLE INFO

#### Keywords:

*Eimeria*  
Indonesia  
Madura cattle  
PCR

### ABSTRACT

Madura cattle, which are native to Indonesia and mainly kept on Madura Island, East Java, are expected to contribute to improving the regional meat self-sufficiency. *Eimeria* spp. are the most pathogenic protozoans among gastrointestinal parasites in livestock but no molecular surveys of *Eimeria* spp. in Madura cattle have been conducted to date. In this study, a total of 183 fecal samples were collected from Madura cattle and 60 (32.8%) were positive for parasites of protozoans and nematodes by the sugar floatation method. Among the samples with parasites, *Eimeria* spp. oocysts were detected in 50 samples (27.3%) with an average OPG value of 1686.1. *Eimeria* spp. were successfully identified to the species level in 26 samples with *Eimeria bovis* being the most prevalent, followed by *E. zuernii* and *E. aubrunensis*. A total of 21 samples showed mixed infection of more than two species of *Eimeria*. *E. bovis* and *E. zuernii* have been recognized as having high virulence and, thus, these parasites are potential sources of severe coccidiosis and the cause of infections in other cattle. Although additional studies are warranted, these results can be helpful for improving the management and productivity of Madura cattle.

### 1. Introduction

Infections by gastrointestinal parasites including protozoans have been reported in livestock worldwide. Parasitic infections are primarily characterized by gastroenteric symptoms such as watery or bloody diarrhea, which often drastically constrain productivity, through reduced growth or milk yield in cattle [1–3]. Among parasites, *Eimeria* spp. are one of the significant protozoan organisms that cause coccidiosis and result in high morbidity and mortality, especially in calves up to 1 year [4]. The prevalence of gastrointestinal parasites differs by country, region and even management systems on farms.

Although more than 12 members of genus *Eimeria* infect cattle [5], five species cause clinical symptoms such as watery or bloody diarrhea followed by loss of appetite, depression, dehydration, and weight loss, which leads to retarded growth [6]. Two species, *E. bovis* and *E. zuernii*, are the greatest impact due to high virulence and high mortality [6]. As it is sometimes difficult to identify the species of *Eimeria* based on oocyst

morphology in routine fecal examinations, molecular characterization is necessary to accurately make identification to the species level.

In Indonesia, the population of 16.5 million beef cattle and 0.5 million dairy cattle [7] is concentrated on Java Island (2.6% and 98.9% of beef and dairy cattle, respectively) [8]. Madura cattle are one of the native livestock breeds in Indonesia and are mainly raised on Madura Island, which is located off the northeastern coast of Java Island and contains four districts of East Java Province. The Madura breed is thought to have been developed by inbreeding Zebu (*Bos indicus*) and Banteng (*Bos javanicus*) and is maintained through careful genetic selection [9]. This breed is reported to possess the advantages of high adaptivity to tropical environments, high reproductive performance, high meat quality, and ease of rearing owing to its small to medium size [10]. Recently, the Ministry of Agriculture in Indonesia launched a revitalization program to reduce beef imports and increase domestic production. Based on policies of the government, Madura cattle is listed as one of beef cattle breeds for supplying high-protein food commodities

<sup>☆</sup> The authors declare that they have no conflicts of interest. All authors contributed equally in writing the manuscript.

\* Corresponding author at: Division of Veterinary Science, Graduate School of Life and Environmental Sciences, Osaka Prefecture University, 1-58 Rinku Orai Kita, Izumisano, Osaka 598-8531, Japan.

E-mail address: [matsubayashi@vet.osakafu-u.ac.jp](mailto:matsubayashi@vet.osakafu-u.ac.jp) (M. Matsubayashi).

<https://doi.org/10.1016/j.parint.2021.102478>

Received 7 August 2021; Received in revised form 20 September 2021; Accepted 3 October 2021

Available online 6 October 2021

1383-5769/© 2021 Elsevier B.V. All rights reserved.

and achieving self-sufficiency in Indonesia [11].

In six reports of infection of cattle by *Eimeria* spp. focused on Java Island, the prevalence of parasite infection ranged from 15.33% to 85.07%, and differences could be observed among provinces and districts and even depending on detection methods [12–17]. In only one of the published reports, identification of *Eimeria* spp. to species level was determined by molecular analysis [14]. In the one paper examining the prevalence of protozoan parasites, 71.4% of Madura cattle were reported to be infected and *Eimeria* spp. accounted for 53.4% of the protozoan parasites [16], but identifications were not made to the species level. In the present study, we parasitologically examined the feces of Madura cattle and used molecular methods to identify *Eimeria* species in order to assess the potential risk of severe coccidiosis.

23

## 2. Materials and methods

### 2.1. Study areas and examined cattle

The study areas were Kamal and Socah subdistricts of the Bangkalan district of Madura Island, East Java, Indonesia. In August 2019 to November 2020, a total of 183 fecal samples were collected from cattle on 50 farms in the Socah subdistrict and 22 farms in the Kamal subdistrict with the following breakdown of age categories: <6 months, 9 and 13, respectively; 6 months to 2 years, 48 and 18, respectively; and > 2 years, 54 and 41, respectively. One to five feces samples were randomly collected on each farm. Most feces were normal, except for 3 diarrheal and 5 soft ones. The collected samples were stored at 4 °C until analysis.

### 2.2. Fecal examination

Feces were examined by the sugar centrifugation method [18,19]. Briefly, 1 g of fecal samples was used and centrifuged at 800 ×g for 5 min. After the supernatant was discarded, sugar solution with a specific gravity of 1.2 was added, followed by centrifugation, then the upper layer was placed on a glass slide. The entire smear was examined by light microscopy. The species of detected parasites were identified based on the published data [20].

### 2.3. Purification of *Eimeria* oocysts

*Eimeria* oocysts were purified from the remaining feces (approximately 5–10 g) by the sugar flotation method as previously described [14]. Briefly, feces were dispersed in distilled water and filtered through a steel mesh or gauze. After centrifugation at 1100 ×g for 5 min, the sugar solution was added, and the sample was centrifuged at 1200 ×g for 10 min. The *Eimeria* oocysts floating on the surface were recovered using a Pasteur pipette and washed with distilled water. The purified oocysts were resuspended with 1–2 ml of PBS, the number of the oocysts was counted under a microscope after putting 15–20 μl of the solution on a slide glass, OPG (oocysts per 1 g feces) values were calculated. Samples were stored at 4 °C until use in molecular identification analysis.

### 2.4. Molecular identification of *Eimeria* spp.

DNA of *Eimeria* oocysts purified as described above was extracted by DNAzol (Molecular Research Center, Cincinnati, Ohio, USA) according to manufacturer recommended protocol but with the addition of five freeze-thaw cycles before lysis. For identification of six *Eimeria* spp., including pathogenic species (*E. bovis*, *E. zuernii*, *E. alabamensis*, *E. aubrunensis*, *E. ellipsoidalis*, and *E. cylindrica*), PCR targeting the internal transcribed spacer 1 (ITS-1) region of the ribosomal RNA gene was carried out as reported previously [21]. PCR products were subjected to electrophoretic separation on agarose gel, stained with ethidium bromide, and visualized on a UV transilluminator.

## 3. Results and discussion

10

Out of a total of 183 fecal samples examined by the sugar flotation method, 60 samples (32.8%) were positive for parasites of protozoans and nematodes. Among 72 farms surveyed, 27 of 50 farms in Socah and 16 of 22 farms in Kamal subdistricts had positive samples, and oocysts of *Eimeria* spp. were detected in 50 fecal samples (27.3%) (Table 1) (Fig. 1). No age-dependent tendency could be identified in the data. The prevalences of *Eimeria* spp. in cattle of Indonesia were reported in six papers to date, ranging 15.33% to 85.07% [12–17], and those of Madura cattle were 66.0% at Socah and 88.2% at Kamal [16]. Compared to previous data, positivity in our study tended to be lower. Although detailed information in some reports could not be confirmed, these differences in prevalences might be attributable to detection methods, sampling seasons, or management strategies of the farms. Especially, the feces were collected only from April to May as ending of rainy season in previous report [16] and our samples were collected in a relatively long period. Because it was implicated that humid climate could play an important role in development to infectious oocysts [22], further studies are needed to resolve the reason for the different prevalences including examined seasons such as rainy or dry ones.

Based on designation of OPG values of 1–499, 500–5000, or more than 5000 as light, moderate, and high infections, respectively [23], the average OPG in the 50 positive samples in the present study of 1686.1 (Table 1) and only five samples with OPG more than 5000 indicates that most of the examined cattle were lightly infected with *Eimeria* spp. although the OPG values could be changed during patent period of the infection. Additionally, because OPG values of the five soft (4 samples infected with *E. bovis*, including 3 samples with mixed infection by *E. zuernii*, and 1 sample with mixed infection by *E. zuernii* and *E. aubrunensis*) and three diarrheal (1 sample infected by *E. bovis*) fecal samples were 500–4000 and 7000–14,500 OPG, respectively, suggesting the possible link between abnormal forms of feces and *Eimeria* infections.

To date, there have been only two reports of molecular investigations of *Eimeria* spp. in cattle in Indonesia and Korea [14,24]. In the present PCR analyses, we identified six *Eimeria* spp. in 26 samples with *E. bovis* being the most prevalent, followed by *E. zuernii* and *E. aubrunensis*. A total of 21 samples appeared to be mixed infections with more than two species present (11; *E. bovis*, *E. zuernii* and *E. aubrunensis*, 6; *E. bovis*, *E. zuernii*, *E. aubrunensis*, and *E. lindrica*, 4; *E. bovis* and *E. zuernii*). All of mixed infections contained *E. bovis* and *E. zuernii*. Generally, *E. bovis* and *E. zuernii* are recognized as being highly virulent. In other surveys, *E. bovis* and *E. zuernii* were reported to be predominant in Indonesia, Brazil or Nigeria based on morphology [15,25,26], and *E. bovis* was detected with high prevalence in Indonesia and Korea by PCR analysis [14,24]. Our results are in agreement with these previous reports. The remaining 23 samples could not be determined to species level. OPG in 17 of 23 samples was less than 10, likely due to there being an insufficient number of the oocysts in feces for successful PCR amplification or contamination by PCR inhibitors.

Of the other parasites, eggs of nematodes were found in 42 of 183 fecal samples, and *Oesophagostomum* spp. were most frequently detected based on being identified in 26 samples (Table 2). These parasites were previously detected in Indonesia and the prevalences were varied among the studies [12–14,17]. These nematode parasites can be sometimes harmful in cattle due to the reduction in productivities such as weight gain [27]. Oocysts of *Eimeria* spp. were found in 32 in 42 samples positive for nematodes. Infections of nematodes and *Eimeria* spp. are transmitted by ingestion of mature eggs and sporulated oocysts, respectively, and infected animals shed newly developed parasites in their feces in forms that are resistant to most disinfectants and environmental conditions. Parasites can survive for long periods and are potential sources of further infection in breeding environments. Thus, presence of these parasites can indicate direct or indirect infection via fecal-oral routes. In our study, the age-dependency of parasite prevalence could not be clarified, and feces with more than 1000 OPG

**Table 1**  
Summary of study areas, ages of cattle and *Eimeria* spp. identification and prevalence.


Subdistrict (number of samples)	Age category	No. of cattle examined	Number positive <sup>a</sup> (%)	Average OPG <sup>b</sup> (range)	Species identification by PCR analysis <sup>c</sup>						Notes
					<i>E. b</i>	<i>E. z</i>	<i>E. aubr</i>	<i>E. alab</i>	<i>E. c</i>	<i>E. e</i>	
Socah (111)	<6 months	9	5 (55.6)	13.2 (1–50)	5	5	3	–	1	–	Coinfected with two ( <i>n</i> = 2), three (2) and four (1) species Coinfected with three (2) and four (2) species Coinfected with three (6) and four (3) species
	6 months to 2 years	48	11 (22.9)	9.6 (1–50)	4	4	4	–	2	–	
	>2 years	54	13 (24.1)	9.8 (1–50)	9	9	9	–	3	–	
Kamal (72)	<6 months	13	3 (23.1)	10,166.7 (5500–14,500)	1	–	–	–	–	–	Coinfected with two species (1) Coinfected with two species (1) and three species (1)
	6 months to 2 years	18	12 (66.7)	3625.3 (1–15,000)	4	1	1	–	–	–	
	>2 years	41	6 (14.6)	1667 (1–4000)	2	2	1	–	–	–	
Total		183	50 (27.3%)	1686.1 (1–15,000)	25 (96.2%)	21 (80.8%)	18 (69.2%)	0 (0%)	6 (23.12%)	0 (0%)	

*E. e.* *Eimeria ellipsoidalis*. Species-level identification was made in 26 of 50 positive samples and percentages indicate the prevalence of each species among the 26 positive samples.

<sup>a</sup> Fecal samples were examined by sugar floatation method.

<sup>b</sup> OPG: oocysts per 20 ram.

<sup>c</sup> Abbreviations: *E. b*; *Eimeria bovis*, *E. z*; *Eimeria zuernii*, *E. aubr*; *Eimeria aubrunensis*, *E. alab*; *Eimeria alabamensis*, *E. c*; *Eimeria cylindrica*.



**Fig. 1.** Photographs of oocyst of *Eimeria* sp. (A) and egg of *Trichuris* sp. (B) from cattle reared in the Socah subdistrict.

**Table 2**  
Summary of study areas, ages and nematodes results.

Subdistrict (number of farms)	Age category	Number of cattle examined	Number positive (%)	Nematodes identified (percent of total number of cattle examined)						
				<i>Oesophagostomum</i> spp.	<i>Strongyloides</i> spp.	<i>Cooperia</i> spp.	<i>Toxocara vitulorum</i>	<i>Trichuris</i> spp.	<i>Mecistocirrus</i> spp.	<i>Capillaria</i> spp.
Socah (50)	<6 months	9	3 (33.3)	2	2	–	–	–	–	–
	6 months to 2 years	48	3 (6.3)	2	2	–	–	1	–	–
	>2 years	54	13 (24.1)	6	7	–	2	1	1	1
Kamal (22)	<6 months	13	3 (23.1)	1	2	–	–	–	–	–
	6 months to 2 years	18	7 (38.9)	5	2	1	–	–	–	–
	>2 years	41	13 (31.7)	10	1	3	–	–	1	–
Total		183	42 (23.0%)	26 (14.2%)	16 (8.7%)	4 (2.2%)	2 (1.1%)	2 (1.1%)	2 (1.1%)	1 (0.5)

values of *Eimeria* spp. were sometimes detected in the same farms (data not shown). Differences in age-dependent prevalence might be induced due to management systems of individual farms for breeding and, therefore, longitudinal and quantitative surveys for parasites are

needed to evaluate the occurrence and sources of the coccidiosis and their productivities.

Although the number of samples in our survey was limited, we showed that samples could be identified to species level for *Eimeria* spp.

by the molecular method. Madura cattle are believed to be resistant to various diseases and tick infestations [10]. While resistance to gastrointestinal parasites remains unknown, further studies are required to clarify pathogenicity in local cattle herds and to improve productivity through improved management practices.

### Ethics statement

All experiments were carried out without sacrificing live animals. Thus, ethical approval for animal experimentation was not required. All examinations in the field study were conducted with permission of the Ministry of the Environment of the Government of Indonesia. Fecal collection was performed in a non-invasive manner. No animals were sacrificed for the purpose of the field study. No human participants were involved in this study.

### References

- [1] H.W. Ploeger, A. Kloosterman, G. Bargeman, L. von Wuijkhuise, R. van den Brink, Milk yield increase after anthelmintic treatment of dairy cattle related to some parameters estimating helminth infection, *Vet. Parasitol.* 35 (1–2) (1990) 103–116, [https://doi.org/10.1016/0304-4017\(90\)90120-z](https://doi.org/10.1016/0304-4017(90)90120-z).
- [2] J. Vercruyse, E. Claerebout, Treatment vs non-treatment of helminth infections in cattle: defining the threshold, *Vet. Parasitol.* 98 (1–3) (2001) 195–214, [https://doi.org/10.1016/s0304-4017\(01\)00431-9](https://doi.org/10.1016/s0304-4017(01)00431-9).
- [3] B. Lassen, T. Lepik, T. Jarvis, Seasonal recovery of *Eimeria* oocysts from soil on naturally contaminated pastures, *Parasitol. Res.* 113 (3) (2014) 993–999, <https://doi.org/10.1007/s00436-013-3731-6>.
- [4] B. Sudhakara Reddy, S. Sivajothi, V.C. Rayulu, Clinical coccidiosis in adult cattle, *J. Parasit. Dis.* 39 (3) (2015) 557–559, <https://doi.org/10.1007/s12639-013-0395-1>.
- [5] R.O. Sánchez, J.R. Romero, R.D. Founroge, Dynamics of *Eimeria* oocyst excretion in dairy calves in the province of Buenos Aires (Argentina), during their first 2 months of age, *Vet. Parasitol.* 151 (2–4) (2008) 133–138, <https://doi.org/10.1016/j.vetpar.2007.11.003>.
- [6] A. Dausgschies, M. Najdrowski, Eimeriosis in cattle: current understanding, *J. Vet. Med. B Infect. Dis. Public Health* 52 (10) (2005) 417–427, <https://doi.org/10.1111/j.1439-0450.2005.00894.x>.
- [7] MoARI, *Livestock and Animal Health Statistic*. Ministry of Agriculture Republic Indonesia, 2018.
- [8] *The Statistical Book on Livestock and Animal Health, Livestock and Animal Health Statistics, Directorate General of Livestock and Animal Health Service, Ministry of Agriculture, 2017* (ISBN: 978-979-628-034-6).
- [9] I.J. Nijman, M. Otsen, E.L. Verkaar, C. de Ruijter, E. Hanekamp, J.W. Ochieng, S. Shamshad, J.E. Rege, O. Hanotte, M.W. Barwegen, T. Sulawati, J.A. Lenstra, Hybridization of banteng (*Bos javanicus*) and zebu (*Bos indicus*) revealed by mitochondrial DNA, satellite DNA, AFLP and microsatellites, *Heredity* (Edinb) 90 (1) (2003) 10–16, <https://doi.org/10.1038/sj.hdy.6800174>.
- [10] V.M.A. Nurgiartiningih, A. Budiarto, Suyadi Kusmartono, Evaluation of performance in female Madura cattle in Madura Island, Indonesia, *Anim. Prod.* 18 (3) (2016) 125–130, <https://doi.org/10.20884/1.anprod.2016.18.3.573>.
- [11] MoARI, *Rencana Strategis Dan Kebijakan Pembangunan Peternakan Nasional Menuju Swasembada Daging*, Ministry of Agriculture Republic Indonesia, 2011.
- [12] S.M. Ananta, Suharno, A. Hidayat, M. Matsubayashi, Survey on gastrointestinal parasites and detection of *Cryptosporidium* spp. on cattle in West Java, Indonesia, *Asian Pac J Trop Med* 7 (3) (2014) 197–201, [https://doi.org/10.1016/S1995-7645\(14\)60020-1](https://doi.org/10.1016/S1995-7645(14)60020-1).
- [13] P.H. Hamid, Y.P. Kristianingrum, J. Prastowo, L.M.R. da Silva, Gastrointestinal parasites of cattle in Central Java, *Am. J. Anim. Vet. Sci.* 11 (2016) 119–124, <https://doi.org/10.3844/ajavsp.2016.119.124>.
- [14] F. Ekawasti, W. Nurcahyo, A.H. Wardhana, T. Shibahara, M. Tokoro, K. Sasai, M. Matsubayashi, Molecular characterization of highly pathogenic *Eimeria* species among beef cattle on Java Island, Indonesia, *Parasitol. Int.* 72 (2019) 101927, <https://doi.org/10.1016/j.parint.2019.101927>.
- [15] P.H. Hamid, Y.P. Kristianingrum, S. Prastowo, Bovine coccidiosis cases of beef and dairy cattle in Indonesia, *Vet. Parasitol. Reg. Stud. Reports* 17 (2019) 100298, <https://doi.org/10.1016/j.vprsr.2019.100298>.
- [16] P. Hastutiek, W.M. Yuniarti, M. Djaeri, N.D.R. Lastuti, E. Suprihati, L.T. Suwanti, Prevalence and diversity of gastrointestinal protozoa in Madura cattle at Bangkalan Regency, East Java, Indonesia, *Vet. World* 12 (2) (2019) 198–204, <https://doi.org/10.14202/vetworld.2019.198-204>.
- [17] D.H. Sawitri, A.H. Wardhana, E. Martindah, F. Ekawasti, D.A. Dewi, B.N. Utomo, T. Shibahara, M. Kusumoto, M. Tokoro, K. Sasai, M. Matsubayashi, Detections of gastrointestinal parasites, including *Giardia intestinalis* and *Cryptosporidium* spp., in cattle of Banten province, Indonesia, *J. Parasit. Dis.* 44 (1) (2020) 174–179, <https://doi.org/10.1007/s12639-019-01179-3>.
- [18] M. Matsubayashi, K. Takami, I. Kimata, T. Nakanishi, H. Tani, K. Sasai, E. Baba, Survey of *Cryptosporidium* spp. and *Giardia* spp. infections in various animals at a zoo in Japan, *J. Zoo Wildl. Med.* 36 (2) (2005) 331–335, <https://doi.org/10.1638/04-032.1>.
- [19] T. Fujino, T. Matsuo, M. Okada, T. Matsui, Detection of a small number of *Cryptosporidium parvum* oocysts by sugar flotation and sugar centrifugation methods, *J. Vet. Med. Sci.* 68 (11) (2006) 1191–1193, <https://doi.org/10.1292/jvms.68.1191>.
- [20] E.J.L. Soulsby, *Helminths, Arthropods and Protozoa of Domesticated Animals*, 7th ed., Baillière Tindall, London, 1982.
- [21] F. Kawahara, G. Zhang, C.N. Mingala, Y. Tamura, M. Koiwa, M. Onuma, T. Nunoya, Genetic analysis and development of species-specific PCR assays based on ITS-1 region of rRNA in bovine *Eimeria* parasites, *Vet. Parasitol.* 174 (1–2) (2010) 49–57, <https://doi.org/10.1016/j.vetpar.2010.08.001>.
- [22] L.B. Cruvinel, L.L. Ferreira, J.E. Nicaretta, L.F.M. Couto, D.M.B. Zapa, A.S. de Assis Cavalcante, L.M. Heller, R.D. de Melo Junior, L.M. de Aquino Gontijo, K. Suzuki, V. E. Soares, W.D.Z. Lopes, *Eimeria* spp. in naturally infected beef cattle: Dynamics of oocysts excretion, prevalence, and comparison between parasitological diagnostics, *Prev. Vet. Med.* 194 (2021) 105447, <https://doi.org/10.1016/j.prevetmed.2021.105447>.
- [23] B. Bangoura, H.C. Mundt, R. Schmäschke, B. Westphal, A. Dausgschies, Prevalence of *Eimeria bovis* and *Eimeria zuernii* in German cattle herds and factors influencing oocyst excretion, *Parasitol. Res.* 110 (2) (2012) 875–881, <https://doi.org/10.1007/s00436-011-2409-1>.
- [24] S.H. Lee, H.Y. Kim, H. Lee, J.W. Kim, Y.R. Lee, M.J. Chae, S.I. Oh, J.H. Kim, M. H. Rhee, O.D. Kwon, Y.K. Goo, T.H. Kim, P.J.L. Geraldino, D. Kwak, *Eimeria* species in cattle with diarrhoea in the Republic of Korea regarding age, season and nature of diarrhoea, *Vet. Rec.* 183 (16) (2018) 504, <https://doi.org/10.1136/vr.104600>.
- [25] L.B. Cruvinel, J.E. Nicaretta, T.S.A. Bastos, L.F.M. Couto, J.B.D. Santos, D.M. B. Zapa, A.S.A. Cavalcante, B.C. Cruz, D.G.L. Borges, F.A. Borges, V.E. Soares, W.D. Z. Lopes, *Eimeria* species in dairy and beef cattle of different ages in Goiás state, Brazil, *Rev. Bras. Parasitol.* 27 (2) (2018) 169–176, <https://doi.org/10.1590/s1984-296120180038>.
- [26] S.D. Ola-Fadunsin, M. Rabiu, K. Hussain, I.M. Sanda, I.A. Ganiyu, Epidemiological studies of *Eimeria* species of cattle in Ilorin, North-Central Nigeria, *Ann. Parasitol.* 66 (3) (2020) 373–384, <https://doi.org/10.17420/ap6603.276>.
- [27] I.A. Sutherland, D.M. Leathwick, Anthelmintic resistance in nematode parasites of cattle: a global issue? *Trends Parasitol.* 27 (2011) 176–181, <https://doi.org/10.1016/j.pt.2010.11.008>.

# Copro-parasitological examinations and molecular determination of *Eimeria* species in Madura cattle reared on Madura Island, Indonesia

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Makoto Matsubayashi, Sayaka Tsuchida, Atsushi Kobayashi, Tomoyuki Shibahara, Hiroshi Nakamura, Koichi Murata, Kazunari Ushida. "Molecular identification of two *Eimeria* species, *E. uekii* and *E. raichoi* as type B, in wild Japanese rock ptarmigans, *Lagopus muta japonica*", *International Journal for Parasitology: Parasites and Wildlife*, 2018  
Publication 2%
- 2 Makoto MATSUBAYASHI, Tomoyuki SHIBAHARA, Tomohide MATSUO, Toshimitsu HATABU, Junya YAMAGISHI, Kazumi SASAI, Takashi ISOBE. "Morphological and molecular identification of *Eimeria* spp. in breeding chicken farms of Japan", *Journal of Veterinary Medical Science*, 2020  
Publication 2%
- 3 Makoto MATSUBAYASHI, Toshimasa KITA, Tsuguto NARUSHIMA, Isao KIMATA, Hiroyuki TANI, Kazumi SASAI, Eiichiroh BABA. 1%

"Coprological Survey of Parasitic Infections in Pigs and Cattle in Slaughterhouse in Osaka, Japan", Journal of Veterinary Medical Science, 2009

Publication

---

4

[www.caas.gov.sg](http://www.caas.gov.sg)

Internet Source

1 %

---

5

Karin Takahashi, Makoto Matsubayashi, Yukio Ohashi, Jun Naohara et al. "Efficacy of ultraviolet light-emitting diodes (UV-LED) at four different peak wavelengths against *Cryptosporidium parvum* oocysts by inactivation assay using immunodeficient mice", *Parasitology International*, 2020

Publication

---

6

Leonardo Bueno Cruvinel, Lorena Lopes Ferreira, João Eduardo Nicaretta, Luiz Fellipe Monteiro Couto et al. "Eimeria spp. in naturally infected beef cattle: Dynamics of oocysts excretion, prevalence, and comparison between parasitological diagnostics", *Preventive Veterinary Medicine*, 2021

Publication

---

7

[www.scielo.br](http://www.scielo.br)

Internet Source

1 %

---

8

Ida, T.. "Cytokine-induced enhancement of calcium-dependent glutamate release from

1 %

astrocytes mediated by nitric oxide",  
Neuroscience Letters, 20080227

Publication

---

9	<a href="http://coek.info">coek.info</a> Internet Source	1 %
10	Yuu Matsuura, Makoto Matsubayashi, Satoko Nukata, Tomoyuki Shibahara et al. "Report of fatal mixed infection with <i>Cryptosporidium parvum</i> and <i>Giardia intestinalis</i> in neonatal calves", <i>Acta Parasitologica</i> , 2017 Publication	1 %
11	<a href="http://www.bioone.org">www.bioone.org</a> Internet Source	1 %
12	Ryusaku Yoshiuchi, Makoto Matsubayashi, Isao Kimata, Masaru Furuya, Hiroyuki Tani, Kazumi Sasai. "Survey and molecular characterization of <i>Cryptosporidium</i> and <i>Giardia</i> spp. in owned companion animal, dogs and cats, in Japan", <i>Veterinary Parasitology</i> , 2010 Publication	<1 %
13	<a href="http://www.e-journal.unair.ac.id">www.e-journal.unair.ac.id</a> Internet Source	<1 %
14	<a href="http://www.e-sciencecentral.org">www.e-sciencecentral.org</a> Internet Source	<1 %
15	Anna Forslid, Dan Christensson, Jan Dahl, Giulio Grandi, Jörg M.D. Enemark. "Bovine	<1 %



eimeriosis in Swedish calves: Epidemiology and insights into sampling procedures", *Veterinary Parasitology: Regional Studies and Reports*, 2015

Publication

---

16

[www.veterinaryworld.org](http://www.veterinaryworld.org)

Internet Source

<1 %

---

17

Abraham Anane, Osman Adamu Dufailu, Francis Addy. "Ascaridia galli and Heterakis gallinarum prevalence and genetic variance of *A. galli* in rural chicken from the Northern Region, Ghana", *Veterinary Parasitology: Regional Studies and Reports*, 2022

Publication

<1 %

---

18

Saroj Kumar, Rajat Garg, P.S. Banerjee, Hira Ram, K. Kundu, Sunil Kumar, M. Mandal. "Genetic diversity within ITS-1 region of *Eimeria* species infecting chickens of north India", *Infection, Genetics and Evolution*, 2015

Publication

<1 %

---

19

Shola David Ola-Fadunsin, Isau Aremu Ganiyu, Musa Rabi, Karimat Hussain et al. "Helminth infections of great concern among cattle in Nigeria: Insight to its prevalence, species diversity, patterns of infections and risk factors", *Veterinary World*, 2020

Publication

---

<1 %

20

ZHU, Wei, Jianbao DONG, Takeshi HAGA, Yoshitaka GOTO, and Masuo SUEYOSHI. "Rapid and Sensitive Detection of Bovine Coronavirus and Group A Bovine Rotavirus from Fecal Samples by Using One-Step Duplex RT-PCR Assay", Journal of Veterinary Medical Science, 2011.

Publication

&lt;1 %

21

[he01.tci-thaijo.org](http://he01.tci-thaijo.org)

Internet Source

&lt;1 %

22

[journal.ipb.ac.id](http://journal.ipb.ac.id)

Internet Source

&lt;1 %

23

[link.springer.com](http://link.springer.com)

Internet Source

&lt;1 %

24

[ugspace.ug.edu.gh](http://ugspace.ug.edu.gh)

Internet Source

&lt;1 %

25

[www.hindawi.com](http://www.hindawi.com)

Internet Source

&lt;1 %

26

Matsubayashi, M.. "Phylogenetic identification of Cystoisospora spp. from dogs, cats, and raccoon dogs in Japan", Veterinary Parasitology, 20110310

Publication

&lt;1 %

27

Terunori Koreeda, Tomo Kawakami, Ayako Okada, Yoshimasa Hirashima et al. "Pathogenic characteristics of a novel

&lt;1 %

intranuclear coccidia in Japanese black calves and its genetic identification as *Eimeria subspherica*", Parasitology Research, 2017

Publication

28

[lib.ugent.be](http://lib.ugent.be)

Internet Source

<1 %

29

[www.jircas.go.jp](http://www.jircas.go.jp)

Internet Source

<1 %

30

"Parasitic Protozoa of Farm Animals and Pets", Springer Science and Business Media LLC, 2018

Publication

<1 %

31

[bmcmgenomdata.biomedcentral.com](http://bmcmgenomdata.biomedcentral.com)

Internet Source

<1 %

32

Aaron S. Lucas, William S. Swecker, David S. Lindsay, Guillermo Scaglia, James P.S. Neel, Francois C. Elvinger, Anne M. Zajac. "A study of the level and dynamics of *Eimeria* populations in naturally infected, grazing beef cattle at various stages of production in the Mid-Atlantic USA", Veterinary Parasitology, 2014

Publication

<1 %

33

Maria Elena Morgoglione, Antonio Bosco, Maria Paola Maurelli, Leucio Camara Alves et al. "A 10-Year Surveillance of *Eimeria* spp. in

<1 %

# Cattle and Buffaloes in a Mediterranean Area", *Frontiers in Veterinary Science*, 2020

Publication

---

34

Sara M. Griffith, Jason Gigley, Jonathan Fox, Berit Bangoura. "Identification and characterization of *Eimeria* spp. in western north American Bison (*Bison bison*) herds and potential risk of cross-species transmission", *Veterinary Parasitology: Regional Studies and Reports*, 2021

Publication

---

<1 %

---

Exclude quotes      Off

Exclude matches      Off

Exclude bibliography      On

# Copro-parasitological examinations and molecular determination of Eimeria species in Madura cattle reared on Madura Island, Indonesia

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---