

Association of Parasite Density and Hematological Parameters of Plasmodium vivax- and Plasmodium vivax and Plasmodium falciparum- infected Patients Attending Merauke General Hospital, Papua, Indonesia

Submission date: 19-Jul-2021 11:42AM (UTC+0800)

Submission ID: 1621373897

File name: Hematological_Parameters_of_Plasmodium_vivax-_and_Plasmodiu.pdf (267.05K)

Word count: 5715

Character count: 33402

by Roy Bagus Kurniawan

40 Association of Parasite Density and Hematological Parameters of *Plasmodium vivax*- and *Plasmodium falciparum*-infected Patients Attending Merauke General Hospital, Papua, Indonesia

Roy Bagus Kurniawan¹, Puspa Wardhani^{2*}, Heny Arwati³, Aryati Aryati⁴, Trieva Verawaty Butarbutar⁵, Christophorus Oetama Adiatmaja⁵, Amarensi Milka Betaubun⁵, Nur Chamidah⁶

¹Medical Doctor Study Programme, Faculty of Medicine, Universitas Airlangga, Jl. Prof. Dr. Moestopo No. 47 Surabaya, Indonesia; ²Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Jl. Prof. Dr. Moestopo No. 47 Surabaya, Indonesia; ³Department of Medical Parasitology, Faculty of Medicine, Universitas Airlangga, Jl. Prof. Dr. Moestopo No. 47 Surabaya, Indonesia; ⁴Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Jl. Prof. Dr. Moestopo No. 47 Surabaya, Indonesia; ⁵Department of Clinical Pathology, Clinical Pathology Specialist Study Programme, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Jl. Prof. Dr. Moestopo No. 47 Surabaya, Indonesia; ⁶Department of Mathematics, Faculty of Sciences and Technology, Universitas Airlangga, Jl. Mulyorejo Surabaya, Indonesia

Abstract

45
Edited by: Ksenija Bogoeva-Kostovska
Citation: Kurniawan RB, Wardhani P, Arwati H, Aryati Aryati, Butarbutar TV, Adiatmaja CO, Betaubun AM, Chamidah N. Association of Parasite Density and Hematological Parameters of *Plasmodium vivax*- and *Plasmodium falciparum*-infected Patients Attending Merauke General Hospital, Papua, Indonesia. Open Access Maced J Med Sci. 2020 Oct 10; 8(B):825-831. https://doi.org/10.3889/oamjms.2020.4881
Keywords: Parasite Density, Hematology, Malaria, *Plasmodium vivax*, *Plasmodium falciparum*
*Correspondence: Puspa Wardhani, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Jl. Prof. Dr. Moestopo No. 47 Surabaya, Indonesia. Phone: +628123176937. E-mail: puspa_pk@yahoo.co.id
Received: 24-May-2020
Revised: 24-Sep-2020
Accepted: 28-Sep-2020
Copyright: © 2020 Roy Bagus Kurniawan, Puspa Wardhani, Heny Arwati, Aryati Aryati, Trieva Verawaty Butarbutar, Christophorus Oetama Adiatmaja, Amarensi Milka Betaubun, Nur Chamidah
Funding: The funding of this research was from the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia with contract number: 544/LN3.141/LT/2019.
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: *Plasmodium falciparum* and *Plasmodium vivax* are frequent causes of malaria. Although they are blood parasites, their biological characteristics are dissimilar, and their species-related consequences on hematological parameters have not been widely investigated. They might be valuable to distinguish both species infection, notably for an endemic region with limited diagnostic resources.

AIM: This study aimed to know the species-specific effect on hematological parameters and its correlation to the parasite density in *P. vivax*- and *P. falciparum*-infected patients attending Merauke General Hospital, Papua, Indonesia.

MATERIALS AND METHODS: Malaria patients confirmed by blood film microscopy from January 1 to July 31, 2019, were recruited, and their hematological parameters were measured using Sysmex XN-1000 instrument. All obtained data were analyzed statistically.

RESULTS: From 100 malaria-positive patients, 87 patients, consisting of 57 *P. vivax* and 30 *P. falciparum* patients, met criteria. Anemia and parasite density >50,000 parasites/ μ L were significantly higher in *P. falciparum* than *P. vivax* patients ($p < 0.05$) though hemoglobin concentration and parasite density were insignificantly different. Interestingly, basophil count was significantly higher in *P. falciparum* compared to *P. vivax* patients ($p = 0.04$). The eosinophil count was significantly higher in *P. vivax* ($p = 0.01$) than *P. falciparum* patients and indicated a significant positive correlation ($p = 0.04$, $r = +0.28$) with the parasite density.

CONCLUSION: There were significant differences between basophil and eosinophil count between *P. vivax* and *P. falciparum* infections. Eosinophil count showed a significant positive correlation with parasite density.

Introduction

Malaria remains one of the deadly infectious diseases that continue as serious public health problems worldwide, particularly in tropical countries. Around 219 million peoples suffered from malaria, and 445 thousand of them died [1]. Indonesia, the country with a high malaria burden, has committed to eradicating malaria in 2030 [2]. Instead, the latest report shows the increased cases of more than 100,000 malaria cases occurred between 2017 and 2016 in Indonesia [1].

Dynamic alterations of hematological parameters are common findings in malaria patients. They might attribute to some changes in significant blood cell types, including erythrocyte, leukocyte, and platelet [3], [4], [5], [6].

Hemoglobin, erythrocyte, total leukocyte, neutrophil, eosinophil, monocyte, and lymphocyte counts seemed significantly lower contrasted with non-malaria patients [3], [4]. Meanwhile, erythrocyte indices, neutrophil-lymphocyte ratio (NLR), and monocyte-lymphocyte ratio (MLR) were higher opposed to non-malaria patients [5], [7]. Leukocyte, neutrophil, monocyte, lymphocyte, erythrocyte, and platelet counts among different levels of parasitemia levels also marked to differ significantly [5].

Plasmodium falciparum and *Plasmodium vivax* are frequent causes of malaria worldwide, including in Indonesia [1]. Although both species are blood parasites, their biological characteristics, such as life cycle, virulence mechanisms, disease manifestation, and management, are dissimilar [8]. Their species-related consequences

on hematological parameters have not been widely investigated. For instance, *P. falciparum* might result in more than 10% parasitemia, whereas *P. vivax* could only reach 2% [8]. Higher parasitemia would undoubtedly result in poor prognosis [9], [10]. However, severe *P. vivax* malaria cases have also been recorded [6], [11]. That evidence inferred that prompt species identification is necessary since both infections require different pharmacotherapeutic approaches. Unfortunately, the microscopic blood smear, the gold standard for diagnosing malaria, needs more extended time, and particular expertise to obtain the result [12]. Hematological parameter changes associated with parasite species-specific effects appear potential to provide a reference in specifically suspecting malaria species, even for estimating the total parasite burden of the disease in the endemic area, due to its availability in the routine examination. The present study aimed to investigate the comparison of hematological parameters between *P. vivax*- and *P. falciparum*-infected patients referred to Merauke General Hospital. The study also tried to correlate the parasite density and hematological parameters in each investigated group. Better understanding in this field could help physicians suspect parasite species in highly suspected malaria patients and predict patient prognosis so that the quick and specific management could be initiated, particularly in the endemic area with limited diagnostic facilities.

27 Materials and Methods

Samples collection

A cross-sectional study was conducted at Merauke General Hospital, Papua, Indonesia, from January to July 2019. Blood samples were collected from patients by vena punctured after signing the informed consent. Blood, then, was transferred to EDTA tubes for parasite and hematology examination. Patients with microscopically confirmed malaria of *P. vivax* or *P. falciparum* were engaged, whereas those with malaria mixed infection, immunocompromised, and comorbid infections were excluded.

Malaria parasite examination

Microscopic examinations were carried out on prepared Giemsa-stained thin and thick blood films to identify malaria species and quantify the parasite density based on the latest WHO guidelines [13]. The formula employed to quantify parasite density is written below.

$$\text{Parasite density} = \frac{\sum \text{counted parasites}}{200 \text{ observed leukocytes}} \times \text{leukocyte count}$$

The examination was accomplished by two trained and experienced field microscopists independently in Merauke clinical pathology laboratory. All obtained data were reassessed by the clinical pathologist at Merauke General Hospital.

Hematology assay

The hematology examination was performed using Sysmex XN-1000 (Sysmex Corporation, Kobe, Japan). The examination included hemoglobin concentration, erythrocyte count, mean corpuscular volume, mean corpuscular hemoglobin (MCH), MCH concentration (MCHC), leukocyte and differential counts, NLR, MLR, and platelet count. All laboratory procedures were conducted based on appropriate standard operating procedures. Quality control of the instrument was completed per manufacturer standards. As parasite examination, all obtained data were reassessed by the clinical pathologist at Merauke General Hospital.

Data analysis

Normality distribution of data was analyzed using the Kolmogorov–Smirnov test if samples exceed 50 and the Shapiro–Wilk if samples were 50 or lesser. Our comparison hypothesis was proved by performing the independent t-test for normally distributed data. Otherwise, the Mann–Whitney U-test would be used. The proportion of data was analyzed using the Z test. The Pearson correlation test was employed to test our correlation hypothesis if data were normally distributed, and the Spearman correlation test in case they did not. The finding was deemed significant at $p < 0.05$ or $|Z| > 1.65$.

Ethical declaration

This study's ethical clearance was issued by the Medical Research Ethics Committee, Faculty of Medicine, Universitas Airlangga (No.169/EC/KEPK/FKUA/2019).

Results

General characteristics

A total of 100 positive malaria samples were collected. However, six patients with mixed infection (*P. vivax*/*P. falciparum*) and seven patients with incomplete data were excluded. The study eventually involved 57 (65.51%) cases of *P. vivax* and 30 (34.48%) cases of *P. falciparum* infections. As many as, 64.37% of involved patients were male and

became the predominant contributor of malaria cases, which comprised 64.91% of *P. vivax* and 63.33% of *P. falciparum* cases. The mean age of *P. vivax* and *P. falciparum* groups was 29.87 ± 16.03 and 29.08 ± 15.33 , respectively.

11 Comparison of parasite density between *P. vivax* and *P. falciparum* infection

The median of *P. vivax* and *P. falciparum* parasite density was 5902.04 (379.84–135,700.00) parasites/ μL and 9061.00 (54.48–1109.675) parasites/ μL , respectively. The median between both species was insignificantly different ($p > 0.05$). Furthermore, parasite density data were grouped into three categories, as shown in Table 1, and noted for their proportion in each malaria species to be compared statistically. The proportion of the parasite density $>50,000$ parasites/ μL was significantly higher in *P. falciparum* compared to *P. vivax* ($|Z| = 1.87$, $p = 0.03$).

Table 1: Parasite density between *Plasmodium vivax* and *P. falciparum*

Parasite density (parasites/ μL)	Number (%)		Z-score (95% CI)	p-value
	<i>P. vivax</i>	<i>P. falciparum</i>		
Category 1 ($<10,000$)	37 (64.91)	16 (53.33)	1.05	0.15
Category 2 (10,000–50,000)	15 (26.32)	7 (23.33)	0.30	0.38
Category 3 ($>50,000$)	5 (8.77)	7 (23.33)	-1.87	0.03*
Total	57 (100.00)	30 (100.00)	N/A	N/A

*Significant value at $|Z| > 1.65$, equal to $p < 0.05$ (one tailed) N/A: Not applicable. *P. falciparum*: *Plasmodium falciparum*; *P. vivax*: *Plasmodium vivax*.

6 Comparison of hematological parameters between *P. vivax* and *P. falciparum* infection

This study found that both eosinophil and basophil count were significantly different between investigated groups. *P. vivax* group's basophil count was substantially lower compared to *P. falciparum* group ($p = 0.04$). Still, *P. vivax* group's eosinophil count was significantly higher contrasted to *P. falciparum* group ($p = 0.01$) (Table 2). Nonetheless, hemoglobin concentration, erythrocyte indices, erythrocyte count, total leukocyte count, neutrophil count, lymphocyte count, monocyte count, platelet count, NLR, and MLR between both groups were considered insignificantly different ($p > 0.05$) (Table 2).

Table 2: Comparison of hematological parameters between *P. vivax* and *P. falciparum* infections

Parameters	Mean \pm SD/Median (Min.–Max.)		p-value
	<i>P. vivax</i>	<i>P. falciparum</i>	
Hemoglobin (g/dL)	12.60 (4.30–16.60)	11.95 \pm 2.78	0.29
Erythrocyte ($\times 10^6/\mu\text{L}$)	4.81 \pm 0.76	4.56 \pm 0.97	0.19 ^b
MCV (fL)	74.93 \pm 8.88	75.82 \pm 8.26	0.65 ^b
MCH (pg/cell)	26.70 (14.50–32.90)	25.50 (21.30–46.10)	0.82
MCHC (g/dL)	34.60 (25.80–38.40)	34.65 (31.20–46.30)	0.72
Platelet ($\times 10^3/\mu\text{L}$)	97.00 (24.00–481.00)	92.00 (13.00–285.00)	0.19
Leukocyte ($\times 10^3/\mu\text{L}$)	7.10 (3.14–17.03)	6.48 (2.19–19.66)	0.92
Eosinophil ($\times 10^3/\mu\text{L}$)	0.10 (0.00–3.48)	0.05 (0.00–0.60)	0.01*
Basophil ($\times 10^3/\mu\text{L}$)	0.02 (0.00–0.11)	0.03 (0.00–0.20)	0.04*
Neutrophil ($\times 10^3/\mu\text{L}$)	5.04 (2.12–14.00)	5.69 \pm 2.83	0.71
Lymphocyte ($\times 10^3/\mu\text{L}$)	1.05 (0.18–3.13)	1.15 (0.33–4.05)	0.20
Monocyte ($\times 10^3/\mu\text{L}$)	0.62 \pm 0.40	0.57 (0.16–2.20)	0.63
NLR	4.95 (1.40–18.27)	4.75 (0.17–24.18)	0.28
MLR	0.55 \pm 0.29	0.47 (0.21–1.13)	0.66

P. falciparum: *Plasmodium falciparum*; *P. vivax*: *Plasmodium vivax*; MCV: Mean corpuscular volume; MCH: Mean corpuscular hemoglobin; NLR: Neutrophil-lymphocyte ratio; MLR: Monocyte-lymphocyte ratio.

Furthermore, the occurrence of anemia was significantly different between both species. Anemia conducted to occur in *P. falciparum* group contrasted to *P. vivax* group ($|Z| = 2.48$, $p = 0.01$) (Table 3). Nevertheless, the occurrence of leukocytosis, leukocytopenia, thrombocytosis, and thrombocytopenia was different insignificantly.

Table 3: Major hematological abnormalities in *P. vivax* and *P. falciparum* infection

Major hematological abnormalities	Number (%)		Z-score (95% CI)	p-value
	<i>P. vivax</i>	<i>P. falciparum</i>		
Anemia (Hb ≤ 11 g/dL)	12 (21.05)	14 (46.67)	-2.48	0.01*
Leukocytosis ($>10,000$ cells/ μL)	7 (12.3)	6 (20.00)	-0.96	0.17
Leukopenia (<4000 cells/ μL)	4 (7.00)	1 (3.00)	0.70	0.24
Thrombocytosis ($>450,000$ cells/ μL)	1 (1.75)	0 (0.00)	0.73	0.23
Thrombocytopenia ($<150,000$ cells/ μL)	41 (71.93)	24 (80.00)	-1.82	0.21

Hb: Hemoglobin; *significant value at $|Z| > 1.65$, equal to $p < 0.05$ (one tailed). *P. falciparum*: *Plasmodium falciparum*; *P. vivax*: *Plasmodium vivax*.

Correlation between parasite density and hematological parameters

The parasite density and eosinophil count in *P. vivax* group indicated a significant positive correlation with $r = +0.28$ ($p = 0.04$) (Table 2). Hemoglobin concentration, erythrocyte count and indices, platelet count, lymphocyte count, total leukocyte count, basophil count, neutrophil count, monocyte count, NLR, and MLR showed insignificant correlation. However, in *P. falciparum* group, neither parasite density nor hematological parameters seemed to correlate significantly.

Discussion

Hematological abnormalities become a prevailing hallmark during malaria and favor to be variable among studies. These diversities appear to be multifactorial in etiology, depending on the parasitic agent, host responses, and environment. *P. vivax* and *P. falciparum* are prevalent causes of malaria in Indonesia [1], while both species possess different characteristics, pathogenesis mechanisms, and therapeutic management [8]. Although the life cycle, biological aspects, and clinical manifestations between both parasites have been studied extensively, the parasite density might differ among endemic areas, and hematological differences resulting from different malaria species seem to lack evidence. This research was conducted based on issues that those species distinctiveness might result in parasite density and hematological parameter abnormalities, which probably own clinical significance in the endemic area. To the best of our knowledge, this was the first study comparing various hematological parameters of *P. vivax* and *P. falciparum* patients and correlating those parameters with the malaria parasite density in Indonesia, particularly in Papua.

The primary cause of malaria in the Merauke was *P. vivax* followed by *P. falciparum*. Six patients with mixed infection (*P. vivax* and *P. falciparum*) were unexpectedly observed. This result contrasts with the study in Jayapura, Papua, which was predominantly caused by *P. falciparum* [14]. Those males were likely to get malaria than females, and insignificant patients' age difference satisfied previous findings [3], [5] and seemed to associate with different occupational exposures.

High parasitemia is likely to occur in *P. falciparum* compared to *P. vivax* due to the parasite indiscrimination to infect all erythrocyte maturation stages, generate multiple infections, and perform rapid schizogony [8]. This study showed that the proportion of parasite density >50,000 parasites/ μ L in *P. falciparum* was significantly higher as opposed to *P. vivax* infection ($|Z| = 1.87$, $p = 0.03$) (Table 1). This study's highest parasite density was 135,700 parasites/ μ L for *P. vivax* and 1109.675 parasites/ μ L for *P. falciparum*. This finding of parasite propensity concurs the previous study reported by Barber *et al.* [10]. Nevertheless, the median of parasite density between both groups was not statistically different. This finding might happen since the evidence of *P. falciparum*-infected erythrocyte performing cytoadherence have been reported [15], [16], resulting in parasite sequestration and preventing circulating parasite to represent the total parasite biomass. Notwithstanding, the present study still confirms the tendency and ability of *P. falciparum* to result in very high parasitemia.

As parasite density relates to the severity of malaria infection [9], [10], it is necessary to correlate the parasite density in each species with common examined hematological parameters. Although, none variables posed a significant correlation, except for eosinophil count in *P. vivax* infection. The parasite density showed significant positive correlation with eosinophil count ($r = 0.28$, $p = 0.04$) (Table 4). These results appear contrary to some studies that indicated several significant correlations. An inverse correlation between the parasite densities with the lymphocyte count in *P. falciparum*

infection and positive correlations of the parasite density with neutrophil and platelet counts in *P. vivax* and *P. falciparum* infection were reported by studies in Ghana and South Ethiopia [3], [7]. A mathematical model study also designated good relationships of the parasite density with neutrophil, lymphocyte, monocyte, and eosinophil count in *P. falciparum* infection [17]. Despite those discordances, insignificant correlations of parasite density, in both species, with hemoglobin concentration, leukocyte count, and monocyte count, in this study, agree with the previous studies [3], [7]. Interestingly, the positive correlation of eosinophil count and parasite density in *P. vivax* group seems becoming a new finding referring to those studies.

Interplays of parasite and hematological parameters still require further elucidation. Intriguing reasons that might underlie insignificant correlations influence malaria transmission, sequestration, and systemic inflammatory reactions [15], [18], [19]. Pro-inflammatory cytokines generated during *P. falciparum* infection promote endothelial activation, leading to leukocyte adherence and sequestration, microvascular thrombosis, and endothelial injury (increased platelet usage) [10], [19], [20], [21], [22]. They might explain why the significant positive correlation of eosinophil and the parasite density, in this study, only occurred in *P. vivax* infection since severe inflammatory reactions during *P. vivax* infection are unlikely to occur [8], [20]. However, the reasons for eosinophil roles during malaria are still controversial. They have been associated with an antiparasitic activity, disease recovery, hypersensitivity, and total parasite burdens [23], [24], [25], [26].

Anemia is frequently observed and associated with malaria severity. However, its exact mechanism seems intricate and poorly understood. As the parasites invade erythrocytes, hemolysis will be occurring and lead to anemia. Nonetheless, the loss of uninfected erythrocytes (UE) appears more influential in the anemia pathogenesis. The previous studies inferred that either mechanical or immune-mediated splenic clearance was principal underlying causes of massive UE loss [27], [28]. Bone marrow suppression was also responsible for the anemia occurrence [29], [30]. The anemia (hemoglobin <11 g/dL) in both *P. vivax* and *P. falciparum* was noted in our study. It favored in *P. falciparum* compared to *P. vivax* ($|Z| = 2.48$, $p = 0.01$) (Table 3). Erythrocyte profiles in *P. falciparum*, however, did not significantly differ from *P. vivax* (Table 2). This result argues the earlier study that found significant differences in erythrocyte count and indices, even if it strengthens the finding of the insignificant hemoglobin concentration difference in both species [5], [11]. These results possibly happen as different research settings might possess different local parasitic strains and malaria transmission intensity. Evidence of *P. falciparum*-infected erythrocyte sequestration [15], [16] makes measured circulating erythrocyte profiles unable to figure out actual states.

Table 4: Correlation between parasite density and hematological parameters in *P. vivax* and *P. falciparum* infection

Parameters	<i>P. vivax</i>		<i>P. falciparum</i>	
	Correlation coefficient	p-value*	Correlation coefficient	p-value*
Parasite density (parasite/ μ L)				
Hemoglobin	-0.20	0.15	0.13	0.50
Erythrocyte	-0.13	0.33	0.07	0.71
MCV	-0.11	0.43	0.14	0.47
MCH	-0.16	0.25	0.18	0.35
MCHC	-0.15	0.26	0.24	0.20
Platelet	-0.13	0.34	-0.08	0.68
Leukocyte	0.14	0.32	0.16	0.40
Eosinophil	0.28	0.04 ^b	-0.16	0.39
Basophil	0.20	0.14	-0.11	0.56
Neutrophil	0.07	0.59	0.26	0.17
Lymphocyte	-0.04	0.76	-0.17	0.37
Monocyte	0.13	0.34	-0.13	0.51
NLR	0.13	0.34	0.29	0.12
MLR	0.18	0.18	-0.05	0.78

*p-value by Spearman's rho correlation test; ^bsignificant value at $p < 0.05$. *P. falciparum*: *Plasmodium falciparum*, *P. vivax*: *Plasmodium vivax*, MCV: Mean corpuscular volume, MCH: Mean corpuscular hemoglobin, NLR: Neutrophil-lymphocyte ratio, MLR: Monocyte-lymphocyte ratio.

The leukocyte deems as the significant element of the immune system that deals with the plasmodia invasion. It prefers to change, meeting on needs and agent properties dynamically. Theoretically, massive schizont ruptures, particularly in *P. falciparum*, will be followed by the extensive exposure with intraerythrocytic materials and parasite antigens, driving to profound immune responses [31]. Our finding has met the theory that leukocytosis (leukocyte >10,000 cells/ μ L) favored occurring in *P. falciparum* rather than *P. vivax* although it was statistically insignificant. That leukocytopenia (leukocyte <4000 cells/ μ L) was found more in *P. vivax* (Table 3) which was in contrast with a report in Thailand during 1998–1999 [32]. Furthermore, leukocyte profiles, such as total leukocyte count, neutrophil count, lymphocyte count, monocyte count, NLR, and MLR, noted insignificant differences between groups. Nevertheless, the significant difference of eosinophil count ($p = 0.01$) and insignificant differences of total leukocyte, lymphocyte, and monocyte counts still followed an investigation in Thailand [5]. That basophil count marked to be significantly different ($p = 0.04$) between observed groups may become a new finding referring to that study. Unimportant differences of NLR and MLR between both species were found, although the previous result showed significant differences compared to non-malaria patients [4].

Inconsistent findings of leukocyte profiles between investigated groups are not unlikely. Microvascular endothelial activation, induced by increased pro-inflammatory cytokines, such as tumor necrosis factor- α and interleukin (IL)-1 β , leads to overexpression of intercellular adhesion molecule-1 and secretion of IL-8, a potent neutrophil attractant [31], [33]. They promote leukocytes marginalization and sequestration [21], resulting in discrepancies between circulating and actual total leukocyte count. That the basophil count appeared higher in *P. falciparum* than in *P. vivax* patients may support the finding of *P. falciparum* ability to produce histamine-releasing factors, belonging to translationally controlled tumor protein (TCTP), suggesting important roles in regulating basophil and eosinophil activity [26], [34]. A study in Senegal reported a higher concentration of *P. falciparum* TCTP in patients with severe malaria than mild malaria and healthy control [35]. That study also identified the enhanced basophil reactivity to stimuli associated with *P. falciparum* TCTP *in vivo* and with malaria severity [36]. TCTP might also reflect total parasite burdens (including sequestered parasite) rather than peripheral parasitemia [26]. This mechanism explains why both parasite densities in our research do not exhibit significant differences regarding the discovery of a considerable difference in eosinophil and basophil count in investigated species. However, the eosinophil count was higher in *P. vivax* than *P. falciparum* might indicate other different *P. vivax* pathogenesis processes that need further elucidation.

Another common complication of malaria is thrombocytopenia. Its pathogenesis has not been certainly

ascertained. Platelet-mediated erythrocyte clumping, von Willebrand factor adherence, sequestration, and immune-mediated destruction are responsible for thrombocytopenic episodes [22], [35], [37], [38], [39]. That thrombocytopenia (<150,000 cells/ μ L) was still a common finding was confirmed in this study (>70% of patients in both groups), and its proportion of both species was roughly similar (Table 3), as reported in another study [4]. A thrombocytosis case was only discovered in *P. vivax* group. Nonetheless, the comparison of platelet count between investigated species was not significantly different, in contrast with the previous investigation in Thailand [5].

This study's limitation was no assessment of primary confounding factors affecting hematological parameters such as the engaged patient's genetic susceptibilities and micronutrient state. That the parasite examination only relied on microscopy also makes submicroscopic malaria unrecognized and could lead to underestimation of parasite density impact on hematological parameters.

Conclusion

Diversities of hematological abnormalities during malaria are common findings. Anemia, leukocyte count abnormalities, and thrombocytopenia are predominant and favor to be in *P. falciparum* infection. Several parameters do not differ significantly between *P. vivax* and *P. falciparum*. However, basophil count is significantly higher in *P. falciparum* infection, whereas eosinophil count is significantly higher in *P. vivax* infection. These findings suggest that basophil and eosinophil count possesses clinical values and might provide a clue, together with clinical presentations, to suspect *P. vivax* or *P. falciparum* infection in the endemic area. The significant positive correlation between the parasite density and eosinophil count in *P. vivax* may be the potential to predict the total parasite burden. These findings offer the opportunity to conduct prompt and specific management, particularly in regions with limited diagnostic facilities. Furthermore, since this report is a pioneer study in Merauke, Papua, researches concerning this topic should be more encouraged.

Acknowledgment

The authors would like to acknowledge Simitabmas Program by KEMENRISTEKDIKTI RI (Ministry of Research, Technology, and Higher Education of the Republic of Indonesia) to support this study (contract number: 544/UN3.14/LT/2019). We

are also grateful to Merauke General Hospital, sample collectors, study participants, and all who gave their hands so that this study could be done.

References

- World Health Organization. Malaria Report, 2018. Geneva: World Health Organization; 2018.
- Kementerian Kesehatan Republik Indonesia. Situasi Terkini Perkembangan Program Pengendalian Malaria di Indonesia Tahun 2018. Jakarta: Kementerian Kesehatan Republik Indonesia; 2018.
- Awoke N, Arota A. Profiles of hematological parameters in *Plasmodium falciparum* and *Plasmodium vivax* malaria patients attending Tercha general hospital, Dawuro Zone, South Ethiopia. *Infect Drug Resist.* 2019;12:521-7. <https://doi.org/10.2147/idr.s184489>
PMid:30881057
- Kotepui M, Phunphuech B, Phiwklam N, Chupeerach C, Duangmano S. Effect of malarial infection on hematological parameters in population near Thailand-Myanmar border. *Malar J.* 2014;13(1):218-24. <https://doi.org/10.1186/1475-2875-13-218>
PMid:24898891
- Kotepui M, Piwkham D, Phunphuech B, Phiwklam N, Chupeerach C, Duangmano S. Effects of malaria parasite density on blood cell parameters. *PLoS One.* 2015;10(3):e0121057. <https://doi.org/10.1371/journal.pone.0121057>
PMid:25807235
- Arévalo-Herrera M, Lopez-Perez M, Medina L, Moreno A, Gutierrez JB, Herrera S. Clinical profile of *Plasmodium falciparum* and *Plasmodium vivax* infections in low and unstable malaria transmission settings of Colombia. *Malar J.* 2015;14(1):154-64. <https://doi.org/10.1186/s12936-015-0678-3>
- Antwi-Baffour S, Kyeremeh R, Buabeng D, Adjei JK, Aryeh C, Kpentey G, et al. Correlation of malaria parasitemia with peripheral blood monocyte to lymphocyte ratio as indicator of susceptibility to severe malaria in Ghanaian children. *Malar J.* 2018;17(1):419-27. <https://doi.org/10.1186/s12936-018-2569-x>
PMid:30419923
- Garcia LS. *Diagnostic Medical Parasitology*. 6th ed. Washington, DC: ASM Press; 2016.
- Argy N, Kendjo E, Augé-Courtoi C, Cojean S, Clain J, Houzé P. Influence of host factors and parasite biomass on the severity of imported *Plasmodium falciparum* malaria. *PLoS One.* 2017;12(4):e0175328. <https://doi.org/10.1371/journal.pone.0175328>
PMid:28410415
- Barber BE, William T, Grigg MJ, Parameswaran U, Piera KA, Price RN, et al. Parasite biomass-related inflammation, endothelial activation, microvascular dysfunction, and disease severity in *Vivax* malaria. *PLoS Pathog.* 2015;11(1):e1004558. <https://doi.org/10.1371/journal.ppat.1004558>
PMid:25569250
- Abdallah TM, Abdeen MT, Ahmed IS, Hamdan HZ, Magzoub M, Adam I. Severe *Plasmodium falciparum* and *Plasmodium vivax* malaria among adults at Kassala hospital, Eastern Sudan. *Malar J.* 2013;12(1):148-54. <https://doi.org/10.1186/1475-2875-12-148>
- Tangpukdee N, Duangdee C, Wilairatana P, Krudsood S. Malaria diagnosis: A brief review. *Korean J Parasitol.* 2009;47(2):93-102. <https://doi.org/10.3347/kjp.2009.47.2.93>
PMid:19488414
- World Health Organization. *Basic Malaria Microscopy*, Part 1. 2nd ed. Geneva: World Health Organization; 2010.
- Sorontou Y, Asih PB, Wanandi SI, Ramelan W, Syafruddin D. Malaria in Jayapura District, Papua Province, Indonesia, and resistance to sulfadoxine-pyrimethamine. *Med J Indones.* 2007;16(1):32-8. <https://doi.org/10.13181/mji.v16i1.254>
- David PH, Hommel M, Miller LH, Udeinya IJ, Oligino LD. Parasite sequestration in *Plasmodium falciparum* malaria: Spleen and antibody modulation of cytoadherence of infected erythrocytes. *Proc Natl Acad Sci U S A.* 1983;80(16):5075-9. <https://doi.org/10.1073/pnas.80.16.5075>
PMid:6348780
- Safeukui I, Correas JC, Brousse V, Hirt D, Deplaine G, Mulé S, et al. Retention of *Plasmodium falciparum* ring-infected erythrocytes in the slow, open microcirculation of the human spleen. *Blood.* 2008;112(6):2520-8. <https://doi.org/10.1182/blood-2008-03-146779>
PMid:18579796
- Evelyn ME, Ezeiruaku FC, Ukaji DC. Experiential relationship between malaria parasite density and some haematological parameters in malaria infected male subjects in port Harcourt, Nigeria. *Glob J Health Sci.* 2012;4(4):139-48. <https://doi.org/10.5539/gjhs.v4n4p139>
PMid:22980350
- Ademolue TW, Aniweh Y, Kusi KA, Awandare GA. Patterns of inflammatory responses and parasite tolerance vary with malaria transmission intensity. *Malar J.* 2017;16(1):145-157. <https://doi.org/10.1186/s12936-017-1796-x>
PMid:28399920
- Coban C, Lee MS, Ishii KJ. Tissue-specific immunopathology during malaria infection. *Nat Rev Immunol.* 2018;18(4):266-78. <https://doi.org/10.1038/nri.2017.138>
PMid:29332936
- Day NP, Hien TT, Schollaardt T, Loc PP, Chuong LV, Chau TT, et al. The prognostic and pathophysiologic role of pro- and anti-inflammatory cytokines in severe malaria. *J Infect.* 1999;180(4):1288-97. <https://doi.org/10.1086/315016>
PMid:10479160
- Hearn J, Rayment N, Landon DN, Katz DR, de Souza JB. Immunopathology of cerebral malaria: Morphological evidence of parasite sequestration in murine brain microvasculature. *Infect Immun.* 2000;68(9):5364-76. <https://doi.org/10.1128/iai.68.9.5364-5376.2000>
PMid:10948166
- Graham SM, Chen J, Chung DW, Barker KR, Conroy AL, Hawkes MT, et al. Endothelial activation, hemostasis, and thrombosis biomarkers in Ugandan children with severe malaria participating in a clinical trial. *Malar J.* 2016;15(1):56-64. <https://doi.org/10.1186/s12936-016-1106-z>
- Waters LS, Taverne J, Tai P, Spry CJ, Targett GA, Playfair JH. Killing of *Plasmodium falciparum* by eosinophil secretory products. *Infect Immun.* 1987;55(4):877-81. <https://doi.org/10.1128/iai.55.4.877-881.1987>
PMid:3549562
- Kurtzhals JA, Reimert CM, Tette E, Dunyo SK, Koram KA, Akanmori BD, et al. Increased eosinophil activity in acute *Plasmodium falciparum* infection-association with cerebral malaria. *Clin Exp Immunol.* 1998;112(2):303-7. <https://doi.org/10.1046/j.1365-2249.1998.00586.x>
PMid:9649195
- Camacho LH, Wilairatana P, Weiss G, Mercader MA, Brittenham GM, Loareesuwan S, et al. The eosinophilic response and haematological recovery after treatment for *Plasmodium falciparum* malaria. *Trop Med Int Health.* 1999;4(7):471-5. <https://doi.org/10.1046/j.1365-3156.1999.00426.x>

- PMid:10470337
26. Mecheri S. Contribution of allergic inflammatory response to the pathogenesis of malaria disease. *Biochim Biophys Acta Mol Basis Dis.* 2012;1822(1):49-56. <https://doi.org/10.1016/j.bbadis.2011.02.005>
 27. Huang S, Amaladoss A, Liu M, Chen H, Zhang R, Preiser PR, et al. *In vivo* splenic clearance correlates with *in vitro* deformability of red blood cells from *Plasmodium yoelii*-infected mice. *Infect Immun.* 2014;82(6):2532-41. <https://doi.org/10.1128/iai.01525-13>
PMid:24686065
 28. Safeukui I, Gomez ND, Adelani AA, Burte F, Afolabi NK, Akondy R, et al. Malaria induces anemia through CD8+ T cell-dependent parasite clearance and erythrocyte removal in the spleen. *mBio.* 2015;6(1):e02493-14. <https://doi.org/10.1128/mbio.02493-14>
PMid:25604792
 29. Casals-Pascual C, Kai O, Cheung JO, Williams S, Lowe B, Nyanoti M, et al. Suppression of erythropoiesis in malarial anemia is associated with hemozoin *in vitro* and *in vivo*. *Blood.* 2006;108(8):2569-77. <https://doi.org/10.1182/blood-2006-05-018697>
PMid:16804108
 30. Panichakul T, Payuhakrit W, Panburana P, Wongborisuth C, Hongeng S, Udomsangpetch R. Suppression of erythroid development *in vitro* by *Plasmodium vivax*. *Malar J.* 2012;11(1):173-80. <https://doi.org/10.1186/1475-2875-11-173>
 31. Abbas A, Lichtman A, Pillai S. Cellular, and Molecular Immunology. 9th ed. Philadelphia, PA: Elsevier; 2018.
 32. McKenzie FE, Prudhomme WA, Magill JA, Forney JA, Permpnich B, Lucas C, et al. White blood cell counts and malaria. *J Infect.* 2005;19(2):323-30. <https://doi.org/10.1086/431152>
PMid:15962228
 33. Chakravorty SJ, Carret C, Nash GB, Ivens A, Szestak T, Craig AG. Altered phenotype and gene transcription in endothelial cells, induced by *Plasmodium falciparum*-infected red blood cells: Pathogenic or protective? *Int J Parasitol.* 2007;37(8-9):975-87. <https://doi.org/10.1016/j.ijpara.2007.02.006>
PMid:17383656
 34. MacDonald SM, Bhisutthibhan J, Shapiro TA, Rogerson SJ, Taylor TE, Tembo M, et al. Immune mimicry in malaria: *Plasmodium falciparum* secretes a functional histamine-releasing factor homolog *in vitro* and *in vivo*. *Proc Natl Acad Sci U S A.* 2001;98(19):10829-32. <https://doi.org/10.1073/pnas.201191498>
PMid:11535839
 35. Coelho HC, Lopes SC, Pimentel JP, Nogueira PA, Costa FT, Siqueira AM, et al. Thrombocytopenia in *Plasmodium vivax* malaria is related to platelets phagocytosis. *PLoS One.* 2013;8(5):e63410. <https://doi.org/10.1371/journal.pone.0063410>
PMid:23723981
 36. Pelleau S, Diop S, Badiane MD, Vitte J, Beguin P, Nato F, et al. Enhanced basophil reactivities during severe malaria and their relationship with the *Plasmodium falciparum* histamine-releasing factor translationally controlled tumor protein. *Infect Immun.* 2012;80(8):2963-70. <https://doi.org/10.1128/iai.00072-12>
PMid:22753372
 37. Pain A, Ferguson DJ, Kai O, Urban BC, Lowe B, Marsh K, et al. Platelet-mediated clumping of *Plasmodium falciparum*-infected erythrocytes is a common adhesive phenotype and is associated with severe malaria. *Proc Natl Acad Sci U S A.* 2001;98(4):1805-10. <https://doi.org/10.1073/pnas.98.4.1805>
PMid:11172032
 38. Grau GE, Mackenzie CD, Carr RA, Redard M, Pizzolato G, Allasia C, et al. Platelet accumulation in brain microvessels in fatal pediatric cerebral malaria. *J Infect.* 2003;187(3):461-6. <https://doi.org/10.1086/367960>
PMid:12552430
 39. O'Sullivan JM, Preston RJ, O'Regan N, O'Donnell JS. Emerging roles for hemostatic dysfunction in malaria pathogenesis. *Blood.* 2016;127(19):2281-8. <https://doi.org/10.1182/blood-2015-11-636464>
PMid:26851291

Association of Parasite Density and Hematological Parameters of Plasmodium vivax- and Plasmodium vivax and Plasmodium falciparum-infected Patients Attending Merauke General Hospital, Papua, Indonesia

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

17%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Esther Felicita Tambayong, Nengah Kuning Atmadjaya, Nyoman Golden, Ketut Wiargitha, Gede Bagus Mahadewa Tjokorda. "Facial Injury Severity Scale Score as a Predictor of Length of Stay for Maxillofacial Fracture at Sanglah General Hospital, Denpasar, Bali, Indonesia", Open Access Macedonian Journal of Medical Sciences, 2020

Publication

2%
- 2 Ferry Efendi, Dwida Rizki Pradiptasiwi, Ilya Krisnana, Tiyas Kusumaningrum et al. "Factors associated with complete immunizations coverage among Indonesian children aged 12–23 months", Children and Youth Services Review, 2020

Publication

1%
- 3 Kotepui, Manas, Bhukdee Phunphuech, Nuoil Phiwklam, Chaowanee Chupeerach, and Suwit Duangmano. "Effect of malarial infection on

1%

haematological parameters in population near Thailand-Myanmar border", *Malaria Journal*, 2014.

Publication

4

Nefsu Awoke, Amsalu Arota. "

1 %

Profiles of hematological parameters in *Plasmodium falciparum* and *Plasmodium vivax* malaria patients attending Tercha General Hospital, Dawuro Zone, South Ethiopia

", *Infection and Drug Resistance*, 2019

Publication

5

repository.usu.ac.id

Internet Source

1 %

6

Sandeep Puri, Omesh Goyal, Subhadra Prashar. "Hepatic dysfunction in falciparum and vivax malaria in Northern India", *Journal of Gastrointestinal Infections*, 2015

1 %

Publication

7

"Comparison of Clinical Profile between *P. vivax*", 'Hindawi Limited', 2014

Internet Source

1 %

8

Wang, Qinghui, Zhenjun Zhao, Xuexing Zhang, Xuelian Li, Min Zhu, Peipei Li, Zhaoqing Yang, Ying Wang, Guiyun Yan, Hong Shang, Yaming Cao, Qi Fan, and Liwang Cui. "Naturally Acquired Antibody Responses to Plasmodium

1 %

vivax and Plasmodium falciparum Merozoite Surface Protein 1 (MSP1) C-Terminal 19 kDa Domains in an Area of Unstable Malaria Transmission in Southeast Asia", PLoS ONE, 2016.

Publication

9

Purnami Nyilo, Manyakori Serafika Permoni Putri. "The Association of Reactive Oxygen Species Levels on Noise Induced Hearing Loss of High Risk Workers in Dr. Soetomo General Hospital Surabaya, Indonesia", Indian Journal of Otolaryngology and Head & Neck Surgery, 2018

Publication

1 %

10

Yohei Kobayashi, Yoshitomo Saita, Hirofumi Nishio, Hiroshi Ikeda et al. "Leukocyte concentration and composition in platelet-rich plasma (PRP) influences the growth factor and protease concentrations", Journal of Orthopaedic Science, 2016

Publication

<1 %

11

indianpediatrics.net

Internet Source

<1 %

12

journals.athmsi.org

Internet Source

<1 %

13

oro.open.ac.uk

Internet Source

<1 %

14

Cho Naing, Maxine A. Whittaker, Victor Nyunt Wai, Joon Wah Mak. "Is Plasmodium vivax Malaria a Severe Malaria?: A Systematic Review and Meta-Analysis", PLoS Neglected Tropical Diseases, 2014

Publication

<1 %

15

Karina Keller Marques da Costa Flaiban. "Hematologic Values of Free-ranging Cebus cay and Cebus nigritus in Southern Brazil", International Journal of Primatology, 10/2008

Publication

<1 %

16

Rodrigues-da-Silva, Rodrigo Nunes, Josué da Costa Lima-Junior, Bruna de Paula Fonseca e Fonseca, Paulo Renato Zuquim Antas, Arlete Baldez, Fabio Luiz Storer, Fátima Santos, Dalma Maria Banic, and Joseli de Oliveira-Ferreira. "Alterations in cytokines and haematological parameters during the acute and convalescent phases of Plasmodium falciparum and Plasmodium vivax infections", Memórias do Instituto Oswaldo Cruz, 2014.

Publication

<1 %

17

D. Mohanty. "Fibrinolysis, inhibitors of blood coagulation, and monocyte derived coagulant activity in acute malaria", American Journal of Hematology, 01/1997

Publication

<1 %

18

Internet Source

<1 %

19

mindamas-journals.com

Internet Source

<1 %

20

Uswatun Hasanah, Ida Leida Maria, Nurhaedar Jafar, Andi Hardianti, Anwar Mallongi, Aminuddin Syam. "Water, Sanitation Dan Hygiene Analysis, and Individual Factors for Stunting among Children Under Two Years in Ambon", Open Access Macedonian Journal of Medical Sciences, 2020

Publication

<1 %

21

journals.plos.org

Internet Source

<1 %

22

pure.tue.nl

Internet Source

<1 %

23

worldwidescience.org

Internet Source

<1 %

24

Beata Stepniak, Sergi Papiol, Christian Hammer, Anna Ramin, Sarah Everts, Lena Hennig, Martin Begemann, Hannelore Ehrenreich. "Accumulated environmental risk determining age at schizophrenia onset: a deep phenotyping-based study", The Lancet Psychiatry, 2014

Publication

<1 %

25 Ngurah, A.A.G.. "H-supermagic labelings of graphs", Discrete Mathematics, 20100428 <1 %
Publication

26 Pramono, C.. "Surgical Technique for Achieving Implant Parallelism and Measurement of the Discrepancy in Panoramic Radiograph", Journal of Oral and Maxillofacial Surgery, 200605 <1 %
Publication

27 www.pjps.pk <1 %
Internet Source

28 Rut Navarro-Martínez, Julio Fernández-Garrido, Cristina Buigues, Elena Torralba-Martínez et al. "Brain-derived neurotrophic factor correlates with functional and cognitive impairment in non-disabled older individuals", Experimental Gerontology, 2015 <1 %
Publication

29 test.dovepress.com <1 %
Internet Source

30 www.balimedicaljournal.org <1 %
Internet Source

31 Barber, Bridget E., Timothy William, Matthew J. Grigg, Uma Parameswaran, Kim A. Piera, Ric N. Price, Tsin W. Yeo, and Nicholas M. Anstey. "Parasite Biomass-Related Inflammation, Endothelial Activation, Microvascular

Dysfunction and Disease Severity in Vivax Malaria", PLoS Pathogens, 2015.

Publication

32

C. Han, Y. Lin, G. Shan, Z. Zhang, X. Sun, Z. Wang, C. Wei, Y. Deng, L. Zhang, L. Bu, D. Shao, H. Wang. "Plasma Concentration of Malaria Parasite-Derived Macrophage Migration Inhibitory Factor in Uncomplicated Malaria Patients Correlates with Parasitemia and Disease Severity", Clinical and Vaccine Immunology, 2010

Publication

<1 %

33

D. E. Ghannam. "Inhaled aminoglycosides in cancer patients with ventilator-associated Gram-negative bacterial pneumonia: safety and feasibility in the era of escalating drug resistance", European Journal of Clinical Microbiology & Infectious Diseases, 03/2009

Publication

<1 %

34

Linda Wati, Ririe Fachrina Malisie, Juliandi Harahap. "Pediatric Observational Priority Score and Early Warning Scoring System to Predict Admission Status in Pediatric Patients in Haji Adam Malik General Hospital", Open Access Macedonian Journal of Medical Sciences, 2021

Publication

<1 %

35 Rick M Fairhurst. "Modulation of malaria virulence by determinants of Plasmodium falciparum erythrocyte membrane protein-1 display", Current Opinion in Hematology, 05/2006
Publication <1 %

36 core.ac.uk
Internet Source <1 %

37 escholarship.org
Internet Source <1 %

38 springerplus.springeropen.com
Internet Source <1 %

39 www.quintessence-publishing.com
Internet Source <1 %

40 www.tandfonline.com
Internet Source <1 %

41 Bernard N. Kanoi, Thomas G. Egwang. "Sex differences in concentrations of HMGB1 and numbers of pigmented monocytes in infants and young children with malaria", Parasitology International, 2021
Publication <1 %

42 F. YILDIZ ZEYREK. "Parasite density and serum cytokine levels in Plasmodium vivax malaria in Turkey", Parasite Immunology, 5/2006
Publication <1 %

43

John Woodford, Ashley Gillman, Peter Jenvey, Jennie Roberts et al. "Positron emission tomography and magnetic resonance imaging in experimental human malaria to identify organ-specific changes in morphology and glucose metabolism: A prospective cohort study", PLOS Medicine, 2021

Publication

<1 %

44

Riccio, Evelyn KP, Paulo RR Totino, Lilian R Pratt-Riccio, Vitor Ennes-Vidal, Irene S Soares, Maurício Rodrigues, José de Souza, Cláudio Daniel-Ribeiro, and Maria Ferreira-da-Cruz. "Cellular and humoral immune responses against the Plasmodium vivax MSP-119 malaria vaccine candidate in individuals living in an endemic area in north-eastern Amazon region of Brazil", Malaria Journal, 2013.

Publication

<1 %

45

Sylvia Jiero, Ayodhia Pitaloka Pasaribu. "Haematological profile of children with malaria in Sorong, West Papua, Indonesia", Malaria Journal, 2021

Publication

<1 %

46

Tienrat Tangchaikeeree, Piamsiri Sawaisorn, Sangdao Somsri, Duangporn Polpanich et al. "Enhanced Sensitivity for Detection of Plasmodium falciparum gametocytes by

<1 %

magnetic nanoparticles combined with enzyme substrate system", Talanta, 2017

Publication

47

Touchchapol Mataradchakul, Chairat Uthaipibull, Francois Nosten, Joel Vega-Rodriguez, Marcelo Jacobs-Lorena, Usa Lek-Uthai. "Plasmodium vivax rhomboid-like protease 1 gene diversity in Thailand", Experimental Parasitology, 2017

Publication

<1 %

48

U. O. Ozojiofor, O. O. Bankole, I. O. Adedeji, K. C. Onuh. "Changes in Liver Function Enzymes in Plasmodium falciparum Infected Malaria Patients in Ajeromi General Hospital, Lagos, Nigeria", Asian Journal of Research in Biochemistry, 2020

Publication

<1 %

49

annalsofian.org

Internet Source

<1 %

50

link.springer.com

Internet Source

<1 %

51

surgicalneurologyint.com

Internet Source

<1 %

52

www.malariajournal.com

Internet Source

<1 %

53

www.tevak.org

Internet Source

<1 %

54

Biomedical Aspects of Histamine, 2011.

Publication

<1 %

55

R. Singh, D. P. Singh, R. Gupta, D. Savargaonkar, O. P. Singh, N. Nanda, R. M. Bhatt, N. Valecha. "Comparison of three PCR-based assays for the non-invasive diagnosis of malaria: detection of Plasmodium parasites in blood and saliva", European Journal of Clinical Microbiology & Infectious Diseases, 2014

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Association of Parasite Density and Hematological Parameters of Plasmodium vivax- and Plasmodium vivax and Plasmodium falciparum-infected Patients Attending Merauke General Hospital, Papua, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
