

In Vitro Antiviral Activity of Morin Compound against Dengue Virus Type 1 in Vero Cells

by Anisa Maharani

Submission date: 19-Jul-2021 11:43AM (UTC+0800)

Submission ID: 1621375193

File name: of_Morin_Compound_against_Dengue_Virus_Type_1_in_Vero_Cells.pdf (954.21K)

Word count: 2564

Character count: 13315

In Vitro Antiviral Activity of Morin Compound against Dengue Virus Type 1 in Vero Cells

Anisa Maharani¹, Teguh H. Sucipto^{2,*}, Harsasi Setyawati³, Yovilianda M. Untoro⁴, Novia F. Sholihah¹, Siti Churrotin², Ilham H. Amarullah², Puspa Wardhani², Aryati², Soegeng Soegijanto²

¹Undergraduate Student, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Campus C UNAIR, Jl. Mulyorejo, Surabaya, East Java, 60115, Indonesia

²Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Campus C UNAIR, Jl. Mulyorejo, Surabaya, East Java, 60115, Indonesia

³Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Campus C UNAIR, Jl. Mulyorejo, Surabaya, East Java, 60115, Indonesia

⁴Master Student, Department of Chemistry, Faculty of Mathematic and Natural Science, Institut Teknologi Bandung, Jl. Ganesha No. 10 Bandung, West Java 40132 Indonesia

*Corresponding Author: Teguh H. Sucipto, Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Campus C UNAIR, Jl. Mulyorejo, Surabaya, Jawa Timur, 60115 Indonesia. Phone: +62-8990484365. E-mail address: teguhharisucipto@staf.unair.ac.id

ABSTRACT

Introduction: Dengue Haemorrhagic Fever (DHF) is an irresistible ailment brought about by dengue infection. Recently, the medicine for dengue fever cannot be disseminated in various countries because of the price was high. One characteristic of a person infected with the virus was thrombocyte decreasing. In previous research it was known that guava can increase the number of thrombocyte patients of dengue fever. Guava was trusted can increase the number of thrombocyte because it contains a group of tannin and flavonoids as anti-bacterial. Some flavonoids in guava include morin, quercetin, and guajavarin.

Objective: Purpose in this study, was to decide the impact of morin compound to the replication of dengue infection.

Methods: This study used to test the inhibition of the virus is by Viral ToxGlo assay to know the value of 50% inhibition of morin compound on dengue virus replication (IC₅₀) and cytotoxicity to determine the toxicity level of the morin compound against body cell (CC₅₀).

Results: This study of IC₅₀ morin for dengue compounds is 9.42 µg/ml and the result of CC₅₀ in Vero cell is 12.46 µg/ml.

Conclusion: In summary, there was an investigation show that morin compound have harmful properties in the body.

Keywords: DENV-1, morin, inhibitor activity, cytotoxicity

Correspondence:

Teguh H. Sucipto
Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Campus C UNAIR, Jl. Mulyorejo, Surabaya, Jawa Timur, 60115 Indonesia.
Phone: +62-8990484365.

E-mail address: teguhharisucipto@staf.unair.ac.id

INTRODUCTION

Dengue shows up in two forms, classic dengue fever and acute dengue fever. In acute dengue fever, Dengue Haemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS) which may cause stomach bleeding, haemorrhage and circulatory collapse. On the off chance that this extreme structure isn't treated with brief and legitimate administration it will prompt the deadly cases.¹ Dengue is a common arbovirus infection transmitted to humans by *Aedes aegypti*.² Dengue Haemorrhagic Fever (DHF) brought about by dengue infections is as yet a noteworthy issue in tropical nations. More than 2.5 billion individuals are at present in danger of DENV disease, with 100 million individuals assessed to be contaminated with DENV every year.³ The main symptoms of Dengue Haemorrhagic Fever are fever, muscle and joint pain, which usually worsens after the first two days and then followed by a decreasing of the amount of thrombocytes. In the most recent decades, dengue has developed as a prime wellbeing worry in tropical and subtropical and until now there has been no effective and affordable drugs circulating.

Morin (2-(2,4-dihydroxyphenyl)-3,5,7-trihydroxychromen-4-one) is a derivative of flavones found in few Moraceae families. Morin has various biological activities, such as antioxidants, antiinflammation, and antivirals for equid herpesvirus 1.⁴ Physical properties of morin include the colour of yellow, water soluble, acid, pyridine and trichloroacetic, alcohol, and acetone solutions, so they can be categorized as water

soluble dyes.⁵ Morin is a polyphenol compound extensively studied in its pharmacological activity to treat human health problems, with few side effects.⁶ Morin was reported anti-CDV (canine distemper virus) activity at the time 0 hour, IC₅₀ 34.02 µg/ml and CC₅₀ 195.90 µg/ml. In the previous research derived of flavonoid, 5-hydroxy-7-methoxy-6-methylflavanone was inhibited DENV-2 infectivity with EC₅₀ 12.31 ± 1.64 µM. Baicalein, fisetin and quercetagenin was reported for powerful inhibition of Chikungunya virus infection, with IC₅₀ of 1.891 µg/ml, 8.444 µg/ml, and 13.85 µg/ml, and with CC₅₀ of 356.3 µg/ml, 194.4 µg/ml, and 226.7 µg/ml, respectively.⁷ Thus, rutin and quercetin on DENV-2 in C₆/36 cells inhibited virus replication with IC₅₀ of 362.68 µg/ml and 500 µg/ml, separately. No cytotoxicity was seen with any of these compounds at up to 1000 µg/ml.⁸

In this research, we have several characterizations that were by doing an activity test in *in vitro* and also the cytotoxicity against body cells. All of the characterizations were performed to determine the inhibitory activity of dengue virus using morin compounds.

MATERIALS AND METHODS

Materials

The chemical reagents used in this research were Morin compound, Ethanol 96% (Merck, Germany), aquades, Dimethyl Sulfoxide (DMSO) (Merck, Germany), Dengue virus serotype 1 (DENV-1) Isolate from Surabaya with Genbank Accession AB915377, Minimum Essential Eagle

Medium (Sigma-Aldrich, Germany), Vero cell (African green monkey kidney), Viral Toxglo Assay (Promega, USA), and Cell Proliferation Reagent WST-1 by Roche Applied Science.

METHODS

Antiviral Activity Assay

Vero Cells with concentration 1.10^6 cells/10 ml were seeded into a 96 well plate and incubated plates containing cells at 37°C in a humidified CO₂ incubator for at least 4 hours (and up to 24 hours) to facilitate attachment and allow cells to recover from seeding stresses. One hundred microliters of dengue virus with concentration 4×10^5 FFU/ml stock was combined with various concentrations of morin (50 µg/ml, 25 µg/ml, 12.5 µg/ml, 6.25 µg/ml, 3.13 µg/ml, 0.78 µg/ml). After one hour of exposure to the dengue virus at room temperature, 100 µl of the morin treated virus was added to Vero cells in individual wells. Cells were infected for 1 hour. Unabsorbed virus was removed and replaced with MEM. After 48 hours post-infection added 100µl of ToxGlo (Promega, Madison, WI). 100 µl of ATP Detection Reagent was added to each well of a 96-well plate (25 µl to each well of a 384-well plate) and

wait at least 10 minutes prior to measuring luminescence. Calculate IC₅₀ values by plotting net RLU (relative luminescence units) values (subtracting the average of blank wells) versus compound concentration. The IC₅₀ value is the compound concentration that caused 50% inhibitory effect on the viral replication or viral cytopathic effect (CPE).

Cytotoxicity Assay

Cytotoxicity used WST-1 cell proliferation reagent by Roche Applied Science, Mannheim, Germany. The assay is very sensitive: it can detect 1,000 cells/well of a 96-well plate reader. Vero cells (1×10^5 cells/ml), 50 µl of serial dilution compound, and a total of 10 µl of Cell Proliferation Reagent was added to each well of a 96-well plate and incubated under 5% CO₂ at 37°C for 1 hour. The plate was read at 450 nm (main filter) and 655 nm (reference filter) using an iMark™ Microplate Absorbance Reader.

RESULTS AND DISCUSSION

To decide the antiviral impact of morin, DENV was treated with morin. The IC₅₀ was analysed from the linear regression equation of the percent infectivity. We found that IC₅₀ of morin was 9.42 µg/ml (Figure 1).




Figure 1. Antiviral activity of Morin for Vero Cells at Varying Concentrations. IC₅₀ (maximal inhibitory concentration) is measure of the effectiveness of a substance in inhibiting a specific DENV-1. In this curve was used 6 concentration; 50 µg/ml, 25 µg/ml, 12.5 µg/ml, 6.25 µg/ml, 3.13 µg/ml, and 0.78 µg/ml. IC₅₀ was calculated from regression linier curve; $y = -7.5429x + 121.07$

Morin has been used in several studies, one of them on the development of sensitized coloured solar cells (*dye sensitized solar cell*). The expansion of morin in previous study as a sensitizer in DSSC material did not cause changes in crystal composition on TiO₂ anatase or TiO₂ synthesis results.⁵ As of late, there have been numerous examinations on dengue antivirals from various organic and crystalline chemicals in TiO₂ anatase and TiO₂ from synthesis. Based on previous research, omega 3/6 fatty acids can be used as potential inhibitors of DENV envelope protein fusion process.⁹ Other studies of antivirals for

dengue have also been performed using curcumin and Pentagamavunon-0.¹⁰ From the results of this study known the concentration of curcumin and Pentagamavunon-0 are good against body cells. An anti-dengue study of organic matter has also been carried out, the results showed that the extract from *Psidium guajava* (Guava), *Euphorbia hirta* (Patikan kerbau), *Piper betel* L. (Sirih), *Carica papaya* (Papaya), *Curcuma longa* L. (Turmeric), *Phyllanthus niruri* L. (Meniran), *Andrographis paniculata* (Sambiloto), and *Cymbopogon citratus* (lemongrass) were considered to have *in vitro* activity dependent on three criteria, for example cell viability >50%, IC₅₀ ≤25 µg/ml and the selectivity index >3.15-17.¹¹ To ensure that morin was not dangerous to the cell, the half cytotoxic concentration (CC₅₀) was measured with the WST-1 test. The CC₅₀ was analysed from the linear regression equation of the percent viability. We found that the CC₅₀ of morin was 12.46 µg/ml (Figure 2).




Figure 2. Cytotoxicity of Morin for Vero Cells at Varying Concentrations. CC_{50} is cytotoxicity level of morin compound to cause death to 50% of Vero cells. CC_{50} was calculated from regression linier curve; $y = 0.2037x + 47.462$ with the axis (x) is concentration of compound and ordinate (y) is %viability.

The results show that from the value IC_{50} and CC_{50} , the CC_{50} value was found to increase with an increasing concentration of the test compound. Morin for Vero cells was 12.46 µg/ml, with an R^2 value of 0.9889. When compared with a previous study, morin is more toxic than plant extracts with an IC_{50} value of 35.7 µg/ml. and CC_{50} value 250 µg/ml.¹¹ The good IC_{50} value has a concentration close to 0, whereas for a good CC_{50} should be above 100 µg/ml so that the results of this study can be seen from the values of IC_{50} and CC_{50} indicating that morin is toxic to Vero cells.

In the previous research, bioflavonoids derived are of intrigue as a result of their natural and health advantages. Baicalein inhibited DENV-2 replication in Vero cells with IC_{50} was 1.55 µg/ml and exhibited cytotoxic effects with CC_{50} was 109 µg/ml¹² and showed potent anti-JEV activity with $IC_{50} = 14.28$ µg/ml when it was acquainted to the Vero cells.¹³ The IC_{50} values for the aqueous extract of *Scutellaria baicalensis* on Vero cells following DENV adsorption ranged from 86.59 to 95.19 µg/ml with CC_{50} of the extract was 912.6 µg/ml. The cytotoxicity suggests that the extract is in general non-cytotoxic and could be well-endured by Vero cells.¹⁴

The mechanisms of how morin exerts it is anti-DENV effects are not known. A possible mechanism of action of intracellular anti-DENV activity is binding of morin to the viral RNA, interaction with DENV structural and non-structural protein, and also prophylactic activity of morin against DENV replication, is about the aggregation of the compound in the cells during the treatment. However, avoidance of infection adsorption to the cells is another plausibility that may prompt hindrance of infection passage to the cells.¹³

CONCLUSION

In conclusion, from screening antiviral activity of morin did not potential against DENV-1 because it was shown to be medium toxic to Vero cells. Further study is expected to affirm antiviral activity through a larvicidal test.

DATA AVAILABILITY

The data used to support the findings of this study are available from the corresponding author upon request.

CONFLICT OF INTEREST STATEMENT

We declare that we have no conflicts of interest.

ACKNOWLEDGEMENT

This study was funded by Program Hibah Riset Mandat Universitas Airlangga (HRMUA) 2021 grant from Universitas Airlangga. Much thanks to Institute of Tropical Disease Universitas Airlangga for research temporary position opportunity in the Dengue Laboratory; and for the Center of Excellence (COE) program by the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia.

REFERENCES

- Gubler DJ. Dengue and Dengue Hemorrhagic Fever. *Clin Microbiol Rev.* 1998; 11(3): 480-96
- Simmons CP, Farrar JJ, Nguyen VC, Wills B. Dengue. *N Engl J Med.* 2012; 366(15): 1423-1432
- Halstead SB. Dengue. *Lancet.* 2007;370(9599): 1644-52
- Gravina HD, Tafuri NF, Junior SJ, et al. In Vitro Assessment of The Antiviral Potential of Trans-cinnamic Acid, Quercetin and Morin against Equid Herpesvirus 1. *Res Vet Sci.* 2011; 91(3): e158-62
- Rahayu D, Bagitaningtyas A, Hidayat A, Suci AP. Pengembangan Sel Surya Berpewarna Tersensitisasi (Dye Sensitized Solar Cell) dengan Senyawa Morin dari Kayu Nangka (*Artocarpus heterophyllus L.*). *Pelita-Jurnal Penelitian Mahasiswa UNY.* 2011; VI (1): 1-11
- Gopal JV. Morin Hydrate: Botanical Origin, Pharmacological Activity and Its Application: a mini review. *Phcog J.* 2013; 5(3): 123-126
- Lani R, Hassandarvish P, Shu MH, et al. Antiviral activity of selected flavonoids against Chikungunya virus. *Antiviral Res.* 2016; 133: 50-61
- Silva ARA, Morais SM, Marques MM, et al. Antiviral activities of extracts and phenolic components of two *Spondias* species against dengue virus. *J Venom Anim Toxins Incl Trop Dis.* 2011; 17(4): 406-413
- Idrus S. Omega 3 Fatty Acid and 6 as Potential Inhibitors of Dengue Virus Fusion. *Majalah Biam.* 2015; 11(2): 49-56
- Marbawati D, Umniyati DR. Effects of Curcumin and Pentagamavunon-0 Against Dengue-2 Virus Infection in Vero Cells; an in Vitro Study. *Procedia Environ Sci.* 2015; 23: 215-221
- Saptawati L, Febrinasari RP, Yudhani RD, et al. In vitro study of eight Indonesian plants extracts as anti-Dengue virus. *Health Sci J Indo.* 2015; 8(1): 12-18
- Zandi K, Teoh BT, Sam SS, et al. Novel antiviral activity of baicalein against dengue virus. *BMC Compl Alternative Med.* 2012; 12(214): 1-9

13. Johari J, Kianmehr A, Mustafa MR, Abubakar S, Zandi K. Antiviral activity of baicalein and quercetin against the Japanese Encephalitis Virus. *Int J Mol Sci.* 2012; 13(12): 16785-95
14. Zandi K, Lim TH, Rahim NA, et al. Extract of *Scutellaria baicalensis* inhibits dengue virus replication. *BMC Compl Alternative Med.* 2013; 13(91): 1-10

In Vitro Antiviral Activity of Morin Compound against Dengue Virus Type 1 in Vero Cells

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Xiaomin Kang, De Jin, Yuqing Zhang, Rongrong Zhou, Yuehong Zhang, Fengmei Lian. "Systematic Elucidation of the Mechanism of Sappan Lignum in the Treatment of Diabetic Peripheral Neuropathy Based on Network Pharmacology", Evidence-Based Complementary and Alternative Medicine, 2021
Publication 1%
- 2 N.S. Aminah, K.N.W. Tun, A.N. Kristanti, H.T. Aung, Y. Takaya, M.I. Choudhary. "Chemical constituents and their biological activities from Taunggyi (Shan state) medicinal plants", Heliyon, 2021
Publication 1%
- 3 Lusida, Maria. "High rate of seronegative HCV infection in HIV-positive patients", Biomedical Reports, 2013.
Publication 1%
- 4 www.mdpi.com
Internet Source 1%

5	Zachary A. Gurard-Levin, Cheng Liu, Andreas Jekle, Ruchika Jaisinghani et al. "Evaluation of SARS-CoV-2 3C-like protease inhibitors using self-assembled monolayer desorption ionization mass spectrometry", Antiviral Research, 2020 Publication	1 %
6	digitalcommons.montclair.edu Internet Source	1 %
7	www.tandfonline.com Internet Source	1 %
8	Nahdlotul Fauziyah, Fermanto, Muhamad Nur Ghoyatul Amin, Mochammad Amin Alamsjah, Sri Subekti, Dwi Yuli Pujiastuti. "Postharvest processing of Sargassum duplicatum for tea products", Journal of Applied Phycology, 2021 Publication	1 %
9	Pawan K. Agrawal, Chandan Agrawal, Gerald Blunden. "Quercetin: Antiviral Significance and Possible COVID-19 Integrative Considerations", Natural Product Communications, 2020 Publication	1 %
10	www.imr.gov.my Internet Source	1 %
11	www.lib.kobe-u.ac.jp Internet Source	1 %

12 A. L. Ivansyah, M.A. Martoprawiro, Buchari. "Computational modeling of inclusion complex of r/s-omeprazole with β -cyclodextrin using oniom2 method", Journal of Physics: Conference Series, 2017 1 %
Publication

13 Sahu, Kapendra, Mohammad Shaharyar, and Anees A. Siddiqui. "Effect of Morin on pharmacokinetics of Piracetam in rats, in vitro enzyme kinetics and metabolic stability assay using rapid UPLC method : Phamacokinetics, metabolic stability, in vitro enzyme kinetics", Drug Testing and Analysis, 2012. 1 %
Publication

14 mdpi.com 1 %
Internet Source

15 Emma Riese, Viggo Kann. "Computer Science Majors' Experiences of Their Distance Education Caused by the COVID-19 Pandemic", 2021 IEEE Global Engineering Education Conference (EDUCON), 2021 1 %
Publication

16 www.biomedcentral.com 1 %
Internet Source

17 www.neliti.com 1 %
Internet Source

www.thieme-connect.com

18

Internet Source

1 %

19

jmscr.igmpublication.org
Internet Source

<1 %

20

www.omicsonline.org
Internet Source

<1 %

21

www.google.com
Internet Source

<1 %

22

www.scitechnol.com
Internet Source

<1 %

23

Élida Cleyse Gomes da Mata, Caroline Barbosa Farias Mourão, Marisa Rangel, Elisabeth Ferroni Schwartz. "Antiviral activity of animal venom peptides and related compounds", Journal of Venomous Animals and Toxins including Tropical Diseases, 2017
Publication

<1 %

24

Heny Arwati, Subagyo Yotopranoto, Etik Ainun Rohmah, Din Syafruddin. "Submicroscopic malaria cases play role in local transmission in Trenggalek district, East Java Province, Indonesia", Malaria Journal, 2018
Publication

<1 %

25

Riss, Terry, Erika Hawkins, Martha O,ÄôBrien, Andrew Niles, and Richard Moravec. "Homogeneous Multiwell Assays for

<1 %

Measuring Cell Viability, Cytotoxicity, and Apoptosis", Drug Discovery Series, 2006.

Publication

26

idpjournal.biomedcentral.com

Internet Source

<1 %

27

Adeel Masood Butt, Mohd Cairul Iqbal Mohd Amin, Haliza Katas, Narong Sarisuta, Wasu Witoonsaridsilp, Ruthairat Benjakul. "

Characterization of Pluronic F127 and D--Tocopheryl Polyethylene Glycol 1000

Succinate Mixed Micelles as Nanocarriers for Targeted Anticancer-Drug Delivery ", Journal of Nanomaterials, 2012

Publication

<1 %

28

Rosmalena Rosmalena, Berna Elya, Beti E. Dewi, Fithriyah Fithriyah et al. "The Antiviral Effect of Indonesian Medicinal Plant Extracts Against Dengue Virus In Vitro and In Silico", Pathogens, 2019

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

In Vitro Antiviral Activity of Morin Compound against Dengue Virus Type 1 in Vero Cells

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
