




# Jurnal KIMIA RISET

Volume 2 No. 2, Desember 2017; ISSN: 2528-0414; e-ISSN: 2528-0422



**Departemen Kimia  
Fakultas Sains dan Teknologi  
Universitas Airlangga  
<http://e-journal.unair.ac.id/index.php/JKR>**



Home > Archives > Vol 2, No 2 (2017)

## VOL 2, NO 2 (2017)

DESEMBER

### TABLE OF CONTENTS

#### ARTICLES

##### REPARASI DAN KARAKTERISASI ALFA-Fe<sub>2</sub>O<sub>3</sub> /ZEOLIT Y UNTUK REAKSI PERENKAHAN ASAM PALMITAT

doi 10.20473/jkr.v2i2.6166

Abstract views = 334 times | PDF views = 768 times  
Abdulloh Abdulloh, Purkan Purkan, Nizar Hardiansyah

PDF  
69-76

##### Membran Blend Kitosan/Poli Vinil Alkohol (PVA): Pengaruh Komposisi material blend, pH, dan Konsentrasi bahan Pengikat Silang

doi 10.20473/jkr.v2i2.6195

Abstract views = 426 times | PDF views = 3951 times  
Bambang Piluharto, A Sjaifullah, I Rahmawati, Erix Nurharianto

PDF  
77-85

##### EFEKTIVITAS KOMBINASI EKSTRAK DAUN SIRIH MERAH (Piper Crocatum) DAN EKSTRAK BIJI ALPUKAT (Persea americana) DALAM MENGHAMBAT PERTUMBUHAN Candida albicans

doi 10.20473/jkr.v2i2.6196

Abstract views = 1121 times | PDF views = 2699 times  
Vini Angraini, Masfufatun Masfufatun

PDF  
86-92

##### Celery herb essential oil in the formulation of anti-dandruff hair tonic against Pityrosporum ovale

doi 10.20473/jkr.v2i2.6197

Abstract views = 535 times | PDF views = 615 times  
Riska Surya Ningrum, Antonius Budi Prasetyo, Alfinda Novi Kristanti

PDF  
93-97

##### OPTIMASI JENIS DAN KADAR SUMBER NITROGEN SERTA pH MEDIUM UNTUK PRODUKSI PROTEASE DARI ISOLAT HtCUM6.2.2 DARI TAUCO SURABAYA

doi 10.20473/jkr.v2i2.6307

Abstract views = 371 times | PDF views = 1012 times  
Monika Rahayu, Evi Susanti

PDF  
98-107

##### POTENSI EKSTRAK METANOL DAUN MANGGA BACANG (Mangifera foetida L.) SEBAGAI ANTIBAKTERI TERHADAP Enterobacter aerogenes DAN IDENTIFIKASI GOLONGAN SENYAWA AKTIFNYA

doi 10.20473/jkr.v2i2.5753

Abstract views = 1080 times | PDF views = 2780 times  
Ely Setiawan, Tien Setyaningtyas, Dwi Kartika, Dian Riana Ningsih

PDF  
108-117

##### The Influence of pH Values on the Crystallite Size of ZnO by Solvothermal Synthesis

doi 10.20473/jkr.v2i2.6430

Abstract views = 225 times | PDF views = 222 times

PDF  
118-122

#### Instruction for Author

Guide for authors

Online Submission

Document Template

#### Journal Policy

Focus and Scope

Publication Ethics

Article Processing Charge

Peer Reviewers

Editorial Team

Open Access Statement

Plagiarism

Copyright

Contact

#### USER

Username

Password

Remember me

Login

#### NOTIFICATIONS

[View](#)  
[Subscribe](#)

#### JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

[By Issue](#)

[By Author](#)

[By Title](#)

[Other Journals](#)

#### INFORMATION

[For Readers](#)

[For Authors](#)



[For Librarians](#)

#### KEYWORDS

Alkaline protease, Chicken feces, Gallus gallus bankiva, Pseudomonas sp Alkaloid turunan furokuinolin, skimmianin, Toddalia asiatica L., antikanker Analytical Chemistry Antifungal preparations, Candida albicans, Betel leaf (SDS), Gellinggang leaf extract (Cassia alata L.) Antioxidant, S. polycystum, T. decurrens, Dutungan island DENV-2, 2-isopropanol, Methanol, Chloroform, Ethanol 96%, RNA Isolation Dioscorea



**Gambaran Histopatologi Hepar Tikus Putih (*Rattus Novergicus*) yang Diinduksi dengan Parasetamol dosis Toksik Pasca Pemberian Ekstrak Etanol Daun Kelor (*Moringa Oleifera*)**

 10.20473/jkr.v2i2.6700

 Abstract views = 536 times |  views = 4803 times  
Noer Kumala I



**PENGEMBANGAN ELEKTRODA PASTA KARBON TERMODIFIKASI MIP DENGAN MONOMER ASAM METAKRILAT SEBAGAI SENSOR PADA ANALISIS KREATIN SECARA POTENSIOMETRI**

 10.20473/jkr.v2i2.6699

 Abstract views = 654 times |  views = 1152 times  
Usreg Sri Handajani, Cahya Nurrahmi H Amilianti, Miratul Khasanah

**Pengaruh pH terhadap Degradasi Pewarna Direct Blue menggunakan Jamur Pelapuk Kayu *Pleurotus flabellatus***

 10.20473/jkr.v2i2.6546

 Abstract views = 376 times |  views = 1157 times  
Niki Rachmanur Husna, Hasri Ummas, Sudding -

 PDF  
123-130

 PDF  
131-139

 PDF  
140-146

hispidia Dennst, Dioscoreaceae, phenolic, 4-ethoxy-3-methoxyphenol, methyl-3,4-dihydroxybenzoic, antioxidant Ganoderma lucidum, senyawa fitokimia, GCMS Inorganic Chemistry RNA, Precipitation Solvent, DMF, EDTA, Ultrapure H2O Selada merah (*Lactuca sativa* var. *Crispa*), sitotoksik, *Artemia salina* Leach. T102 ZSM-5, metakaolin, terdealuminasi, tanpa templat, pengaturan suhu dan waktu antibakteri, *Enterobacter aerogenes*, *Mangifera foetida* L., metanol asam amino, besi, nanomagnetit, dan mineral enzym, biodegradation, fungus enzyme fenil propanoid, skopoletin, dan *Ipomoea batatas* L. fotokatalitik methanol, Phytochemical screening, Piper betle L. var *Nigra*. methyloleate, catalyst, Ni/H5NZ, hydrocracking remazol red 198

**CURRENT ISSUE**


ATOM	1.0
RSS	2.0
RSS	1.0



View JKR Stats  
00044598



Pusat Pengembangan Jurnal dan Publikasi Ilmiah  
**Universitas Airlangga**  
copyright@2017 Template PPJPI



[Home](#) > [About the Journal](#) > [Editorial Team](#)

## EDITORIAL TEAM

### EDITOR-IN-CHIEF

[Dr. Purkan Purkan](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

### VICE EDITOR-IN-CHIEF

[Dr. Abdulloh Abdulloh](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

### EDITORIAL BOARD

[Prof. Ni Nyoman Tri Puspaningsih](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

[Prof. Chun-Hu Chen](#), Department of Chemistry, National Sun Yat-sen University, Kaohsiung, 80424, Taiwan

[Prof. Punnapayak Hunsa](#), Faculty of Science, Chulalongkorn University, Bangkok, 10330, Thailand

[Dr. rer.nat Ganden Supriyanto](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

[Dr. Hery Suwito](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

### MANAGING EDITOR

[Dr. Imam Siswanto](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

[Rico Ramadhan, Ph.D](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

[Qurrota A'yuni, M.Si](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

[Fatiha Khairunnisa, M.Si](#), Department of Chemistry, Faculty of Science and Technology, Airlangga University

### ADMINISTRATION SUPPORT

[Fendi Kristanto](#), Department of Chemistry Faculty of Science and Technology Universitas Airlangga

[Oktavia Wahyu Ningsih](#), Department of Chemistry Faculty of Science and Technology Universitas Airlangga

#### Instruction for Author

[Guide for authors](#)

[Online Submission](#)

[Document Template](#)

#### Journal Policy

[Focus and Scope](#)

[Publication Ethics](#)

[Article Processing Charge](#)

[Peer Reviewers](#)

[Editorial Team](#)

[Open Access Statement](#)

[Plagiarism](#)

[Copyright](#)

[Contact](#)

#### USER

Username

Password

Remember me

#### NOTIFICATIONS

[View](#)  
[Subscribe](#)

#### JOURNAL CONTENT

Search

Search Scope

All

Browse

[By Issue](#)

[By Author](#)

[By Title](#)

[Other Journals](#)

#### INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

#### KEYWORDS

Alkaline protease, Chicken feces, Gallus gallus bankiva, Pseudomonas sp  
Alkaloid turunan furokuinolin,  
skimmianin, Toddalia asiatica L.,  
antikanker Analytical Chemistry  
Antifungal preparations, Candida  
albicans, Betel leaf (SDS), Gellinggang  
leaf extract (Cassia alata L.)  
Antioxidant, S. polycystum, T.  
deccurens, Dutungan island DENV-2, 2-  
isopropanol, Methanol, Chloroform,  
Ethanol 96%, RNA Isolation Dioscorea



View JKR Stats  
00044596



hispidia Dennst, Dioscoreaceae, phenolic, 4-ethoxy-3-methoxyphenol, methyl-3,4-dihydroxybenzoic, antioxidant Ganoderma lucidum, senyawa fitokimia, GCMS Inorganic Chemistry RNA, Precipitation Solvent, DMF, EDTA, Ultrapure H2O Selada merah (Lactuca sativa var. Crispa), sitotoksik, Artemia salina Leach. TiO2 ZSM-5, metakaolin, terdealuminasi, tanpa templat, pengaturan suhu dan waktu antibakteri, Enterobacter aerogenes, Mangifera foetida L., metanol asam amino, besi, nanomagnetit, dan mineral enzym, biodegradation, fungus enzyme fenil propanoid, skopoletin, dan Ipomoea batatas L. fotokatalitik methanol, Phytochemical screening, Piper betle L. var Nigra. methyloleate, catalyst, Ni/H5NZ,hydrocracking remazol red 198



Pusat Pengembangan Jurnal dan Publikasi Ilmiah  
**Universitas Airlangga**  
copyright@2017 Template PPJPI

## CELERY HERB ESSENTIAL OIL IN THE FORMULATION OF ANTIDANDRUFF HAIR TONIC AGAINST *PITYROSPORUM OVALE*

Riska Surya Ningrum<sup>1</sup>, Antonius Budi Prasetyo<sup>2</sup>, Alfinda Novi Kristanti<sup>2\*</sup>

<sup>1</sup>Student of Undergraduate Chemistry Study Program, Faculty of Science and Technology

<sup>2</sup>Department of Chemistry, Faculty of Science and Technology

Universitas Airlangga

\*email: alfinda-n-k@fst.unair.ac.id

Received 16 Oktober 2017

Accepted 28 Nopember 2017

### Abstract

Essential oil from *Apium graveolens* (celery), a species belonged to *Apiaceae*, was isolated, identified, and tested its activity against fungi *Pityrosporum ovale*, fungi that cause dandruff problem. Essential oil isolation was conducted by steam distillation. The identification with GC-MS showed that the major compound of celery essential oil was 3-isobutylidenphalide. The anti-fungal activity test was performed using pure celery oil and celery oil added with virgin coconut oil (VCO) and paraffin oil that are usually used as components in hair tonic formulation. The bioactivity test showed that celery oil, whether in pure form or in hair tonic formulation exhibited excellent growth inhibition activity of *Pityrosporum ovale*

**Keywords:** essential oil, celery, *Apium graveolens*, *Pityrosporum ovale*

### Introduction

Celery belongs to *Apiaceae*, is aromatic herb growing on lowland or highland, and is well known of its essential oil. According to the history, celery has been used as a vegetable since 1640, but it was only recognized as a medicinal plant in 1942. As a medicinal plant, celery is able to stimulate digestive enzymes in increasing the appetite, use as diuretic agent, to maintain blood vessel flexibility, to lower blood pressure and wide peripheral blood vessels (Sudarsono, 1996). A study showed that celery can used as hypocholesterolemia and hypolipidemia agents even though the dosages used in that study not decrease the total cholesterol and lipid significantly (Juheini, 2002). In addition, celery is also efficacious to nourish hair. The juice of celery herb showed an activity in promoting hair growth and increasing hair thickness (Kuncari, 2015). Previous studies mentioned that the ethanol extract of celery herb showed inhibitory activity against

*Pityrosporum ovale*, a fungi causing dandruff (Sukandar, 2006), *Streptococcus mutans*, the most important bacteria in the process of dental caries (Majidah, 2014), and *Candida albicans*, one microorganism that causes candidiasis (Rachmawati, 2014). Meanwhile, the essential oils of celery can inhibit the growth of *Staphylococcus aureus*, *Listeria monocytogenes* and *L. ivanovii* (Misis, 2008).

In this study we tested the antifungal activity of the essential oil of celery against *Pityrosporum ovale*, a fungus that causes dandruff. This fungus is a naturally occurring yeast-like organism found on the scalp and other parts of human skin. It's mostly found on skin areas with plenty of sebaceous glands, such as on scalp, face, and upper part of the body (Niharika, 2010). Several synthetic fungicide have been produced to fight dandruff problems, besides the herbal ingredients which assumed to have good anti-dandruff activity. This research is a study on the

efficacy of anti-dandruff substances from herbal origin, in this case was celery essential oil, in the formulation of anti-dandruff hair tonic against *Pityrosporum ovale* in vitro. The celery essential oil was also used due to expectation of its flavor properties. Ketoconazole was used as positive control due to its best known of anti-*Pityrosporum* activity (Prabhamanju, 2009).

## Materials and Methods

### Equipment and materials

Celery herbs were procured from the local market of Surabaya, East-Java, as well as ketoconazole. *Pityrosporum ovale* was cultured in the dermatitis laboratory of local hospital. The chemicals used in this study were n-hexane, CaCl<sub>2</sub> anhydrous, distilled water and the media Sabouraud Dextrose Agar (SDA) for fungus. Apparatus and instrument used in this study was a set of steam distillation and GC-MS GC2010 MSQP 2010S Shimadzu.

### Research procedure

#### Isolation and identification of celery essential oil

The celery essential oil was obtained by steam distillation from celery herbs (*Apium graveolens* L.) which were first cleaned and cut into small pieces. The essential oil obtained was placed in a dark glass bottle, closed tightly, and stored in a refrigerator. Furthermore, celery essential oil was analyzed using GC-MS

#### Preparation of fungus

*Pityrosporum ovale* was taken from seborrhoeic dermatitis patient and then was cultured on agar SDA (*Sabouraud Dextrose Agar*) media which was contained 20 g of glucose, 5 g of pepton, and 9 g of agar. The final volume of medium was adjusted to 500 mL by adding aquades (Funder, 1968). Preparation of inoculum was done in this way. Sterilized water (10 mL) was added to *P.ovale* stock culture and then was shaken with vortex. OD was measured using Spectrophotometer at 425

nm to get 0.45. This suspension would be used to activity test (Khan, 1992).

#### Determination of Antifungal properties of celery essential oil

Various concentration of celery essential oil (2 mL, 4 mL, 6 mL, 8 mL, 10 mL) were prepared. Ten ml of VCO and 2.5 ml of paraffin liquid were added into each concentration. The inhibiting capacity of each concentration of essential oil was tested using agar disc method. The control positive using for this test was a solution of ketoconazole 2% (w/w). As comparative, the anti-*pytyrosporum* of pure essential oil experiment was also conducted.


Twenty ml of SDA media which had been diluted was mixed with 1 mL of inoculum and then poured in a petri dish, and allowed to solid. Three sterile paper discs were placed on agar medium in a petri dish. Thirty ml of each test solution was added to the disc paper drop wise. Each petri dish was incubated for 3-4 days at 25 °C. Growth inhibition around the paper discs was measured. The experiment was conducted in triplicate

## Results and Discussion

The yield of celery essential oil obtained from steam distillation process was 0,13 %. The analysis by GC-MS showed that the main component of this essential oil was  $\beta$ -selinene (time retention 22.025, % Area 0.77) and 3-isobutylidenphthalide (time retention 25.933, % Area 30.16). The structure and the mass spectrum of these compounds are described in Figure 1 and Figure 2 respectively. This result differed from those published before, which stated that limonene was the major constituent of *Apium graveolens* essential oil (Van Wassenhove, 1990; Hassanen, 2015). However, it has been said that terpenes and phtalides are the key volatile components found in various cultivars of celery (van Wassenhove, 1990). Comparison result of antifungal activity between celery


essential oil and ketoconazole against *P. ovale* are presented in the Table 1. The activity test of pure essential oils and ketokenazol 2% indicates that diameter of inhibition was 16.7 cm and 25.7 cm respectively. From this result showed that celery oil exhibited an antifungal activity against *P. ovale*. However, its activity is lower than ketokenazole Thus, celery essential oil showed as alternative herbal

substance to be used in the formulation of anti-dandruff hair tonic. Furthermore, the anti-fungal activity test of celery essential oil mixed with VCO and paraffin liquid as usually found in hair tonic formulations was done. The result of the bioactivity test was displayed in Table 2




**Figure 1.** The structure of main compounds in celery oil

Hit#:1 Entry:70939 Library:WILEY229.LIB  
 SI:93 Formula:C15 H24 CAS:17066-67-0 MolWeight:204 RetIndex:0  
 CompName:.beta.-Selinene \$\$ Naphthalene, decahydro-4a-methyl-1-methylene-7-(1-methylethenyl)-, [4aR-(4a.alpha.,7.alpha.,8a.b



Hit#:1 Entry:57372 Library:WILEY229.LIB  
 SI:94 Formula:C12 H12 O2 CAS:0-0-0 MolWeight:188 RetIndex:0  
 CompName:3-Isobutyldenphthalide \$\$



**Figure 2.** Mass spectrums of main compounds


**Table 1.** Antifungal activity of celery essential oil and ketoconazole

Sample	Diameter (mm)			Average
	Replication			
	1	2	3	
1 mL essential oil	9	24	17	16.7
Ketokenazole 2%	19	23	35	25.7



**Table 2.** Bioactivity test result of sample (celery essential oil in various concentrations mixed with VCO and paraffin liquid)

No.	Essential oil (mL)	VCO (mL)	Paraffin Liquid (mL)	Diameter (cm)			Average
				Replication			
				1	2	3	
1	0	10	2,5	0	0	0	0
2	2	10	2,5	0	0	0	0
3	4	10	2,5	0	0	0	0
4	6	10	2,5	1,8	0,9	0	0,9
5	8	10	2,5	1,7	0,6	1,6	1,3
6	10	10	2,5	2,0	0,9	1,5	1,47

**Figure 3.** Curve of the relationship between the concentration of celery oil in hair tonic with diameter of growth inhibition of *P.ovale*

At a concentration of 2 and 4 mL of essential oil, no anti-fungal activity was observed. The anti-fungal activity was observed started from the concentration of 6, 8, and 10 ml, whereas the mixture of VCO with liquid paraffin (no essential oil) did not exhibit anti-fungal activity.

### Conclusion

Based on the result, it can be stated that the essential oil of celery may inhibit the growth of *Pityrosporum ovale*. Therefore, the use of this essential oil as an ingredient in hair tonic or anti-dandruff shampoo formulation can be recommended. In addition, it can also serve as a fragrance.

### References

Funder, S., 1968, *Practical Micology Manual For Identification Of Fungi*, Hafner Publishing Company Inc., New York, USA

Hassanen, N.H.M.; Eissa, A.M.F.; Hafez, S.A.M.; Mosa, E.A.M. 2015. Antioxidant and antimicrobial activity of celery (*Apium graveolens*) and coriander (*Coriandrum sativum*) herb and seed essential oil. *International Journal Current Microbiology and Applied Sciences*, 4(3), 284-296

- Juheini. 2002. Pemanfaatan herba seledri (*Apium graveolens* L.) untuk menurunkan kolesterol dan lipid dalam darah tikus putih yang diberi diet tinggi kolesterol dan lemak. *Makara, Sains*, 6(2), 65-69
- Khan, M.Y.; Dahot, M.U.; Khan, M.Y. 1992. Single cell protein production by *Penicillium javanicum* from pretreated rice husk, *Journal of Islamic Academy of Sciences*, 5(1), 39-43.
- Kuncari, E.S.; Iskandarsyah; Praptiwi. 2015. The irritation and hair growth activity test of male S-D mice : Effect of gel containing apigenin and celery (*Apium graveolens* L.) juice, *Media Litbangkes*, 25(1), 15-22
- Majidah, D.; Fatmawati, D.W.A.; Gunadi, A. 2014. Antibacterial activity of celery leaves extract (*Apium graveolens* L.) against *Streptococcus mutans* as an alternative mouthwash, *Artikel ilmiah hasil penelitian mahasiswa*, Universitas Jember
- Misic, D.; Zizovic, I.; Stamenic, M.; Asanin, R.; Ristic, M.; Petrovic, S.D.; Skala, D. 2008. Antimicrobial activity of celery fruit isolates and SFE process modeling, *Biochemical Engineering Journal*, 42, 148-152.
- Niharika, A.; Aquicio, J.M.; Anand, A. 2010. Antifungal properties of neem (*Azadirachta indica*) leaves extract to treat hair and dandruff, *E-International Scientific Research Journal*, 2(3), 244-252.
- Prabhmanju, M.; Shankar, S.G.; Babu, K.; Ranjith, M.S. 2009. Herbal vs chemical substances as antidandruff ingredients : which are more in effective the management of dandruff ?-An overview, *Egyptian Dermatology Online Journal*, 5(2): 8
- Rachmawati, I. 2014. Pengaruh konsentrasi ekstrak etanol daun seledri (*Apium graveolens*) terhadap hambatan pertumbuhan *Candida albicans* in vitro, *Skripsi*, Universitas Muhammadiyah Surakarta
- Sudarsono; Pudjoanto, A.; Gunawan, D.; Wahyuono, S.; Donatus, I. A.; Drajad, M.; Wibowo, S.; Ngatidjan. 1996. *Tumbuhan Obat, Hasil Penelitian, Sifat-sifat dan Penggunaan*, Pusat Penelitian Obat Tradisional, UGM, Yogyakarta
- Sukandar, E. Y.; Suwendar; Ekawati, E. 2006. Aktivitas ekstrak etanol herba seledri (*Apium graveolens*) dan daun urang aring (*Eclipta prostate* L.) terhadap *Pytirosporum ovale*, *Majalah Farmasi Indonesia*, 17(1), 7-12.
- van Wassenhove, F.; Dirinck, P.; Vulsteke, G.; Schamp, N. 1990. Aromatic volatile composition of celery and celeriac cultivars, *Hort Science*, 25(5), 556-559.