

Dengue Virus Serotype 4 Is Responsible for the Outbreak of Dengue in East Java City of Jember, Indonesia

by Aryati Aryati

Submission date: 19-Jul-2021 11:48AM (UTC+0800)





Submission ID: 1621379439

File name: he_Outbreak_of_Dengue_in_East_Java_City_of_Jember,_Indonesia.pdf (4.72M)

Word count: 10190

Character count: 60585

Dengue Virus Serotype 4 Is Responsible for the Outbreak of Dengue in East Java City of Jember, Indonesia


Aryati Aryati ^{1,2}, Billy J. Wrahatnala ³, Benediktus Yohan ⁴ , May Fanny ^{5,6}, Faradila K. N. Hakim ³, Eka Putri Sunari ⁷, Nelly Zuroidah ⁷, Puspa Wardhani ^{1,2}, Marsha S. Santoso ⁴, Dominicus Husada ⁸, Ali Rohman ⁹ , Siti Nadia Tarmizi ¹⁰, Justus T. O. Sievers ⁴  and R. Tedjo Sasmono ^{4,*} 

¹ Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya 60286, Indonesia; dr_aryati@yahoo.com (A.A.); puspa_pk@yahoo.co.id (P.W.)

² Institute of Tropical Diseases, Universitas Airlangga, Surabaya 60115, Indonesia

³ Master Program of Tropical Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya 60286, Indonesia; drbillyjordanw@gmail.com (B.J.W.); fara.faradilahakim@gmail.com (F.K.N.H.)

⁴ Eijkman Institute for Molecular Biology, Jakarta 10430, Indonesia; yohan@eijkman.go.id (B.Y.); marshasantoso@eijkman.go.id (M.S.S.); jto.sievers@gmail.com (J.T.O.S.)

⁵ Clinical Pathology Sub-Specialization Program, Universitas Airlangga, Surabaya 60286, Indonesia; dr.mayfannytanzilia@gmail.com 

⁶ Department of Clinical Pathology, Faculty of Medicine, Universitas Ciputra, Surabaya 60219, Indonesia

⁷ Clinical Pathology Specialization Program, Universitas Airlangga, Surabaya 60286, Indonesia; siputrisunari@yahoo.com (E.P.S.); nelly.zuroidah-2016@fk.unair.ac.id (N.Z.)

⁸ Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya 60286, Indonesia; dominicushusada@yahoo.com

⁹ Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115, Indonesia; alirohman@fst.unair.ac.id

¹⁰ Directorate of Vector-Borne and Zoonotic Disease, Ministry of Health of Indonesia, Jakarta 12950, Indonesia; nadiawiweko@gmail.com

* Correspondence: sasmono@eijkman.go.id; Tel.: +62-21-391-7131

³ Received: 29 June 2020; Accepted: 18 August 2020; Published: 20 August 2020



Abstract: Outbreaks of dengue virus (DENV) in Indonesia have been mainly caused by the DENV serotype-1; -2; or -3. The DENV-4 was the least-reported serotype in Indonesia during the last five decades. We recently conducted a molecular epidemiology study of dengue in the Jember regency, East Java province, Indonesia. Dengue is endemic in the region and outbreaks occur annually. We investigated the clinical characteristics and etiology of dengue-like febrile illness in this regency to understand the disease dynamics. A total of 191 patients with clinical symptoms similar to dengue were recruited during an 11-month study in 2019–2020. Children accounted for the majority of cases and dengue burden was estimated in 41.4% of the cases based on NS1 antigen, viral RNA, and IgG/IgM antibody detection with the majority (73.4%) being primary infections. Secondary infection was significantly associated with a higher risk of severe dengue manifestation. All four DENV serotypes were detected in Jember. Strikingly, we observed the predominance of DENV-4, followed by DENV-3, DENV-1, and DENV-2. Genotype determination using Envelope gene sequence revealed the classification into Genotype I, Cosmopolitan Genotype, Genotype I, and Genotype II for DENV-1, -2, -3, and -4, respectively. The predominance of DENV-4 in Jember may be associated with a new wave of DENV infections and spread in a non-immune population lacking a herd-immunity to this particular serotype.

Keywords: dengue; molecular epidemiology; serotype; Java; Indonesia

47

1. Introduction

The dengue virus (DENV) is a member of the *Flaviviridae* family that causes a systemic infection simply known as dengue. DENV maintains arboreal-human transmission cycles via the *Aedes* mosquito vector, with historical records of cases matching the symptoms of infection dating as far back as 265 AD in China [1]. DENV's ~10.7 kb single-stranded positive-sense RNA genome encodes three structural (C, prM/M, E) and seven non-structural (NS1, NS2A, NS2B, NS3, NS4A, NS4B, NS5) proteins [2,3]. The four serotypes (DENV-1, -2, -3, and -4) are able to cause repeated and simultaneous infections with varying clinical manifestations, most commonly an acute febrile illness, accompanied by myalgia, arthralgia, headache, and a rash, in the case of classical dengue fever (DF) [4,5]. In more extreme cases, the infection can progress to dengue hemorrhagic fever (DHF), and the frequently fatal dengue shock syndrome (DSS), making dengue a significant public health problem in tropical and subtropical areas of the world, including Indonesia, driven by the global distribution of mosquito vectors, among other factors [4,5].

The serotypes of DENV are antigenically different, enabling repeated and simultaneous infections, as well as being genetically clearly distinct, with separate genotypes within each serotype, which demonstrate varying pathogenicity and infectivity [6–8]. Phylogenetic analyses of DENV genomes have recently also allowed an expanded understanding of the factors driving the virus' evolution and pathogenicity, as well as its historical origins [8,9].

Indonesia has a long history of dengue, with metropolitan centers Jakarta and Surabaya reporting cases since 1968 [10]. Infections are widespread, with all provinces and 85.6% of cities in the country reporting at least one case of DHF in 2018 [11]. In the same year, East Java province recorded the second highest number of DHF cases (8449 in total) and the highest number of total fatalities (84 in total) in Indonesia, with an incidence rate of 21.4/100,000 [11]. However, while the history and development of DENV is well-characterized in the province's capital Surabaya [12–14] and Madura [15], there is a distinct lack of studies elucidating the nature of dengue in the Jember regency, the third largest city in East Java after Surabaya and Malang.

The expansion of the monitoring of DENV evolution in less-studied areas such as Jember is essential in implementing effective public health strategies to mitigate and avoid future dengue epidemics. Such epidemics can occur when new genotypes of DENV emerge in or are introduced to naïve populations, as has been reported previously [16,17]. As DENV genotypes have been reported to show antigenic differences [18], new genotypes present a significant risk, even to populations with a history of exposure to other DENV strains.

The spread of a rare DENV serotype would have a similar potency as an epidemic. Previous analyses of dengue outbreaks in Indonesia have shown these to be overwhelmingly caused by DENV-1–3 [14,19,20], with cases of serotype shift over time [12,13]. While DENV-4 has circulated in East Asia and other parts of South-East Asia over the past 60 years [9], there have been no reports of outbreaks associated with DENV-4 in Indonesia, except for an isolated report from Jakarta based on a limited number of partial genomic sequences [21]. Due to this rarity of infections, DENV-4 is also the least studied of the serotypes, with five identified genotypes (I–V), one of which is exclusively sylvatic (IV), with genotype-II being split into -IIa and -IIb [9,18].

The process of monitoring dengue cases in Indonesia must also account for ambiguity in clinical symptoms and cross-reactive antibodies affecting the identification of patients. Infection with other *Aedes*-transmitted viruses often show very similar symptoms to those of dengue and will often share spatio-temporal distribution with dengue. One such disease, chikungunya, caused by the Alphavirus chikungunya virus (CHIKV), not only shares a large amount of clinical symptoms with DF, but is also widespread in Indonesia [22]. While point-of-care rapid antibody testing kits for DENV, such as those developed for IgG and IgM, can exclude chikungunya patients, these immunoglobulins may be cross-reactive for DENV if the patient is infected with another *Flaviviridae* virus, such as the Zika virus (ZIKV) [23] or the Japanese encephalitis virus (JEV) [24]. Zika, a member of the same *Flaviviridae* family as DENV, causes clinical symptoms which may easily be confused with DF and

there is evidence to support the notion that it is widespread in Indonesia, with a large number of undiagnosed/misdiagnosed cases due to the low severity of symptoms [25].

In this study, we performed a molecular epidemiology study of DENV and other arboviruses circulating in Jember regency, East Java. Our major aims were to inspect both clinical epidemiology and virological aspects of dengue in Jember. Here, we highlight the role of DENV-4 in causing an outbreak in the region and describe the genetic and epidemiological make up that it may contribute.

1

2. Materials and Methods

2.1. Ethical Considerations for the Commencement of the Study

The study was initiated after obtaining ethical approval from the Research Ethics Committee of the Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia (Ethical Approval No. 156/EC/KEPK/FKUA/2019, 17 May 2019). The voluntary involvement into the study was ensured by obtaining written informed consent from all patients and/or children's legal guardians.

100

2.2. Study Design and Patient Recruitment

The study was designed as a cross-sectional study performed in the Jember regency, East Java Province, Indonesia. Jember is the third largest urban area in East Java after Surabaya and Malang, located at $-8^{\circ}10'8''$ S latitude, $113^{\circ}42'8''$ E longitude with an area of 3306 km^2 and an average altitude of 0–500 m above sea level (masl). The population consists of 2,450,668 inhabitants with 0.41% population growth in 2019, living in 31 districts and 248 villages [26]. Jember is located approximately 200 km to the southeast from Surabaya, the capital city of East Java.

Study subjects' recruitment was conducted for 11 months from May 2019 to March 2020, covering a monsoon season occurring typically from November to April, peaking in January, which is also the dengue peak season in the area. Patients suspected of having dengue were recruited at six hospitals/healthcare centers, namely RS Jember Klinik, RSD Dr. Soebandi, RS Citra Husada, Klinik Dr. M. Suherman, Klinik Dokterku Taman Gading, and Puskesmas Sumbersari, Jember. The inclusion criteria were febrile patients willing to participate in the study, aged up to 80 years old with a body temperature of $\geq 38^{\circ}\text{C}$ for less than 5 days of fever, accompanied by at least two of the clinical symptoms of dengue, i.e., headache, retro-orbital pain, myalgia, arthralgia/bone pain, and rash, as described in the World Health Organization, South-East Asia Regional Office (WHO-SEARO) 2011 guidelines [5]. The exclusion criteria were patients with fever and upper respiratory tract infections and/or those diagnosed as non-dengue. An interview session was performed by a doctor or research nurse to record demographic data and study participants' medical and relevant clinical information. During the interview, study participants were asked to provide written informed consent signed by the patients or their legal guardian. Upon consent given, single 3 to 5 mL blood during an acute phase was drawn from each patient for serum processing. Sera were separated by centrifugation and stored frozen (at -20°C) until further processing and laboratory analyses were done at the Eijkman Institute for Molecular Biology, Jakarta. The fraction of blood sample from each study participant was analyzed for routine/common blood parameters, i.e., hemoglobin, hematocrit, erythrocyte, platelet, and leucocyte count.

The determination of dengue clinical manifestation was based on WHO-SEARO 2011 guidelines [5]. Patients were classified as having either dengue fever (DF), dengue hemorrhagic fever (DHF), or dengue shock syndrome (DSS). Moreover, patients having unusual manifestations and severe organ involvement, such as of the liver, kidneys, brain, or heart were considered as if they had expanded dengue syndrome. In terms of dengue confirmation, patients with clinical symptoms of dengue and who tested positive for dengue IgM and/or IgG were classified as "probable dengue", while patients who tested positive for the dengue NS1 antigen and/or RT-PCR were categorized as "confirmed dengue" [5]. Data on the dengue fever IR in Jember, East Java, and Indonesia were compiled from published reports by Jember Regency Health Office, East Java Provincial Health Office and the Ministry

of Health of the Republic of Indonesia (*Profil Kesehatan Indonesia/Indonesia Health Profiles 2014–2019*, accessible as data repository at <https://pusdatin.kemkes.go.id>).

2.3. Dengue NS1 Antigen and Serology Tests

Acute serum samples were first screened for the presence of the DENV-encoded Non Structural Protein 1 (NS1) antigen using lateral flow immunochromatographic assays by Standard Q Dengue NS1 Ag Rapid Test (SD Biosensor, Gyeonggi-do, Korea) for dengue confirmation [5]. The presence of anti-dengue IgM and IgG was determined by Standard Q Dengue IgM/IgG Rapid (SD Biosensor). Results from the IgM/IgG tests were used as criteria in defining the infection status (primary or secondary infection). A positive IgM result was indicative of active primary or secondary infection, while an IgG-positive result was indicative of active secondary infection. Primary infection was determined by positive IgM and negative IgG results, while secondary infection was determined by positive IgG, which could be accompanied by a positive IgM result.

2.4. Molecular Detection for DENV, Other Arboviruses, and Typhoid Fever

A total amount of 140 µL of serum sample was used for nucleic acid extraction using QIAamp Viral RNA mini kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The extracted RNA was then used as a template in the simultaneous DENV detection and serotyping as well as CHIKV detection performed using abTES DEN/CHIKU 5 qPCR (AIT Biotech, Singapore), according to the method detailed by the manufacturer. The possible presence of flavivirus group pathogens (such as ZIKV and JE viruses) was also screened for using pan-flavivirus real-time qRT-PCR, using primers as described elsewhere [27].

Further screening was done for samples which tested negative for dengue and other arbovirus infection to detect other possible causative pathogens. Analysis for typhoid fever was performed using Tubex TF (IDL Biotech, Bromma, Sweden) based on the Inhibition Magnetic Binding Immunoassay (IMBI) method to detect *Salmonella typhi* IgM anti-O9 antibodies in serum. The identification of typhoid fever was done following the manufacturer's recommendations.

2.5. DENV Isolation in Cell Culture

Virus isolation approach was performed on serum samples using the cell culture technique. African green monkey (*Cercopithecus aethiops*) kidney cells (Vero CCL-81 (ATCC)) were seeded at 2×10^5 cells/well of 24-well plate (Corning, Corning, NY, USA) in MEM medium supplemented with 10% of fetal bovine serum (FBS), 2 mM of L-glutamine and 1% antibiotic/antimycotic (all from Gibco, Thermo Scientific, Carlsbad, CA, USA). Plates were incubated overnight at 37 °C inside a humidified incubator with 5% CO₂ supplementation to generate cell monolayers of 80–100% confluency. An amount of 50 µL of serum sample was prepared in a total of 200 µL of MEM – 2% FBS and inoculated into the designated well containing monolayer of cells. Virus adsorption was allowed for 1 h at 37 °C, 5% CO₂ before inoculant aspiration and replenishment with fresh medium. Plates were incubated at 37 °C, 5% CO₂ with daily checking for the presence of cell's cytopathic effect (CPE), typically for 7–9 days. Supernatant was harvested from wells, centrifuged 3000× g for cells and debris separation and stored frozen at –80 °C until use. The presence of DENV was tested for using qRT-PCR molecular detection of extracted RNA.

2.6. Determination of DENV Genotype by Envelope Gene Sequencing

The determination of the DENV Genotype was based on the full length envelope (E) gene sequencing using the protocol described earlier [14]. Briefly, cDNA was generated from DENV RNA using Superscript III Reverse Transcriptase (Invitrogen-Thermo Scientific, Carlsbad, CA, USA) and DENV-specific primers. PCR amplification was done using cDNA template and *Pfu* Turbo DNA Polymerase (Stratagene-Agilent Technologies, Santa Clara, CA, USA) to generate PCR products of approximately 2.5 kb in size that were then purified on 0.8% agarose gel using the QIAquick gel

extraction kit (Qiagen). Purified amplicons were used as templates in cycle sequencing reactions using BigDye Dideoxy Terminator kits v.3.1 (Applied Biosystems-Thermo Scientific, Foster City, CA, USA) employing overlapping primers for each serotype from both strands [20]. Sequencing was performed on purified DNA using capillary sequencing implemented on a 3130 xl Genetic Analyzer (Applied Biosystems). SeqScape v.2.5 software (Applied Biosystems) was used to assemble the sequence reads, with manual inspection to clarify sequence ambiguities. Sequence alignment was done using a Multiple Sequence Comparison by Log-Expectation (MUSCLE) program inside MEGA X [28] software (v.10.1.1, Penn State University, State College, PA, USA). The classification of the genotypes in each serotype was based on classifications by Goncalvez et al. [29], Twiddy et al. [30], Lanciotti et al. [31], and Lanciotti et al. [32] for DENV-1, -2, -3, and -4, respectively. Initial DENV genotype classification trees are available as Supplementary Figures S1–S4.

2.7. DENV Phylogenetic Reconstruction and Evolutionary Analysis

The genetic distance method was used to filter all closely related DENV sequences available in the GenBank repository as of 30 April 2020 implemented in the BLAST website (<https://blast.ncbi.nlm.nih.gov>). A number of 100 closely related strains from each DENV serotype were used as initial datasets along with Jember isolate sequences. Further filtering was applied to retrieve sequences of the DENV E gene from each serotype with more than 1400 nt. From this, we selected only 60 of the most closely related strains to achieve clarity of the tree view.

The resulting alignment of 1485 nt (1479 nt for DENV-3) was then used in a robust phylogenetic and evolutionary analysis using the Bayesian Markov Chain Monte Carlo (MCMC) algorithm, as implemented in BEAST v.2.6.1 [33]. Datasets were uploaded to the BEAUti v.2.6.1 graphical interface with the isolation year used for the calibration of each taxon. A phylogenetic tree was inferred based on selection of the statistical model for likelihood calculation, using MEGA X [28]. A statistical model was selected based on the Bayesian Information Criterion (BIC) results and the best selected model for all four serotypes datasets was the TrN (TN93) algorithm with four gamma parameters (G4). Molecular clock measurement was set using a relaxed uncorrelated lognormal molecular clock and a Bayesian skyline prior. The analysis was set to generate 100 million chains sampled for every 1000 chains with an initial estimated evolutionary rate set at 7.6×10^{-4} substitutions per site per year [34]. The MCMC trace was analyzed using Tracer v.1.7.1 to monitor the adequate Effective Sampling Size (ESS) for all the parameters after 10% burn-in. A maximum clade credibility (MCC) tree was created using TreeAnnotator v.2.6.1 and visualized using FigTree v.1.4.3. The evolutionary parameters were estimated as a median number with 95% Highest Posterior Density (HPD).

2.8. Statistical Analysis

Univariate testing was conducted using Pearson's Chi-square tests for categorical data, while age and hematological data were compared using Kruskal–Wallis tests. Multivariate testing was also conducted using binomial regression analysis for reported symptoms with age, gender, immunologic status, and serotype as potential covariates. All statistical analysis was done using R statistical software (<http://www.r-project.org>). A probability value of $p < 0.05$ was considered as statistically significant.

3. Results

3.1. Sample Collection and Demographical Data

From May 2019 to March 2020, a total of 191 febrile patients with dengue-like illness were enrolled at the six collection sites located within the Jember city center (Figure 1). These sites included reference hospitals where people all over Jember regency seek medication for their illness. In general, the annual dengue IR in Jember was lower than that in East Java province and on the national level. The annual IR of dengue in Jember in 2019 was documented at 40.32/100,000 population (Figure 2A). Historically, annual IRs in Jember were relatively stable within the period of three years between 2016 and 2018,

a downward-sloping trend from previous years. However, an increasing IR was observed in 2019 and continued with a marked increase from January to March 2020, culminating in a spike in February 2020 with 240 cases (equivalent to an IR of 117.08/100,000 population if extrapolated to 12 months). This sudden increase in dengue incidence was covered in the study period where 55 (or 28.8%) samples were collected during these three months (data not shown).



Figure 1. Map of study sites in Jember, East Java province, Indonesia.




Figure 2. Dengue cases in Jember regency. (A) Annual incidence rate (IR) of dengue in Jember regency, East Java province and Indonesia national data, with additional monthly incidence for Jember regency in early 2020 (extrapolated for 12 months). (B) Distribution of patient age by groups in Jember in 2019–2020.

Based on demographical data collected during patient recruitment, the recruited study participants were aged between 13 months and 64 years with a median age of 20 years (IQR: 9.5–29.0). A total of 98 patients were male (51.3%), with a male-to-female ratio of 1:0.95. Among all of the suspected dengue patients, children aged 0–10 years account for the highest proportion (27.2%), followed by adolescents 11–20 years old (25.7%) and young adults 21–30 y.o (23.0%) (Figure 2B). More children than adults were suspected of having dengue in Jember.

3.2. Clinical Examination of Dengue-Suspected Patients

We used clinical examination data to study the most common symptoms suffered by all dengue-suspected patients. Among all 191 patients, headache was the most common symptom reported (83.2%), followed by myalgia (52.4%) and vomiting (41.9%) (Table 1). Bleeding manifestation was observed in 14.1% of patients. However, none were reported to have other signs of severe dengue such as pleural effusion, ascites, and altered consciousness. Moreover, there was no significant correlation observed between reported symptoms and potentially affecting covariates, such as age, gender, infection status, and the infecting DENV serotype (Supplementary Table S1).

Table 1. Reported symptoms of acutely febrile dengue-suspected patients.

Symptoms (n = 191)	n (%)
Headache	159 (83.2)
Myalgia	100 (52.4)
Vomiting	80 (41.9)
Stomachache	73 (38.2)
Arthralgia	69 (36.1)
Retro-orbital Pain	47 (24.6)
Bleeding manifestation	27 (14.1)
Rash	13 (6.8)
Positive Tourniquet Test	10 (5.2)
Altered Consciousness	0 (0)
Ascites	0 (0)
Pleural Effusion	0 (0)

3.3. Dengue Serological Testing and Case Confirmation

Among all 191 dengue-suspected patients identified by symptoms, we found that 17 samples (8.9%) were probable dengue cases while another 62 samples (32.5%) were confirmed as dengue by NS1 antigen detection and/or molecular testing (Figure 3). Combined together, the burden of dengue in Jember was 41.4% (79/191). Out of the 62 confirmed dengue patients, 25 (40.3%) tested positive for both NS1 and via RT-PCR. The other 36 (58.1%) samples tested positive via RT-PCR only, and 1 sample (1.6%) tested positive for NS1 only.

When looking at the clinical manifestations of 79 probable and confirmed dengue patients, the majority (58 or 73.4%) manifested as the milder DF clinical manifestation, while the other 21 (26.6%) patients had the more severe DHF manifestation. Based on the infection status, a higher percentage of primary infection (73.4%) was recorded than secondary infections (26.6%). There was no correlation between patients' age and gender and infection status. However, secondary infections were more likely to present with DHF compared to primary infection ($p = 0.005$) (Table 2). Examining the laboratory data, the median platelet count of patients with secondary infections is significantly lower ($p < 0.001$) than that of primary infections. Out of 58 primary dengue infections, 10 were categorized as DHF due to severe thrombocytopenia of $50 \times 10^3/\mu\text{L}$ counts or less, rapidly decreasing platelet counts, or an increase in hematocrit levels of 20% or more.




Figure 3. The samples processing flow implemented in Jember dengue study.

Table 2. Characteristics of dengue among probable and confirmed patients in term of infection status.

Parameters	Infection Status (n = 79)		p-Value ^a
	Primary (n = 58)	Secondary (n = 21)	
Age, Median (IQR)	19.5 (8–28.5)	19 (9–26)	0.665 ^b
Age groups			1.000
Children (<20 y.o)	28	10	
Adults (≥20 y.o)	30	11	
Gender			0.111
Male	33	7	
Female	25	14	
Clinical manifestation			0.005
DF	48	10	
DHF	10	11	
Hematology data, Median (IQR)			
Platelet (×10 ³ /μL)	172 (105.3–223)	57 (35–85)	<0.001 ^b
Hematocrit (%)	41.1 (38.3–44.8)	42.9 (41.3–46.0)	0.052 ^b
WBC (×10 ³ /μL)	4.3 (3.2–7.3)	3.6 (2.7–5.6)	0.502 ^b

IQR, interquartile range; WBC, white blood count ^a Pearson’s Chi-squared test ^b Kruskal–Wallis test. Statistically significant values are shown in bold.

3.4. DENV Circulating in Jember

Among 62 dengue-confirmed samples, the serotype of DENV could be determined from 61 samples using RT-PCR, leaving one sample with a negative RT-PCR result but positive for the NS1 antigen. All four serotypes of DENV were detected with the predominance of DENV-4 in 43 (70.5%) of the samples, followed by 12 DENV-3 (19.7%), and three samples (4.9%) for both DENV-1 and DENV-2 (Figure 3). Within the time period between January and March 2020, where a sudden increase in cases was observed, a total number of 12 samples tested positive by RT-PCR and serotyping determined the presence of DENV-1, -2, -3, and -4 in two (16.7%), three (25.0%), three (25.0%), and four (33.3%) samples, respectively.

3.5. DENV Serotypes and Their Clinical Correlates

There was no correlation between the infecting DENV serotypes and patient's age, gender, clinical manifestation, and infection status. Platelet count was found to be correlated with DENV serotypes ($p = 0.008$) with post hoc analysis showing the lower platelet count in DENV-1 samples ($p = 0.019$) (Table 3). The median platelet count was also significantly lower in patients with secondary infections than in those with primary infections ($p < 0.001$, data not shown). This was in line with the result among probable and confirmed patients (Table 2).

Table 3. Jember DENV serotypes and their clinical correlates.

Parameters	DENV Serotype ($n = 61$)				p -Value
	DENV-1 ($n = 3$)	DENV-2 ($n = 3$)	DENV-3 ($n = 12$)	DENV-4 ($n = 43$)	
Age, Median (IQR)	8 (7–12.5)	11 (7.2–20)	20 (11.8–22.3)	20 (11.5–30.5)	0.336 ^a
Gender					
Male	3	1	4	25	0.143 ^b
Female	0	2	8	18	
Clinical manifestation					
DF	0	1	10	33	0.011 ^{b,c}
DHF	3	2	2	10	
Infection status					
Primary	1	1	12	36	0.004 ^{b,d}
Secondary	2	2	0	6	
Hematology, Median (IQR)					
Platelet ($\times 10^3/\mu\text{L}$)	20 (15.5–20.5)	73 (53.5–86)	185 (118.8–194)	143 (76–227.5)	0.008 ^{a,e}
Hematocrit (%)	2.90 (2.65–3.25)	2.60 (2.50–3.20)	3.65 (2.90–4.60)	4.64 (2.95–8.50)	0.529 ^a
WBC ($\times 10^3/\mu\text{L}$)	47.5 (43.2–47.8)	44.4 (41.6–45.2)	42.0 (38.3–45.6)	41.5 (38.6–45.1)	0.204 ^a

IQR, interquartile range. ^a Kruskal–Wallis test. ^b Pearson's Chi-squared test. ^c Post-hoc Bonferroni: $p = 0.112$ (DENV-1—others), $p = 1.000$ (DENV-2—others), $p = 1.000$ (DENV-3—others), $p = 1.000$ (DENV-4—others). ^d Post-hoc Bonferroni: $p = 0.338$ (DENV-1—others), $p = 0.338$ (DENV-2—others), $p = 0.776$ (DENV-3—others), $p = 1.000$ (DENV-4—others). ^e Post-hoc Dunn: $p = 0.019$ (DENV-1—others), $p = 0.438$ (DENV-2—others), $p = 0.660$ (DENV-3—others), $p = 1.000$ (DENV-4—others).

3.6. Non-DENV Arboviruses and Other Pathogens Circulating Jember

Testing for viruses other than DENV was conducted for the possible CHIKV and other flavivirus infections. From all 191 samples tested, we did not find any CHIKV infection. Further testing using pan-flavivirus RT-PCR yielded 11 (5.8%) positively detected samples (data not shown). To determine the identity of those flavivirus PCR amplicons, Sanger capillary sequencing was performed using pan-flavivirus primers. All pan-flavivirus-positive samples were confirmed as DENVs.

Tubex testing for typhoid fever diagnosis was also done on 169 samples, 32 (18.9%) of which were positive. Out of those 32 samples, 11 samples also tested positive for DENV and 4 samples also tested positive for dengue antibodies. The remaining 95 samples that were negative for DENV, CHIKV, and had negative or unavailable Tubex results were categorized under “fever of unknown origin (FUO)”. Patients with dengue, typhoid fever, and fever of unknown origin report very similar symptoms. However, when comparing hematology results, cases of dengue infections present with lower platelet counts (Table 4).

Table 4. Clinical features of dengue, typhoid fever, double infections, and fever of unknown origin (FUO).

<i>n</i> (%)	Dengue (<i>n</i> = 64)	Typhoid (<i>n</i> = 17)	Double Dengue-Typhoid (<i>n</i> = 15)	FUO (<i>n</i> = 95)	Pearson's Chi Square Test
Headache	50 (78.1)	14 (82.4)	14 (93.3)	81 (85.3)	0.404
Retro-orbital Pain	17 (26.6)	4 (23.5)	1 (6.7)	25 (26.3)	0.402
Myalgia	35 (54.7)	9 (52.9)	4 (26.7)	52 (54.7)	0.218
Arthralgia	25 (39.1)	5 (29.4)	2 (13.3)	37 (38.9)	0.225
Rash	3 (4.7)	0 (0)	0 (0)	10 (10.5)	0.182
Stomachache	27 (42.2)	7 (41.2)	4 (26.7)	35 (36.8)	0.713
Vomiting	23 (35.9)	10 (58.8)	6 (40.0)	41 (43.2)	0.382
Bleeding	8 (12.5)	4 (23.5)	3 (20.0)	12 (12.6)	0.577
Positive Tourniquet	11 (17.2)	0 (0)	2 (13.3)	7 (7.4)	0.097
					One-Way ANOVA
Platelet ($\times 10^3/\mu\text{L}$)	140 (71–223)	174 (135–223)	136 (74–181)	171 (138–254)	0.005 *
Hematocrit (%)	42 (38.9–45.0)	36 (34.2–41.0)	42 (35.5–45.6)	40 (36.6–44.0)	0.249
WBC ($\times 10^3/\mu\text{L}$)	4.4 (2.85–7.13)	5.6 (4.63–7.10)	4.0 (3.25–6.05)	5.6 (4.22–8.35)	0.119

* Post-hoc Dunn Test: $p = 0.701$ (Dengue vs. Double Infection), $p = 0.016$ (Dengue vs. FUO), $p = 0.062$ (Double Infection vs. FUO), $p = 0.299$ (Dengue vs. Typhoid), $p = 0.304$ (Double Infection vs. Typhoid), $p = 0.779$ (FUO vs. Typhoid).

3.7. DENV Genotypes in Jember

We have successfully generated 22 full-length E gene sequences from Jember DENV isolates. The sequences were deposited in the GenBank repository, with accession numbers shown in Table 5. The full-length E gene sequence is 1485 nt for DENV-1, -2, and -4, and 1479 nt for DENV-3.

Table 5. Samples with sequenced DENV E genes and their genotype grouping.

No.	Sample ID	Serotype	Genotype	Age (y)	Manifestation	Accession No.
1.	JBB-140	DENV-1	I	8	DHF	MT377728
2.	JBB-145	DENV-2	Cosmopolitan	4	DHF	MT377729
3.	JBB-015	DENV-3	I	1	DF	MT377730
4.	JBB-021	DENV-3	I	8	DF	MT377731
5.	JBB-022	DENV-3	I	2	DF	MT377732
6.	JBB-028	DENV-3	I	21	DF	MT377733
7.	JBB-138	DENV-3	I	35	DHF	MT377734
8.	JBB-004	DENV-4	II	40	DF	MT377735
9.	JBB-007	DENV-4	II	16	DHF	MT377736
10.	JBB-008	DENV-4	II	26	DF	MT377737
11.	JBB-023	DENV-4	II	38	DF	MT377738
12.	JBB-024	DENV-4	II	7	DF	MT377739
13.	JBB-037	DENV-4	II	62	DF	MT377740
14.	JBB-042	DENV-4	II	19	DHF	MT377741
15.	JBB-044	DENV-4	II	18	DF	MT377742
16.	JBB-045	DENV-4	II	12	DF	MT377743
17.	JBB-049	DENV-4	II	18	DF	MT377744
18.	JBB-050	DENV-4	II	11	DHF	MT377745
19.	JBB-051	DENV-4	II	8	DHF	MT377746
20.	JBB-055	DENV-4	II	42	DF	MT377747
21.	JBB-072	DENV-4	II	5	DF	MT377748
22.	JBB-136	DENV-4	II	38	DF	MT377749

We were only able to sequence one DENV-1 isolate from Jember. The isolate was grouped into the Genotype I of DENV-1 and showed close-relatedness to strains from Samarinda, East Kalimantan [35], and other strains from cities in Indonesia, i.e., Bali [36], Yogyakarta [37], Jambi [38], and Purwokerto, Central Java [39], along with strains from imported cases to Taiwan and China and an isolate from Singapore (Figure 4). The Jember DENV-1 isolate was likely becoming the basal taxon for

clade-consisting strains from Samarinda, Bali, and Singapore, with the time to the most recent common ancestor (tmrca) estimated in 2014. Within the larger clade, Surabaya strains were also grouped. The Genotype I was estimated to emerge circa 2004. Unlike the previous studies by our group and others in Surabaya, the presence of Genotype IV in Jember was not detected.




Figure 4. The Maximum Clade Credibility (MCC) phylogenetic tree of DENV-1 genotype I and IV strains generated by BEAST Bayesian inference method with TrN + G evolution model calculated using E gene sequences. The red labels indicate isolates from Jember, while blue labels indicate strains from other cities in Indonesia and green labels indicate strains from the nearby Surabaya city. The number in the node indicates the posterior probability of that particular cluster, with values higher than 0.5 shown. Time to the most recent common ancestor (tmrca) is indicated as median year (95% Highest Posterior Density (HPD)).

Similarly, only one isolate of DENV-2 was successfully sequenced and genetically grouped to the Cosmopolitan genotype of DENV-2. This isolate from Jember was forming a monophyletic group with Indonesian strains from Balikpapan, East Kalimantan [35] with the tmrca estimated at 2011 (Figure 5). However, we observed that the isolate was relatively distinct to DENV-2 from other cities in Indonesia. Moreover, the Jember isolate was quite distinct and evolutionarily separated for years from Surabaya strains.




Figure 5. The Maximum Clade Credibility (MCC) phylogenetic tree of DENV-2 Cosmopolitan genotype generated by BEAST Bayesian inference method with TrN + G evolution model calculated using E gene sequences. The red labels indicate isolates from Jember while blue labels indicate strains from other cities in Indonesia and green labels indicate strains from nearby Surabaya city. The number in the node indicates the posterior probability of that particular cluster, with values higher than 0.5 shown. Time to the most recent common ancestor (tmrca) is indicated as median year (95% Highest Posterior Density (HPD)).

Among 12 samples detected as DENV-3, we successfully sequenced the E gene of five (41.7%) isolates. The DENV-3 isolates from Jember were classified as Genotype I (Figure 6). Further grouping into clades were detected with the first clade consisting of four Jember isolates grouped together in a monophyletic clade supported by a strong posterior, which emerged circa 2013 to 2015. These isolates were closely related to strains from the Indonesian cities of Samarinda and Balikpapan, East Kalimantan [35] and Makassar, South Sulawesi [17], and strains related to dengue cases in Taiwan. The second clade consists of one isolate of Jember DENV-3, which also shared a relationship with strains from East Kalimantan and a strain from an imported case in Taiwan with a tmrca of around the year 2013. A strain from Surabaya is located in the basal position of the first clade and shared the same common ancestor as a strain which emerged around 2012. Another Surabaya strain was also identified

in the basal position, with a tmrca estimated to be in 2009, sharing the common ancestor with both clades where Jember isolates were located.




Figure 6. The Maximum Clade Credibility (MCC) phylogenetic tree of DENV-3 Genotype I generated by BEAST Bayesian inference method with a TrN + G evolution model calculated using E gene sequences. The red labels indicate isolates from Jember, while blue labels indicate strains from other cities in Indonesia and green labels indicate strains from nearby Surabaya city. The number in the node indicates the posterior probability of that particular cluster, with values higher than 0.5 shown. Arabic numbers depict further grouping into DENV lineages. Time to the most recent common ancestor (tmrca) is indicated as median year (95% Highest Posterior Density (HPD)).

As the predominant serotype, all DENV-4 isolates from Jember were grouped into Genotype II (Figure 7). We have generated sequences from 15 out of 43 (34.9%) isolates confirmed as DENV-4.

Further grouping into three monophyletic clades were observed. Clade 1 consists of nine Jember DENV-4 isolates grouped together with a strain of imported cases to Taiwan and other strains from cities in Indonesia, i.e., Makassar, Bali [40], and Sukabumi, West Java [41]. These taxa were estimated to have a tmrca around 2012. The second clade grouped four Jember DENV-4 isolates along with strains from Taiwan, and the Indonesian cities of Bali and Jambi. This clade was estimated to have a younger tmrca in 2015. The other clade—Clade 3—showed a relatively distinct grouping with two Jember isolates displaying a close relationship with strains from Jambi. The estimated tmrca for this particular clade is around 2008.




Figure 7. The Maximum Clade Credibility (MCC) phylogenetic tree of DENV-4 Genotype II generated by BEAST Bayesian inference method with TrN + G evolution model calculated using E gene sequences. The red labels indicate isolates from Jember, while blue labels indicate strains from other cities in Indonesia and green labels indicate strains from nearby Surabaya city. The number in the node indicates the posterior probability of that particular cluster, with values higher than 0.5 shown. Arabic numbers depict further grouping into DENV lineages. Time to the most recent common ancestor (tmrca) is indicated as median year (95% Highest Posterior Density (HPD)).

The molecular clock analysis inferred the evolutionary parameters for DENV isolates from Jember. The age for each DENV serotype tree was estimated at 83, 41, 24, and 29 years for DENV-1, DENV-2, DENV-3, and DENV-4, respectively. Higher evolutionary rates were recorded for DENV-3 and DENV-4 trees (Table 6). However, the rates were still within the normal range of DENV evolutionary rates [34]. Looking at Jember isolates, the evolutionary rate of the DENV isolates were similar, or within the range of, the median rate of the trees. However, DENV-3 isolates from Jember showed distinctively high evolutionary rates above the range of the DENV-3 tree.

Table 6. Evolutionary rates of DENV from Jember.

Serotype	DENV Tree			Jember DENV Isolate(s)	
	<i>n</i>	Tree Age (Years)	Median Rate $\times 10^{-4}$ (95% HPD)	<i>n</i>	Mean Rate ($\times 10^{-4}$) \pm STDEV
DENV-1	60	83	7.7 (5.9–9.7)	1	7.7
DENV-2	60	41	6.5 (4.2–9.4)	1	7.2
DENV-3	60	24	11.2 (9.0–13.4)	5	15.5 \pm 4.2
DENV-4	60	29	10.0 (8.2–11.8)	15	9.9 \pm 0.17

4. Discussion

Dengue and other arbovirus molecular surveillance studies were performed in Jember regency, the third largest city in East Java Province, Indonesia. This 11-month study aimed to monitor the transmission of DENV and other arboviruses in the region, also describing the characteristics of dengue outbreak based on clinical, serological, and viral genetic data. Such monitoring and detailed analysis of clinical cases is essential in less-developed, and thus under-studied, areas such as Jember, as there is lack of historical data documenting dengue in this urban city.

Within the 11-month period, a total of 191 samples of dengue-suspected patients were collected. Although lower dengue IR in Jember was documented compared to East Java province and national rates, there was a sudden increase of dengue incidence in the early months of 2020 (Figure 2A). Looking deeper into this time period, the dengue cases peaked in February 2020. Nearly 30% of our samples were collected in the 3-month period around this peak (from January to March). Moreover, from those 55 samples, there are 12 (21.8%) samples which tested positive using RT-PCR with all four DENV serotypes detected. The persistent incidence of dengue in Jember, the routine annual outbreak happening in the regency, and the sudden increase of cases during the monsoon rainy season, need further investigation.

The majority of patients attending six sample collection sites were children under 10 y.o (Figure 2B). This is similar to our previous dengue disease data gathered in Surabaya [14]. The patients were seeking clinical attention mostly due to symptoms of headache, myalgia, and vomiting, with some patients having bleeding manifestations. Looking at the dengue burden in Jember, we observed a percentage of 41.4% burden with confirmed cases making up 32.5%, reflecting the relatively high burden imposed on the Jember population. Out of DENV-negative samples, we did not detect any other arbovirus infections. This may be due to the low prevalence of Chikungunya [22,42] and Zika [25] infections in Indonesia. However, we detected cases of typhoid fever with a positivity of 18.9%, including those also infected with DENV, with very similar reported symptoms and laboratory results. This finding confirms that typhoid fever is considered to be one of the causative agents of febrile illness and should be considered in the clinical setting as a differential or an additional diagnosis of dengue [43].

Exploring the clinical data documented during the study, we observed that more patients presented with milder DF compared to DHF manifestations. Furthermore, when looking at the infection status, the higher percentage of primary infection was evident. The pattern of milder dengue manifestations being correlated with primary infection was also reported in a dengue surveillance study in Jambi in 2015 [38]. On the other hand, secondary infections were more likely to present with DHF manifestation in Jember (Table 2). This is consistent with reports describing the correlation between secondary

infection and the more severe manifestation of dengue in patients [44]. Moreover, secondary infection is correlated with a lower platelet count in Jember patients. It has been reported that platelet-associated IgG (PAIgG) formation involving anti-dengue virus IgG induces transient thrombocytopenia during the acute phase of secondary dengue virus infection [45], which may explain the correlation observed.

All four serotypes of DENV were found to be circulating in Jember. Strikingly, we found a predominance of DENV-4 with some additional DENV-3 cases, along with a few cases of DENV-1 and DENV-2 (Figure 3). While DENV-4 is known to circulate in East and South East Asia [9], analysis of previous dengue outbreaks in Indonesia have almost exclusively highlighted DENV-1 to -3 as the causative pathogens [12,14,17,19,20], with only one report of an outbreak in Jakarta based on a limited number of cases [21]. Furthermore, the predominant serotype in the nearby city of Surabaya in 2012 was documented as DENV-1 [14,15]. Here, we observed a different serotype distribution in Jember. This is the first data of serotype distribution in Jember, since there is no historical published data for this regency.

The epidemic of DENV-4 in Jember may be caused by several possible considerations. First, this serotype may have been newly introduced to the area from other regions in Indonesia [46]. The low herd immunity in Jember (26.6%) may easily have facilitated the local transmission of this “benign” serotype. Secondly, this serotype may have re-emerged from local endemic viruses, as previously described in Jakarta in the case of DENV-3 [46]. Another alternative is the occurrence of a serotype switch, as has happened in the nearby city of Surabaya [12,13]. The introduction of a new DENV serotype in a population with a low herd immunity would lead to a high ratio of primary to secondary infections, even in an area in which other serotypes commonly circulate. This seems to be the case here, as there is an overrepresentation of primary infections in this outbreak (73.1%), demonstrating a relatively low dengue herd immunity in the area. By comparison, other studies of DENV outbreaks in Java, such as those in Semarang and Jakarta, found the rates of primary infections to be only 23% and 18.7% respectively [20,47], showing the hyperendemicity of DENV across the island. All four DENV serotypes were detected in our study, which may lead to future outbreaks due to serotype-shift. Since secondary infections are known to lead to more severe clinical manifestations than primary infections due to antibody-dependent enhancement (ADE), possible future outbreaks in Jember may lead to larger proportions of the population with severe disease. Interestingly, despite the widespread nature of DENV-3 in Java, all of the confirmed cases of this serotype in this study were primary infections (Table 3). This indicates that there may not have been a consistent circulation of this serotype in the regency for the past few years, long enough for existing immunity to be lost [48,49]. Following this absence of DENV-3, a recent reintroduction of the virus would provide an explanation for the new primary infections.

Molecular surveillance of DENV is essential in a case such as this, as the emergence of new serotypes, genotypes, or lineages of DENV or the introduction of one of these into a naïve population has been shown to significantly increase the incidence of dengue, as well as the severity of the symptoms [16,29]. There is also some data to suggest that different serotypes are linked to a varying severity of the disease [50–52]. While our data did not show a significant correlation between serotype and disease severity (Table 3), this may be attributable to the low number of cases of DENV-1 and -2. However, the fact that DENV-4 was the predominant serotype during the epidemic of mild dengue in Jember is in line with the opinion that considered DENV-4 as the most mild of the four DENV serotypes [4].

Phylogenetic analyses revealed the close relationship between Jember isolates and strains from other cities in Indonesia. Surprisingly, the genetic distance analysis revealed that there was no direct relationship between Jember isolates and strains from Surabaya, the nearest large urban city to Jember in the East Java province. The DENV isolates from Jember were likely to be more related to strains from Samarinda and Balikpapan, both cities in East Kalimantan, from a dengue surveillance study conducted in recent years [35]. Further connections between these cities may need to be explored more extensively. However, traces of transmission and the influence of Surabaya city on Jember may be seen

in the presence of Surabaya strains in the basal position of clades leading to Jember isolates. Together, the data shows that the Jember DENV isolates were local endemic viruses circulating in the region, with DENV-4 re-emerging, likely due to a low herd immunity.

The limitations of this study were related to the sampling sites, which were focused in the Jember city center area and were not spread throughout all the areas of the regency. The distinct proportion of DENV serotypes also lowered the power of the statistical analysis, especially for DENV-1 and DENV-2 samples. Moreover, we did not assess the contribution of mosquito vectors on the transmission of dengue in Jember. Nevertheless, our data provides important epidemiological information that will be useful for future surveillance.

In summary, we reveal the first comprehensive clinical and virological data of the DENV-4 outbreak in Indonesia in the Jember regency. Our study highlights the correlation of this outbreak with primary infections and milder forms of dengue. Continuous dengue surveillance is important to have more complete data on dengue in Jember and surrounding areas and to increase preparedness for potential outbreaks in the future.

Supplementary Materials: The following are available online at <http://www.mdpi.com/1999-4915/12/9/913/s1>, Table S1: Multivariate analysis of dengue symptoms and the potentially affecting covariates, Figure S1: Phylogenetic tree and genotype classification of DENV-1 isolate from Jember, East Java (marked with red dot) with other reference strains, Figure S2: Phylogenetic tree and genotype classification of DENV-2 isolate from Jember, East Java (marked with red dot) with other reference strains, Figure S3: Phylogenetic tree and genotype classification of DENV-3 isolates from Jember, East Java (marked with red dots) with other reference strains, Figure S4: Phylogenetic tree and genotype classification of DENV-4 isolates from Jember, East Java (marked with red dot) with other reference strains.

Author Contributions: Conceived and designed the study: A.A., R.T.S. Performed sample collection and clinical data analysis: P.W., M.F., E.P.S., B.J.W., F.K.N.H., M.S.S., D.H., A.R., N.Z., S.N.T. Performed laboratory experiments: B.Y., J.T.O.S., M.F., E.P.S., B.J.W., F.K.N.H. Data analysis: B.Y., J.T.O.S., M.S.S. Wrote the paper: B.Y., J.T.O.S., M.S.S., R.T.S. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by a research grant from the Universitas Airlangga (Grant number 965/UN3.14/LT/2019) to A.A.

Acknowledgments: We would like to thank study participants and medical staff from hospitals/healthcare centers involved in the study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Wei, K.; Li, Y. Global evolutionary history and spatio-temporal dynamics of dengue virus type 2. *Sci. Rep.* **2017**, *7*, 45505. [[CrossRef](#)] [[PubMed](#)]
2. Henchal, E.A.; Putnak, J.R. The dengue viruses. *Clin. Microbiol. Rev.* **1990**, *3*, 376–396. [[CrossRef](#)] [[PubMed](#)]
3. Guzman, M.G.; Halstead, S.B.; Artsob, H.; Buchy, P.; Farrar, J.; Gubler, D.J.; Hunsperger, E.; Kroeger, A.; Margolis, H.S.; Martínez, E.; et al. Dengue: A continuing global threat. *Nat. Rev. Microbiol.* **2010**, *8*, S7–S16. [[CrossRef](#)] [[PubMed](#)]
4. Gubler, D.J. Dengue and dengue hemorrhagic fever. *Clin. Microbiol. Rev.* **1998**, *11*, 480–496. [[CrossRef](#)] [[PubMed](#)]
5. World Health Organization. *Comprehensive Guidelines for Prevention and Control of Dengue and Dengue Haemorrhagic Fever*; Revised and Expanded; WHO Regional Office for South-East Asia: New Delhi, India, 2011; ISBN 978-92-9022-387-0.
6. Holmes, E.C.; Twiddy, S.S. The origin, emergence and evolutionary genetics of dengue virus. *Infect. Genet. Evol.* **2003**, *3*, 19–28. [[CrossRef](#)]
7. Rico-Hesse, R. Microevolution and virulence of dengue viruses. *Adv. Virus Res.* **2003**, *59*, 315–341.
8. Weaver, S.C.; Vasilakis, N. Molecular Evolution of Dengue Viruses: Contributions of Phylogenetics to Understanding the History and Epidemiology of the Preeminent Arboviral Disease. *Infect. Genet. Evol.* **2009**, *9*, 523–540. [[CrossRef](#)]
9. Sang, S.; Liu-Helmerson, J.; Quam, M.B.M.; Zhou, H.; Guo, X.; Wu, H.; Liu, Q. The evolutionary dynamics of DENV 4 genotype I over a 60-year period. *PLoS Negl. Trop. Dis.* **2019**, *13*. [[CrossRef](#)]

10. Sumarmo. Dengue haemorrhagic fever in Indonesia. *Southeast Asian J. Trop. Med. Public Health* **1987**, *18*, 269–274.
11. Ministry of Health of the Republic of Indonesia *Indonesia Health Profile 2018*; Ministry of Health: Jakarta, Indonesia, 2019.
12. Kotaki, T.; Yamanaka, A.; Mulyatno, K.C.; Churrotin, S.; Labiqah, A.; Sucipto, T.H.; Soegijanto, S.; Kameoka, M.; Konishi, E. Continuous dengue type 1 virus genotype shifts followed by co-circulation, clade shifts and subsequent disappearance in Surabaya, Indonesia, 2008–2013. *Infect. Genet. Evol.* **2014**, *28*, 48–54. [[CrossRef](#)]
13. Kotaki, T.; Yamanaka, A.; Mulyatno, K.C.; Churrotin, S.; Sucipto, T.H.; Labiqah, A.; Ahwanah, N.L.F.; Soegijanto, S.; Kameoka, M.; Konishi, E. Divergence of the dengue virus type 2 Cosmopolitan genotype associated with two predominant serotype shifts between 1 and 2 in Surabaya, Indonesia, 2008–2014. *Infect. Genet. Evol.* **2016**, *37*, 88–93. [[CrossRef](#)] [[PubMed](#)]
14. Wardhani, P.; Aryati, A.; Yohan, B.; Trimarsanto, H.; Setianingsih, T.Y.; Puspitasari, D.; Arfijanto, M.V.; Bramantono, B.; Suharto, S.; Sasmono, R.T. Clinical and virological characteristics of dengue in Surabaya, Indonesia. *PLoS ONE* **2017**, *12*, e0178443. [[CrossRef](#)] [[PubMed](#)]
15. Sucipto, T.H.; Kotaki, T.; Mulyatno, K.C.; Churrotin, S.; Labiqah, A.; Soegijanto, S.; Kameoka, M. Phylogenetic Analysis of Dengue Virus in Bangkalan, Madura Island, East Java Province, Indonesia. *J. Trop. Med.* **2018**, *2018*, 8127093. [[CrossRef](#)] [[PubMed](#)]
16. Klungthong, C.; Putnak, R.; Mammen, M.P.; Li, T.; Zhang, C. Molecular genotyping of dengue viruses by phylogenetic analysis of the sequences of individual genes. *J. Virol. Methods* **2008**, *154*, 175–181. [[CrossRef](#)] [[PubMed](#)]
17. Sasmono, R.T.; Wahid, I.; Trimarsanto, H.; Yohan, B.; Wahyuni, S.; Hertanto, M.; Yusuf, I.; Mubin, H.; Ganda, I.J.; Latief, R.; et al. Genomic analysis and growth characteristic of dengue viruses from Makassar, Indonesia. *Infect. Genet. Evol.* **2015**, *32*, 165–177. [[CrossRef](#)] [[PubMed](#)]
18. Gallichotte, E.N.; Baric, T.J.; Nivarthi, U.; Delacruz, M.J.; Graham, R.; Widman, D.G.; Yount, B.L.; Durbin, A.P.; Whitehead, S.S.; de Silva, A.M.; et al. Genetic Variation between Dengue Virus Type 4 Strains Impacts Human Antibody Binding and Neutralization. *Cell Rep.* **2018**, *25*, 1214–1224. [[CrossRef](#)]
19. Harapan, H.; Michie, A.; Yohan, B.; Shu, P.Y.; Mudatsir, M.; Sasmono, R.T.; Imrie, A. Dengue viruses circulating in Indonesia: A systematic review and phylogenetic analysis of data from five decades. *Rev. Med. Virol.* **2019**, *29*, e2037. [[CrossRef](#)]
20. Lestari, C.S.W.; Yohan, B.; Yunita, A.; Meutiawati, F.; Hayati, R.F.; Trimarsanto, H.; Sasmono, R.T. Phylogenetic and evolutionary analyses of dengue viruses isolated in Jakarta, Indonesia. *Virus Genes* **2017**, *53*, 778–788. [[CrossRef](#)]
21. Dewi, B.E.; Naiggolan, L.; Putri, D.H.; Rachmayanti, N.; Albar, S.; Indriastuti, N.T.; Sjamsuridzal, W.; Sudiro, T.M. Characterization of dengue virus serotype 4 infection in Jakarta, Indonesia. *Southeast Asian J. Trop. Med. Public Health* **2014**, *45*, 53–61.
22. Harapan, H.; Michie, A.; Mudatsir, M.; Nusa, R.; Yohan, B.; Wagner, A.L.; Sasmono, R.T.; Imrie, A. Chikungunya virus infection in Indonesia: A systematic review and evolutionary analysis. *BMC Infect. Dis.* **2019**, *19*, 243. [[CrossRef](#)]
23. Priyamvada, L.; Hudson, W.; Ahmed, R.; Wrammert, J. Humoral cross-reactivity between Zika and dengue viruses: Implications for protection and pathology. *Emerg. Microbes Infect.* **2017**, *6*, e33. [[CrossRef](#)] [[PubMed](#)]
24. Vaughn, D.W.; Nisalak, A.; Kalayanarooj, S.; Solomon, T.; Dung, N.M.; Cuzzubbo, A.; Devine, P.L. Evaluation of a Rapid Immunochromatographic Test for Diagnosis of Dengue Virus Infection. *J. Clin. Microbiol.* **1998**, *36*, 234–238. [[CrossRef](#)] [[PubMed](#)]
25. Sasmono, R.T.; Dhenni, R.; Yohan, B.; Pronyk, P.; Hadinegoro, S.R.; Soepardi, E.J.; Ma'roef, C.; Satari, H.; Menzies, H.; Hawley, W.; et al. Endemicity of Zika virus in Indonesia: Serological evidence in pediatric population. *Int. J. Infect. Dis.* **2019**, *79*, 13–14. [[CrossRef](#)]
26. BPS-Statistics of Jember Regency. *Jember Regency in Figures 2020 Delivering Data to Inform Development Planning*; BPS-Statistics of Jember Regency: Jember, Indonesia, 2020; ISBN 978-602-71310-3-3.
27. Kuno, G.; Chang, G.J.; Tsuchiya, K.R.; Karabatsos, N.; Cropp, C.B. Phylogeny of the genus *Flavivirus*. *J. Virol.* **1998**, *72*, 73–83. [[CrossRef](#)] [[PubMed](#)]
28. Kumar, S.; Stecher, G.; Li, M.; Nnyaz, C.; Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* **2018**, *35*, 1547–1549. [[CrossRef](#)]

29. Goncalvez, A.P.; Escalante, A.A.; Pujol, F.H.; Ludert, J.E.; Tovar, D.; Salas, R.A.; Liprandi, F. Diversity and evolution of the envelope gene of dengue virus type 1. *Virology* **2002**, *303*, 110–119. [[CrossRef](#)]
30. Twiddy, S.S.; Farrar, J.J.; Vinh Chau, N.; Wills, B.; Gould, E.A.; Gritsun, T.; Lloyd, G.; Holmes, E.C. Phylogenetic relationships and differential selection pressures among genotypes of dengue-2 virus. *Virology* **2002**, *298*, 63–72. [[CrossRef](#)]
31. Lanciotti, R.S.; Lewis, J.G.; Gubler, D.J.; Trent, D.W. Molecular evolution and epidemiology of dengue-3 viruses. *J. Gen. Virol.* **1994**, *75*, 65–75. [[CrossRef](#)]
32. Lanciotti, R.S.; Gubler, D.J.; Trent, D.W. Molecular evolution and phylogeny of dengue-4 viruses. *J. Gen. Virol.* **1997**, *78*, 2279–2284. [[CrossRef](#)]
33. Bouckaert, R.; Vaughan, T.G.; Barido-Sottani, J.; Duchêne, S.; Fourment, M.; Gavryushkina, A.; Heled, J.; Jones, G.; Kühnert, D.; De Maio, N.; et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS Comput. Biol.* **2019**, *15*, e1006650. [[CrossRef](#)]
34. Costa, R.L.; Voloch, C.M.; Schrago, C.G. Comparative evolutionary epidemiology of dengue virus serotypes. *Infect. Genet. Evol.* **2012**, *12*, 309–314. [[CrossRef](#)] [[PubMed](#)]
35. Sasmono, R.T.; Kalalo, L.P.; Trismasih, S.; Denis, D.; Yohan, B.; Hayati, R.F.; Haryanto, S. Multiple introductions of dengue virus strains contribute to dengue outbreaks in East Kalimantan, Indonesia, in 2015–2016. *Virol. J.* **2019**, *16*, 93. [[CrossRef](#)] [[PubMed](#)]
36. Masyeni, S.; Yohan, B.; Somia, I.K.A.; Myint, K.S.A.; Sasmono, R.T. Dengue infection in international travellers visiting Bali, Indonesia. *J. Travel. Med.* **2018**, *25*. [[CrossRef](#)] [[PubMed](#)]
37. Utama, I.M.S.; Lukman, N.; Sukmawati, D.D.; Alisjahbana, B.; Alam, A.; Murniati, D.; Utama, I.M.G.D.L.; Puspitasari, D.; Kosasih, H.; Laksono, I.; et al. Dengue viral infection in Indonesia: Epidemiology, diagnostic challenges, and mutations from an observational cohort study. *PLoS Negl. Trop. Dis.* **2019**, *13*, e0007785. [[CrossRef](#)] [[PubMed](#)]
38. Haryanto, S.; Hayati, R.F.; Yohan, B.; Sijabat, L.; Sihite, I.F.; Fahri, S.; Meutiawati, F.; Halim, J.A.N.; Halim, S.N.; Soebandrio, A.; et al. The molecular and clinical features of dengue during outbreak in Jambi, Indonesia in 2015. *Pathog. Glob. Health* **2016**, *110*, 119–129. [[CrossRef](#)] [[PubMed](#)]
39. Kusmintarsih, E.S.; Hayati, R.F.; Turnip, O.N.; Yohan, B.; Suryaningsih, S.; Pratiknyo, H.; Denis, D.; Sasmono, R.T. Molecular characterization of dengue viruses isolated from patients in Central Java, Indonesia. *J. Infect. Public Health* **2017**. [[CrossRef](#)]
40. Megawati, D.; Masyeni, S.; Yohan, B.; Lestari, A.; Hayati, R.F.; Meutiawati, F.; Suryana, K.; Widarsa, T.; Budiyasa, D.G.; Budiyasa, N.; et al. Dengue in Bali: Clinical characteristics and genetic diversity of circulating dengue viruses. *PLoS Negl. Trop. Dis.* **2017**, *11*, e0005483. [[CrossRef](#)]
41. Nusa, R.; Prasetyowati, H.; Meutiawati, F.; Yohan, B.; Trimarsanto, H.; Setianingsih, Y.; Sasmono, R.T. Molecular surveillance of dengue in Sukabumi, West Java province, Indonesia. *J. Infect. Dev. Ctries* **2014**, *8*, 733–741. [[CrossRef](#)]
42. Kosasih, H.; de Mast, Q.; Widjaja, S.; Sudjana, P.; Antonjaya, U.; Ma'roef, C.; Riswari, S.F.; Porter, K.R.; Burgess, T.H.; Alisjahbana, B.; et al. Evidence for endemic chikungunya virus infections in Bandung, Indonesia. *PLoS Negl. Trop. Dis.* **2013**, *7*, e2483. [[CrossRef](#)]
43. Capeding, M.R.; Chua, M.N.; Hadinegoro, S.R.; Hussain, I.I.H.M.; Nallusamy, R.; Pitisuttithum, P.; Rusmil, K.; Thisyakorn, U.; Thomas, S.J.; Huu Tran, N.; et al. Dengue and other common causes of acute febrile illness in Asia: An active surveillance study in children. *PLoS Negl. Trop. Dis.* **2013**, *7*, e2331. [[CrossRef](#)]
44. Guzman, M.G.; Alvarez, M.; Halstead, S.B. Secondary infection as a risk factor for dengue hemorrhagic fever/dengue shock syndrome: An historical perspective and role of antibody-dependent enhancement of infection. *Arch. Virol.* **2013**, *158*, 1445–1459. [[CrossRef](#)] [[PubMed](#)]
45. Oishi, K.; Inoue, S.; Cinco, M.T.D.D.; Dimaano, E.M.; Alera, M.T.P.; Alfon, J.A.R.; Abanes, F.; Cruz, D.J.M.; Matias, R.R.; Matsuura, H.; et al. Correlation between increased platelet-associated IgG and thrombocytopenia in secondary dengue virus infections. *J. Med. Virol.* **2003**, *71*, 259–264. [[CrossRef](#)] [[PubMed](#)]
46. Ong, S.H.; Yip, J.T.; Chen, Y.L.; Liu, W.; Harun, S.; Lystianingsih, E.; Heriyanto, B.; Beckett, C.G.; Mitchell, W.P.; Hibberd, M.L.; et al. Periodic re-emergence of endemic strains with strong epidemic potential—a proposed explanation for the 2004 Indonesian dengue epidemic. *Infect. Genet. Evol.* **2008**, *8*, 191–204. [[CrossRef](#)] [[PubMed](#)]

47. Fahri, S.; Yohan, B.; Trimarsanto, H.; Sayono, S.; Hadisaputro, S.; Dharmana, E.; Syafruddin, D.; Sasmono, R.T. Molecular Surveillance of Dengue in Semarang, Indonesia Revealed the Circulation of an Old Genotype of Dengue Virus Serotype-1. *PLoS Negl. Trop. Dis.* **2013**, *7*, e2354. [[CrossRef](#)] [[PubMed](#)]
48. Forshey, B.M.; Reiner, R.C.; Olkowski, S.; Morrison, A.C.; Espinoza, A.; Long, K.C.; Vilcarrero, S.; Casanova, W.; Wearing, H.J.; Halsey, E.S.; et al. Incomplete Protection against Dengue Virus Type 2 Re-infection in Peru. *PLoS Negl. Trop. Dis.* **2016**, *10*, e0004398. [[CrossRef](#)]
49. Waggoner, J.J.; Balmaseda, A.; Gresh, L.; Sahoo, M.K.; Montoya, M.; Wang, C.; Abeynayake, J.; Kuan, G.; Pinsky, B.A.; Harris, E. Homotypic Dengue Virus Reinfections in Nicaraguan Children. *J. Infect. Dis.* **2016**, *214*, 986–993. [[CrossRef](#)]
50. Soo, K.M.; Khalid, B.; Ching, S.M.; Chee, H.Y. Meta-Analysis of Dengue Severity during Infection by Different Dengue Virus Serotypes in Primary and Secondary Infections. *PLoS ONE* **2016**, *11*, e0154760. [[CrossRef](#)]
51. Vicente, C.R.; Herbinger, K.H.; Fröschl, G.; Malta Romano, C.; de Souza Areias Cabidelle, A.; Cerutti Junior, C. Serotype influences on dengue severity: A cross-sectional study on 485 confirmed dengue cases in Vitória, Brazil. *BMC Infect. Dis.* **2016**, *16*, 320. [[CrossRef](#)]
52. Yung, C.F.; Lee, K.S.; Thein, T.L.; Tan, L.K.; Gan, V.C.; Wong, J.G.X.; Lye, D.C.; Ng, L.C.; Leo, Y.S. Dengue serotype-specific differences in clinical manifestation, laboratory parameters and risk of severe disease in adults, Singapore. *Am. J. Trop. Med. Hyg.* **2015**, *92*, 999–1005. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Dengue Virus Serotype 4 Is Responsible for the Outbreak of Dengue in East Java City of Jember, Indonesia

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	Marsha S. Santoso, Benediktus Yohan, Dionisius Denis, Rahma F. Hayati et al. "Diagnostic accuracy of 5 different brands of dengue virus non-structural protein 1 (NS1) antigen rapid diagnostic tests (RDT) in Indonesia", <i>Diagnostic Microbiology and Infectious Disease</i> , 2020 Publication	1%
2	www.ncbi.nlm.nih.gov Internet Source	1%
3	www.openagrar.de Internet Source	1%
4	doi.org Internet Source	1%
5	s3.amazonaws.com Internet Source	1%
6	link.springer.com Internet Source	<1%

stacks.cdc.gov

7

Internet Source

<1 %

8

"Treated Plasmodium berghei infected pregnant mice by Andrographis paniculata tablet (AS201-01) decreasing the TLR-4 expression and apoptosis index of placental tissue", Journal of Applied Pharmaceutical Science, 2018

Publication

<1 %

9

www.repository.cam.ac.uk

Internet Source

<1 %

10

www.scdiagnostics.com.au

Internet Source

<1 %

11

res.mdpi.com

Internet Source

<1 %

12

boris.unibe.ch

Internet Source

<1 %

13

Berta Nelly Restrepo, Leidy Diana Piedrahita, Ivony Yireth Agudelo, Gabriel Parra-Henao, Jorge E. Osorio. "Frequency and Clinical Features of Dengue Infection in a Schoolchildren Cohort from Medellin, Colombia", Journal of Tropical Medicine, 2012

Publication

<1 %

14

www.nature.com

Internet Source

<1 %

15 Silvia Riccetti, Alessandro Sinigaglia, Giovanna Desole, Norbert Nowotny, Marta Trevisan, Luisa Barzon. "Modelling West Nile Virus and Usutu Virus Pathogenicity in Human Neural Stem Cells", *Viruses*, 2020
Publication

16 pubmed.ncbi.nlm.nih.gov
Internet Source

17 repository.ubaya.ac.id
Internet Source

18 R. Tedjo Sasmono, Anne-Frieda Taurel, Ari Prayitno, Hermin Sitompul et al. "Dengue virus serotype distribution based on serological evidence in pediatric urban population in Indonesia", *PLOS Neglected Tropical Diseases*, 2018
Publication

19 Rico-Hesse, R.. "Origins of Dengue Type 2 Viruses Associated with Increased Pathogenicity in the Americas", *Virology*, 19970414
Publication

20 edoc.unibas.ch
Internet Source

21 studentsrepo.um.edu.my
Internet Source

- | | | |
|----|--|------|
| 22 | oro.open.ac.uk
Internet Source | <1 % |
| 23 | raw.githubusercontent.com
Internet Source | <1 % |
| 24 | Jiaqi Cao, Hong Deng, Lei Ye, Xuezheng Ma et al. "Epidemiological and clinical characteristics of Dengue virus outbreaks in two regions of China, 2014 – 2015", PLOS ONE, 2019
Publication | <1 % |
| 25 | Soroy Lardo, Yaldiera Utami, Benediktus Yohan, Seri MMU. Tarigan, Widayat Djoko Santoso, Leonard Nainggolan, R. Tedjo Sasmono. "Concurrent infections of dengue viruses serotype 2 and 3 in patient with severe dengue from Jakarta, Indonesia", Asian Pacific Journal of Tropical Medicine, 2016
Publication | <1 % |
| 26 | Isabel Cristina Guerra-Gomes, Bruna Macêdo Gois, Rephany Fonseca Peixoto, Camila Alves Oliveira et al. "Molecular and clinical epidemiological surveillance of dengue virus in Paraíba, Northeast Brazil", Revista da Sociedade Brasileira de Medicina Tropical, 2017
Publication | <1 % |
| 27 | R. Tedjo Sasmono, Isra Wahid, Hidayat Trimarsanto, Benediktus Yohan et al. | <1 % |

"Genomic analysis and growth characteristic of dengue viruses from Makassar, Indonesia", *Infection, Genetics and Evolution*, 2015

Publication

28

Cheng-Fen Yang, Shu-Fen Chang, Tung-Chien Hsu, Chien-Ling Su et al. "Molecular characterization and phylogenetic analysis of dengue viruses imported into Taiwan during 2011-2016", *PLOS Neglected Tropical Diseases*, 2018

Publication

<1 %

29

Himawan Bayu Patriadi. "Chapter 3 Surviving in the Globalized World Through Local Perspectives: Pesantrens and Sustainable Development", Springer Science and Business Media LLC, 2018

Publication

<1 %

30

Mandary, Masomian, Poh. "Impact of RNA Virus Evolution on Quasispecies Formation and Virulence", *International Journal of Molecular Sciences*, 2019

Publication

<1 %

31

Unai Perez-Sautu, Se Hun Gu, Katie Caviness, Dong Hyun Song et al. "A Model for the Production of Regulatory Grade Viral Hemorrhagic Fever Exposure Stocks: From Field Surveillance to Advanced Characterization of SFTSV", *Viruses*, 2020

Publication

<1 %

32	repositori.uji.es Internet Source	<1 %
33	scholarbank.nus.edu.sg Internet Source	<1 %
34	www.dovepress.com Internet Source	<1 %
35	www.jove.com Internet Source	<1 %
36	www.radioprotection.org Internet Source	<1 %
37	www.tm.nagasaki-u.ac.jp Internet Source	<1 %
38	Susan H. Eshleman. "Analysis of <i>pol</i> Integrase Sequences in Diverse HIV Type 1 Strains Using a Prototype Genotyping Assay", AIDS Research and Human Retroviruses, 03/2009 Publication	<1 %
39	academicjournals.org Internet Source	<1 %
40	www.ajtmh.org Internet Source	<1 %
41	Juthamas Phadungsombat, Marco Yung-Cheng Lin, Narinee Srimark, Atsushi Yamanaka et al. "Emergence of genotype Cosmopolitan of dengue virus type 2 and	<1 %

genotype III of dengue virus type 3 in Thailand", PLOS ONE, 2018

Publication

42

Ming Te Yeh, Sara Capponi, Adam Catching, Simone Bianco, Raul Andino. "Mapping Attenuation Determinants in Enterovirus-D68", *Viruses*, 2020

Publication

<1 %

43

Wen-Ting Yeh, Rong-Fu Chen, Lin Wang, Jien-Wei Liu, Men-Fang Shaio, Kuender D. Yang. "Implications of previous subclinical dengue infection but not virus load in dengue hemorrhagic fever", *FEMS Immunology & Medical Microbiology*, 2006

Publication

<1 %

44

cmjournal.biomedcentral.com

Internet Source

<1 %

45

en.wikipedia.org

Internet Source

<1 %

46

www.dgspj.de

Internet Source

<1 %

47

Hoa Thi My Vo, Veasna Duong, Sowath Ly, Quan-Zhen Li, Philippe Dussart, Tineke Cantaert. "Autoantibody Profiling in Plasma of Dengue Virus-Infected Individuals", *Pathogens*, 2020

Publication

<1 %

48

hdl.handle.net

Internet Source

<1 %

49

www.mbaa.com

Internet Source

<1 %

50

Ankita Agarwal, Sudheer Gupta, Tanvi Chincholkar, Vivek Singh et al. "Co-circulation of dengue virus serotypes in Central India: Evidence of prolonged viremia in DENV-2", *Infection, Genetics and Evolution*, 2019

Publication

<1 %

51

Ayu Rahayu, Utari Saraswati, Endah Supriyati, Dian Aruni Kumalawati et al. "Prevalence and Distribution of Dengue Virus in Aedes aegypti in Yogyakarta City before Deployment of Wolbachia Infected Aedes aegypti", *International Journal of Environmental Research and Public Health*, 2019

Publication

<1 %

52

Capeding, Maria Rosario, Mary Noreen Chua, Sri Rezeki Hadinegoro, Ismail I. H. M. Hussain, Revathy Nallusamy, Punnee Pitisuttithum, Kusnandi Rusmil, Usa Thisyakorn, Stephen J. Thomas, Ngoc Huu Tran, Dewa Nyoman Wirawan, In-Kyu Yoon, Alain Bouckenoghe, Yanee Hutagalung, Thelma Laot, and Tram Anh Wartel. "Dengue and Other Common Causes of Acute Febrile Illness in Asia: An

<1 %

Active Surveillance Study in Children", PLoS
Neglected Tropical Diseases, 2013.

Publication

53

Egon A. Ozer, Lacy M. Simons, Olubusuyi M.
Adewumi, Adeola A. Fowotade et al.

"Coincident rapid expansion of two SARS-CoV-
2 lineages with enhanced infectivity in
Nigeria", Cold Spring Harbor Laboratory, 2021

Publication

54

Jéssica Barreto Lopes Silva, Debora
Magalhães Alves, Vanessa Bottino-Rojas,
Thiago Nunes Pereira et al. "Wolbachia and
dengue virus infection in the mosquito *Aedes
fluviatilis* (Diptera: Culicidae)", PLOS ONE,
2017

Publication

55

Pooja Rao, Achappa Basavaprabhu, Suchitra
Shenoy, Nikhil Victor Dsouza, Basavaiah
Sridevi Hanaganahalli, Vaman Kulkarni.
"Correlation of Clinical Severity and
Laboratory Parameters with Various
Serotypes in Dengue Virus: A Hospital-Based
Study", International Journal of Microbiology,
2020

Publication

56

citeseerx.ist.psu.edu

Internet Source

<1 %

<1 %

<1 %

<1 %

57

Internet Source

<1 %

58

reproductive-health-
journal.biomedcentral.com

Internet Source

<1 %

59

www.jbiomedsci.com

Internet Source

<1 %

60

www.kohtaoems.com

Internet Source

<1 %

61

"Proceedings of the 1st International
Conference on Electronics, Biomedical
Engineering, and Health Informatics", Springer
Science and Business Media LLC, 2021

Publication

<1 %

62

Groen, J.. "Diagnostic value of dengue virus-
specific IgA and IgM serum antibody
detection", Microbes and Infection, 199911

Publication

<1 %

63

Leandra Barcelos Figueiredo, Tetsu Sakamoto,
Luiz Felipe Leomil Coelho, Eliseu Soares de
Oliveira Rocha et al. "Dengue Virus 2
American-Asian Genotype Identified during
the 2006/2007 Outbreak in Piau , Brazil
Reveals a Caribbean Route of Introduction
and Dissemination of Dengue Virus in Brazil",
PLoS ONE, 2014

Publication

<1 %

- 64 M. OhAinle, A. Balmaseda, A. R. Macalalad, Y. Tellez et al. "Dynamics of Dengue Disease Severity Determined by the Interplay Between Viral Genetics and Serotype-Specific Immunity", *Science Translational Medicine*, 2011
Publication <1 %
-
- 65 Sneha Singh, Dustin Farr, Ashok Kumar. "Ocular Manifestations of Emerging Flaviviruses and the Blood-Retinal Barrier", *Viruses*, 2018
Publication <1 %
-
- 66 Xiaoying Jia, Jiao Guo, Weirong Yuan, Lingling Sun et al. "Mechanism through which retrocyclin targets flavivirus multiplication", *Cold Spring Harbor Laboratory*, 2020
Publication <1 %
-
- 67 archive.org
Internet Source <1 %
-
- 68 bmccresnotes.biomedcentral.com
Internet Source <1 %
-
- 69 www.ejurnal-analiskesehatan.web.id
Internet Source <1 %
-
- 70 www.lib.kobe-u.ac.jp
Internet Source <1 %
-

71

Angela Ferreira Lopes de Teive e Argolo, Valéria Christina de Rezende Féres, Marli Tenório Cordeiro, Lucimeire Antonelli da Silveira et al. "High frequency of pre-existing neutralizing antibody responses in patients with dengue during an outbreak in Central Brazil", BMC Infectious Diseases, 2016

Publication

<1 %

72

Arindam Sarkar, Debjani Taraphdar, Shyamalendu Chatterjee. "Molecular Typing of Dengue Virus Circulating in Kolkata, India in 2010", Journal of Tropical Medicine, 2012

Publication

<1 %

73

Corbett, K. S., L. Katzelnick, H. Tissera, A. Amerasinghe, A. D. de Silva, and A. M. de Silva. "Preexisting Neutralizing Antibody Responses Distinguish Clinically Inapparent and Apparent Dengue Virus Infections in a Sri Lankan Pediatric Cohort", The Journal of Infectious Diseases, 2014.

Publication

<1 %

74

Jeyanthi Suppiah, Siew-Mooi Ching, Syafinaz Amin-Nordin, Lailatul-Akmar Mat-Nor et al. "Clinical manifestations of dengue in relation to dengue serotype and genotype in Malaysia: A retrospective observational study", PLOS Neglected Tropical Diseases, 2018

<1 %

75

Karina Rocha Dutra, Betânia Paiva Drumond, Izabela Maurício de Rezende, Maurício Lacerda Nogueira et al. "Molecular surveillance of dengue in Minas Gerais provides insights on dengue virus 1 and 4 circulation in Brazil", *Journal of Medical Virology*, 2017

Publication

<1 %

76

Letícia Borges da Silva HEINEN, Nayara ZUCHI, Belgath Fernandes CARDOSO, Marcelo Adriano Mendes dos SANTOS et al. "DENGUE OUTBREAK IN MATO GROSSO STATE, MIDWESTERN BRAZIL", *Revista do Instituto de Medicina Tropical de São Paulo*, 2015

Publication

<1 %

77

Mansyur Arif, Patricia Tauran, Herman Kosasih, Ninny Meutia Pelupessy et al. "Chikungunya in Indonesia: Epidemiology and diagnostic challenges", *PLOS Neglected Tropical Diseases*, 2020

Publication

<1 %

78

Syed Fazil Ahamed, Rosario Vivek, Shalini Kotabagi, Kaustuv Nayak, Anmol Chandele, Murali-Krishna Kaja, Anita Shet. "Enhancing the sensitivity of Dengue virus serotype detection by RT-PCR among infected children in India", *Journal of Virological Methods*, 2017

Publication

<1 %

79 Umesh Jayarajah, Pradeep K de Silva, Priyankara Jayawardana, Upul Dissanayake et al. "Pattern of dengue virus infections in adult patients from Sri Lanka", Transactions of The Royal Society of Tropical Medicine and Hygiene, 2018
Publication

<1 %

80 Wellington Sun. "Phase 2 clinical trial of three formulations of tetravalent live-attenuated dengue vaccine in flavivirus-naïve adults", Human Vaccines, 01/01/2009
Publication

<1 %

81 apps.searo.who.int
Internet Source

<1 %

82 core.ac.uk
Internet Source

<1 %

83 ecancer.org
Internet Source

<1 %

84 hal-pasteur.archives-ouvertes.fr
Internet Source

<1 %

85 jidc.org
Internet Source

<1 %

86 parasitesandvectors.biomedcentral.com
Internet Source

<1 %

87 profiles.stanford.edu
Internet Source

<1 %

88 tel.archives-ouvertes.fr <1 %
Internet Source

89 test.dovepress.com <1 %
Internet Source

90 worldwidescience.org <1 %
Internet Source

91 www.scielo.br <1 %
Internet Source

92 Ericel Hernández-García, Maria de Lourdes Muñoz, Randy E. David, Gerardo Pérez-Ramírez et al. "Epidemiological implications of the genetic diversification of dengue virus (DENV) serotypes and genotypes in Mexico", *Infection, Genetics and Evolution*, 2020
Publication

93 Harapan Harapan, Amanda Yufika, Samsul Anwar, Haypheng Te, Hamzah Hasyim, Roy Nusa, Pandji Wibawa Dhewantara, Mudatsir Mudatsir. "Effects of El Niño Southern Oscillation and Dipole Mode Index on Chikungunya Infection in Indonesia", *Tropical Medicine and Infectious Disease*, 2020
Publication

94 A. Imrie, Z. Zhao, S. N. Bennett, P. Kitsutani, M. Laille, P. Effler. "Molecular epidemiology of dengue in the Pacific: introduction of two

distinct strains of dengue virus type-1 into Hawaii", *Annals of Tropical Medicine & Parasitology*, 2013

Publication

95

Daniel H. Libraty. "A Comparative Study of Leptospirosis and Dengue in Thai Children", *PLoS Neglected Tropical Diseases*, 12/26/2007

Publication

96

Harapan Harapan, Alice Michie, Mudatsir Mudatsir, Roy Nusa et al. "Chikungunya virus infection in Indonesia: a systematic review and evolutionary analysis", *BMC Infectious Diseases*, 2019

Publication

97

Jiang, Liyun, Xinwei Wu, Yejian Wu, Zhijun Bai, Qinglong Jing, Lei Luo, Zhiqiang Dong, Zhicong Yang, Yang Xu, Yimin Cao, Biao Di, Yulin Wang, and Ming Wang. "Molecular epidemiological and virological study of dengue virus infections in Guangzhou, China, during 2001–2010", *Virology Journal*, 2013.

Publication

98

Kaifa Wei, Yuhan Li. "Global evolutionary history and spatio-temporal dynamics of dengue virus type 2", *Scientific Reports*, 2017

Publication

99

Meity Ardiana, Anwar Santoso, Hanestya O. Hermawan, Ricardo A. Nugraha, Budi S. Pikir, I

<1 %

<1 %

<1 %

<1 %

<1 %

Gde Rurus Suryawan. "Acute Effects of Cigarette on Endothelial Nitric Oxide Synthase, Vascular Cell Adhesion Molecule 1 and Aortic Intima Media Thickness "Cigarette smoke-induced pro-atherogenic changes"", Cold Spring Harbor Laboratory, 2021

Publication

100

Somphavanh Somlor, Ludovic BROSSAULT, Marc Grandadam. "Evaluation of VIDAS diagnostic assay prototypes detecting dengue virus NS1 antigen and anti-dengue virus IgM and IgG antibodies", Cold Spring Harbor Laboratory, 2021

Publication

<1 %

101

Zompi, Simona, and Eva Harris. "Animal Models of Dengue Virus Infection", Viruses, 2012.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Dengue Virus Serotype 4 Is Responsible for the Outbreak of Dengue in East Java City of Jember, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19
