

The incidence of Clostridium Difficile infection in diarrhea patients after receiving antibiotics at Dr. Soetomo Hospital Surabaya

by Rima Hayyu Chrisnanda

Submission date: 19-Jul-2021 12:56PM (UTC+0800)

Submission ID: 1621426952

File name: after_receiving_antibiotics_at_Dr._Soetomo_Hospital_Surabaya.pdf (725.69K)

Word count: 2896

Character count: 15161



Published by DiscoverSys

The incidence of *Clostridium Difficile* infection in diarrhea patients after receiving antibiotics at Dr. Soetomo Hospital Surabaya



CrossMark

Rima Hayyu Chrisnanda, Puspa Wardhani*

ABSTRACT

Introduction: Improper use of antibiotics is a risk factor for *C. difficile* infection. The increasing incidence of *C. difficile* infection is one of the indications of failure in infection prevention and control in hospitals. The aim of this study was to analyze the incidence of *C. difficile* infection in diarrhea patients after receiving antibiotics in the ICU and internal medicine ward of Dr. Soetomo Hospital Surabaya.

Methods: Stool samples were taken from 31 diarrhea patients with at least 2 x 24-hours of antibiotic use who were admitted to the ICU and hospitalized at Dr. Soetomo Hospital Surabaya from August 2017 to May 2018. Each sample was examined for glutamate dehydrogenase (GDH) and A and B toxins of *C. difficile*. The results of this study were analyzed descriptively.

Results: The average age of respondents was 48.94 (21 – 86) years old. In addition, based on demographic data, it was predominant by females (65%), diabetes mellitus and fever (16%), and ceftriazone usage (22%). The results showed one sample (3%) with positive results and 30 samples (97%) with negative results for both A/B toxins and GDH enzyme.

Conclusion: It can be concluded that the incidence of diarrhea in these patients was not due to *C. difficile* infection. The cause of diarrhea in these patients could be due to other causes such as a virus, fungus, or other bacterial infections. More sample collection is needed to provide more accurate data about *C. difficile* infection, so it can help to fulfill data for infection prevention and control in hospitals.

Keywords: *Clostridium difficile*, glutamate dehydrogenase, A/B toxins

Cite this Article: Chrisnanda, R.H., Wardhani, P. 2019. The incidence of *Clostridium Difficile* infection in diarrhea patients after receiving antibiotics at Dr. Soetomo Hospital Surabaya. *Bali Medical Journal* 8(2): 342-346. DOI: [10.15562/bmj.v8i2.1404](https://doi.org/10.15562/bmj.v8i2.1404)

Department of Clinical Pathology,
Faculty of Medicine Airlangga
University, Dr. Soetomo Hospital,
Surabaya, Indonesia

INTRODUCTION

Clostridium difficile (*C. difficile*) is an obligate anaerobe, gram-positive rod bacteria, which can cause diseases with clinical manifestations ranging from pseudomembranous colitis to severe fulminant colitis.¹ Irrational use of antibiotics is the most important risk factor for *C. difficile* infection.² The description of pseudomembranous colitis was first reported in 1893, and the association between the administration of clindamycin and pseudomembranous colitis was reported in 1974.¹ *C. difficile* infection is suspected to cause antibiotic-associated diarrhea in 10 – 35% of cases, and is the most common cause of nosocomial diarrhea with significant morbidity and mortality. About 80% of *C. difficile* infection are healthcare-associated infections (HAI), while 20% of cases are community-acquired.³

Several researchers in developed countries have reported the incidence of *C. difficile* infection, but it is still very limited in developing countries.³ The increasing incidence of *C. difficile* infection in hospitals is an indication of failure in infection prevention and control in hospitals. In the past two decades, the incidence of *C. difficile* infection, both healthcare-associated infections and community-acquired, have increased. This problem has an impact on mortality, hospital care, and health

costs.^{3,4} Carriage of *C. difficile* occurs in around 5 - 15% of healthy adults, 84.4% of healthy infants and neonates, and 57% of hospitalized patients.⁵

The Joint Commission International (JCI) standard includes the number of *C. difficile* infections as the number of infections that must be monitored continuously. The increasing incidence of *C. difficile* infection in hospitals is one of the indications of failure in infection prevention and control in hospitals. This could be caused by several things, such as the lack of hand hygiene, inadequate decontamination of medical equipment, low quality of environmental cleaning, low awareness of human resources in the practice of preventing infection, and the use of antibiotics that are not indicated. Dr. Soetomo Hospital Surabaya has not routinely performed *C. difficile* checks as a parameter to evaluate HAI, which may be due to the difficulty of doing *C. difficile* culture.

The aim of this study was to analyze the incidence of *C. difficile* infection in diarrhea patients who received antibiotics in the intensive care unit (ICU) and the internal medicine ward of Dr. Soetomo Hospital Surabaya. It is hoped that knowing the rate of infection due to *C. difficile* can increase the awareness of infection prevention and control in the hospital.

*Correspondence to:
Puspa Wardhani, Department
of Clinical Pathology, Faculty of
Medicine Airlangga University,
Dr. Soetomo Hospital, Surabaya,
Indonesia
puspa_pk@yahoo.co.id

Received: 2018-12-01
Accepted: 2018-12-19
Published: 2019-08-01

METHOD

This study was conducted at ²² Dr. Soetomo Hospital Surabaya. Samples were taken from the ICU and internal medicine ward from August 2017 to May 2018. Samples were collected from male and female adult patients who experienced diarrhea after receiving antibiotics for at least two days in the inpatient room. The examination was carried out at the Clinical Pathology Laboratory using the Immuno Chromatography Test (ICT) method quantitatively to determine the presence of toxins and GDH enzymes of *C. difficile*.

A/B toxins were checked using the IMMUNOQUICK® Tox A/B, which is designed as a rapid lateral flow immunoassay to detect the presence of antigen A and B toxins in the form of fresh, frozen and stool specimens that have been stored. The IMMUNOQUICK® Tox A/B rapid test with nitrocellulose zone part of the test line coated with anti-toxin A and B antibodies. Anti-toxin A and B antibodies also bind to red latex particles inserted in the test strip below the zone of nitrocellulose. The presence of A and B toxins in a combination of samples with antibodies will be a complex form.

Complex migration to the nitrocellulose strip will bind to antibodies in the test region and will be seen as a pink/red line. The appearance of two red/pink lines with varying intensity on the device in the test line position and the control line shows reactive results that are interpreted as positive results on the IMMUNOQUICK® Tox A/B. The appearance of one red/pink line with varying intensity on the device is in the position of the control line are interpreted as negative results. No line that appears on the device in the control line position indicates an invalid result and cannot be interpreted. If this condition occurs, the test must be repeated with a new kit.

Glutamate dehydrogenase (GDH) is a metabolic enzyme that is encoded by the *gluD* gene. All *C. difficile* isolates produce this antigen in high concentrations, for both toxigenic and non-toxigenic strains. *C. difficile* GDH IMMUNOQUICK is a qualitative lateral flow immunoassay to detect GDH antigens in feces. This test uses specific antibodies for GDH in the membrane on the test line. During the examination, the presence of GDH in fecal specimens will react with anti-GDH antibodies that are attached to gold particles. The mixture will move upward towards the membrane capillary and react with the anti-GDH contained in the test line. The appearance of a colored line on the line test (T) shows a positive result, whereas no line indicates a negative result. The results are declared invalid if a line does not appear in the control, so the test must be repeated.

Data were presented descriptively in tables and diagrams. Patients were descriptively presented based on inpatient rooms, gender, diagnosis, therapy, and *C. difficile* infection status.

RESULT

A total of 31 diarrhea patients with at least 2 x 24-hours of antibiotic use after admission to ICU and internal medicine ward were included in this study. Demographic data and clinical characteristics can be seen in Table 1.

As shown in Figure 1, two patients were obtained (6%) from the ICU, and 29 patients (94%) patients came from the internal medicine ward.

As shown in Figure 2, there were 20 females (65%) patients and 11 males (35%). The average age of patients from both the ICU and the inpatient ward was 48.94 years, with the youngest being 21 years old and the oldest being 86 years old. The shortest duration of antibiotic administration was four days, while the longest was 18 days. (Table 1)

As shown in Figure 3, patient diagnosis included chronic kidney disease, diabetes mellitus, febrile

¹⁵ **Table 1** Demographic data and clinical characteristics of patients (n = 31)

| Parameter | Mean | N | % |
|---|-----------------|----|-----|
| Age (years) | 48.94 (21 – 86) | | |
| Sex | | | |
| Male | | 11 | 35% |
| Female | | 20 | 65% |
| Diagnosis | | | |
| Chronic kidney disease | | 2 | 7% |
| Diabetes mellitus | | 5 | 16% |
| Fever | | 5 | 16% |
| Typhoid fever | | 3 | 10% |
| Hepatitis B | | 3 | 10% |
| Pneumonia | | 4 | 13% |
| Urinary tract infection | | 2 | 6% |
| Abdominal colic | | 2 | 6% |
| Other | | 5 | 16% |
| Therapy | | | |
| Ceftriaxone | | 22 | 71% |
| Cefotaxime | | 5 | 16% |
| Ciprofloxacin | | 3 | 10% |
| Meropenem | | 1 | 3% |
| Length of stay (days) | 7.65 (5 – 21) | | |
| Duration of antibiotic administration (Days) | 7.46 (4 – 18) | | |

observation, typhoid fever, hepatitis B, pneumonia, urinary tract infections, abdominal colic, and as many as 5 people (16%) suffered from other diseases, which were systemic lupus erythematosus, Ca caecum, pulmonary TB, HELLP syndrome and hypertension.

In Figure 4, 71% of the samples were treated with ceftriaxone, 16% received cefotaxime, 10% received ciprofloxacin, and one patient (3%) received meropenem therapy; all drugs were given through intravenous injection. The length of stay varied, with an average of 7.65 days. The shortest length of stay was five days from inpatient care, while the longest was 21 days from the ICU (Table 1).

All samples were examined for A/B toxins and GDH enzymes using ICT. The results showed only




Figure 1 Diagram of patients based on inpatient rooms




Figure 2 Diagram of patients based on gender




Figure 3 Diagram of patients based on diagnosis




Figure 4 Diagram of patients based on therapy




Figure 5 C. difficile infection percentage




Figure 6 The positive result of C. difficile infection

one sample with positive results for A/B toxins and GDH enzyme. Therefore, it can be concluded that the incidence of diarrhea in this patient was due to C. difficile infection. A total of 30 samples (97%) had negative results for both A/B toxins and GDH enzyme. (Figure 5)

DISCUSSION

In humans, most of the normal flora bacteria are anaerobic bacteria. The most commonly found bacterium in the digestive tract is *C. difficile*. *C. difficile* is found in about 3% of adults, 20 – 50% of infants, and 50 – 70% of neonates.⁶ Rolfe et al. and Fekety et al. stated that ampicillin, penicillin, clindamycin, chloramphenicol, tetracycline, and cephalixin are antibiotics that were often associated with the onset of *C. difficile* infection. However, Allen et al. stated that almost all antibiotics, except vancomycin, bacitracin, and metronidazole, were often associated with *C. difficile* infection.

In 2015, the Centers for Disease Control (CDC) estimated that *C. difficile* infected nearly half a million Americans and caused 29,000 deaths in 2011. Forty percent of cases came from home care and community health services, while 24% occurred in hospitals.⁷ In developing countries, *C. difficile* infection report is still uncommon. A study conducted by a hospital in Malaysia found that out of 175 fecal samples from patients suspected of being infected with *C. difficile*, 13.7% had positive results, with the majority occurring in patients over 50 years old.⁸

During the period of sample collection (August 2017 – May 2018), we obtained 31 samples. All samples were examined for A/B toxins and GDH enzymes using ICT. Only one sample (3%) was found to have positive results for A/B toxins and GDH enzyme, which meant that the incidence of diarrhea in this patient was due to *C. difficile* infection. A total of 30 samples (97%) had negative results for both for A/B toxins and GDH enzyme; hence it can be concluded that the incidence of diarrhea in these patients was not due to *C. difficile* infection. The cause of diarrhea in these patients could be due to a virus or other bacterial infections.

The results obtained in this study are lower than the study conducted in four hospitals in Semarang from 2014 to 2015. The study used fecal samples taken from 340 patients suspected of being infected with *C. difficile*. As many as 19 (5.6%) samples had positive results for toxins and GDH examination.⁹ Diagnosing *C. difficile* infection is difficult because of the lack of laboratories that can provide anaerobic culture facilities.

The patient who had positive results in this study was admitted to the hospital with a diagnosis of hepatitis B. The patient was 54 years old. The patient was treated in the inpatient of the internal medicine ward and received a treatment of 3rd generation cephalosporin injections twice per day along with other therapies according to the

treatment protocol for hepatitis B. On the third day of antibiotic administration, the patient began to have diarrhea, yet the antibiotic treatment continued until the seventh day. The patient's fecal sample was examined on the fifth day of antibiotic administration and showed positive results. Two days after the antibiotic was stopped, the patient no longer had diarrhea. Three days later, or on the 12th day, the patient was discharged from the hospital. The results of the examination for A/B toxins and GDH enzymes have also been reported to the clinician. This supports the theory that diarrhea in hospitalized patients who received at least 2 x 24-hour antibiotic therapy can be a sign of *C. difficile* infection.

Further research examining *C. difficile* infection should be conducted in a wider range of hospitals with longer periods, considering how difficult it is to collect the samples. It is expected that with more sample collection and anaerobic culture examination, more accurate data on *C. difficile* infection can be obtained to help fulfill data for infection prevention and control in hospitals.

CONCLUSION

Only one sample showed positive results, while the remaining 30 samples showed negative results. Therefore, it can be concluded that diarrhea in these patients was not due to *C. difficile* infection, but could be due to other causes such as viruses, fungi or other bacterial infections. Stool examination in patients suffering from diarrhea after receiving antibiotics for at least two days can help establish a diagnosis of whether the patient is suffering from *C. difficile* infection. Controlling *C. difficile* infection is very critical because the increasing incidence of *C. difficile* infection in hospitals is one of the indicators of failure in infection prevention and control in hospitals.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding manuscript

ETHICAL CLEARANCE

This study has been received ethical approval by the Ethics Committee prior study was carried out.

FUNDING

The authors are responsible for the funding without involvement of grant, sponsorship, or any other sources of funding.

AUTHOR CONTRIBUTION

All authors are contributed equally to the content of study from data preparation, statistical analysis, results, and data synthesis.

REFERENCES

- Goudarzi M, Seyedjavadi SS, Goudarzi H, Mehdizadeh Aghdam E, Nazeri S. Clostridium difficile Infection: Epidemiology, Pathogenesis, Risk Factors, and Therapeutic Options. *Scientifica (Cairo)*. 2014; 2014: 916826. DOI: [10.1155/2014/916826](https://doi.org/10.1155/2014/916826).
- Burnham CA, Carroll KC. Diagnosis of Clostridium difficile infection: an ongoing conundrum for clinicians and for clinical laboratories. *Clin Microbiol Rev*. 2013; 26(3): 604 – 30. DOI: [10.1128/CMR.00016-13](https://doi.org/10.1128/CMR.00016-13).
- Khan FY, Elzouki AN. Clostridium difficile infection: a review of the literature. *Asian Pac J Trop Med*. 2014; 7S1: S6 – S13. DOI: [10.1016/S1995-7645\(14\)60197-8](https://doi.org/10.1016/S1995-7645(14)60197-8).
- Vindigni SM, Surawicz CM. C. difficile Infection: Changing Epidemiology and Management Paradigms. *Clin Transl Gastroenterol*. 2015; 6: e99. DOI: [10.1038/ctg.2015.24](https://doi.org/10.1038/ctg.2015.24).
- Surawicz CM, Brandt LJ, Binion DG, Ananthkrishnan AN, Curry SR, Gilligan PH et al. Guidelines for diagnosis, treatment, and prevention of Clostridium difficile infections. *Am J Gastroenterol*. 2013; 108(4): 478 – 98. DOI: [10.1038/ajg.2013.4](https://doi.org/10.1038/ajg.2013.4).
- Rolfe RD, Finegold SM. Purification and characterization of Clostridium difficile toxin. *Infect Immun*. 1979; 25(1): 191 – 201.
- Rupnik M, Wilcox MH, Gerding DN. Clostridium difficile infection: new developments in epidemiology and pathogenesis. *Nat Rev Microbiol*. 2009; 7(7): 526 – 36. DOI: [10.1038/nrmicro2164](https://doi.org/10.1038/nrmicro2164).
- Collins DA, Hawkey PM, Riley TV. Epidemiology of Clostridium difficile infection in Asia. *Antimicrob Resist Infect Control*. 2013; 2(1): 21. DOI: [10.1186/2047-2994-2-21](https://doi.org/10.1186/2047-2994-2-21).
- Collins DA, Gasem MH, Habibie TH, Arinton IG, Hendriyanto P, Hartana AP, Riley TV. Prevalence and molecular epidemiology of Clostridium difficile infection in Indonesia. *New Microbes New Infect*. 2017; 18: 34 – 37. DOI: [10.1016/j.nmnl.2017.04.006](https://doi.org/10.1016/j.nmnl.2017.04.006).



This work is licensed under a Creative Commons Attribution

The incidence of Clostridium Difficile infection in diarrhea patients after receiving antibiotics at Dr. Soetomo Hospital Surabaya

ORIGINALITY REPORT

14%

SIMILARITY INDEX

8%

INTERNET SOURCES

12%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Ave Olivia Rahman, Anati Purwakanthi, Hasna Dewi. "Antifertility effect of betel nut (Areca catechu L) in male rat", MEDISAINS, 2020
Publication 1%
- 2** eprints.skums.ac.ir
Internet Source 1%
- 3** S. Mayor. "Drugs are as good as PCI in stable coronary artery disease, study shows", BMJ, 4/7/2007
Publication 1%
- 4** en.wikipedia.org
Internet Source 1%
- 5** gazeta.neurologia.com.pl
Internet Source 1%
- 6** www.singulexclarity.eu
Internet Source 1%
- 7** Daniel Curcio, Alejandro Cané, Francisco Andrés Fernández, Jorge Correa. "Clostridium

difficile-associated Diarrhea in Developing Countries: A Systematic Review and Meta-Analysis", Infectious Diseases and Therapy, 2019

Publication

8

www.ajmb.org

Internet Source

1 %

9

Yingchao Cui, Danfeng Dong, Lihua Zhang, Daosheng Wang, Cen Jiang, Qi Ni, Chen Wang, Enqiang Mao, Yibing Peng. "Risk factors for Clostridioides difficile infection and colonization among patients admitted to an intensive care unit in Shanghai, China", Research Square, 2019

Publication

1 %

10

Petru A. Pop, Liviu Lazar, Florin M. Marcu. "Benefic Effect of Conservative Treatments on Patients Diagnosed With Osteoporosis", Volume 2: Biomedical and Biotechnology, 2012

Publication

1 %

11

Marwan S. Abougergi, John H. Kwon. "Intravenous Immunoglobulin for the Treatment of Clostridium difficile Infection: A Review", Digestive Diseases and Sciences, 2010

Publication

<1 %

12

Internet Source

<1 %

13

www.researchgate.net

Internet Source

<1 %

14

Na Liu, Yu-xia Deng. "Observations on treatment of piriformis injury syndrome by a technique of green tortoise exploring point", *Journal of Acupuncture and Tuina Science*, 2008

Publication

<1 %

15

www.ihsanalur.com

Internet Source

<1 %

16

"Infectious Diseases in the Intensive Care Unit", Springer Science and Business Media LLC, 2020

Publication

<1 %

17

Burnham, C.-A. D., and K. C. Carroll. "Diagnosis of *Clostridium difficile* Infection: an Ongoing Conundrum for Clinicians and for Clinical Laboratories", *Clinical Microbiology Reviews*, 2013.

Publication

<1 %

18

Mansoor Kodori, Zohreh Ghalavand, Abbas Yadegar, Gita Eslami et al. "Molecular Characterization of Pathogenicity Locus (PaLoc) and *tcdC* Genetic Diversity Among

<1 %

tcdA+B+ Clostridioides Difficile Clinical Isolates in Tehran, Iran", Research Square, 2020

Publication

| | | |
|----|---|------|
| 19 | d-nb.info Internet Source | <1 % |
| 20 | tel.archives-ouvertes.fr Internet Source | <1 % |
| 21 | www.dovepress.com Internet Source | <1 % |
| 22 | "Moderated Poster Presentations", International Journal of Urology, 8/2006 Publication | <1 % |
| 23 | R. V. FENGER, A. LINNEBERG, M. TVEDE, C. ØSTERGAARD. " Increasing seroprevalence of in an adult Danish general population ", Epidemiology and Infection, 2008 Publication | <1 % |
| 24 | teknolabjournal.com Internet Source | <1 % |
| 25 | www.ncbi.nlm.nih.gov Internet Source | <1 % |
| 26 | www.researchsquare.com Internet Source | <1 % |
| 27 | "Updates on Clostridium difficile in Europe", Springer Science and Business Media LLC, 2018 | <1 % |

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off

The incidence of Clostridium Difficile infection in diarrhea patients after receiving antibiotics at Dr. Soetomo Hospital Surabaya

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
