

Batatasin III a derivative of dihydrostilbene compound from Yam Peel of Uwi Tuban and Its Antioxidant Activity

by Alfinda Novi Kristanti

Submission date: 09-Mar-2020 05:58PM (UTC+0800)

Submission ID: 1272174468

File name: 2018-IOP_Conf._Series-Wiwik.pdf (437.15K)


Word count: 1910


Character count: 8971

PAPER · OPEN ACCESS

Batatasin III a derivative of dihydrostilbene compound from Yam Peel of Uwi Tuban and Its Antioxidant Activity

To cite this article: W Agustina *et al* 2018 *J. Phys.: Conf. Ser.* **1116** 042003

 View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Batatasin III a derivative of dihydrostilbene compound from Yam Peel of Uwi Tuban and Its Antioxidant Activity

W Agustina¹, A N Kristanti², Y Takaya³, E Fitriana¹ and N S Aminah^{2*}

¹ Student of Magister Chemistry Program, Dept. Of Chemistry, Faculty of Science and Technology, Universitas Airlangga, C-Campus, Jl. Ir Soekarno, Surabaya, 60115

² Chemistry, Faculty of Science and Technology, Universitas Airlangga, C-Campus, Jl. Ir Soekarno, Surabaya, 60115

³ Faculty of Pharmacy, Meijo University, 468-8503, Tempaku Nagoya, Japan

*email address : nanik-s-a@fst.unair.ac.id

Abstract A dihydrostilbene derivative compound, namely 3,3-dihydroxy-5-methoxybenzyl or batatasin III (**1**) was isolated from the yam peel of *Dioscorea alata* L. The structure of compound **1** has been elucidated based on spectroscopy UV-Vis, 1D and 2D NMR Analysis. The IC₅₀ of DPPH radical scavenging of this compound (206.82 µg/mL) lower than ethyl acetate extract (109.99 µg/mL), but higher than methanol extract (893.59 µg/mL).

1. Introduction

The genus *Dioscorea* (Dioscoreaceae) comprises more than 600 species that are widely distributed in tropical and subtropical region such as Indonesia. Wild yam or *Dioscorea alata* occur in several part of Indonesia and the plants is commonly known as Uwi tuban. Many of these species are used local food crop rich in starch of Indonesia but the bioactivity of this species is still poorly understood. This genus has been shown to produce a number of secondary metabolite such as terpenoid [1], saponin [2], steroid [3] and phenolic compounds [4]. Previous research from other country reported that *Dioscorea* are used indigenously as traditional medicines to leprosy, tumor in Bangladesh [5], inflammatory diseases such as asthma, rheumatoid arthritis and bronchitis in Taiwan [6].

In continuation of the research of the phenolic compounds in this medicinal plant, our research group already reported from two species: *D. esculenta* L. successfully isolated two phenanthrene derivatives namely confusarin and nudol [7], methyl-3,4-dihydroxybenzoate and 9,10-dihydrophenanthrene from *D. alata* L.[8]. In this research, reported of 3,3-dihydroxy-5-methoxybenzyl or batatasin III is a phenolic compound isolated from the methanol extract of the yam peel of *Dioscorea alata* L.(uwi tuban). Uwi Tuban is one of *D. alata* species which mostly cultivated in Tuban, East Java. The chemical structure of compound **1** was established by UV, IR, 1D and 2D NMR. The antioxidant activity against DPPH radical scavenging the isolated compound **1**, methanol and ethyl acetate extract are also briefly described.

2. Materials and Methods

2.1 General experimental

NMR spectra were recorded on JEOL 600 ECA spectrometer using CDCl₃ at 600 (¹H) and 125 (¹³C) MHz. UV and FTIR spectrum recorded in KBr powder with Shimadzu series 1800 spectrophotometer.



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Gravimetry Column Chromatography (GCC) and radial chromatography were carried out using Si gel 60 GF254 and Si gel PF254 for TLC analysis and pre-coated silica gel plates (Merck, Darmstadt, Germany, Kieselgel 60 GF254 0,25 mm thickness) were used.

2.2 Plant Material

Sample of yam peels of uwi Tuban were collected in March 2015 from the district Tuban, East Java, Indonesia, and the specimen was deposited at the Department of Biology, Fac. Of Science and Technology, Universitas Airlangga. The yam peels were cleaned, dried under the shade, cut into small pieces and milled.

2.3 Extraction and Isolation

The dried yam peels of uwi Tuban (2.1 kg) were macerated in methanol at room temperature (3x24 hours). The methanol extract was evaporated under reduced pressure to give a dark brown residue (88 g). The crude methanol extract (88 g) was partitioned respectively with n-hexane and ethyl acetate. The ethyl acetate extract (16 g) was separated by vacuum column chromatography on silica gel using eluent the mixture of n-hexane and ethyl acetate with increasing polarity gradient and gravity column chromatography with the same method of eluent. The subfraction that showed two spot on TLC test, then purified using radial chromatography using the mixture of chloroform-aceton (increasing polarity gradient) yielded pure compound that showed one spot on TLC test (15 mg)

3.4 DPPH Radical Scavenging

The antioxidant assay of compound **1** against DPPH (2,2-diphenyl-1-picrihidrazil) radical was measured by UV-Vis spectrophotometer λ 517 nm as described previously [10]. The inhibition percentage (%) of radical scavenging activity was calculated using the following equation

$$\text{Inhibition (\%)} = \left(\frac{A_0 - A_s}{A_0} \right) \times 100\%$$

Where A_0 is the absorbance of the control (containing all reagents except the test compound), and A_s is the absorbance of the test compound.

3. Results and Discussions

Extraction of the dried milled of yam peel of uwi Tuban (2.1 kg) was carried out using methanol and then methanol extract was portioned successively with n-hexane and ethyl acetate. The ethyl acetate extract (16 g) was separated by gravity column chromatography on silica gel and radial chromatography yielded 3,3-dihydroxy-5-methoxybibenzyl or batatasin III (15mg) (**Figure 1**)

The compound of 3,3-dihydroxy-5-methoxybibenzyl (**1**) was isolated as a brown liquid (15 mg). UV spectrum (MeOH) λ_{max} (log ϵ) 226.4 nm(3.67) and 274.2 nm (2.42). IR ν_{max} cm^{-1} 3334.93, 2937.59, 2858.51, 1697.36, 1494.83, 1149.57, 1058.92, 692.44. $^1\text{H-NMR}$ (600 MHz, CDCl_3) δ 2.84 (4H, m, H-6' a,b), 3.77 (1H, s, 5'-OMe), 6.28 (1H, s, H-2', H-4'), 6.35 (1H, s, H-6'), 6.66 (1H, s, H-2), 6.70 (1H, d, J = 12, H-4), 6.77 (2H, d, J = 6, H-6), 7.17 (3H, t, J = 6, 12, H-5). $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) 160.79 (C 5'-OMe), 144.45 (C-1'), 143.57 (C-1), 129.56 (C-5), 120.92 (C-6), 115.43 (C-2), 112.94 (C-4), 108.09 (C-2'), 106.88 (C-6'), 99.17 (C-4'), 37.63 (C-6a), 37.29 (C-6b). The $^1\text{H-NMR}$ (Table 1) spectrum of **1** exhibited the presence seven aromatic proton at δ 6.28 (1H, s, H-2', H-4'), 6.35 (1H, s, H-6'), 6.66 (1H, s, H-2), 6.70 (1H, d, J = 12, H-4), 6.77 (2H, d, J = 6, H-6), 7.17 (3H, t, J = 6, 12, H-5), two methylene proton at δ 2.84 (4H, m, H-6' a,b). The $^{13}\text{C-NMR}$ spectrum showed the presence seven aromatic ring, two hydroxy aromatic ring (δ 156.51 and 155.45), one methoxy aromatic ring (δ 160.79), one methoxy (δ 55.31) and two methylene carbon (δ 37.29 and 37.62). Base on NMR data, it is predicted that compound **1** is a dihydrostilbene derivative (**Figure 1**). The methoxy position were confirmed to be C-5' by HMBC. And compound **1** contain two hydroxy (C-3' and C-3) by HSQC spectrum data. Based on further comparison with published data (Table 2)[9], the structure of **1** was identified as 3,3-dihydroxy-5-methoxybibenzyl with the trivial name batatasin III. The complete of HMBC correlations consistent with the structure **1** are shown in Table 1 and Figure 2.

Based on the literature study, this compound has never been reported yet from *D. alata* especially from Indonesia.




Figure 1. Structure of 3,3-dihydroxy-5-methoxybibenzyl




Figure 2. Selected HMBC correlation for compound 1.

Table 1. NMR spectroscopic data of 3,3-dihydroxy-5-methoxybibenzyl in CDCl₃

| No | ¹ H (m, J in Hz) | ¹³ C | HMBC |
|-------|-----------------------------|-----------------|------------------------|
| 1 | 8 | 143.45 | |
| 2 | 6.66 (1H, s) | 115.43 | |
| 3 | - | 155.45 | |
| 4 | 6.70(1H, d, J=12) | 112.94 | 5 |
| 5 | 7.17 (1H, t, J=6;12) | 129.56 | C-1, C-3 |
| 6 | 6.77 (1H, d, J=6) | 120.92 | C-6'a,b , C-4, C-1 |
| 1' | 7 | 144.45 | |
| 2' | 6.28 (1H, s) | 108.09 | C-6' a,b , C-4', 3'OH |
| 3' | - | 154.45 | |
| 4' | 6.28(1H, s) | 99.71 | C-6' , 2'-OH, C-5' OMe |
| 5' | - | 160.79 | |
| 6' | 6.35 (1H, s) | 106.88 | C-2' , C-4' , C-5' OMe |
| a | 2.84 (2H, m) | 37.29 | C-6' , C-1, C-1' |
| b | 2.84 (2H, m) | 37.62 | C-6' , C-1, C-1' |
| 3-OH | 5.22 (1H, sbr) | - | |
| 3'-OH | 5.22 (1H, sbr) | - | |

5'-OMe 3.77 (3H, s) 55.31

Table 2. The comparison of chemical shift data of 3,3-dihydroxy-5-methoxybenzyl from isolated compound *B. striata* [9]

| Isolated compound from <i>D. alata</i> | | | 3,3-dihydroxy-5-methoxybenzyl rom <i>B. striata</i> [9] | | |
|--|-----------------------------|-----------------|--|-----------------------------|-----------------|
| No | ¹ H (m, J in Hz) | ¹³ C | No | ¹ H (m, J in Hz) | ¹³ C |
| 1 | 6.66 (1H, s) | 143.45 | 1 | - | 143.5 |
| 2 | 6.70 (1H, d, J=12) | 115.43 | 2 | 6.65 | 121.2 |
| 3 | 7.17 (1H, t, J=6;12) | 155.45 | 3 | - | 129.7 |
| 4 | 6.77 (1H, d, J=6) | 112.94 | 4 | 6.70 | 113.1 |
| 5 | 6.28 (1H, s) | 129.56 | 5 | 7.14 (1H, m) | 155.7 |
| 6 | 6.28 (1H, s) | 120.92 | 6 | 6.75 (1H, d, J=8) | 115.5 |
| 1' | 6.35 (1H, s) | 144.45 | 1' | - | 144.6 |
| 2' | 2.84 (2H, m) | 108.09 | 2' | 6.25 | 108.1 |
| 3' | 2.84 (2H, m) | 154.45 | 3' | - | 156.7 |
| 4' | 5.22 (1H, sbr) | 99.71 | 4' | 6.28 | 99.33 |
| 5' | 5.22 (1H, sbr) | 160.79 | 5' | - | 161.1 |
| 6' | 3.77 (3H, s) | 106.88 | 6' | 6.32 | 107.0 |
| a | 3.77 (3H, s) | 37.29 | a | 2.84 (2H, m) | 37.8 |
| b | 3.77 (3H, s) | 37.62 | b | 2.79 (2H, m) | 37.5 |
| 3-OH | - | - | 3-OH | - | - |
| 3'-OH | - | - | 3'-OH | - | - |
| 5'-OMe | 3.77 (3H, s) | 55.31 | 5'-OMe | 3.75(3H, s) | 55.4 |

4. Conclusions

On antioxidant evaluation against DPPH radical scavenging of compound **1** exhibited IC₅₀ values of 206.82 µg/mL suggested that compound **1** has moderate activity. This compound more active than IC₅₀ of methanol extract (893.59 µg/mL) and less active than ethyl acetate extract (109.66 µg/mL).

References

- [1] Teponno R B *et al* 2013 *Phytochem Let.*, 310-314
- [2] Kim K H *et al* 2011 *Biorg & Med Chem Lett.* 2075-2078
- [3] Taponjhou L A *et al* 2013, *Phytochem Lett.* 341-350
- [4] Yang M H *et al* 2009 *J. Bioorg & Med Chem.* 2689-2694
- [5] Murray R D H *et al* 1984 *Phytochemistry* **23** 623
- [6] Chen H *et al* 2003 *Nutrition* **19** 646
- [7] Aminah S A *et al* 2017 *J. of Chemical Technology and Metallurgy* **52** 1135
- [8] Aminah N S *et al* 2017 *AIP Conference Proceedings* **1888** 020050
- [9] Woo W K *et al* 2014 *Natural Product Sciences* **20** 91
- [10] Brand-Williams W *et al* 1995 *Lebensmittel-Wissenschaft und-Technologie/Food Science and Technology* **28** 25

Batatasin III a derivative of dihydrostilbene compound from Yam Peel of Uwi Tuban and Its Antioxidant Activity

ORIGINALITY REPORT

20%
SIMILARITY INDEX

10%
INTERNET SOURCES

18%
PUBLICATIONS

0%
STUDENT PAPERS

PRIMARY SOURCES

- 1** Philippe Colomban. "Understanding the nano- and macromechanical behaviour, the failure and fatigue mechanisms of advanced and natural polymer fibres by Raman/IR microspectrometry", *Advances in Natural Sciences: Nanoscience and Nanotechnology*, 2012
Publication **3%**
- 2** Iqbal Musthapa. "An oxepinoflavone from *Artocarpus elasticus* with cytotoxic activity against P-388 cells", *Archives of Pharmacal Research*, 02/2009
Publication **2%**
- 3** vdocuments.site
Internet Source **2%**
- 4** Chapman, C.J.. "Peptide modification through site-selective residue interconversion: application of the rhodium-catalysed 1,4-addition of aryl siloxanes and boronates", *Tetrahedron*, 20080929
Publication **2%**

| | | |
|----|--|-----|
| 5 | Yoshiteru Ida, Yohko Satoh, Mariko Katoh, Masumi Katsumata (nee Ohtsuka) et al. "Achyranthosides A and B, novel cytotoxic saponins from <i>Achyranthes fauriei</i> root", <i>Tetrahedron Letters</i> , 1994 Publication | 1 % |
| 6 | www.ifrj.upm.edu.my Internet Source | 1 % |
| 7 | Sunazuka, T.. "Absolute stereochemistries and total synthesis of (+)-arisugacins A and B, potent, orally bioactive and selective inhibitors of acetylcholinesterase", <i>Tetrahedron</i> , 20040830 Publication | 1 % |
| 8 | www.allindianpatents.com Internet Source | 1 % |
| 9 | www.thieme-connect.com Internet Source | 1 % |
| 10 | www.myanmar-education.edu.mm Internet Source | 1 % |
| 11 | Ali Loukaci, Michèle Guyot, Angèle Chiaroni, Claude Riche. " A New Indole Alkaloid from the Marine Tunicate ", <i>Journal of Natural Products</i> , 1998 Publication | 1 % |
| 12 | www.freepatentsonline.com Internet Source | |

1 %

13

P.L. Majumder, Mausumi Basak. "Two bibenzyl derivatives from the orchid *Cirrhopetalum andersonii*", *Phytochemistry*, 1991

Publication

1 %

14

Eveen Al-Shalabi, Muzn Alkhaldi, Suhair Sunoqrot. "Development and evaluation of polymeric nanocapsules for cirsiol isolated from Jordanian *Teucrium polium* L. as a potential anticancer nanomedicine", *Journal of Drug Delivery Science and Technology*, 2020

Publication

<1 %

15

Lamberto Tomassini, Jianjun Gao, Sebastiano Foddai, Mauro Serafini, Antonio Ventrone, Marcello Nicoletti. " Iridoid glucosides from ", *Natural Product Research*, 2006

Publication

<1 %

16

cirworld.org

Internet Source

<1 %

17

Allan M. Jordan, Helen M. I. Osborn, Petra M. Stafford, George Tzortzis, Robert A. Rastall. "Synthesis of Unsaturated Aminopyranosides as Possible Transition State Mimics for Glycosidases", *Journal of Carbohydrate Chemistry*, 2007

Publication

<1 %

18

Munekazu Inuma, Hideki Tosa, Naeko Toriyama, Toshiyuki Tanaka, Tetsuro Ito, V. Chelladurai. "Six xanthenes from *Calophyllum austroindicum*", *Phytochemistry*, 1996

Publication

<1 %

19

S. A. Emami, J. Asili, Z. Mohagheghi, M. K. Hassanzadeh. "Antioxidant Activity of Leaves and Fruits of Iranian Conifers", *Evidence-Based Complementary and Alternative Medicine*, 2007

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Batatasin III a derivative of dihydrostilbene compound from Yam Peel of Uwi Tuban and Its Antioxidant Activity

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
