

Immature Granulocyte and Mean Platelet Volume as a Predictor of 30-Day Postoperative Mortality in Patients with Sepsis Caused by Peritonitis

by Rudi Hartono

Submission date: 19-Jul-2021 01:11PM (UTC+0800)

Submission ID: 1621435882

File name: tive_Mortality_in_Patients_with_Sepsis_Caused_by_Peritonitis.pdf (514.69K)

Word count: 5070

Character count: 25587

Immature Granulocyte and Mean Platelet Volume as a Predictor of 30-Day Postoperative Mortality in Patients with Sepsis Caused by Peritonitis

Rudi Hartono Sinaga, Arie Utariani, Puspa Wardhani¹, Hardiono Hardiono

Departments of Anesthesiology and Reanimation and ¹Clinical Pathology, Faculty of Medicine Universitas Airlangga, Dr. Soetomo Hospital, Surabaya, Indonesia

Abstract

Background: Prompt and reliable identification and risk stratification in sepsis patients are needed to reduce the risk of mortality. Immature granulocytes (IG) and mean platelet volume (MPV) are considered as the predictors of 30-day mortality in sepsis patients. This study aims to analyze the relationship between IG and MPV with 30-day mortality following emergency laparotomy in patients with sepsis due to peritonitis. **Materials and Methods:** In this observational retrospective study, IG, MPV value, and 30-day mortality were obtained from the medical records of sepsis patients due to peritonitis who underwent an emergency laparotomy that met the inclusion criteria. We recorded the patients' data that met the inclusion criteria from the medical records that consisted of age, sex, diagnosis, sequential organ failure assessment score, and routine laboratory examination at the time of admission. Then, we analyzed each variable to determine the valid predictors of mortality. **Results:** From a total of 107 patients, the mortality rate was 34.58%. IG of day 1 (cutoff = 1.05), MPV of day 3 (cutoff = 10.35), and mean difference of platelet volume between day-0 and day-3 (cutoff = 0.35) were valid predictors for 30-day mortality ($P = 0.004$, $P = 0.006$, and $P < 0.001$, respectively). The mean difference of platelet volume day-0 and day-3 had the highest sensitivity and specificity, which was 67.6% and 72.9%, respectively. **Conclusion:** The number of IG on day-1, MPV on day-3, and mean difference of platelet volume between day-0 and day-3 are the valid predictors of mortality in sepsis patients due to peritonitis who underwent emergency surgery within 30 days.

Keywords: Immature granulocyte, mean platelet volume, peritonitis, sepsis

INTRODUCTION

Sepsis is a life-threatening state of organ dysfunction caused by immune dysregulation against infection.^[1] The epidemiological burden of sepsis globally is difficult to be determined, but sepsis is estimated to affect more than 30 million people worldwide each year and potentially causes around 6 million deaths.^[2] A research conducted in seven countries by collecting the data from 1979 to 2015 showed 288 cases of sepsis and 148 cases of severe sepsis per 100,000 people/year with in-hospital mortality rates were 17% for sepsis and 26% for severe sepsis.^[3] Another study found that 27.08% of all patients with sepsis had severe sepsis with 40%–60% mortality.^[4]

Peritonitis is one of the most common causes of sepsis. Research on critical surgical patients with severe sepsis showed that the most infection site (72.3%) was the abdomen.^[5] A study on 675 sepsis patients that needed surgical site control showed

389 patients (57.6%) experiencing intra-abdominal sepsis.^[6] The high mortality rate in sepsis patients can be caused by several factors, including low awareness, late identification, and inappropriate management of the disease.^[7-9]

Some biomarkers of organ dysfunction occur in patients with sepsis, especially in septic shock, and these markers are related to the severity and mortality of sepsis.^[9-13] Thrombocytopenia is a common condition and a multifactorial phenomenon

Address for correspondence: Dr. Arie Utariani,

Department of Anesthesiology and Reanimation, Faculty of Medicine Airlangga University, Dr. Soetomo General Hospital, Surabaya, Indonesia.
E-mail: arie.utariani@fk.unair.ac.id

Submitted: 19-Jun-2020

Revised: 16-Jul-2020

Accepted: 16-Aug-2020

Published: 05-Oct-2020

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Sinaga RH, Utariani A, Wardhani P, Hardiono H. Immature granulocyte and mean platelet volume as a predictor of 30-day postoperative mortality in patients with sepsis caused by peritonitis. *Bali J Anaesthesiol* 2020;4:166-71.

Access this article online

Quick Response Code:



Website:
www.bjoaonline.com

DOI:
10.4103/BJOA.BJOA_114_20

that occurs during sepsis.^[14] When thrombocytopenia occurs, younger platelets are released into the blood, and these platelets are larger and more active, so mean platelet volume (MPV) increases. An increase in MPV indicates a platelet diameter that can be used as a marker for platelet activation, endothelial damage, and thrombotic and inflammatory conditions.^[15-18] Previous studies have shown that patients who died of sepsis had a more considerable increase in MPV values and increased MPV from baseline is an independent risk factor for 28-day mortality in patients with sepsis.^[15] MPV has the potential to be used as an easily accessible prognostic marker of sepsis.^[18]

Although several markers have been investigated, only a few can be applied clinically due to the complexity of sepsis, and easy identification of markers is needed to provide adequate treatment for patients with sepsis.^[19] There were no studies that analyzed the MPV and immature granulocytes (IG) as the predictors of mortality in patients with sepsis caused by peritonitis who underwent surgery. This study set goals to compare the significance of IG and MPV by diagnostic testing to predict the sepsis prognosis due to peritonitis in patients undergoing emergency surgery.

MATERIALS AND METHODS

This was a retrospective study conducted in Dr. Soetomo General Hospital in Surabaya, Indonesia. After the approval of our Institutional Ethics Committee (1903/KEPK/III/2020 dated on March 20, 2020), we collected the data of the patients diagnosed with peritonitis and presented with sepsis who underwent emergency laparotomy from the medical records between May and December 2019.

We employed all-inclusive sampling for subject selection. Patients who met the following criteria were enrolled in the study protocol: aged ≥ 18 years old, diagnosed with peritonitis who underwent emergency laparotomy, presented with sepsis at the time of admission with the suspected source of infection (peritonitis), and ≥ 2 quick sequential organ failure assessment (SOFA) criteria based on the 3rd International Consensus Definition for Sepsis and Septic Shock 3. The exclusion criteria were the patient with incomplete medical record and conditions that cause the increase of IG and MPV, such as blood cell malignancy, an autoimmune disease, an immunocompromised condition, and history of drugs consumption (anti-platelet such as clopidogrel) before admission.

We recorded the patients' data that met the inclusion criteria from the medical records that consisted of age, sex, early diagnosis, infection site, SOFA score, routine laboratory examination at the time of admission in the emergency room (d0), day-1 (d1) postlaparotomy, and day-3 (d3) postlaparotomy, and patient's outcome between day-3 and day-30 postlaparotomy. Then, patients were divided into two groups: the survivor group (Group S) and the nonsurvivor group (Group NS). SOFA score was used to determine the outcome and correlate with IG and MPV as a predictor of mortality in sepsis patients.

Statistical analysis was performed with the SPSS statistics software version 25 (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, version 25.0. Armonk, NY: IBM Corp.). Continuous variables were expressed as mean \pm standard deviation or median (range) where appropriate, and categorical variables were expressed as the absolute number and proportions. Logistic regression was performed to see the most influential subvariables and receiver operating characteristic (ROC) curve plotting to determine IG and MPV cutoff value as a predictor of mortality. The cutoff value was calculated to get the best sensitivity and specificity. $P < 0.05$ was considered statistically significant.

RESULTS

A total of 107 participants were enrolled in this study, with a mortality rate of 34.6%. Demographic, clinical, and laboratory characteristics in each group are shown in Table 1. In this study, we did not find any significant differences in sex, age, and BMI between nonsurvivor and survivor group. The median of days in mechanical ventilation was longer in nonsurvivor than in the survivor group.

Logistic regression analysis to analyze the correlation between SOFA score, IG, and MPV with 30 days-mortality is shown in Table 2. IG day 0, day 1, and day 3, Δ IG day 3-day 0, MPV day 3, and Δ MPV day 3-day 0 were further analyzed with ROC curves to determine the cutoff value, as shown in Figure 1 and Table 3. According to the ROC curve, all of these parameters were significant as a predictor of 30 days-mortality. Upon diagnostic testing [Table 4], only IG-d1, MPV-d3, and Δ MPV d3-d0 were valid as a predictor for 30-days mortality in sepsis patients, which were shown with McNemar test and κ -value of >0.05 . In this study, Δ MPVd3-d0 >0.35 ($\times 10^3/\text{mm}^3$) was the best predictor for sepsis-related mortality in 30 days with the highest sensitivity (67.6%) and specificity (72.9%).




Figure 1: Receiver operating characteristic curves for immature granulocytes and mean platelet volume to predict 30 days-mortality in sepsis patients with peritonitis who underwent emergency laparotomy

Table 1: Demographic, clinical, and laboratory characteristics comparison

	Group NS	Group S	P
Number of patients	37 (34.6)	70 (65.4)	
Sex, n (%)			0.174*
Male	29 (78.4)	46 (65.7)	
Female	8 (21.6)	24 (34.3)	
Age (years)	60 (18-80)	48 (18-83)	0.076**
BMI (kg/m ²)	24.2 (17.3-31.2)	23.9 (14.5-45.4)	0.639**
Infection site, n (%)			
Perforated appendicitis	8 (21.6)	44 (62.9)	0.015*
Gastric perforation	15 (40.5)	16 (22.9)	
Others	14 (37.8)	10 (14.3)	
Comorbidities, n (%)			
Diabetes mellitus	9 (24.3)	11 (15.7)	0.277*
Hypertension	11 (29.7)	19 (27.1)	0.777*
COPD	4 (10.8)	4 (5.7)	0.340*
Osteoarthritis	4 (10.8)	2 (2.9)	0.089*
Gastritis	1 (2.7)	3 (4.3)	0.681*
Malignancy	3 (8.1)	3 (2.9)	0.221*
Coronary disease	0	1 (1.4)	0.465*
Asthma	1 (2.7)	0	0.167*
No comorbidity	11 (29.7)	31 (44.3)	0.142*
LOS (days)	7 (4-15)	8 (4-20)	0.165**
MV (days)	7 (0-14)	1 (0-8)	<0.001**
SOFA score	11 (6-12)	5 (0-8)	<0.001**
Laboratory values			
Hb d0 (g/dL)	10.8 (±2.3)	11.9 (±2.7)	0.041***
WBC d0 (10 ³ /mm ³)	11.2 (4.3-36.5)	13.4 (0.9-38.4)	0.130**
Platelet d0 (10 ³ /mm ³)	324.0 (87.0-643.0)	271.0 (37.0-801.0)	0.911**
Albumin d0 (10 ³ /mm ³)	2.8 (±0.5)	3.1 (±0.5)	0.001***
IG d0 (10 ³ /mm ³)	1.5 (0.2-9.0)	0.6 (0.2-5.5)	0.002**
IG d1 (10 ³ /mm ³)	1.7 (0.2-15.0)	0.7 (0.3-14.0)	0.005**
IG d3 (10 ³ /mm ³)	1.9 (0.5-9.6)	0.8 (0.3-5.9)	0.001**
ΔIG d3-d0	0.3 ([-1.9]-2.9)	0.1 ([-1.1]-1.3)	0.043**
MPV d0 (10 ³ /mm ³)	10.04±0.94	9.93±0.96	0.568***
MPV d1 (10 ³ /mm ³)	10.16±0.69	9.94±1.01	0.242***
MPV d3 (10 ³ /mm ³)	10.69±0.80	10.02±0.94	<0.001***
ΔMPV d3-d0	0.6 ([-0.7]-2.0)	0.05 ([-8.5]-2.0)	0.001**

Data for age, body mass index, length of stay, days on mechanical ventilation, SOFA score, and laboratory values are expressed as mean±standard deviation or median (range). *Chi-square test, **Mann-Whitney U-test, ***T2-independent test. BMI: Body mass index, COPD: Chronic obstructive pulmonary disease, LOS: Length of stay, MV: Mechanical ventilation, SOFA: Sequential organ failure assessment, Hb: Hemoglobin, WBC: White blood cells, IG: Immature granulocytes, MPV: Mean platelet volume, Δ: Difference

Table 2: Logistic regression analysis for the independent predictors of 30 days mortality

	Odds ratio	P
SOFA score	2.309	<0.001*
IG d0	1.378	0.036*
IG d1	0.867	0.120*
IG d3	1.271	0.049*
ΔIG d3-d0	2.216	0.032*
MPV d0	1.131	0.566*
MPV d1	0.766	0.241*
MPV d3	2.361	0.001*
ΔMPV d3-d0	2.567	0.001*

*Logistic regression analysis. SOFA: Sequential organ failure assessment, IG: Immature granulocytes, MPV: Mean platelet volume, Δ: Difference

DISCUSSION

Sepsis is a health problem that affects all age groups worldwide and is a complication of peritonitis which can cause multiple organ failure and death.^[20] This mortality rate varies and depends on the severity of the disease, incidence rate, and organ failure, so that early detection of severe sepsis or septic shock is needed not only for risk stratification but also in monitoring the efficacy of treatment and disease progression.^[15]

In this study, mortality in patients with sepsis due to peritonitis who underwent emergency laparotomy was 34.58%. This mortality rate is in line with previous studies with settings that similar to this study. In another study, a 30-day mortality rate during hospital stay was 40%, and this mortality rate was

Table 3: The cutoff point for immature granulocytes and mean platelet volume in predicting 30 days mortality in sepsis patients with peritonitis

	AUROC	P	Cutoff
IG d0	0.678 (0.568-0.787)	0.003*	0.95
IG d1	0.665 (0.555-0.774)	0.005*	1.05
IG d3	0.703 (0.602-0.804)	0.001*	1.15
ΔIG d3-d0	0.619 (0.496-0.741)	0.044*	0.15
MPV d3	0.711 (0.607-0.815)	<0.001*	10.35
ΔMPV d3-d0	0.688 (0.579-0.798)	0.001*	0.35

AUROC: Area under the receiver operating characteristic curve, IG: Immature granulocytes, MPV: Mean platelet volume, Δ: Difference, *significant if $P < 0.05$

Table 4: Immature granulocytes and mean platelet volume predictive performance for 30-day mortality in sepsis patients with peritonitis

	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	McNemar test	κ
IG d0	67.6	62.9	51.0	78.6	0.034	0.279
IG d1	64.9	64.3	49.0	77.6	0.073	0.271
IG d3	67.6	62.9	51.0	78.6	0.034	0.279
ΔIG d3-d0	62.2	57.1	43.4	74.1	0.023	0.175
MPV d3	62.2	65.7	48.9	76.7	0.143	0.262
ΔMPV d3-d0	67.6	72.9	56.8	81.0	0.143	0.262

PPV: Positive predictive value, NPV: Negative predictive value, IG: Immature granulocytes, MPV: Mean platelet volume, Δ: Difference

significantly increased in patients with septic shock.^[21] Death in sepsis patients that occurred in the initial phase is caused by multiple organ failure due to primary infection, while in the final phase is mostly caused by secondary infection due to immunosuppression in the clinical course of sepsis.^[22,23]

In this study, older age was significantly associated with mortality within 30 days ($P = 0.015$). Some studies suggest that old age (geriatrics) can be an independent risk factor for predisposing to severe sepsis.^[24] There were several reasons why older people were more likely to develop infections. It has been established that immune function decreases with age; also known as immunosenescence, which makes older people vulnerable to an increased risk of infection and more severe and prolonged infections. The elderly were more susceptible to infection due to the process of body changes and decreased organ function and the presence of comorbid diseases. The diagnosis of sepsis in the elderly is even more difficult because the elderly provide responses and clinical symptoms of sepsis that are less clear, and sometimes accompanied by delirium.^[25]

The median SOFA score in the nonsurvivor group was significantly higher than in the survivor group. This finding was in accordance with previous research that stated that significantly high-SOFA scores were found in many of the nonsurvivor groups.^[11,26] SOFA scores at initial hospital admission and changes in SOFA scores during treatment have good accuracy

in predicting hospital mortality. Furthermore, SOFA scores are an accepted method for determining risk stratification and prognosis for severe sepsis patients at hospital admission.^[27,28]

The mean Hb at admission in the nonsurvivor group was significantly lower than in the survivor group, and both groups had anemia. These results were similar to the study by Jung *et al.*,^[29] which states that the percentage of patients with 90-day mortality increases with decreasing initial hemoglobin levels and initial hemoglobin levels independently associated with 90-day mortality and mortality increases proportionately with decreasing hemoglobin levels. Low hemoglobin levels at hospital admission indicate inadequate tissue oxygenation. They may reflect more severe inflammation compared to patients with septic shock with normal hemoglobin levels.^[29] Anemia in sepsis can be caused by several mechanisms, including a systemic inflammatory response process. It may cause a decrease in red blood cell production, and destruction of red blood cells due to hemolysis and bleeding due to disseminated intravascular coagulation. The combination of anemia and changes in oxygen consumption induced by sepsis can increase tissue oxygenation disorders, resulting in cellular hypoxia, cell dysfunction, and can end in multiple organ dysfunction syndromes, so that mortality increases.^[30]

The mean albumin at admission in the nonsurvivor group was lower than the survivor group. This result was in accordance with the previous study, where albumin results significantly lower in the group of nonsurvived patients.^[15] Takegawa *et al.*^[31] reported that the value and change of albumin in a short time reflects the risk of death in clinical and surgical patients undergoing treatment in the intensive care unit (ICU). Progressive decreases in serum albumin caused by the decreased synthesis in the liver have effects on colloidal osmotic pressure and carriers of endogenous or exogenous compounds. Therefore, they can dynamically change the permeability of blood vessels, which may contribute to death.

The IG count on day 0, 1, and 3 in the nonsurvivor patients was significantly higher than the survivors. Similarly, the Δ IG day 3-day 0 was significantly higher in the nonsurvivor group. This result was similar to the study by Nierhaus *et al.*^[32] in sepsis patients treated at the ICU. They reported a difference of IG in between survivor and nonsurvivor groups, in the period 15 (days 10–14 treatment; area under curve [AUC] = 0.617, $P = 0.042$) and the period 20 (days 15–21 treatment; AUC = 0.682, $P < 0.001$), but they stated that IG did not have prognostic power in predicting mortality despite the significant difference. This discrepancy occurred because the number of patients with infection relatively increased in the final period (day 10–21) due to several noninfectious patients who left the ICU early, so the balance of the two groups studied was leaning toward the infected population.

We found that the cutoff value of first day's IG as a predictor of mortality in septic patients due to peritonitis was 1.05, with a sensitivity of 64.9% and specificity of 64.3%. Previous studies had shown that IG could significantly distinguish the sepsis

group with complicated severe sepsis with a cutoff of 0.5% and can differentiate better than leukocytes, C-reactive protein (CRP), and procalcitonin.^[33]

This study shows that the MPV value of day 3 was valid as a predictor of mortality in sepsis patients due to peritonitis performed by emergency laparotomy. This finding was in line with a study by Vardon-Bounes *et al.*^[14] which showed that MPV values were significantly correlated with mortality, particularly the MPV value on the 10th day can predict 90-day survival of sepsis patients. Several previous studies have also shown that MPV increments are statistically significant in the first 3 days in Gram-positive sepsis patients and can predict 28-day mortality in septic shock.^[15,34]

This study shows that the value of Δ MPV d3-d0 ≥ 0.35 is a valid predictor of mortality in sepsis patients due to peritonitis underwent emergency laparotomy. This finding was in line with a study by Kim *et al.*^[15] based on the multivariate analysis shown that Δ MPV was an independent predictor of 28-day mortality after confounding factors were adjusted (hazard ratio = 1.44; 95% confidence interval = 1.01–2.06; $P = 0.044$), so an increase in MPV in the first 72 h of hospital admission was a poor predictor of 28-day mortality in sepsis patients.^[15] Daily monitoring of MPV values can stratify the risk of death in sepsis patients due to the changes in MPV during the 1st week after the onset of sepsis due to MPV values that do not return to baseline or normal values correlated with unfavorable outcomes.^[14]

Increased MPV indicates platelet diameter, which can be used as a marker for platelet activation, endothelial damage, and thrombotic and inflammatory conditions.^[15,17,18] The size of the MPV is a picture of the proinflammatory and prothrombotic conditions of the body. Large platelets are more functional, metabolic, and enzymatic active than platelets with smaller MPVs, so large platelets have more significant prothrombotic potential. Furthermore, the inflammatory process can induce procoagulant changes and cause embolization which is the most common cause of death in patients with systemic infections. MPV can be considered as an integrative measure of the inflammatory process and the destructive state of hypercoagulation in critical illness. Thus, the relationship between increased MPV and mortality in patients with sepsis can occur partly explained by this idea.^[15]

This study still has several limitations that might affect the results of the study. Patients were selected from a single health center with a retrospective design, so it may not cover the general population. Since this is a retrospective study, no further investigation was carried out on the confounding factors that may influence the IG and MPV values. The serological marker evaluations such as CRP, calcitonin, or lactate are not evaluated because they are not routinely carried out in our hospital.

CONCLUSION

We found that the IG on day-1, MPV on day-3, and mean difference of platelet volume between day-0 and day-3 are valid

predictors of mortality in sepsis patients due to peritonitis who underwent emergency surgery within 30 days. IG and MPV are easy, fast screening tool in stratifying the risk of mortality in sepsis patients due to peritonitis underwent emergency surgery.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, *et al.* Surviving sepsis campaign: International guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med* 2017;43:304-77.
- World Health Organization. Sepsis. Available from: <https://www.who.int/sepsis/en/>. [Last accessed on 2020 Feb 04].
- Fleischmann C, Scherag A, Adhikari NK, Hartog CS, Tsaganos T, Schlattmann P, *et al.* Assessment of global incidence and mortality of hospital-treated sepsis. Current estimates and limitations. *Am J Respir Crit Care Med* 2016;193:259-72.
- Zharfan R, Hakim A, Purba A, Sulistiawan S, Semedi B. Albumin, leukosit, and protrombin as predictor of sepsis mortality among adult patients in soetomo general hospital, Surabaya, Indonesia. *Indones J Anesthesiol Reanim* 2019;1:8-12.
- Cheng B, Xie G, Yao S, Wu X, Guo Q, Gu M, *et al.* Epidemiology of severe sepsis in critically ill surgical patients in ten university hospitals in China. *Crit Care Med* 2007;35:2538-46.
- Green S, Kong VY, Clarke DL, Sartorius B, Odendaal J, Bruce JL, *et al.* The spectrum and outcome of surgical sepsis in pietermaritzburg, South Africa. *S Afr Med J* 2017;107:134-6.
- Shukri KH. The burden of sepsis; a call to action in support of world sepsis day 2013. *Bull Emerg Trauma* 2013;1: 52-5.
- Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, *et al.* Surviving Sepsis campaign: International guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med*. 2013;39:165-228.
- Sweeney T, Wong H. Risk Stratification and prognosis in sepsis. *Clin Chest Med* 2016;37:209-18.
- Moore LJ, McKinley BA, Turner KL, Todd SR, Sucher JF, Valdivia A, *et al.* The epidemiology of sepsis in general surgery patients. *J Trauma* 2011;70:672-80.
- Kim JW, Park JH, Kim DJ, Choi WH, Cheong JC, Kim JY. The delta neutrophil index is a prognostic factor for postoperative mortality in patients with sepsis caused by peritonitis. *PLoS One* 2017;12:e0182325.
- Ostrand-Rosenberg S, Fenselau C. Myeloid-derived suppressor cells: Immune-suppressive cells that impair antitumor immunity and are sculpted by their environment. *J Immunol* 2018;200:422-31.
- Liu Y, Wei G, Cheng WA, Dong Z, Sun H, Lee VY, *et al.* Targeting myeloid-derived suppressor cells for cancer immunotherapy. *Cancer Immunol Immunother* 2018;67:1181-95.
- Vardon-Bounes F, Gratacap MP, Groyer S, Ruiz S, Georges B, Seguin T, *et al.* Kinetics of mean platelet volume predicts mortality in patients with septic shock. *PLoS One* 2019;14:e0223553.
- Kim CH, Kim SJ, Lee MJ, Kwon YE, Kim YL, Park KS, *et al.* An increase in mean platelet volume from baseline is associated with mortality in patients with severe sepsis or septic shock. *PLoS One* 2015;10:e0119437.
- Korniluk A, Koper-Lenkiewicz OM, Kamińska J, Kemona H, Dymicka-Piekarska V. Mean platelet volume (MPV): New perspectives for an old marker in the course and prognosis of inflammatory conditions. *Mediators Inflamm* 2019;2019:1-14.
- Budak YU, Polat M, Huysal K. The use of platelet indices, plateletcrit, mean platelet volume and platelet distribution width in emergency non-traumatic abdominal surgery: A systematic review. *Biochem Med (Zagreb)* 2016;26:178-93.
- Eberhardt A, Lessig F, Schreiter K, Fuchs M, Sablitzki A, Ludewig M, *et al.* Mean platelet volume (MPV) is an outcome marker in sepsis

- patients. *Int J Infect Dis* 2012;16:218.
19. Huang Z, Fu Z, Huang W, Huang K. Prognostic value of neutrophil-to-lymphocyte ratio in sepsis: A meta-analysis. *Am J Emerg Med* 2020;38:641-7.
 20. Riché FC, Dray X, Laisné MJ, Matéo J, Raskine L, Sanson-Le Pors MJ, *et al.* Factors associated with septic shock and mortality in generalized peritonitis: Comparison between community-acquired and postoperative peritonitis. *Crit Care* 2009;13:R99.
 21. Mačiulienė A, Maleckas A, Kriščiukaitis A, Mačiulis V, Vencius J, Macas A. Predictors of 30-day in-hospital mortality in patients undergoing urgent abdominal surgery due to acute peritonitis complicated with sepsis. *Med Sci Monit* 2019;25:6331-40.
 22. Otto GP, Sossdorf M, Claus RA, Rödel J, Menge K, Reinhart K, *et al.* The late phase of sepsis is characterized by an increased microbiological burden and death rate. *Crit Care* 2011;15:R183.
 23. Daviaud F, Grimaldi D, Dechartres A, Charpentier J, Geri G, Marin N, *et al.* Timing and causes of death in septic shock. *Ann Intensive Care* 2015;5:16.
 24. Nasa P, Juneja D, Singh O. Severe sepsis and septic shock in the elderly: An overview. *World J Crit Care Med* 2012;1:23-30.
 25. Wardani IS. Tatalaksana sepsis berat pada pasien lanjut usia. *Jurnal Kedokteran Unram* 2017;7:33-9.
 26. Park BH, Kang YA, Park MS, Jung WJ, Lee SH, Lee SK, *et al.* Delta neutrophil index as an early marker of disease severity in critically ill patients with sepsis. *BMC Infect Dis* 2011;11:299.
 27. Jones AE, Trzeciak S, Kline JA. The sequential organ failure assessment score for predicting outcome in patients with severe sepsis and evidence of hypoperfusion at the time of emergency department presentation. *Crit Care Med* 2009;37:1649-54.
 28. Karakike E, Kyriazopoulou E, Tsangaris I, Routsis C, Vincent JL, Giamarellos-Bourboulis EJ. The early change of SOFA score as a prognostic marker of 28-day sepsis mortality: Analysis through a derivation and a validation cohort. *Crit Care* 2019;23:387.
 29. Jung SM, Kim YJ, Ryoo SM, Kim WY. Relationship between low hemoglobin levels and mortality in patients with septic shock. *Acute Crit Care* 2019;34:141-7.
 30. Muady GF, Bitterman H, Laor A, Vardi M, Urin V, Ghanem-Zoubi N. Hemoglobin levels and blood transfusion in patients with sepsis in Internal Medicine Departments. *BMC Infect Dis* 2016;16:569.
 31. Takegawa R, Kabata D, Shimizu K, Hisano S, Ogura H, Shintani A, *et al.* Serum albumin as a risk factor for death in patients with prolonged sepsis: An observational study. *J Crit Care* 2019;51:139-44.
 32. Nierhaus A, Klatt S, Linssen J, Eismann NM, Wichmann D, Hedke J, *et al.* Revisiting the white blood cell count: Immature granulocytes count as a diagnostic marker to discriminate between SIRS and sepsis—a prospective, observational study. *BMC Immunol* 2013;14:8.
 33. Ha SO, Park SH, Park SH, Park JS, Huh JW, Lim CM, *et al.* Fraction of immature granulocytes reflects severity but not mortality in sepsis. *Scand J Clin Lab Invest* 2015;75:36-43.
 34. Aydemir H, Piskin N, Akduman D, Kokturk F, Aktas E. Platelet and mean platelet volume kinetics in adult patients with sepsis. *Platelets* 2015;26:331-5.

Immature Granulocyte and Mean Platelet Volume as a Predictor of 30-Day Postoperative Mortality in Patients with Sepsis Caused by Peritonitis

ORIGINALITY REPORT

23%

SIMILARITY INDEX

15%

INTERNET SOURCES

19%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Fanny Vardon-Bounes, Marie-Pierre Gratacap, Samuel Groyer, Stéphanie Ruiz et al. "Kinetics of mean platelet volume predicts mortality in patients with septic shock", PLOS ONE, 2019
Publication 1%
- 2** Surya Kanta Das, Ashish Kumar Barik, Choudhury Sabyasachi Swain. "Guiding arm template for manual set-up", Journal of Indian Orthodontic Society, 2019
Publication 1%
- 3** Jong Wan Kim, Jun Ho Park, Doo Jin Kim, Won Hyuk Choi, Jin Cheol Cheong, Jeong Yeon Kim. "The delta neutrophil index is a prognostic factor for postoperative mortality in patients with sepsis caused by peritonitis", PLOS ONE, 2017
Publication 1%
- 4** Theresa A. Rowe, June M. McKoy. "Sepsis in Older Adults", Infectious Disease Clinics of 1%

North America, 2017

Publication

-
- | | | |
|----|---|------|
| 5 | Ryosuke Takegawa, Daijiro Kabata, Kentaro Shimizu, Saya Hisano, Hiroshi Ogura, Ayumi Shintani, Takeshi Shimazu. "Serum albumin as a risk factor for death in patients with prolonged sepsis: An observational study", <i>Journal of Critical Care</i> , 2019
Publication | 1 % |
| 6 | link.springer.com
Internet Source | 1 % |
| 7 | Tuncay Kırış, Selcuk Yazici, Zeki Yüksel Günaydin, Şükrü Akyüz et al. "The Prognostic Impact of In-Hospital Change in Mean Platelet Volume in Patients With Non-ST-Segment Elevation Myocardial Infarction", <i>Angiology</i> , 2016
Publication | 1 % |
| 8 | www.e-sciencecentral.org
Internet Source | 1 % |
| 9 | mafiadoc.com
Internet Source | 1 % |
| 10 | www.scielo.br
Internet Source | 1 % |
| 11 | Fábio M. de Castilho, Antonio Luiz P. Ribeiro, José Luiz P. da Silva, Vandack Nobre, Marcos R. de Sousa. "Heart rate variability as | <1 % |

predictor of mortality in sepsis: A prospective cohort study", PLOS ONE, 2017

Publication

12	www.ajol.info Internet Source	<1 %
13	apm.amegroups.com Internet Source	<1 %
14	twitter.com Internet Source	<1 %
15	www.avicennajmed.com Internet Source	<1 %
16	www.dovepress.com Internet Source	<1 %
17	www.ijphrd.com Internet Source	<1 %
18	www.turkishjournalpediatrics.org Internet Source	<1 %
19	repository.ubaya.ac.id Internet Source	<1 %
20	seq.es Internet Source	<1 %
21	www.haievent.com Internet Source	<1 %

22

Leonardo Lorente, Alejandro Jiménez, María M. Martín, José Luis Iribarren, Juan José Jiménez, María L. Mora. "Clinical cure of ventilator-associated pneumonia treated with piperacillin/tazobactam administered by continuous or intermittent infusion", *International Journal of Antimicrobial Agents*, 2009

Publication

<1 %

23

Noor Al-Bader, Austin Meier, Matthew Geniza, Yamid Sanabria Gongora, James Oard, Pankaj Jaiswal. " Loss of premature stop codon in the () gene confers sheath blight disease resistance in rice ", *Cold Spring Harbor Laboratory*, 2019

Publication

<1 %

24

SEVİNÇ, Mert Mahsuni, KINACI, Erdem, ÇAKAR, Ekrem, BAYRAK, Savaş, ÖZAKAY, Abdulkerim, AREN, Acar and SARI, Serkan. "Diagnostic value of basic laboratory parameters for simple and perforated acute appendicitis: an analysis of 3392 cases", *Ulusal Travma ve Acil Cerrahi Derneği*, 2016.

Publication

<1 %

25

Ariani Arista Putri Pertiwi, Dan Fraczkowski, Sheryl L. Stogis, Karen Dunn Lopez. "Using Heuristic Evaluation to Improve Sepsis Alert

<1 %

Usability", Critical Care Nursing Clinics of North America, 2018

Publication

26

Syed Mustafa Karim, Jamal Zekri, Ehab Abdelghany, Reyad Dada, Husna Munsoor, Imran Ahmad. "Time from last chemotherapy to death and its correlation with the end of life care in a referral hospital", Indian Journal of Medical and Paediatric Oncology, 2021

Publication

<1 %

27

W. Strayer, Robert Walsh, Carl Livadas, David Lapsley. "Detecting Botnets with Tight Command and Control", Proceedings. 2006 31st IEEE Conference on Local Computer Networks, 2006

Publication

<1 %

28

synapse.koreamed.org

Internet Source

<1 %

29

www.apsr2018.com

Internet Source

<1 %

30

www.tevak.org

Internet Source

<1 %

31

Belousova, Hassan, Lampkin. "Mixed-Method Study of Utilizing Portfolios to Document and Assess Co-Curricular Activities: Student and Advisor Perceptions", Pharmacy, 2019

Publication

<1 %

32 Haytham Wali, William Malik, Juliane Mayette. "Efficacy of Using High-Dose Vitamin C, Thiamine, and Hydrocortisone in Sepsis and Septic Shock: A Retrospective Before-After Study", Research Square Platform LLC, 2021
Publication <1 %

33 Yuxing Zhang, Jian Chen, Chunyan Bai, Dawei Zhang, Qiwen Zhan. "Dynamical generation of multiple focal spot pairs with controllable position and polarization", Optics Express, 2020
Publication <1 %

34 ejournal.unsri.ac.id
Internet Source <1 %

35 academic.hep.com.cn
Internet Source <1 %

36 ccforum.biomedcentral.com
Internet Source <1 %

37 jag.journalagent.com
Internet Source <1 %

38 repozitorij.svkst.unist.hr
Internet Source <1 %

39 www.thoughtco.com
Internet Source <1 %

40 Cihan Bedel, Mustafa Korkut, Ali Avcı, Ahmet Uzun. "Immature Granulocyte Count and <1 %

Percentage as New Predictors of Mortality in Patients with Upper Gastrointestinal Bleeding", Indian Journal of Critical Care Medicine, 2020

Publication

41

Elvira Grandone, Raffaele Pesavento, Giovanni Tiscia, Antonio De Lorenzo et al. "Mortality and Transfusion Requirements in COVID-19 Hospitalized Italian Patients According to Severity of the Disease", Journal of Clinical Medicine, 2021

Publication

<1 %

42

Varsha Wilson, Kunnothara Kantan Velayudhan, Harshavardhan Rao, Sheejamol Velickakathu Sukumaran. "Low Absolute Eosinophil Count Predicts In-Hospital Mortality in Cirrhosis With Systemic Inflammatory Response Syndrome", Cureus, 2021

Publication

<1 %

43

bmcimmunol.biomedcentral.com

Internet Source

<1 %

44

epublications.uef.fi

Internet Source

<1 %

45

www.ajp.eg.net

Internet Source

<1 %

46

www.bjoaonline.com

Internet Source

<1 %

47

Gao, Yanxia, Yi Li, Xuezhong Yu, Shigong Guo, Xu Ji, Tongwen Sun, Chao Lan, Valery Lavergne, Marc Ghannoum, and Li Li. "The Impact of Various Platelet Indices as Prognostic Markers of Septic Shock", PLoS ONE, 2014.

Publication

<1 %

48

Guillermo Rodriguez-Nava, Goar Egoryan, Tianyu Dong, Qishuo Zhang et al. "Comparison of Clinical Characteristics and Outcomes of Hospitalized Patients with Seasonal Coronavirus Infection and COVID-19: A Retrospective Cohort Study", Research Square Platform LLC, 2021

Publication

<1 %

49

Gönül Açıksarı, Mehmet Koçak, Yasemin Çağ, Lütfiye Nilsun Altunal et al. "Prognostic Value of Inflammatory Biomarkers in Patients with Severe COVID-19: A Single-Center Retrospective Study", Biomarker Insights, 2021

Publication

<1 %

50

Hayato Go, Hitoshi Ohto, Kenneth E. Nollet, Shunya Takano et al. "Using Platelet Parameters to Anticipate Morbidity and Mortality Among Preterm Neonates: A

<1 %

Retrospective Study", *Frontiers in Pediatrics*,
2020

Publication

51

Jian-Feng Wu. "Changes of monocyte human leukocyte antigen-DR expression as a reliable predictor of mortality in severe sepsis", *Critical Care*, 2011

Publication

<1 %

52

Pravin Ambure, Arantxa Ballesteros, Francisco Huertas, Pau Camilleri, Stephen J. Barigye, Rafael Gozalbes. "Development of Generalized QSAR Models for Predicting Cytotoxicity and Genotoxicity of Metal Oxides Nanoparticles", *International Journal of Quantitative Structure-Property Relationships*, 2020

Publication

<1 %

53

Xiao-Fei Shen, Ke Cao, Jin-peng Jiang, Wen-Xian Guan, Jun-Feng Du. "Neutrophil dysregulation during sepsis: an overview and update", *Journal of Cellular and Molecular Medicine*, 2017

Publication

<1 %

54

bmcinfectdis.biomedcentral.com

Internet Source

<1 %

55

etheses.bham.ac.uk

Internet Source

<1 %

56	medinform.jmir.org Internet Source	<1 %
57	repositorio.unicamp.br Internet Source	<1 %
58	www.roaic.eg.net Internet Source	<1 %
59	www.smj.org.sa Internet Source	<1 %
60	www.thieme-connect.com Internet Source	<1 %
61	"Day 4 - Tuesday 31 August 2004", European Heart Journal, 09/02/2004 Publication	<1 %
62	Seong Yeob Lee, Jin Joo Kim, Jae Ho Jang, In Cheol Hwang. "Prognostic Factors in Septic Shock Patients on Arrival at Emergency Department", Annals of Geriatric Medicine and Research, 2017 Publication	<1 %
63	Gu-Lung Lin, Joseph P. McGinley, Simon B. Drysdale, Andrew J. Pollard. "Epidemiology and Immune Pathogenesis of Viral Sepsis", Frontiers in Immunology, 2018 Publication	<1 %
64	I Re Lee, Jae Il Shin, Se Jin Park, Ji Young Oh, Ji Hong Kim. "Mean platelet volume in young	<1 %

children with urinary tract infection", Scientific Reports, 2015

Publication

65

Rahul Chaturvedi, Brittany N. Burton, Suraj Trivedi, Ulrich H. Schmidt, Rodney A. Gabriel. "The Association of Preoperative Hematocrit With Adverse Events Following Exploratory Laparotomy in Septic Patients: A Retrospective Analysis", Journal of Intensive Care Medicine, 2020

Publication

<1 %

66

Yasemin Ustundag Budak, Murat Polat, Kagan Huysal. "The use of platelet indices, plateletcrit, mean platelet volume and platelet distribution width in emergency non-traumatic abdominal surgery: a systematic review", Biochemia Medica, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Immature Granulocyte and Mean Platelet Volume as a Predictor of 30-Day Postoperative Mortality in Patients with Sepsis Caused by Peritonitis

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
