



## About Us

**Bukti C.5**

**EurAsian Journal of BioSciences (Eurasia J Biosci) has ceased publication.**

**EurAsian Journal of BioSciences (Eurasia J Biosci)** e-ISSN 1307-9867 (online) is an international, refereed electronic journal. It publishes the results of original research in the field of biological sciences especially related to morphology, physiology, genetics, taxonomy, ecology and biogeography of both prokaryotic and eukaryotic organisms.

**EurAsian Journal of BioSciences** is indexed by AGRICOLA, EBSCOhost, ProQuest, SCOPUS, Zoological Record (Thomson ISI) etc. All published articles are assigned to Digital Object Identifier (DOI)-CrossRef.

**EurAsian Journal of BioSciences** is an Open Access Journal.

**Eurasia J Biosci** is published by Foundation for Environmental Protection and Research (FEPR) is a non-profit, volunteer organization. FEPR was founded by young academicians in 1991 who, instead of complaining about the environmental problems, thought that it was a duty to do something to protect the environment.



**Development of single nucleotide polymorphism markers for bacterial leaf blight resistance in rice**

Shahril Ab Razak, Rahiniza Kamaruzaman, Kogeethavani Ramachandran, Nor Helwa Ezzah Nor Azman, Siti Norhayati Ismail, Muhammad Fairuz Mohd Yusof, Asfaliza Ramli

[Download Full Article - PDF format](#)   [Abstract](#)

**Non-thermal carbon dioxide-mediated inactivation process of *Escherichia coli* using bespoke system**

Ali R. Mulakhudair

[Download Full Article - PDF format](#)   [Abstract](#)

**Phytochemicals screening and antibacterial activity of silver nanoparticles, phenols and alkaloids extracts of conocarpus lancifolius**

Rana Hussein Raheema, Roaa M. H. Shoker

[Download Full Article - PDF format](#)   [Abstract](#)

**The clean and healthy lifestyle behaviors and healthy housing influenced the incident of acute respiratory infection in childhood**

Muhammad El Kautsar, Nilam Anggraeni, Vivi Permata Sari, Ribka B. A Mboe, Anindya Widyasari, Zahwa Afdhilani Azwan, Athiyah Amatillah, Priscillia Ivana Jie, **Sulistiawati**, R Bambang

[Download Full Article - PDF format](#)   [Abstract](#)

**Isolation of different trematodes cercariae spp. of medical and veterinary importance from Shatt-Al-Arab water south of Iraq**

Shaimaa A. Alsamir

[Download Full Article - PDF format](#)   [Abstract](#)

**Evaluate the effect of periodontally accelerated osteogenic orthodontics on gingivae and alveolar bone: a systematic review and meta-analysis**

Meysam Haghghat, Mohammad Ghasemirad, Maryam Sarbaz, Kimiya Sezavar

[Download Full Article - PDF format](#)   [Abstract](#)



## The clean and healthy lifestyle behaviors and healthy housing influenced the incident of acute respiratory infection in childhood

Muhammad El Kautsar<sup>1</sup>, Nilam Anggraeni<sup>1</sup>, Vivi Permata Sari<sup>1</sup>, Ribka B. A Mboe<sup>1\*</sup>, Anindya Widyasari<sup>1</sup>, Zahwa Afdhilani Azwan<sup>1</sup>, Athiyah Amatillah<sup>1</sup>, Priscillia Ivana Jie<sup>1</sup>, Sulistiawati<sup>2</sup>, R Bambang<sup>3</sup>

<sup>1</sup> Faculty of Medicine, Universitas Airlangga, Surabaya, INDONESIA

<sup>2</sup> Department Public Health and Prevention Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, INDONESIA

<sup>3</sup> Technical Service Unit of Latkesmas Mumajati, Lawang, Malang, INDONESIA

\*Corresponding author: ribka.brillyand.angeli-2018@fk.unair.ac.id

### Abstract

**Introduction:** Based on preliminary surveys in this study, it was found 86.7% of respondents suffered from acute respiratory infection experienced 4 times reinfection within 3 months and 50% of respondents rarely consumed balanced nutritious food. **Aims:** to find the correlation between clean and healthy lifestyle behaviours, healthy housing and the frequency of acute respiratory infection in children under the age of 1-4 years. **Method:** a cross-sectional study was used by involving mothers or caregivers with infants aged 1-4 years in Malang Regency. A questionnaire was applied as research instrument. Data obtained were conducted by a nonparametric contingency coefficient test with IBM SPSS v17. **Result:** Statistical analysis shows that there is a significant correlation between family members who smoke ( $p\text{-value}=0.021$ ) and the level of home density ( $p\text{-value}=0.03$ ) with the frequency of acute respiratory tract infections in toddlers. While the relationship of other variables such as the sex of children under five ( $p\text{-value}=0.799$ ), clean and healthy lifestyle behaviour: delivery at primary health care ( $p\text{-value}=0.084$ ), exclusive breastfeeding ( $p\text{-value}=0.940$ ), routinely visiting Integrated Healthcare Center ( $p\text{-value}=0.396$ ), hand washing behaviour ( $p\text{-value}=0.523$ ), consuming healthy food ( $p\text{-value}=0.247$ ), and infant's activity ( $p\text{-value}=0.096$ ), healthy housing: ventilation ( $p\text{-value}=0.396$ ) and lighting ( $p\text{-value}=0.767$ ) have no a significant correlation with incident of acute respiratory infection in infants. **Conclusion:** There is a correlation between family members who smoke and the level of home density with the frequency of acute respiratory infection in children under the age of 1-4 years.

**Keywords:** clean and healthy lifestyle behaviours, healthy housing, frequently of acute respiratory infection, children, health behavior

Kautsar ME, Anggraeni N, Sari VP, Mboe RBA, Widyasari A, Azwan ZA, Amatillah A, Jie PI, Sulistiawati, Bambang R (2020) The clean and healthy lifestyle behaviors and healthy housing influenced the incident of acute respiratory infection in childhood. Eurasia J Biosci 14: 4837-4842.

© 2020 Kautsar et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

### INTRODUCTION

Acute Respiratory Infection is a disease mostly affected on children in the age of 5-14 years, the prevalence of infected between men and women seems almost the same, namely 510,714 and 506,576 respectively. Housing is an influential factor, where more people in urban areas are infected than in rural areas (Health 2018). This disease is caused by a small substance (smaller than bacteria) - called virus - that causes infection and transmitted from one person to another through coughing or sneezing (Ellita 2013).

Acute Respiratory Infection is a disease mostly affected on children in the age of 5-14 years, the prevalence of infected between men and women seems almost the same, namely 510,714 and 506,576

respectively. Housing is an influential factor, where more people in urban areas are infected than in rural areas (Health 2018). This disease is caused by a small substance (smaller than bacteria) - called virus - that causes infection and transmitted from one person to another through coughing or sneezing (Ellita 2013).

According to WHO (2001), a house is a physical structure or building for shelter, where the environment is useful for physical and spiritual health and social conditions both for the health of families and individuals (World Health Organization 2008). Thus it can be said

Received: October 2019

Accepted: March 2020

Printed: October 2020

**Table 1.** The distribution of clean and healthy lifestyle behaviors, healthy housing, and acute respiratory infection

|   | Yes |      | No |      |
|---|-----|------|----|------|
|   | n   | %    | n  | %    |
| Clean and healthy lifestyle Behaviors                   |     |      |    |      |
| Delivery at primary health care                         | 50  | 98   | 1  | 2    |
| Exclusive breastfeeding                                 | 24  | 47.1 | 27 | 52.9 |
| Regularly visiting Integrated Healthcare Center         | 43  | 84.3 | 8  | 15.7 |
| Wash hand behavior                                      | 35  | 68.6 | 16 | 31.4 |
| Consuming healthy food                                  | 51  | 86.3 | 7  | 13.7 |
| Exercise  | 44  | 86.3 | 7  | 13.7 |
| Smoking at home   | 39  | 76.5 | 12 | 23.5 |
| Healthy Housing   |     |      |    |      |
| Good Lighting   | 46  | 90.2 | 5  | 9.8  |
| Ceramic floor   | 48  | 94.1 | 3  | 5.9  |
| Room Ventilation >10%                                   | 8   | 15.7 | 43 | 84.3 |
| Large of home >8 m <sup>2</sup> / person                | 15  | 29.4 | 36 | 70.6 |
| Acute respiratory infection in children within 3 months | 38  | 74.5 | 13 | 25.5 |

that a healthy housing is a building for shelter and rest as well as a means of family coaching that fosters healthy living physically, mentally and socially, hence all family members can work productively. Healthy housing have criteria to prevent the occurrence of acute respiratory tract infections consisting of: the walls of the house, the floor of the house, air ventilation, residential density and lighting must be in accordance with the requirements of a healthy home so that humidity does not occur. Healthy housing factors can prevent the growth of a variety of bacteria and pneumonia viruses (Nurjazuli 2009). Based on preliminary survey, approximately 86.7% of respondents suffered from acute respiratory infections 4 times in the last 3 months, and 50% of them consumed balanced nutritious food in low levels.

The study aims to find the relationship between clean and Healthy Behavior, Healthy Housing and the frequency of acute respiratory infections in children under the age of 1-4 years.

## METHOD

### Participant

A cross sectional with observational analytic approach was used in this study. The population was children aged under 5 years in Sumber Mlaten sub-village, Kalirejo sub-district, Lawang sub-district, East Java in 2019. The sample was taken using the total sampling method and meets the inclusion criteria which include: families of children aged 1-4 years who are willing to become respondents (mother/caregiver/other family), present during data collection

### Data Collection and Analysis

The research instrument was a questionnaire that had been tested for validity in previous studies. Questionnaire questions included data on parental education, parental occupation, and sex of children under five years, infection status, clean and healthy lifestyle behaviours, and criteria for a healthy home referring to the Republic of Indonesia Ministry of Health (2016). The research period was conducted for 7 days, starting from 8 December 2019 to 14 December 2019.

The data collected was performed by processing stages which include cleaning, editing, coding and entry. After the data entered, we analyzed using a bivariate test with chi-square test and contingency coefficient test to determine the correlation between clean and healthy lifestyle behaviours, Healthy Homes and acute respiratory infection in Children aged under five years. The data was analysed using statistical software program Statistical Package for the Social Science (SPSS) version 17.0 (SPSS.Inc., Chicago, IL).

### Ethics Statement

This study was approved by the Ethics Committee of the Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. Written informed consent was obtained from participants prior to their enrolment in this study.

## RESULT

Based on inclusion criteria's, overall participants in this study were 51 participants divided by mother and caregiver of children. The socio-demographic data included basic variables of mother or caregiver and children. Participants who have aged between 18-35 years and having primary school history have the percentage as much as 60.8% and 41.2%, respectively. Meanwhile, the number of participants were mostly unemployed status about 82.4%. The highest children age in this study was 2 years, namely 35.3% and female's children were the majority of children which counted as many as 56.9%. The immunization status were dominated by children who have completed immunization status, namely 94.1% (Fig. 1).

Table 1 represented three variable which divided into clean and healthy lifestyle behaviors, healthy housing and incidence of acute respiratory infection in children within 3 months. According to clean and healthy lifestyle behaviors, it revealed that about 98% of participants visited primary health care as the method to deliver their children, 52.9% of participants have no given exclusive breastfeeding to their children. Moreover, approximately 84.3% of participants were routinely visiting integrated healthcare center, 68.8% of participants had gotten used to washing their hands




Fig. 1. Socio Demography Participants

using clean water. All participants declared that they had provided enough nutrition for their children (100%) and almost all participants have a settled job (86.3%). The smoking at home was done by 76.5% of participants. Healthy housing consisted by several factors, including natural lighting to the room which experienced by 90.2% of participant, floor of housing using tiles or ceramics had have by 94.1%. Furthermore, as much as 84.3% of house has ventilation area <10%, and density <8m<sup>2</sup>/person was 70.6%.

Bivariate analysis shows several factors associated with incident of acute respiratory infection in children aged 1-4 years (Table 2). Parents who have smoked habit at home were more likely to increase the incident of acute respiratory infection in children compared to parents who had no smoked (p-value=0.021). Further,

Table 2. Statistical Analysis between variables

|   | Experiencing acute respiratory infection in infant within last 3 months |    | P Value |
|---|---|----|---------|
|   | Yes   | No |         |
| Total family members                            |   |    |         |
| >4  | 4   | 4  | 0.083   |
| <4  | 34  | 9  |         |
| Immunization status                             |   |    |         |
| Complete  | 36  | 12 | 0.748   |
| uncomplete                                      | 2   | 1  |         |
| Exclusive Breastfeeding                         |   |    |         |
| Yes   | 18  | 6  | 0.940   |
| No  | 20  | 7  |         |
| Regularly visiting Integrated Healthcare Center |   |    |         |
| Yes   | 33  | 10 | 0.396   |
| No  | 5   | 3  |         |
| Parents smoking behavior                        |   |    |         |
| Yes   | 26  | 13 | 0.021   |
| No  | 12  | 0  |         |
| Delivery at primary health care                 |   |    |         |
| Yes   | 38  | 12 | 0.084   |
| No  | 0   | 1  |         |
| Wash hand behavior                              |   |    |         |
| Yes   | 27  | 8  | 0.523   |
| No  | 11  | 5  |         |
| Infant activity                                 |   |    |         |
| Yes   | 31  | 13 | 0.096   |
| No  | 7   | 0  |         |
| Consuming healthy food                          |   |    |         |
| Yes   | 38  | 13 | 0.247   |
| No  | 0   | 0  |         |
| Large of home                                   |   |    |         |
| >8 m <sup>2</sup> / person                      | 7   | 8  | 0.003   |
| <8 m <sup>2</sup> / person                      | 31  | 5  |         |
| Room Ventilation                                |   |    |         |
| <10%  | 33  | 10 | 0.396   |
| >10%  | 5   | 3  |         |
| Room lighting                                   |   |    |         |
| Sunshine  | 34  | 12 | 0.767   |
| no  | 4   | 1  |         |
| Kind of floor                                   |   |    |         |
| Ceramic   | 35  | 13 | 0.296   |
| None  | 3   | 0  |         |

population density which was <8m<sup>2</sup>/person, was more likely to increase the incident of acute respiratory infection in children compared to population density under 8m<sup>2</sup>/person (p-value=0.003). Whereas, variables related to children, such as total family members, immunization status, excusive breastfeeding and regularly visiting integrated healthcare center showed no correlation with the incident of acute respiratory infection as the p-values were above the minimum standard, namely 0.083, 0.748, 0.940, 0.396, respectively. With regard to clean and healthy lifestyle behaviors variables, included: delivery at primary healthcare, washing hand, children activity, and consuming healthy food were less likely affected on the incident of acute respiratory infection, which recorded p-value as much as 0.084, 0.523, 0.096, 0.247, respectively. While, other healthy housing components, such as: room ventilation, room lighting, kind of floor have less likely significant correlation he incident of acute respiratory infection, which the p-value were 0.396, 0.767, 0.296, respectively.

## DISCUSSION

Bivariate analysis showed that there was no significant relationship between hands washing behavior with the incidence of acute respiratory infections in children aged 1-4 years. This is in accordance with previous studies which produced a significant relationship between the incidences of acute respiratory infections with hand washing behavior. However, there was no significant relationship between exclusive breastfeeding ( $p=0.01$ ;  $OR=0.113$ ), use of latrines ( $p=0.00$ ;  $OR=0.93$ ), daily giving of vegetables and fruit ( $p=0.01$ ;  $OR=0.311$ ), and not smoking in the house ( $p=0.00$ ;  $OR=0.113$ ) with acute respiratory infection occurrence in children under the age of 12-59 months (Prayitno 2019). Acute respiratory infections is the leading cause of death for children under five and hand washing habits can reduce respiratory infections by more than 50% (Kementerian Kesehatan RI 2014; Merk et al. 2014).

We found as many as 31 children under five (61%) had acute respiratory infections in the last 3 months who actively played both outside and inside the house. Chi Square analysis results obtained  $p$  value of 0.096. These results indicate that there is no positive effect of playing activity with the incidence of acute respiratory infections in children under the age of 1-4 years. This is contrast with study that revealed a relationship between physical activity and morbidity, especially in the respiratory tract. The results of the study explained that subjects who did less activity were more susceptible to illness or often experienced pain compared to those who did many activities (Tomatala et al. 2019). Regular physical activity is one way to maintain the immune system. Various studies have confirmed the beneficial effect of exercise activities on the components of the immune system. Regular physical activity also teaches the body to distribute blood better to muscles during activity (Purwanto 2011). Physical activity and regular exercise and sufficient quantities, can help maintain optimal health for those concerned (Kepmenkes RI 1999).

The results showed that there was no significant influence between regularly visiting integrated healthcare center and acute respiratory infection in children under five in Sumber Mlaten Hamlet. This contradicts with other studies to 51 respondents that the incidence of acute respiratory infection in infants and regular visiting to integrated healthcare center have a signification of attachment (Fatmawati 2017). Visiting integrated healthcare center is a form of Community-Based Health Efforts which manage and organize from, by, for and with the community in organizing health development, in order to empower the community and provide facilities to the community in obtaining basic health services. To accelerate the reduction in maternal

and infant mortality rates (Direktorat Jendral P2PL 2011).

In this study, we found that the correlation between breastfeeding and acute respiratory infection was not significant with a significance value of 0.94 ( $p> 0.05$ ). The results of this study are consistent with research in 2011 which showed that there was no relationship between exclusive breastfeeding and the incidence of acute respiratory infection in children aged 6 months to 24 months (Lestari 2011). However, this study differs from other studies that there is a relationship between duration of exclusive breastfeeding exclusively with acute respiratory infections (Prameswari, G. 2009). Breastfeeding has a protective effect of acute respiratory infection during the first year, but with increasing age of children under five years the level of immunoglobulin in breast milk will decrease so that if breastfeeding is given longer does not have much effect on body resistance to overcome microorganisms that cause acute respiratory infection (Endiyani et al. 2009).

Smoking habits of parents showed highly likely affected the incident of acute respiratory infection in children as it exposure to cigarette smoke. This data is supported by previous studies that shows that there is a positive influence between exposure to cigarette smoke to acute respiratory infection in children under five (Riyanto & Kusumawati 2016; Hidayangsih et al. 2017; Mahendra & Farapti 2018). Basically, smoking is one of the risk factors for acute respiratory infection in children under five, especially in East Java. Based on the results of Basic Health Research (Risikesdas) in 2007, showed the prevalence of acute respiratory infection at 6.4%, while in Malang Regency the prevalence of acute respiratory infection was 4.3% (Department of Health in Malang 2013). According to WHO (2008), the adverse effects of cigarette smoke on Passive smokers are greater than active smokers. Side stream smoke or side smoke caused by the activity of passive smokers, is proven to contain more tobacco burners than mainstream smoke. This side smoke is proven to contain carbon monoxide 5 times larger, tar and nicotine 3 times, ammonia 46 times, nickel 3 times, nitrosamine as a cause of cancer levels reached 50 times greater in side smoke compared with the main smoke levels (World Health Organization 2008).

These results indicate no significant effect on childbirth in health professionals and the incidence of acute respiratory infection in children under five. Women treated by traditional healers do not tend to get good early breastfeeding initiation so that neonates can lose important nutrients in colostrum which may cause health problems in the future (Virarisca et al. 2019). Childbirth performed by health professionals, doctors, midwives or paramedics has a stake in the prevention of infections and other hazards that can threaten the safety of babies and mothers (Direktorat Jendral P2PL 2011) placenta does not come out, bleeding and infant death. However,

due to ignorance of information, participants did not realize that the impact was very dangerous to their health (Lestari 2014). In neonatal births, the first hour of life is very important in neonates' life. There are interventions to be done by health workers who understand the condition of the neonate, including paying attention to nutrition, seeing whether there are congenital abnormalities, providing support to the respiratory tract, cardiovascular system, and preventing infection. If the treatment has not done properly, it can interfere with the growth, development and health of children in the future (Sharma 2017).

Population density in house has significant association with incident of acute respiratory infection in children and another have proven the similar result as this finding (Firdaus & Ahmad 2013). Mostly of respiratory problems, such as acute respiratory infection, were affected with overcrowding, indoor and poor condition of the houses (Firdaus & Ahmad 2013).

The results of the bivariate analysis interpreted that there is no significant correlation between lighting with acute respiratory infection incident in children under five. This finding is contrast with previous study in Indonesia which showed the significant correlation between indoor air pollution including lighting source in home, and incident of acute respiratory infection in children under five years (Hidayangsih et al. 2017). A healthy housing requires sufficient light, both artificial lighting and natural lighting from the sun (Agustina & Mado 2018).

The type of floor has no associated with acute respiratory infection in children under five as it showed from bivariate analysis which contained  $<0.005$ . This is

different from study that revealed the variable of the type of house floor which cumulatively can be interpreted to have a significant relationship between the type of floor with the incidence of acute respiratory infection in children under five (Agustina & Mado 2018). The same thing was expressed in study that showed the condition of the house floor with the occurrence of acute respiratory infection in children under five years old in the Work Area of the Salibabu Public Health Center in Talaud Islands Regency had a significant relationship with the value ( $p = 0,000$ ). The type of house floor affects the incidence of acute respiratory infection in children under five as houses that have ceramic or tile floor types tend to be better because they are easy to clean and not damp. Conversely, only casted floors tend to be damp, not waterproof, and can be a breeding ground for bacteria or viruses that cause acute respiratory infection (Bee et al. 2015).

### CONCLUSION

In the criteria for a healthy home, there is a relationship between the points of population density and the incidence of acute respiratory infection in children under five, while in other criteria (giving birth in health workers, giving exclusive breastfeeding, routinely bringing to the primary care, washing hands, giving nutritious food) is not found a correlation. The correlation between smoking points in the house and the incidence of acute respiratory infection in children under five years old have proven, while in the other criteria (Ventilation Area, Natural Lighting, and Floor Type) is no correlation.

### REFERENCES

- Agustina and Mado, F. G. (2018) 'Analysis of Housing Environmental Factors with Arrival Disease Events in Communities in Kelurahan Oenesu District Kupang in 2018', in 1 international conference. Kupang: Health Polytechnic of Ministry of Health in Kupang.
- Bee, L. W., Akili, R. H. and Sinolungan, J. V. S. (2015) 'Hubungan Antara Kondisi Lingkungan Fisik Rumah Dengan Kejadian Infeksi Saluran Pernapasan Akut (ISPA) Pada Anak Balita Di Wilayah Kerja Puskesmas Salibabu Kabupaten Kepulauan Talaud', *Kesmas*, 4(2). Available at: <https://ejournal.unsrat.ac.id/index.php/kesmas/article/view/12692>.
- Department of Health in Malang (2013) Health Profile. Malang.
- Direktorat Jendral P2PL (2011) Rencana Aksi Program Pengendalian Penyakit dan Penyehatan Lingkungan tahun 2010-2014. Jakarta.
- Ellita (2013) Hubungan antara Tingkat Pengetahuan Ibu dan Pemberian ASI dengan Kejadian ISPA Pada Anak balita di Rumah Sakit BLUD Ibu dan Anak di Pemerintah Aceh Tahun 2013. Universitas Aceh.
- Endiyani, B. et al. (2009) 'Infeksi Saluran napas Akut pada Anak balita di Daerah Urban Jakarta', *Seri Pediatri*, 11(4). Available at: <http://www.idai.or.id/saripediatri/pdf/11-4-%0A1.pdf>.
- Fatmawati, T. Y. (2017) Pengaruh Pendidikan Kesehatan dengan Media Leaflet terhadap Pengetahuan Ibu tentang Penatalaksanaan ISPA pada Anak balita di Posyandu. Universitas Batanghari Jambi.
- Firdaus, G. and Ahmad, A. (2013) 'Relationship between housing and health: A cross-sectional study of an urban centre of India', *Indoor and Built Environment*, 22(3), pp. 498–507. doi: 10.1177/1420326X12443846.
- Health, M. O. (2018) 'Riset Kesehatan Dasar (Riskesdas)', Ministry of Health Indonesia.

- Hidayangsih, P. S. et al. (2017) 'The Relationship between Respiratory Infections and Healthy Homes in Children Under Five, Indonesia 2013', *Media Penelitian dan Pengembangan Kesehatan*, 27(3). doi: 10.22435/mpk.v27i3.6663.153-160.
- Kementerian Kesehatan RI (2011) *Buku Pedoman PHBS, Kementerian Kesehatan Republic Indonesia*. Available at: <http://promkes.kemkes.go.id/phbs> (Accessed: 8 December 2019).
- Kementerian Kesehatan RI (2014) *Perilaku Mencuci Tangan Pakai Sabun di Indonesia*. Jakarta.
- Kepmenkes RI (1999) No: 829/MENKES/SK/VII/1999 tentang Persyaratan Kesehatan Perumahan.
- Lestari, P. (2011) Hubungan antara Pemberian ASI Eksklusif dengan kejadian ISPA pada Anak Usia di atas 6 bulan sampai 24 bulan di Dusun Gondang, Desa Randuagung, Kecamatan Singosari, Kabupaten Malang. Surabaya. Available at: [http://penelitian.unair.ac.id/detail\\_dosen\\_2153](http://penelitian.unair.ac.id/detail_dosen_2153).
- Lestari, R. D. (2014) 'Analisis Persalinan Dengan Tenaga Non Kesehatan Di Wilayah Kerja Puskesmas Punggur Kabupaten Kubu Raya Tahun 2014', *Jurnal proners*, 1(1).
- Mahendra, I. G. A. P. and Farapti, F. (2018) 'Relationship between Household Physical Condition with The Incidence of ARI on Todler at Surabaya', *Jurnal Berkala Epidemiologi*, 6(3), p. 227. doi: 10.20473/jbe.v6i32018.227-235.
- Merk, H. et al. (2014) 'Associations of hand-washing frequency with incidence of acute respiratory tract infection and influenza-like illness in adults: A population-based study in Sweden', *BMC Infectious Diseases*, 14(1). doi: 10.1186/1471-2334-14-509.
- Nurjazuli, W. R. (2009) 'Faktor Risiko Dominan Kejadian pneumonia pada Anak balita. Jakarta', *Jurnal Respirologi Indonesia*, 29(2).
- Prameswari, G., N. (2009) 'Hubungan Lama Pemberian ASI Eksklusif dengan Terjadinya ISPA', *Kemas*, 5(1). Available at: <http://journal.unnes.ac.id/index.php/kemas>.
- Prayitno, A. (2019) 'Hubungan Perilaku Hidup Bersih Dan Sehat Rumah Tangga Terhadap Kejadian Infeksi Saluran Pernafasan Akut Dan Diare Pada Anak Balita', *Respiratory Unair*. Available at: <http://repository.unair.ac.id/84884/Repository.unair.ac.id>.
- Purwanto (2011) 'Dampak Senam Aerobik terhadap Daya Tahan Tubuh dan Penyakit.', *Jurnal Media Ilmu Keolahragaan Indonesia*, 1.
- Riyanto, R. and Kusumawati, A. (2016) 'Pengaruh asap rokok terhadap terjadinya penyakit ISPA pada anak balita di puskesmas Kedung Banteng Banyumas', *Medisains*, 14(3). Available at: <http://jurnalnasional.ump.ac.id/index.php/medi%0Asains/artic%0Ale/view/1614>.
- Sharma, D. (2017) 'Golden hour of neonatal life: Need of the hour', *Maternal Health, Neonatology and Perinatology*. doi: 10.1186/s40748-017-0057-x.
- Tomatala, S. et al. (2019) 'Hubungan Antara Aktivitas Fisik dengan Kekambuhan ISPA Pada Anak Usia Sekolah di Kecamatan Bringin Kabupaten Semarang', *Jurnal Keperawatan Respati Yogyakarta*, 6(1), p. 537. doi: 10.35842/jkry.v6i1.289.
- Virarisca, S., Dasuki, D. and Sofowan, S. (2019) 'Metode persalinan dan hubungannya dengan inisiasi menyusui dini di RSUP Dr. Sardjito Yogyakarta'. Available at: <https://journal.ugm.ac.id/jgki/article/view/177%0A44/11518>.
- World Health Organization (2008) 'Pencegahan dan Pengendalian ISPA', *Jurnal Kesmasindo*, 6, pp. 35-42. Available at: <http://www.who.int/csr/resources/publications/AMpandemicbahasa.pdf>.





### Nursing Research and Practice

Hindawi's Academic Journals Cover A Wide Range of D  
Submit With Us.

Hindawi

## EurAsian Journal of BioSciences

**COUNTRY**

Turkey

Universities and research institutions in Turkey

**SUBJECT AREA AND CATEGORY**

Agricultural and Biological Sciences  
Agricultural and Biological Sciences (miscellaneous)

Biochemistry, Genetics and Molecular Biology  
Biochemistry, Genetics and Molecular Biology (miscellaneous)

Environmental Science  
Environmental Science (miscellaneous)

**PUBLISHER**

Foundation for Environmental Protection and Research

**H-INDEX**

7

**PUBLICATION TYPE**

Journals

**ISSN**

13079867

**COVERAGE**

2013-2020

**INFORMATION**

Homepage  
How to publish in this journal  
ejobios@ejobios.org

## Best Minecraft Server Hosting

Start Your Own Minecraft Server With Instant Setup  
And 24/7 Live Chat.

Open

### SCOPE

EurAsian Journal of BioSciences (Abbrev. Eurasia J Biosci or EJOB IOS) is an international, refereed electronic journal. It publishes the results of original research in the field of biological sciences restricted to morphology, physiology, genetics, taxonomy, ecology and biogeography of both prokaryotic and eucaryotic organisms. The journal encourages submission of manuscripts dealing with plant biology, animal biology, plant physiology, microbiology, hydrobiology, ecology and environmental science, ethnobiology, biodiversity and conservation biology. EurAsian Journal of BioSciences publishes original articles in the following areas: -Agriculture, Fisheries & Food -Anatomy & Morphology -Behavioural Sciences -Biology, Biochemistry and Biotechnology -Biophysics -Biology Education -Cellular Biology and Anatomical Sciences -Ecology, Evolution & Environment -Entomology -Forestry -General Biology -Genetics & Heredity -Life Sciences - Other topics -Microbiology and Immunology -Molecular Biology -Mycology -Palaeontology -Parasitology -Pharmacology & Pharmacy -Physiology and Related Sciences -Plant Sciences -Toxicology -Veterinary Sciences -Virology -Zoology

 Join the conversation about this journal

# Best Minecraft Server Hosting

Start Your Own Minecraft Server With Instant Setup And 24/7 Live Ch

Shockbyte



Quartiles

## FIND SIMILAR JOURNALS

1  
**Bioscience Research**

PAK

**35%**  
similarity

2  
**Plant Archives**

IND

**34%**  
similarity

3  
**Pertanika Journal of Tropical  
Agricultural Science**

MYS

**33%**  
similarity

4  
**Saudi Journal of  
Sciences**

SAU

**32%**  
simila

SJR



Citations per document

