

Implementation of Hygiene and Environmental Sanitation in Under Five Years Old Diarrhea Patients at Surabaya Primary Health Center

by Anindita Rehana

Submission date: 30-Nov-2021 10:49AM (UTC+0800)

Submission ID: 1715891047

File name: ears_Old_Diarrhea_Patients_at_Surabaya_Primary_Health_Center.pdf (956.88K)

Word count: 7433

Character count: 39649

Research Articles

Implementation of Hygiene and Environmental Sanitation in Under Five Years Old Diarrhea Patients at Surabaya Primary Health Center Penerapan Higiene dan Sanitasi Lingkungan pada Pasien Diare Balita di Puskesmas Surabaya

Anindita Rehana¹, Rebekah Juniati Setiabudi^{2,4}, Sulistiawati^{3,4},
Manik Retno Wahyunitisari^{2,4}

¹ Faculty of Medicine, Airlangga University

² Department of Microbiology, Faculty of Medicine Airlangga University

³ Department of Public Health and Preventive Medicine, Faculty of Medicine
Airlangga University

⁴ Dr. Soetomo General Hospital Surabaya

Abstract

Surabaya has the most cases of diarrhea in East Java, with the highest detection targets in Gading Health Center (2346 cases) and Mojo Health Center, with diarrhea cases handled by 1322 cases. This study aimed to describe hygiene and environmental sanitation in under-five diarrhea patients in the working area of Surabaya Health Center. This study used a survey method, descriptive approach, interviews, and questionnaires in October 2019-March 2020. The results showed 62,2% of under-fives 'drinking water sources were ineligible, 86,5% of mothers' handwashing behavior were ineligible, 67,6% of the use of serving hoods were ineligible, 100% of food ripeness level were eligible, 59,5% of the distance between well and septic tank were eligible, 100% of latrines were eligible, 89,2% of sewerages were ineligible, and 83,8% of waste disposal facilities were ineligible. This research concluded that drinking water sources, handwashing behavior, serving hoods, sewerages, and waste disposal facilities were ineligible. The problems can be solved by consuming branded gallons of water / well water or refillable water that is boiled before consumption, improving how to wash hands with soap and running water, covering food using a serving hood, use the cover to close the sewerage, and throwing the trash into the temporary shelter (TPS) as much as two times a week or less than 3x24 hours.

Keywords: under five years old, diarrhea, facilities

Abstrak


Kota Surabaya memiliki kasus diare terbanyak di Jawa Timur dengan target penemuan terbanyak di Puskesmas Gading sebanyak 2346 kasus dan Puskesmas Mojo dengan kasus diare ditangani sebanyak 1322 kasus. Tujuan penelitian ini adalah mengetahui gambaran higiene dan sanitasi lingkungan pasien diare balita di wilayah kerja Puskesmas Surabaya. Penelitian ini menggunakan metode survei, pendekatan deskriptif, wawancara dan kuesioner pada bulan Oktober 2019-Maret 2020. Hasil penelitian menunjukkan 62,2% sumber air minum balita tidak memenuhi syarat, 86,5% perilaku cuci tangan ibu tidak memenuhi syarat, 67,6% penggunaan tudung saji tidak memenuhi syarat, 100% kematangan makanan memenuhi syarat, 59,5% jarak sumur dengan *septic tank* memenuhi syarat, 100% jamban memenuhi syarat, 89,2%, saluran pembuangan air limbah tidak memenuhi syarat, dan 83,8% sarana pembuangan sampah tidak memenuhi syarat. Simpulan penelitian ini adalah sumber air minum, perilaku cuci tangan, penggunaan tudung saji, saluran pembuangan air limbah, dan sarana pembuangan sampah masing-masing tidak memenuhi syarat. Permasalahan tersebut dapat diatasi dengan mengonsumsi air galon bermerk/air sumur atau air isi

22
ulang yang direbus sebelum dikonsumsi, memperbaiki cara mencuci tangan dengan sabun dan air mengalir, menutup makanan dengan tudung saji, memasang penutup untuk saluran pembuangan air limbah, dan membuang sampah ke TPS sebanyak 2 kali dalam seminggu atau kurang dari 3x24 jam.

Kata Kunci: balita, diare, fasilitas

*Korespondensi:

Anindita Rehana, email: anindita.rehana-2017@fk.unair.ac.id



This is an open-access article under the CC-BY license

INTRODUCTION

3
Diarrheal disease is an endemic disease in Indonesia and is a potential disease of outbreaks (KLB), often accompanied by death. In 2017, there were 21 outbreaks of diarrhea in Indonesia, spread across 12 provinces in 17 districts/cities. KLB (Kejadian Luar Biasa) is the emergence or increase in the incidence of illness or death, which is epidemiologically significant in an area within a certain period and is a condition that can lead to an outbreak (Kemenkes RI, 2018). In East Java, Surabaya ranks first in the most cases of diarrhea to be precise in the working area of Gading Health Center with a target number of finding 2346 cases and the working place of Mojo Health Center with 1322 cases of diarrhea handled (Dinas Kesehatan Kota Surabaya, 2019). Based on Riskekdas (2007), the age group of diarrhea with the highest prevalence was children under five (1-4 years), specifically 16.7%, therefore this study used under-fives as research subjects.

Diarrhea is described as the occurrence of watery stools at least three times or more a day (Nemeth and Pflieger, 2020). The most common cause of diarrhea in under-fives is Rotavirus infection. Low awareness of hygiene and implementation of environmental sanitation was also found to be risk factors for diarrhea outbreaks (Kemenkes RI, 2011). The number of people who do not have access to clean water in cities has increased sharply in developing countries due to urbanization, so they are looking for alternative water sources in the form of drilled wells and dug wells (Sari et al, 2018). The well is not free from *E. coli* contamination if the water source is less than 10 meters away from sewage disposal facilities, waste storage, and garbage collection or disposal sites (Kemenkes RI, 2018). *E. coli* is used as an indicator of water contamination by feces because the bacteria cannot survive long outside the intestine. *E. coli* itself is a normal flora in the large intestine. However, some strains of *E. coli* can produce toxins that can cause diarrhea (Thani et al, 2016).

The practice of promoting hygiene that is most effective as a form of overcoming diarrhea problems is washing hands properly because it can cut off the transmission of pathogens that cause diarrhea (Ndisika and Solomon, 2019). Processing food until it is ripe, washing fruits and vegetables before eating, washing tableware in a sterile manner, covering food with a serving hood have also been shown to reduce the incidence of diarrhea (Zyoud et al, 2019). The correct implementation of environmental sanitation in households is that they must have a gooseneck or embankment latrine with a lid, have a septic tank or wastewater drainage, and can be used alone or together (Kemenkes RI, 2018). A trash can with a closed condition is also essential because flies are said to be a vector of digestive tract diseases like damp and dirty places such as piles of household waste (Fitri et al, 2020).

Similar research has been conducted by Suda et al. (2019) and Magdalena et al.

(2019) regarding the factors associated with the incidence of children under five. This study contributed to the discussion because it contained variables that were also discussed in this study, including the relationship between mother's knowledge, drinking water, latrines, handwashing behavior, and the cleanliness of toddler food utensils. Research by Gloria et al. (2019) also contributed to the discussion of food storage which refers to the serving hood in this study. Research by Fitri et al. (2020) reveals the relationship between contamination of well water and the incidence of diarrhea in children under five related to this study, that the distance of the wells also affects the quality of the drinking water source for toddlers when it comes from well water. The relationship between sewerage and diarrhea is also discussed in research by Langit (2016).

Researchers are interested in conducting research on the implementation of hygiene and environmental sanitation in under-five diarrhea patients at the Surabaya Primary Health Center to provide information to the public about implementing hygiene and environmental sanitation.

METHODS

This research used a survey method with a descriptive approach which was carried out in October 2019-March 2020 in Gading Health Center and Mojo Health Center working areas. This study population was under five who visit Gading Health Center and Mojo Health Center in October 2019-March 2020 and obtained 37 under-fives who had met the inclusion criteria through the purposive sampling technique. Researchers set specific criteria in this study, so the results are more representative of the purpose and research subjects. Therefore, the researchers determine inclusion criteria in obtaining samples, so it can be said that this research uses the purposive sampling technique. The inclusion criteria in this study were (1) under-fives aged 12-59 months, (2) Suffering from diarrhea based on the diagnosis from the doctor, (3) Coming to Gading and Mojo Health Center, (4) Mother was willing to be research subjects for interviews and home surveys.

Data collection techniques were carried out by interview and field survey. The instrument used in this study was a questionnaire. The data source used is primary data obtained through interviews and questionnaires. The data were analyzed using the IBM SPSS Statistics 25 application. The variables in this study were hygiene and environmental sanitation. Hygiene researched included drinking water sources for under-fives, mothers' handwashing behavior, use of a serving hood, and the food ripeness level. Environmental sanitation variables studied included the distance between the septic tank and the well, latrines, sewerages, waste disposal facilities. Ethics has been obtained from the KEPK Faculty of Medicine, Universitas Airlangga no. 253/EC/KEPK/FKUA/2020.

RESULTS AND DISCUSSION

Characteristics of research subjects

The descriptive characteristics of the analyzed subjects are shown in Table 1 which shows that the mean age of mothers is $32,89 \pm 5,425$ years, with the minimum age 21 years and the maximum age 46 years and the mean age of under-five is $23,32 \pm 13,597$ months, with the minimum age 12 months and the maximum age 54 months.

Table 1 shows the mean age of the mother age. Research by Haswari et al. (2019) said that there is a relationship between a mother's education and the incidence of diarrhea with a p-value of 0,000 (p-value < 0,05). Adult age will affect a more systematic frame of thinking. This level of maturity will affect their parenting, caring for their children, so that the more mature the mother's age, the more experienced she will be in caring, including in implementing hygiene and environmental sanitation, providing food intake, and maintaining children's health (Haswari et al, 2019).

Table 1. Characteristics of research subjects

	Minimal	Maximum	Mean	Standard Deviation
Mother's Age (years)	21	46	32,89	5,425
Toddler Age (months)	12	54	23,32	13,597
		Frequency		Presentation (%)
Gender of Toddler				
Male		19		51,4
Female		18		48,6
Mother's highest attainment	Educational			
Elementary School		6		16,2
Junior High School		14		37,8
High School		13		35,1
Bachelor		3		8,1
Diploma		1		2,7
Mother's work				
Housewives		33		89,2
Private Sector Worker		4		10,8

Source: Primary Data, 2019

Table 1 shows that most of the mother's education level is junior high school as many as 14 (37.8%) research subjects. Most occupations of mothers of children under-five are housewives, 33 (89.2%) research subjects. Research by Sumampouw et al. (2019) and Santika et al. (2020) said that the level of education has a significant effect on diarrhea cases in children under five. The level of education is a major determining factor for the survival of children, especially those under five who cannot carry out activities independently (Omona et al, 2020). They depend on their mother or caregiver for activities such as eating, drinking, defecating, and urinating, which will have a major impact on their health if negligent.

People with low education pay less attention to or ignore environmental hygiene and sanitation behavior towards themselves and their families (Ugboko et al, 2020). People who have a higher education level prefer to work and have a greater chance of getting health information from their work environment than mothers who do not work or housewives (Ainsyah and Lusno, 2018). Therefore, it is necessary to consider providing counseling on parenting, the application of good hygiene, and environmental sanitation, especially for mothers with low education and housewives to be better prepared to educate and care for children properly (Santika et al, 2020).

Under-fives drinking water source

Table 2 shows the analysis of water sources for children under-five hygiene application. The results showed that only 14 (37,8%) under-fives consumed water that met the hygiene requirements by consuming branded gallon bottled water or drinking PDAM / well water that was boiled and 23 (62,2%) under-fives consumed water that was not meet the hygiene requirements by not boiling refill water before consumption.

Table 2. Characteristics of under-fives drinking water source

Under-fives Drinking Water Source	Frequency	Presentation (%)
Eligible	14	37,8
Ineligible	23	62,2
Total	37	100

Source: Primary Data, 2019

Research by Asnel and Sari (2019) said that there is a significant relationship between the drinking water source and the incidence of diarrhea with the p-value 0,003 (p-value < 0,05) and POR 4,054 (95% CI = 1,678-9,798). It means that toddlers with ineligible drinking water sources are at risk 4 times to had diarrhea than those who had eligible drinking water sources (Asnel and Sari 2019). Research by Suda et al. (2019) said that there is a significant relationship between drinking water sources for under-fives and acute diarrhea in under-fives. Safe (proper) drinking water for health is drinking water that meets physical, microbiological, chemical, and radioactive requirements (Kemenkes RI, 2018).

This study has limitations that microbiological, chemical, and radioactive examinations as required for safe (proper) drinking water for health stipulated by Kementerian Kesehatan RI were not carried out. Still, the researchers used the standards of previous studies which stated that the water was boiled first until it boils before consumption is an effective way to reduce the risk of diarrhea that is easily done by every individual (Dinkel et al, 2020). Boiling water is mainly carried out to refill drinking water sources prone to contamination. This statement is supported by Wahyuningsih and Kamaningroem (2019) research, which said that the total coliform test results show that 21 out of 25 refill drinking water facilities in Gubeng District Surabaya City, do not meet the standard threshold.

Research by Dewanti and Sulistryorini (2017) regarding the bacteriological quality analysis of refill drinking water in Sememi Village, Benowa District, Surabaya City shows all research subjects or owners and handlers of Refill Drinking Water Depot (DAMIU) do not wash their hands with running water and soap before filling consumer gallons of water. The results of the examination of the raw water samples at DAMIU also showed the growth of *E. coli*, but after processing, there were 2 out of 6 refill water samples that still contained *E. coli*. As for Bottled Drinking Water (AMDK), research by Monikayani et al. (2020) said that bottled drinking water is safer from the side of the Most Probable Number (MPN) test.

The technique of washing milk bottles also affects the sterilization of water that under-fives drink. Research by Lanida and Farapti (2018) said that there is a relationship between washing milk bottles and the incidence of under-five diarrhea. The risk of diarrhea in children who consume breast milk by the bottle is 1.95 times, the risk of diarrheain children who consume formula milk using bottles is 2.12 times, while the risk of diarrhea in children who consume exclusive breastfeeding, supplementary food other than breast milk, breastfeeding using a milk bottle, and bottle feeding formula is 1.77 times. This shows that milk bottles are very susceptible to contamination by bacteria, therefore it is necessary to wash bottles properly (Masood, 2017).

Techniques for cleaning and storing feeding bottles in infants have been proposed by the CDC. Before washing milk bottles, mothers are required to wash their hands with soap and running water for 20 seconds. Then start rinsing the parts of the bottle under running water. Parts of baby milk bottles should not be placed in the sink during the

washing process because they have the potential to contain germs that can stick to any part of the feeding bottle. After rinsing, the bottle is placed in a basin filled with hot water and soap. Then do the rubbing of the bottles and other parts using a clean brush. The rinse is carried out again with running water and placed in a separate basin dedicated for baby food items in clean and dust-free conditions and then allowed to dry. For a better bottle sterilization process, it is recommended to boil for 5 minutes. Washing with soap and running water only physically removes germs from the bottle, whereas the boiling process is an extra step to kill more germs and extra protection from all infections (CDC, 2020). If not cleaned properly, the fat and protein content in milk will stick to the bottle and form white spots which, if allowed to accumulate for a long time and dry out, will be difficult to clean and become a breeding ground for bacteria. If bacteria enter the digestion, it will increase the risk of diarrhea (Kosapilawan et al, 2019).

Mothers’ hand washing behavior

Table 3 shows the analysis of hygiene application of mothers' hand washing behavior. The results showed that only 5 (13.5%) research subjects met the hygiene requirements by washing their hands properly with the criteria of washing hands before eating, after defecating and urination, after changing under-five diapers, washing hands with running water, and always using soap.

Table 3. Characteristics of mothers’ hand washing behavior

	Frequency	Presentation (%)
Mothers’ Hand Washing Behavior		
Eligible	5	13,5
Ineligible	32	86,5
Total	37	100

Source: Primary Data, 2019

Alamsyah and Marianthi (2020) said that there is a relationship between the behavior of hand washing with soap with the incidence of diarrhea, with the p-value 0,03 (p-value < 0,05) and OR 3,80 (95% ci 1,25-11,50). It means that the mothers who don’t wash their hands properly have a risk of a 3,80 times greater occurrence of diarrhea in their children than mothers who behave well (Alamsyah and Marianthi 2020). Research by Radhika (2020) said a significant relationship between mothers’ hand washing behavior and cases of diarrhea in under-fives.

Research by Fuhrmeister et al. (2020) said that there are enteric bacteria, an average of 1.7% on the palms. Enteric pathogens are transmitted fecal-orally via various reservoirs, including flies, fomites, hands, soil, food, and water. Therefore, one effective way to block pathogen transmission is hand washing. The practice of washing hands properly has requirements, namely washing hands with soap and running water, carried out at least 30 seconds, and done at least before eating, after changing diapers, and after defecating (Ndisika and Solomon, 2019). Based on the interview results, respondents did not always wash their hands with running water and even used finger bowl and rarely used soap.

Research by Ndisika and Solomon (2019) said that the practice of washing hands with soap could reduce the threat of severe diarrhea by up to 48% and the likelihood of diarrhea by 47%. Even though awareness, knowledge, attitude, and practice have been carried out, the method of washing hands is not correct, the desired results will not be

achieved. This can occur because the information provided is inadequate or the mother of the under-five does not understand the information. Therefore there is a need for counseling on the importance of getting used to washing hands properly.

Use of serving hood

Table 4 shows an analysis of hygiene applications in using a serving hood. The results showed that only 12 (32,4%) research subjects had met the correct hygiene requirements by always using a serving hood or another food cover.

Table 4. Characteristics of serving hood use

	Frequency	Presentation (%)
Use of Serving Hood		
Eligible	12	32,4
Ineligible	25	67,6
Total	37	100

Source: Primary Data, 2019

Research by Gloria et al. (2019) said that there is a significant relationship between the use of a serving hood and cases of diarrhea in under-fives. Research by Yanti et al. (2018) said that there is a significant relationship between poor food management and storage and incidence of diarrhea with the p-value 0,029 (p-value < 0,05) and OR = 4,278, it means that poor food management and storage is four times more likely to be at risk of diarrhea than good food management and storage.

Food is one of the most important routes for children to be exposed to pathogens that come from feces. Exposure to pathogens from feces from both humans and animals can lead to diarrheal disease, or even continuous exposure will lead to infection without symptoms (Freeman et al, 2020). Research by Zyoud et al. (2019) said that 50% of cases of food poisoning are related to improper storage of food or reheating of food. The diarrhealdisease can occur by consuming food or water that is contaminated by infective organisms (bacteria and viruses), toxic chemicals, radioactive substances, and other harmful substances. The process of cooking, serving, and storing food greatly affects contamination by toxic chemicals or pathogens that cause disease. A good way to store food before consumption is to use a food cover or serving hood (Zyoud et al, 2019). However, the use of a serving hood presents socio-cultural challenges, including the belief about food spoilage when covered with a serving hood (Freeman et al, 2020).

The purpose of using a serving hood as a food cover is that flying animals do not catch food, especially flies, which can transmit pathogens. Cockroaches, lizards, and mice may also approach eating a small portion of the food served and transmit pathogens as well (Zyoud et al, 2019). Hygienic food storage is defined as food inaccessible to animals, food inaccessible to young children, covered food, and fly-free food (Freeman et al, 2020). An alternative if you don't have a serving hood is to store food in a special food cabinet and always keep the cupboard and create a clean environment around the food. In addition, people can also use wire netting on windows, use curtains, or put fly-resistant adhesive paper around the food to avoid flies or other flying animals (Yanti et al, 2018).

Food ripeness level

Table 5 shows the analysis of hygiene applications of food ripeness level that are usually carried out by the research subjects. The results showed all research subjects had met the correct hygiene requirements, namely cooking the food until it is fully cooked.

Table 5. Characteristics of food ripeness level

	Frequency	Presentation (%)
Food Ripeness Level		
Eligible	37	100
Total	37	100

Source: Primary Data, 2019

Cooked food has an important role in reducing the level of bacteria in food and beverages (Fitri et al, 2020). Cooking, serving, and storing food greatly affects contamination by toxic chemicals or pathogens that cause disease. Symptoms of toxic food poisoning mostly appear within 24 hours of consuming contaminated food, while foodborne infections may not appear until 2-3 days later. Symptoms include nausea, vomiting, diarrhea, abdominal pain, headache, and fever (Zyoud et al, 2019).

Research by Fiona (2017) said that raw meat could be contaminated with certain pathogenic bacteria spores that are not easily destroyed. Therefore cooking food until it is cooked is highly recommended to destroy pathogenic bacteria in raw meat. Spores develop into vegetative cells that can reproduce rapidly in food at room temperature. Therefore, processing food until it is ripe, washing fruits and vegetables before eating, washing tableware in a sterile manner, covering food with a serving hood are believed to reduce the incidence of diarrhea due to *E. coli* infection (Zyoud et al, 2019).

CDC has appealed to prevent food poisoning with four principles, namely Clean-Separate-Cook-Chill. Cleaning is an appeal to wash hands and cooking utensils, including cutting boards and kitchen tables before, during, and after preparing food, because microorganisms can survive around the kitchen, including hands, cooking utensils, cutting boards, and tables. Fresh fruits and vegetables are also encouraged to be rinsed under running water before consumption. Then Separate is an appeal to separate raw meat, poultry, seafood, and eggs from ready-to-eat food. The use of cutting boards is also advised to separate raw materials and ready-to-eat food ingredients. The next principle is Cook, the appeal to cooking food at the right temperature to kill harmful microorganisms. Then the Chill principle is an appeal to freeze food properly. The bacteria that cause food poisoning multiply at the fastest between 4°C-60°C, so make sure the refrigerator temperature is 4°C or lower (CDC, 2020).

Research by Wainaina et al. (2020) said that norovirus is found among food handlers in densely populated, low-income, and urban areas. As many as 87% of the study participants answered, having a dirty cutting board can cause food contamination. One of the most common risk factors for norovirus infection in food makers is a lack of hand washing behavior. Increasing hand washing compliance with soap and running water can reduce and eliminate viral contamination.

Research by Sutoko et al. (2019) shows that the results of colony calculations on a cutting board have the highest average value of 7.92 CFU/cm² while glass has the smallest average value, namely 0.01 CFU/cm². The liquid residue from raw meat or poultry on the cutting board's surface has the potential to transmit pathogens to other foods that use the same cutting board. When the cutting board is contaminated, pathogens can survive and multiply. Foodborne pathogens such as *Salmonella* spp., *Campylobacter* spp., *Bacillus cereus*, *Staphylococcus aureus*, and *Listeria monocytogenes* are found in household kitchen appliances (Deza et al, 2007).

The surface of the cutting board has scratches that cause gaps due to the pressure from the knife. The gaps make it difficult to clean the remaining food ingredients. Based on previous research, *S. aureus* and *E. coli* were able to survive on polyethylene material for a long time. Therefore, the recommended use of a cutting board is a cutting board made from closed-grained wood. Woodcutting boards are known to have porous materials that can absorb moisture quickly, so bacteria cannot grow on these surfaces (Sutoko et al, 2019). Other studies have also suggested that some woods may contain antibacterial properties, and the hygroscopic characteristics of these materials can cause bacterial drying (Deza et al, 2007).

Well distance to septic tank

The results showed that the majority of research subjects had met the environmental sanitation requirements by not having a well with a distance of <10 m from the septic tank, namely 25 (59,5%) research subjects.

Table 6. Characteristics of good distance to septic tank

	Frequency	Presentation (%)
Well Distance to Septic Tank		
Eligible	22	59,5
Ineligible	15	40,5
Total	37	100

Source: Primary Data, 2019

Research by Fitri et al. (2020) and Bangun et al. (2020) said there is a relationship between contaminated well water sources and diarrhea in under-fives. Research by Dangiran and Dharmawan (2020), which analyzed the relationship between the incidence of diarrhea and the presence of dug wells in Jabung Village, Semarang City, proved that the distribution of diarrhea was found in houses that have dug wells with bacteriological water quality that do not meet the requirements, namely >50 CFU/100 ml and distance from latrines and/or septic tanks <11 meters (Dangiran and Dharmawan, 2020).

In China, an epidemic of infectious diarrheal disease has occurred in schools due to consuming contaminated well water. The location of the well was found near a toilet or septic tank, a damaged sanitary sewer, and wastewater that spilled into the well. Broken toilets or septic tanks and sewers near water wells, improper water well construction, and a lack of sterilization facilities are the main causes of water contamination by *E. coli* (Ding et al, 2017).

The septic tank must be ≥ 10 meters from the water source to avoid contamination of *E. coli* in the feces (Kemenkes RI, 2018). Soil contamination causes contamination of clean water sources to become a good growth medium for *E. coli* and increases soil concentration (Fitri et al. 2020). In areas with high population density, the ideal 10-meter distance between water sources and septic tanks seems difficult to implement due to lack of land. Another alternative can be done to overcome limited land. One of them is by knowing the direction of groundwater flow so that the direction of the flow does not lead to a well or water source, with the septic tank being free from contamination (Sidik et al. 2020). If groundwater or well water is consumed without boiling it first or not boiling, it can cause diarrhea (Fitri et al, 2020).

Latrine

The results showed all research subjects had met the environmental sanitation requirements by having latrines with the criteria that feces did not pollute the soil and

surface water, the latrines were closed, and free from unpleasant odors.

Table 7. Characteristics of latrine

	Frequency	Presentation (%)
Latrine		
Eligible	37	100
Total	37	100

Source: Primary Data, 2019

Research by Magdalena et al. (2019) said that there is a significant relationship between the type of latrine and diarrhea in children under five. The toddler who uses latrines that do not meet requirements had a two times greater chance of having diarrhea based on research by Sidabalok et al. (2019). Attitudes will determine a person in choosing latrine, as evidenced by Theresiana et al. (2020) with a p-value of 0,000 (p-value <0,05) and OR = 23, which means that there is a significant relationship between attitude and choice of latrine by the respondent. Respondents with positive attitudes were 23 times more likely to choose healthy latrines than those with negative attitudes. Therefore, the community needs information about the benefits, disadvantages, and consequences obtained if the latrine used does not meet the requirements (Theresiana et al, 2020).

Under-fives can experience diarrhea even though they have a healthy latrine because the water used does not meet the requirements, namely well water that has not been tested for healthy water requirements. In addition, under-fives and mothers do not practice hygiene after defecating by washing their hands with running water and soap (Magdalena et al., 2019). A suitable method of disposal of feces is that feces should not pollute the soil and surface water because this results in soil contamination and clean water sources, becoming a good growth medium for *E. coli* (Fitri et al, 2020). The latrine must also be closed, free from odors or unsightly conditions, so as not to invite flies and other animals to transmit the pathogens that cause diarrhea. Feces should not be touched with hands directly, and when cleaning, it is also mandatory to wash hands with soap and run water (Kemenkes RI, 2018).

Sewerage

The results showed that most research subjects 33 (89,2%) did not meet the environmental sanitation requirements by having open sewerage.

Table 8. Characteristics of sewerage

	Frequency	Presentation (%)
Sewerage		
Eligible	4	10,8
Ineligible	33	89,2
Total	37	100

Source: Primary Data, 2019

Research by Langit (2016) and Ikhwan (2013) said that there is a significant relationship between sewerage and diarrhea in children under five. Research by Mirsiyanto et al. (2020) said that there is a significant relationship between sewerage and the incidence of chronic diarrhea in children under five with OR = 3,202, which means that respondents with a bad wastewater treatment system are 3.202 times more likely to

suffer from diarrhea than respondents with a good wastewater treatment system.

According to the Badan Pelatihan Kesehatan Kementerian Kesehatan (2011), sewerage is a facility in the form of excavated soil or pipes made of cement or pipe which function to dispose of washing water, used bathing water, dirty/other used water to the disposal site. The wastewater generally contains materials or substances harmful to health and cause an unpleasant smell that can invite disease vectors. Good sewerage is sewerage that does not pollute wells or other water sources; not a breeding ground for mosquitoes, flies and centipedes; closed tightly using a board or other non-permanent cover for easy cleaning; cannot cause accidents, especially in children; and does not interfere with the aesthetics (Kemenkes RI, 2011).

Population density has a consequence of decreasing environmental quality that can affect health, for example, the potential for water to overflow when it rains. During the interview, the researcher asked whether there was a frequent overflow of wastewater into the road and the research subjects answered the overflow occurred when it rained. Shallow sewerage that is rarely cleaned can trigger this. The environment that becomes dirty has become a preferred place for flies and mice, which are vectors of various diseases such as diarrhea (Fitri et al, 2020).

Waste Disposal Facilities

The results showed that most research subjects, as many as 31 (83.8%), did not meet the environmental sanitation requirements by having an open trash can and throwing the trash into the temporary shelter (TPS) less than two times a week or more than 3x24 hours.

Table 9. Characteristics of waste disposal facilities

Waste Disposal Facilities	Frequency	Presentation (%)
Eligible	6	16,2
Ineligible	31	83,8
Total	37	100

Source: Primary Data, 2019

Research by Mustika (2019) and Yuniar et al. (2020) said a significant relationship between trash cans and under-five diarrhea is said. Disposal of household waste, especially trash cans that are open and allowed to accumulate for over three days, can cause an odor that invites disease-transmitting vectors. Garbage must be disposed of in a temporary shelter (TPS) at least two times a week or a maximum of every 3x24 hours.

Research by Jamaluddin and Zarnila (2020) said that there is a significant relationship between waste management and the incidence of diarrhea with a p-value = 0,000 (p-value < 0,05). The act of separating and storing waste between organic and non-organic waste can reduce the risk of diarrhea because the separation and storage of organic and organic waste and properly covered will reduce the frequency of flies (Jamaluddin and Zarnila, 2020).

Garbage is closely related to public health because garbage can spread disease through vectors (Yuniar et al, 2020). Uncovered (open) trash bins invite disease-transmitting vectors into the house and reach humans. The vector can transmit diseases, such as flies perching on the trash, and can cause diarrhea. Garbage also it needs to be separated between (organic) food scraps and inorganic waste because food scraps attract lots of flies and have the potential to breed in the trash, which then flies off and land on food or drinks so it can cause diarrhea (Fitri et al. 2020). For this reason, it is necessary to

increase public knowledge about procedures for the proper disposal of household wastewater.

CONCLUSION

It can be concluded that the drinking water source for children under five, the behavior of washing hands of mothers, the use of serving hoods, sewerages, and waste disposal facilities do not meet the requirements, only food ripeness level, the distance between the well and the septic tank, and latrines that have been qualify. The problems can be solved by consuming branded gallons of water / well water or refillable water that is boiled before consumption, improving how to wash hands with soap and running water, covering food using a serving hood, use the cover to close the sewerage, and throwing the trash into the temporary shelter (TPS) as much as two times a week or less than 3x24 hours.

ACKNOWLEDGMENTS

We would like to express our gratitude to the staff of Gading Health Center, Mojo Health Center, and the staff of the Surabaya City Health Office who assisted this research.

CONFLICT OF INTEREST STATEMENT

The authors declare that there is no conflict of interest.

REFERENCES

- Ainsyah RW, Lusno MFD. 2018. The Protective Factor of Diarrhea Incidence in Toddler in Surabaya. *Jurnal Berkala Epidemiologi*. 6(1):51.
- Alamsyah T, Marianthi D. 2020. Correlation between Incidence of Diarrhea in Toddlers and Housewife's Clean and Healthy Lifestyle. *J-Kesmas: Jurnal Fakultas Kesehatan Masyarakat (The Indonesian Journal of Public Health)*. 7(1):57.
- Asnel R, Sari LA. 2019. Correlation Family's Healthy Lifting Behavior with Diarrhea in Toddlers. *KnE Life Sciences*. 4(10):197.
- Bangun HA, Nababan D, Hestina. 2020. Hubungan Sanitasi Dasar Dengan Kejadian Diare Pada Balita Di Desa Durian Kecamatan Pantai Labu Kabupaten Deli Serdang. *Jurnal Tekesnos*. 2(1):57-66.
- CDC [Center for Disease Control and Prevention]. 2020a. Food Safety. Centers for Disease Control and Prevention. USA: CDC.
- CDC [Center for Disease Control and Prevention]. 2020b. How to Clean, Sanitize, and Store Infant Feeding Items. Centers for Disease Control and Prevention. USA: CDC.
- Dangiran HL, Dharmawan Y. 2020. Analisis Spasial Kejadian Diare Dengan Keberadaan Sumur Gali Di Kelurahan Jabungan Kota Semarang. *Jurnal Kesehatan Lingkungan Indonesia*. 19(1):68-75.
- Dewanti RA, Sulistyorini L. 2017. Analisis Kualitas Bakteriologis Air Minum Isi Ulang Di Kelurahan Sememi, Kecamatan Benowo. *The Indonesian Journal of Public Health*. 12(1):39.
- Deza MA, Araujo M, Garrido MJ. 2007. Efficacy of Neutral Electrolyzed Water to Inactivate *Escherichia Coli*, *Listeria Monocytogenes*, *Pseudomonas Aeruginosa*, and *Staphylococcus Aureus* on Plastic and Wooden Kitchen Cutting Boards. *Journal of*

- Food Protection. 70(5):1070.
- Dinas Kesehatan Kota Surabaya. 2019. Profil Kesehatan 2018. Surabaya: Dinas Kesehatan Kota Surabaya.
- Ding Z, Zhai Y, Wu C, Wu H, Lu Q, Lin J, He F. 2017. Infectious Diarrheal Disease Caused by Contaminated Well Water in Chinese Schools: A Systematic Review and Meta Analysis. *Journal of Epidemiology*. 27:274–28.
- Dinkel KA, Costa M, Kraft T, Stieglitz J, Cummings D, Gurven M, Kaplan H, Trumble B. 2020. Relationship of Sanitation, Water Boiling, and Mosquito Nets to Health Biomarkers in a Rural Subsistence Population. *American Journal of Human Biology*.
- Fiona. 2017. Bacteria in Raw Meat vs Cooked Meat. Hong Kong: Centre for Food Safety.
- Fitri IS, Kusnopranto H, Soesilo TEB. 2020. The Source of Potential Pollution and Diarrhea on Toddlers at Populous Area (a Study at Johar Baru Subdistrict, Central Jakarta). *E3S Web of Conferences*. 153:1–13.
- Freeman MC, Ellis AS, Ogutu EA, Caruso BA, Linabarger M, Micek K, Muga R, Girard AW, Wodnik BK, Arriola KJ. 2020. Impact of a Demand• Side Integrated WASH and Nutrition Community-Based Care Group Intervention on Behavioural Change : A Randomised Controlled Trial in Western Kenya. *BMJ Global Health*: 5:e002806.
- Fuhrmeister ER, Ercumen A, Grembi JA, Islam M, Pickering AJ, Nelson KL. 2020. Shared Bacterial Communities between Soil, Stored Drinking Water, and Hands in Rural Bangladeshi Households. *Water Research X*. 9:100056.
- Gloria CV, Maharani R, Hayana, Dewi S. 2019. Determinants of Diarrhea Events on under Five Children in Harapan Raya Public Health Center Working Area. *Avicenna*. 14(3):52–110.
- Haswari GT, Wijayanti Y, Laksono B. 2019. Analysis Factors of Diarrhea Incidentin Toddlers At Purwodadi District Health Centre, Grobogan. *Public Health Perspective Journal*. 4(3):232–39.
- Ikhwan Z. 2013. Faktor Individu Dan Keadaan Saluran Pembuangan Air Limbah (SPAL) Rumah Tangga Dengan Kejadian Diare Di RT 01 RW 09 Kelurahan Sei Jang Kecamatan Bukit Bestari Kota Tanjungpinang. *Jurnal Kesehatan*. 4(2):416–419.
- Jamaluddin, Zarnila E. 2020. Hubungan Pengelolaan Sampah Rumah Tangga dan Kualitas Air Bersih dengan Kejadian Diare pada Balita di Desa Suka Mulia dan Desa Alur Manis Kecamatan Rantau Kabupaten Aceh Tamiang. *Jurnal Edukes*. 3(1):40–45.
- Kemenkes RI. 2011. *Jurnal Buletin Jendela Data & Informasi Kesehatan Situasi Diare Di Indonesia*. Jakarta: Kementerian Kesehatan RI.
- Kemenkes RI. 2018. 1227 Profil Kesehatan Indonesia 2018. Jakarta: Kementerian Kesehatan RI.
- Kosapilawan MM, Gunawan DCD, Nofiantika F. 2019. Hubungan Praktik Penggunaan Botol Susu dengan Kejadian Diare pada Balita di Wilayah Kerja Puskesmas Umbulharjo 1 , Kota Yogyakarta. *Ilmu Gizi Indonesia*. 03(01):45–52.
- Langit SL. 2016. Hubungan Kondisi Sanitasi Dasar Rumah dengan Kejadian Diare pada Balita di Wilayah Kerja Puskesmas Rembang 2. *Jurnal Kesehatan Masyarakat*. 4(2):160–65.
- Lanida BP, Farapti F. 2018. Prevention of the Incidence of Diarrhea in Infants through Hygiene of Milk Bottles. *Jurnal Berkala Epidemiologi*. 6(3):244.
- Magdalena I, Rantetampang AL, Pongtiku A, Mallongi A. 2019. The Risk Factors Environment and Behavior Influence Diarrhea Incidence to Child in Abepura Hospital Jayapura City. *International Journal of Science and Healthcare Research*. 4(March):171–80.
- Masood MSA. 2017. The Relationship between Diarrhea and Feeding Indicators among

- Children in the Capital Sana'a Yemen. *Nutrition and Food Toxicology*. 1(3):98–105.
- Mirsiyanto E, Sitorus RJ, and Misnaniarti. 2020. Analysis of Environmental Factors with Chronic Diarrhea in Toddlers in Jambi City in 2019. *International Journal of Science and Society*. 2(4):300–310.
- Monikayani R, Khatimah H, Muthmainah N, Rahmiati, Oktaviyanti IK. 2020. Gambaran Most Probable Number Air Galon Bermerek Dan Isi Ulang Di Banjarmasin. *Homeostasis*. 3(1):105–10.
- Mustika DA. 2019. Hubungan Pemberian MP-ASI dan Sanitasi Lingkungan dengan Kejadian Diare pada Anak Usia 6-24 Bulan di Desa Kedungkendo Kecamatan Candi Kabupaten Sidoarjo. [Skripsi]. Surabaya: Universitas Airlangga.
- Ndisika AOO, Solomon T. 2019. Knowledge, Attitude and Practice of Hand Washing among Mothers of Children 0-59 Months of Age in Lagos Nigeria. *Universal Journal of Public Health*. 7(2):52–58.
- Nemeth V, Pfliegerhaer N. 2020. Diarrhea. USA: StatPearls.
- Omona S, Malinga G, Opoke R, Openy G, Opiro R. 2020. Prevalence of Diarrhoea and Associated Risk Factors among Children under Five Years Old in Pader District, Northern Uganda. *BMC Infectious Diseases*. 20(1):1–9.
- Radhika A. 2020. Hubungan Tindakan Cuci Tangan Pakai Sabun Dengan Kejadian Diare Pada Balita Di Rw Xi Kelurahan Sidotopo, Kecamatan Semampir, Kota Surabaya. *Medical Technology and Public Health Journal*. 4(1):16–24.
- Santika NKA, Efendi F, Rachmawati PD, Has EMM, Kusnanto K, Astutik E. 2020. Determinants of Diarrhea among Children under Two Years Old in Indonesia. *Children and Youth Services Review*. 111(2020):104838.
- Sari SYI, Sunjaya DK, Furusawa HS, Watanabe C, Raksanagara AS. 2018. Water Sources Quality in Urban Slum Settlement along the Contaminated River Basin in Indonesia: Application of Quantitative Microbial Risk Assessment. *Journal of Environmental and Public Health*. 2018.
- Sidabalok DL, Samsudin, Djaja IM. 2019. Relationship between Environmental Factors and Personal Hygiene with Diarrhea among Children under Five in West Kotawaringin, Central Kalimantan. 6th International Conference on Public Health 2019.:31–39.
- Sidik M, Wijayanti SPM, Iqbal A. 2020. The Determinants of Diarrhea Disease Incidence in Densely Populated Area of West Nusa Tenggara, Indonesia. *Jurnal Kesehatan Lingkungan*. 12(2):107.
- Suda ED, Nabuasa E, Hinga IAT. 2019. Faktor-Faktor Yang Berhubungan Dengan Kejadian Diare Pada Balita Di Desa Buru Kaghu Kecamatan Wewewa Selatan Kabupaten Sumba Barat Daya. *Lontar : Journal of Community Health*. 1(4):119–26.
- Sumampouw OJ, Nelwan JE, Rumayar AA. 2019. Socioeconomic Factors Associated with Diarrhea among Under-Five Children in Manado Coastal Area, Indonesia. *Journal of Global Infectious Diseases*. 11(4):140–46.
- Sutoko A, Hapsari R, Hadi P. 2019. Kualitas Bakteriologi Peralatan Masak Dan Makan Di Rumah Sakit Nasional Diponegoro. *Jurnal Kedokteran Diponegoro*. 8(4):1327–37.
- Thani TS, Symekher SML, Boga H, Oundo J. 2016. Isolation and Characterization of Escherichia Colipathotypes and Factors Associated with Well and Boreholes Water Contamination in Mombasa County. *Pan African Medical Journal*. 23:1–11.
- Theresiana Y, Triyana L, Clara S. 2020. Determinants of Healthy Latrines Ownership in Working Area at Public Health Center of Suak Tapeh in Banyuasin Regency

- South Sumatra 2019. *Health Science Research*. 25:268–276.
- Ugboko HU, Nwinyi OC, Oranusi SU, Oyewale JO. 2020. Childhood Diarrhoeal Diseases in Developing Countries. *Heliyon*. 6(4):e03690.
- Wahyuningsih I, Karnaningroem N. 2019. Minimizing Failure of Production Quality from Refill Drinking Water in Gubeng District, Surabaya City, Using Failure Mode and Effect Analysis. *IOP Conference Series: Earth and Environmental Science*. 259(1).
- Wainaina E Otieno CA, Kamau J, Nyachieo A, Lowther SA. 2020. Norovirus Infections and Knowledge, Attitudes and Practices in Food Safety among Food Handlers in an Informal Urban Settlement, Kenya 2017. *BMC Public Health*. 20(1):1–8.
- Yanti CA, Ediana D, Rizki M. 2018. Hubungan Perilaku Dan Tingkat Kepadatan Lalat Dengan Kejadian Diare Di Pasar Sarilamak. *Human Care Journal*. 3(1).
- Yuniar N, Majid R, Karimuna SR. 2020. Analysis of Factors Associated with the Incidence of Diarrhea in Toddlers in the Working Area of Puuwatu Health Center, Kendari in 2019. *Indian Journal of Public Health Research & Development*. 11(05):733–38.
- Zyoud S, Shalabi J, Imran K, Ayaseh L, Radwany N, Salameh R, Sa'dalden Z, Sharif L, Sweileh W, Awang R, Al-Jabi S. 2019. Knowledge, Attitude and Practices among Parents Regarding Food Poisoning: A Cross-Sectional Study from Palestine. *BMC Public Health*. 19(1):1–10.

Implementation of Hygiene and Environmental Sanitation in Under Five Years Old Diarrhea Patients at Surabaya Primary Health Center

ORIGINALITY REPORT

13%

SIMILARITY INDEX

11%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	jurnal.utu.ac.id Internet Source	1%
2	www.cldcollege.com Internet Source	1%
3	Tuhu Pindari, Suparji Suparji. "Handwashing Behavior Using Soap, Physical Conditions of Cooking Food Storage and Deare Incidence", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	1%
4	bmcpublikealth.biomedcentral.com Internet Source	1%
5	Ika Syiami Fitri, Haryoto Kusnoputranto, Tri Edhi Budhi Soesilo. "The source of potential pollution and diarrhea on toodlers at populous area (a study at Johar Baru Subdistrict, Central Jakarta)", E3S Web of Conferences, 2020 Publication	1%

6	Nurhaedah Nurhaedah. "HUBUNGAN ANTARA SANITASI LINGKUNGAN DENGAN KEJADIAN DIARE PADA LANJUT USIA", JURNAL ILMIAH KESEHATAN SANDI HUSADA, 2019 Publication	<1 %
7	eprints.ums.ac.id Internet Source	<1 %
8	www.cdc.gov Internet Source	<1 %
9	www.grafiati.com Internet Source	<1 %
10	escholarship.org Internet Source	<1 %
11	assets.researchsquare.com Internet Source	<1 %
12	www.hrpub.org Internet Source	<1 %
13	www.neliti.com Internet Source	<1 %
14	repository.uinjkt.ac.id Internet Source	<1 %
15	M. A. DEZA, M. ARAUJO, M. J. GARRIDO. "Efficacy of Neutral Electrolyzed Water To Inactivate Escherichia coli, Listeria monocytogenes, Pseudomonas aeruginosa,	<1 %

and Staphylococcus aureus on Plastic and Wooden Kitchen Cutting Boards", Journal of Food Protection, 2007

Publication

16

core.ac.uk

Internet Source

<1 %

17

ejurnal.poltekkes-tjk.ac.id

Internet Source

<1 %

18

www.myfoodresearch.com

Internet Source

<1 %

19

mafiadoc.com

Internet Source

<1 %

20

Opara, Peace, Balafama Alex-Hart, and Tamunoiyowuna Okari. "Hand-washing practices amongst mothers of under-5 children in Port Harcourt, Nigeria", Paediatrics and International Child Health, 2016.

Publication

<1 %

21

Matthew C Freeman, Anna S Ellis, Emily Awino Ogutu, Bethany A Caruso et al. "Impact of a demand-side integrated WASH and nutrition community-based care group intervention on behavioural change: a randomised controlled trial in western Kenya", BMJ Global Health, 2020

Publication

<1 %

dizifood.com

22	Internet Source	<1 %
23	digilib2.unisayogya.ac.id Internet Source	<1 %
24	erepository.mkuit.ac.rw Internet Source	<1 %
25	ejournal.undana.ac.id Internet Source	<1 %
26	eprints.umm.ac.id Internet Source	<1 %
27	journal.uin-alauddin.ac.id Internet Source	<1 %
28	opendentistryjournal.com Internet Source	<1 %
29	www.research-system.siam.edu Internet Source	<1 %
30	Niu Di, Shanshan Li, Hao Xiang, Yinyu Xie et al. "Associations of Residential Greenness with Depression and Anxiety in Rural Chinese Adults", The Innovation, 2020 Publication	<1 %
31	docplayer.info Internet Source	<1 %
32	downloads.hindawi.com Internet Source	<1 %

33	e-journal.unair.ac.id Internet Source	<1 %
34	hdl.handle.net Internet Source	<1 %
35	jgpt.co.in Internet Source	<1 %
36	pubs.acs.org Internet Source	<1 %
37	www.dovepress.com Internet Source	<1 %
38	www.internationaljournalsrsg.org Internet Source	<1 %
39	Sa'ed Zyoud, Jawad Shalabi, Kathem Imran, Lina Ayaseh et al. "Knowledge, attitude and practices among parents regarding food poisoning: a cross-sectional study from Palestine", BMC Public Health, 2019 Publication	<1 %
40	"The Effect of Environmental Factors on the Event of Acute Diarrhea", International Journal of Innovative Technology and Exploring Engineering, 2020 Publication	<1 %
41	Myra Mesnuath Kosapilawan, Delima Citra Dewi Gunawan, Fera Nofiartika. "Correlation between milk bottle practice and diarrhea on	<1 %

children under five at Puskesmas Umbulharjo 1, Yogyakarta City", Ilmu Gizi Indonesia, 2019

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Implementation of Hygiene and Environmental Sanitation in Under Five Years Old Diarrhea Patients at Surabaya Primary Health Center

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15
