

A mathematical model of tuberculosis (TB) transmission with children and adults groups: A fractional model

by Fatmawati Fatmawati

Submission date: 02-Jun-2020 03:54PM (UTC+0800)

Submission ID: 1336399293

File name: 6..pdf (1.09M)

Word count: 10491

Character count: 45031



84

Research article

A mathematical model of tuberculosis (TB) transmission with children and adults groups: A fractional model

Fatmawati¹, Muhammad Altaf Khan^{2,*}, Ebenezer Bonyah³, Zakia Hammouch⁴ and Endrik Mifta Shaiful¹

31

¹ Department of Mathematics, Faculty of Science and Technology, Universitas Airlangga Surabaya 60115, Indonesia

² Faculty of Natural and Agricultural Sciences, University of the Free State, South Africa

³ Department of Mathematics Education, University of Education Winneba, Kumasi Campus, Kumasi, Ghana

⁴ Département de Mathématiques, Faculté des Sciences et Techniques Errachidia, Université Moulay Ismail, Morocco

* Correspondence: Email: altafdir@gmail.com.

Abstract: We consider a novel fractional model to investigate the (tuberculosis) TB model dynamics with two age groups of human, that is, the children and the adults. First, we formulate the model and present the basic results associated to the model. Then, using the fractional operators, Caputo and the Atangana-Baleanu and obtain a generalized model. Further, we give a novel numerical approach for the solution of the fractional model and obtain their approximate solution. We show graphical results with various values of the fractional order. A comparison of the two operators are shown graphically. The results obtained through Atangana-Baleanu operator is flexible than that of Caputo derivative. The infection in tuberculosis (TB) infected people decreases fast when decreasing the fractional order.

Keywords: tuberculosis; two-age-class transmission; fractional differential equations; Caputo derivative; Atangana-Baleanu (A-B) derivative

Mathematics Subject Classification: 34A34

1. Introduction

The human population has not been spared by infectious diseases irrespective of the medical facilities. In spite of great achievement in the advancement of medicine, the world is still struggling to combat infectious diseases [1]. *Mycobacterium tuberculosis* (MTB) is responsible for Tuberculosis (TB) infection which primarily affects the lungs (pulmonary TB). However, the disease can also affect

some other systems in the human body such as the circulatory system, the central nervous system, the genital-urinary system. TB can be transmitted through a medium including the kiss, speak, cough, sneeze [1, 2]. It can even affect individuals through the utilization of infected persons unsterilized eating utensils. In some instance, pregnant women who have an active TB can affect the baby in the womb. The actively infected individuals can cause infection but not the latent TB patient. The transmission of TB depends on the number of infectious droplets expelled by a carrier, the quality of ventilation available and finally, the duration of the individuals exposed to virulence of the MTB strain. The spread of TB can be reduced by isolating the active infectious individual and immediately introducing viable anti-tuberculosis therapy [3–5]. At the moment, 95% of the world total TB patients of 8 million each year are found in the developing countries. The age bracket that most affected by TB is 15–59. In the developing countries, TB has been identified as the leading death causes in the majority of the Sub-Sahara counties [6]. TB has been established as the leading cause of death in sub-Saharan Africa which account for at least 2 million deaths [7]. TB in children under the age of fifteen (15) is very critical public health problem since it account for most of the current TB transmission [8].

Adaption of modern practice help improves TB control because of factors such as endogenous reactivation, the presence of multi-drug resistant TB, the high rate of HIV incidence in the last two decades. Exogenous re-infection of TB is another concern and very common in Africa. A comprehensive understanding of treatment with respect to the regional transmission of TB is crucial in the fighting of this disease [9].

Mathematical modeling in recent times has been identified as an essential tool that provides qualitative information about the epidemiology of many diseases and strategies in controlling them. Several theoretical studies have been undertaken with regard to TB infection. Several integer-order TB models have been constructed and investigated by some authors [10–14]. The mentioned integer TB models do not predict accurately because they do not have memory effect in order to provide an accurate prediction. However, non-integer models possess memory effect and most of the operators have crossover properties that enhance accurate predictions.

Fractional calculus (FC) is a branch of applied mathematics that generalizes the integer calculus to fractional order [15, 16]. In recent times, the use of fractional calculus in the scientific community has gained enough attention because of its numerous applications including epidemiology, engineering, water resource management [17–24]. The use of fractional derivative in modeling epidemiology has become important because the result takes into consideration the memory effect which naturally occurs in several biological models. The reason is that fractional derivative models give the real phenomena associated with the given problem. The fractional of derivatives and integrals give a vivid description of the memory and hereditary characteristics inherent in many materials and other processes [25]. Usually, the integer order can provide information about two points which realistically may not be true. The concept of classical fractional-order derivative was proposed by Riemann and Liouville. It was well known that Caputo and Riemann-Liouville have singular kernels. Next, Caputo and Fabrizio in [26] presented a new definition of fractional derivative without singular kernel which proved to be good and many researchers are applying. Caputo and Fabrizio [27] also investigated the notion of fractional derivative and applications to the hysteresis phenomena. The fractional calculus is now used in mathematical modeling of impulsive fractional functional differential equations. A more interesting study regarding the impulsive fractional functional differential equations and their

applications is studied in [28].

Recently, Atangana and Baleanu developed a new operator which is based on generalized Mittag-Leffler function where the kernel is non-singular and non-local kernel [29]. Numerous studies on non-integer order models in the sense Atangana and Baleanu have been carried out by researchers. For instance, Atangana and Koca in [30] studied chaos in a simple nonlinear system using Atangana-Baleanu derivatives in Caputo sense. Atangana and Owolabi in [31] developed a new numerical approach for fractional differential equations for solving the system of fractional derivative equations which proved to be efficient. Atangana and Gomez [32] presented a detailed analysis of the decolonization of fractional calculus rules: breaking commutativity and associativity to capture more natural phenomena. Khan et al. [33] studied the TB dynamics under the framework of the work of Atangana-Baleanu derivative with relapse cases. Bonyah in [34] studied a new 5-D hyperchaotic system with four wings by Atangana-Baleanu operator. Khan and Atangana [35] studied the Ebola disease with different fractional operators. The comparison of Atangana-Baleanu derivative with Caputo-Fabrizio for the dengue model are studied in [36]. Bank data through fractional modeling and their analysis is studied in [37]. The dynamics of the chaotic system through Caputo-Fabrizio and the Atangana-Baleanu derivative is studied in [38]. The authors studied in [39] through different fractional operators of fractional calculus for bank data. Using the Atangana-Baleanu derivative the authors studied the TB dynamics [40]. The previously defined fractional operators have some deficiencies which may or may not correctly investigate the dynamics of such mathematical models that come from science and engineering fields. The Caputo derivative has a singular and local kernel which may not exactly studied the dynamics of a particular problem but these operators are still well known to the researchers and uses for the modeling dynamics. The Caputo-Fabrizio operator was defined without singular kernel. Many researchers used this operator for modeling purposes and still popular among researchers. To have a more precise definition of fractional operator a new operator called Atangana-Baleanu derivative was defined which has a non-singular and non-local kernel and has been used widely for many problems of science and engineering and found it interesting. This new definition is now much famous among researchers around the world. Here, in this work, we consider this newly operator to formulate a new mathematical model for TB dynamics in which we consider the adults and the children as a two groups and present to discuss deeply its mathematical as well as numerical results. This is the first attempt of the authors to have a new mathematical that exploring the children and adults TB dynamics.

It is well known that integer order model does not describes well the dynamics of the disease because there involve history of the disease and their memory. The fractional order models well address the memory effect involve in the disease. At each value of the fractional order parameters we can have a solution but in integer order model we don't have except at the integer case. The difference between two points can be effectively estimated through fractional order models. It is also well known that fractional order models are good for data fitting, where we have variety of choice's for the fractional order parameter. The Caputo operator is singular and local while the AB operator is non-singular and nonlocal. The cross over behavior in the TB model can be addressed well through the AB derivative. Therefore, we considered both the operators and its comparison through graphical results.

To date, there has not been anyone who has investigated the dynamics of TB model with children and adults population using Caputo and Atangana-Baleanu derivative. So, this study investigates a fractional order model in the Atangana-Baleanu sense to explore the spread of TB diseases with in

children and adults population. We classify the population into children and adult classes. Based on the risk of transmission, most TB in children is usually not contagious compared to TB in adults [8,41]. Hence, the TB model that involves two-age-classes that distinguishes children and adults can indicate a new understanding of TB transmission.

This paper begins by mathematical preliminaries of the fractional derivative. The description of the model with the two-age-class transmission is given in Section 3. The TB model in Caputo derivative with basic properties, calculation of the basic reproduction number, and the numerical result are done in Section 4. The expression of the TB model in Atangana–Baleanu derivative with the existence and uniqueness of the solution, the numerical iterative, and also the numerical simulation are presented in Section 5. The comparison results of both derivatives with various fractional order parameters are shown in Section 6. Finally, the conclusion is shown in Section 7.

2. Essentials of fractional calculus

We present the details of the fractional operators in the following sections that should be used later in the proposed study.

Definition 2.1. (see [16]) For $\alpha > 0$, with $n - 1 < \alpha < n$, $n \in \mathbb{N}$, the fractional derivative in the sense of Caputo is defined to be:

$${}^c D_t^\alpha f(t) = \frac{d^\alpha f(t)}{dt^\alpha} := \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{(n-\alpha-1)} f^{(n)}(s) ds, \quad (2.1)$$

where the symbol $\Gamma(\cdot)$ denotes the gamma function.

The Riemann-Liouville fractional integral is defined by

$$I_t^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{(\alpha-1)} f(s) ds. \quad (2.2)$$

Definition 2.2. (see [29]) Suppose $f \in H^1(0, b)$, $b > 0$, $\alpha \in [0, 1]$, then the Atangana-Baleanu fractional operator in Caputo sense (ABC) can be written as follows:

$${}^{ABC} D_t^\alpha f(t) = \frac{B(\alpha)}{1-\alpha} \int_0^t f'(s) E_\alpha \left[-\alpha \frac{(t-s)^\alpha}{1-\alpha} \right] ds, \quad (2.3)$$

where $B(\alpha) = 1 - \alpha + \frac{\alpha}{\Gamma(\alpha)}$ is a normalized function with $B(0) = B(1) = 1$ and E_α is Mittag-Leffler function

$$E_\alpha(z) = \sum_{k=0}^{\infty} \frac{(z)^k}{\Gamma(\alpha k + 1)}, \alpha > 0.$$

The integral associated to the ABC derivative in Riemann-Liouville sense is given by,

$$I_t^\alpha f(t) = \frac{1-\alpha}{B(\alpha)} f(t) + \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_0^t f(s)(t-s)^{\alpha-1} ds. \quad (2.4)$$

3. Description of the model

The present section describes the formulation of the mathematical model of TB transmission with two groups of human that is, the children and the adults. We design the model by splitting the population into children (C) and the adults class (A). Furthermore, each of the classes is divided into three classes, namely, the susceptible classes (S_C, S_A), the latent TB classes (L_C, L_A), and the active TB classes (I_C, I_A), where S_C, L_C and I_C represent the susceptible, latent and the active population at any time t respectively, whereas the adults sub-population are respectively shown by S_A, L_A and I_A . Hence, the total population is $N = S_C + L_C + I_C + S_A + L_A + I_A$. Individuals who are in the latent stage are not infectious and are thus unable to transmit bacteria. We also assumed that the natural death rate of the children and adults population is equal as we consider the average natural death rate of the total population.

The majority of the children with tuberculosis are not infectious to others [8,41]. Hence, we assume that only the active TB adults could spread TB within the population. In this model, it is assumed that children got TB infection by doing contacts with active TB adults. Therefore, the latent TB and active TB children do not spread TB within the population. A small percentage of newly infected individuals from the sub-population is assumed to undergo fast progress directly into the class of active TB, while the rest are latently infected and enter the latent class [42]. Furthermore, the TB patients both children and adults cannot recover completely but will enter the latent stage again. Considering these assumptions and facts the following system of evolutionary dynamics of the TB individuals is described through the following systems where the description of the parameters flows and the variables are given in Figure 1.




Figure 1. Tuberculosis transmission diagram.

$$\begin{aligned}\frac{dS_C}{dt} &= \Lambda - \beta_C I_A S_C - (g + \mu) S_C, \\ \frac{dL_C}{dt} &= \beta_C (1 - p_C) I_A S_C + \gamma_C I_C - k_C (1 - r_C) L_C - (g + \mu) L_C,\end{aligned}$$

$$\begin{aligned}
 \frac{dI_C}{dt} &= \beta_C p_C I_A S_C + k_C(1 - r_C)L_C - \gamma_C I_C - (g + \mu + d_C)I_C, \\
 \frac{dS_A}{dt} &= gS_C - \beta_A I_A S_A - \mu S_A, \\
 \frac{dL_A}{dt} &= (1 - p_A)\beta_A I_A S_A + gL_C + \gamma_A I_A - k_A(1 - r_A)L_A - \mu L_A, \\
 \frac{dI_A}{dt} &= p_A \beta_A I_A S_A + k_A(1 - r_A)L_A + gI_C - (\gamma_A + \mu + d_A)I_A,
 \end{aligned} \tag{3.1}$$

where the initial values for the model variables are given by

$$S_C(0) = S_{C0}, L_C(0) = L_{C0}, I_C(0) = I_{C0}, S_A(0) = S_{A0}, L_A(0) = L_{A0}, \text{ and } I_A(0) = I_{A0}.$$

In the above model (3.1), the population of healthy people is generated by the birth rate Λ and the growing up rate of children is shown by g . The natural death rate of children and adults is denoted by μ . The effective contact rate for both the children and adults is shown by β_C and β_A . The proportion of fast progression of latent class children to infected class children is given by p_C while for adults from latent adult class to infected adult class is by p_A . The rate of progression from latent to infected class for the children compartment is shown by k_C while for the adults infection progression from latent to infected class is given by k_A . The rate of effective therapy for children and adults are shown respectively by γ_C and γ_A . The rate of effective chemoprophylaxis for children and adults are shown by r_C and r_A respectively. Natural death from TB disease in children infected class is given by d_C while for the adults with TB infection by d_A . The description of the parameters for the TB model is set out in Table 1.

Table 1. Parameters with descriptions of the TB model.

Description	Parameter	
Populations growth rate	Λ	
Growing-up rate for children	g	
Natural death rate	μ	
	Children population	Adult population
Infection rate	β_C	β_A
Proportion of fast progression	p_C	p_A
Rate of progression from latent to the infectious	k_C	k_A
Rate of effective therapy	γ_C	γ_A
Rate of effective chemoprophylaxis	r_C	r_A
TB death rate	d_C	d_A

The model (3.1) has the biologically feasible region on Ω_1 with

$$\Omega_1 = \left\{ (S_C, L_C, I_C, S_A, L_A, I_A) \in \mathbb{R}_+^6 : 0 \leq N \leq \frac{\Lambda}{\mu} \right\},$$

which is feasible biologically and all the results associated holds for it. It should be noted that the parameters involved in the model (3.1) are associated to humans so it is non-negative and the region shown by Ω_1 is positive invariant for the described model (3.1) above. The region shown for the model (3.1) below is well-posed and the entire solutions for the initial values belonging to Ω_1 , remains in Ω_1 for every time $t \geq 0$.

Next, in the following section, we explore the TB model in Caputo sense. We apply the definitions of Caputo operator described in above section, and apply it on our model (3.1) and present the results associated to it in the following section in details.

4. A Caputo model

This section present a TB model with two groups of humans which is shown in (3.1), by applying the Caputo derivative on it. The fractional order model of TB transmission with two-age-class in the Caputo sense is given by:

$$\begin{aligned}
 {}^C D_t^\alpha S_C &= \Lambda - \beta_C I_A S_C - (g + \mu) S_C, \\
 {}^C D_t^\alpha L_C &= \beta_C (1 - p_C) I_A S_C + \gamma_C I_C - k_C (1 - r_C) L_C - (g + \mu) L_C, \\
 {}^C D_t^\alpha I_C &= \beta_C p_C I_A S_C + k_C (1 - r_C) L_C - \gamma_C I_C - (g + \mu + d_C) I_C, \\
 {}^C D_t^\alpha S_A &= g S_C - \beta_A I_A S_A - \mu S_A, \\
 {}^C D_t^\alpha L_A &= (1 - p_A) \beta_A I_A S_A + g L_C + \gamma_A I_A - k_A (1 - r_A) L_A - \mu L_A, \\
 {}^C D_t^\alpha I_A &= p_A \beta_A I_A S_A + k_A (1 - r_A) L_A + g I_C - (\gamma_A + \mu + d_A) I_A,
 \end{aligned} \tag{4.1}$$

where α represents the fractional order and $0 < \alpha \leq 1$. The initial values for the model variables in fractional order model (4.1) are given by,

$$S_C(0) = S_{C0}, L_C(0) = L_{C0}, I_C(0) = I_{C0}, S_A(0) = S_{A0}, L_A(0) = L_{A0}, \text{ and } I_A(0) = I_{A0}.$$

4.1. Non-negative solutions

This section deals with the positivity of the model solution in the Caputo derivative. We assume that the initial condition of the model (4.1) is non-negative. Let us consider

$$\Omega_2 = \{x = (S_C, L_C, I_C, S_A, L_A, I_A) \in \mathbb{R}_+^6 : S_C, L_C, I_C, S_A, L_A, I_A \geq 0\}.$$

Here, recalling the important concept given in [43] to prove our result.

Lemma 4.1. (see [43]) Suppose that $f(t) \in C[a, b]$ and ${}^C D_t^\alpha f(t) \in C[a, b]$, for $0 < \alpha \leq 1$, then we have

$$f(t) = f(a) + \frac{1}{\Gamma(\alpha)} {}^C D_t^\alpha f(\xi)(t - a)^\alpha$$

with $a \leq \xi \leq t, \forall t \in (a, b)$.

Corollary 4.2. Consider that $f(t) \in C[a, b]$ and ${}^C D_t^\alpha f(t) \in C[a, b]$, for $0 < \alpha \leq 1$. Then if

(1) ${}^C D_t^\alpha f(t) \geq 0, \forall t \in (a, b)$, then $f(t)$ is non-decreasing.

(2) ${}^C D_t^\alpha f(t) \leq 0, \forall t \in (a, b)$, then $f(t)$ is non-increasing.

Now, we state the main theorem.

Theorem 4.3. A unique solution exists for $x(t) = (S_C, L_C, I_C, S_A, L_A, I_A)^T$ of the model (4.1) and the solution will be in Ω_2 . Additionally, the solution is non-negative.

Proof. Using Theorem 3.1. and Remark 3.2. of [44], it is easily to prove the solution existence and also the uniqueness of the solution of the model (4.1) for all $t > 0$. Next, we will prove the positivity of the solution of the model (4.1). In order to do, we have to prove for every hyperplane bounding the nonnegative orthant, of the field points of the vector Ω_2 . It follows from the model given by (4.1), we have

$$\begin{aligned} {}^C D_t^\alpha S_C |_{S_C=0} &= \Lambda \geq 0, \\ {}^C D_t^\alpha L_C |_{L_C=0} &= \beta_C(1 - p_C)I_A S_C + \gamma_C I_C \geq 0, \\ {}^C D_t^\alpha I_C |_{I_C=0} &= \beta_C p_C I_A S_C + k_C(1 - r_C)L_C \geq 0, \\ {}^C D_t^\alpha S_A |_{S_A=0} &= g S_C \geq 0, \\ {}^C D_t^\alpha L_A |_{L_A=0} &= (1 - p_A)\beta_A I_A S_A + g L_C + \gamma_A I_A \geq 0, \\ {}^C D_t^\alpha I_A |_{I_A=0} &= k_A(1 - r_A)L_A + g I_C \geq 0. \end{aligned}$$

By using Corollary 1, the model solution remains in Ω_2 . \square

Thus, it follows from the results described above the feasible given by Ω_2 is positively invariant for model (4.1).

4.2. Model fixed points

We present here the possible fixed points of the fractional model given by (4.1) and then, we obtain at these possible equilibrium points the stability analysis of the model. For the given model (4.1), there exists two possible equilibrium points that is, the disease free and the endemic equilibrium. We can obtain these equilibrium points by setting the time rate of change equal to zero of the model (4.1) and is given by:

$$\begin{aligned} {}^C D_t^\alpha S_C |_{S_C=0} &= 0, \\ {}^C D_t^\alpha L_C |_{L_C=0} &= 0, \\ {}^C D_t^\alpha I_C |_{I_C=0} &= 0, \\ {}^C D_t^\alpha S_A |_{S_A=0} &= 0, \\ {}^C D_t^\alpha L_A |_{L_A=0} &= 0, \\ {}^C D_t^\alpha I_A |_{I_A=0} &= 0. \end{aligned}$$

We have the following expression,

$$\begin{aligned} \Lambda - \beta_C I_A S_C - (g + \mu)S_C &= 0, \\ \beta_C(1 - p_C)I_A S_C + \gamma_C I_C - k_C(1 - r_C)L_C - (g + \mu)L_C &= 0, \end{aligned}$$

$$\begin{aligned}
\beta_C p_C I_A S_C + k_C(1-r_C)L_C - \gamma_C I_C - (g + \mu + d_C)I_C &= 0, \\
gS_C - \beta_A I_A S_A - \mu S_A &= 0, \\
(1-p_A)\beta_A I_A S_A + gL_C + \gamma_A I_A - k_A(1-r_A)L_A - \mu L_A &= 0, \\
p_A \beta_A I_A S_A + k_A(1-r_A)L_A + gI_C - (\gamma_A + \mu + d_A)I_A &= 0.
\end{aligned} \tag{4.2}$$

At the disease free equilibrium, denoted by E_{01} , we have from (4.2),

$$E_{01} = \left(\frac{\Lambda}{g + \mu}, 0, 0, \frac{g\Lambda}{\mu(g + \mu)}, 0, 0 \right).$$

The endemic equilibrium of the model (4.1) is given by $E_{22} = (S_C^*, L_C^*, I_C^*, S_A^*, L_A^*, I_A^*)$ where

$$\begin{aligned}
S_C^* &= \frac{\Lambda}{I_A^* \beta_C + g + \mu} \\
L_C^* &= \frac{\beta_C(1-p_C)I_A^* S_C^* + \gamma_C I_C^*}{\mu + g + k_C(1-r_C)} \\
I_C^* &= \frac{\beta_C p_C I_A^* S_C^* + k_C(1-r_C)L_C^*}{\gamma_C + g + \mu + d_C} \\
S_A^* &= \frac{gS_C^*}{\beta_A I_A^* + \mu} \\
L_A^* &= \frac{(1-p_A)\beta_A I_A^* S_A^* + gL_C^* + \gamma_A I_A^*}{k_A(1-r_A) + \mu} \\
I_A^* &= \frac{k_A(1-r_A)L_A^* + gI_C^*}{\gamma_A + \mu + d_A - \beta_A p_A S_A^*}.
\end{aligned}$$

The endemic equilibrium E_{22} exists if $\gamma_A + \mu + d_A > \beta_A p_A S_A^*$.

In order to find the basic reproduction number \mathcal{R}_0 , we consider the method in [45] and the matrices are given by

$$F = \begin{pmatrix} 0 & 0 & 0 & P_9 \\ 0 & 0 & 0 & P_{10} \\ 0 & 0 & 0 & P_{12} \\ 0 & 0 & 0 & P_{13} \end{pmatrix}, \quad V = \begin{pmatrix} P_2 & -\gamma_C & 0 & 0 \\ -P_5 & P_3 & 0 & 0 \\ -g & 0 & P_4 & -r_A \\ 0 & -g & -P_6 & P_7 \end{pmatrix}. \tag{4.3}$$

The basic reproduction number is the spectral radius of $\rho(FV^{-1})$ and is given by

$$\mathcal{R}_0 = \mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3 + \mathcal{R}_4, \tag{4.4}$$

where

$$\mathcal{R}_1 = \frac{P_6 P_3 (g P_9 + P_2 P_{12})}{(P_4 P_7 - P_6 r_A)(P_2 P_3 - P_5 \gamma_C)},$$

$$\begin{aligned}
\mathcal{R}_2 &= \frac{P_4 P_2 (g P_{10} + P_3 P_{13})}{(P_4 P_7 - P_6 r_A) (P_2 P_3 - P_5 \gamma_C)}, \\
\mathcal{R}_3 &= \frac{P_6 \gamma_C (g P_{10} - P_5 P_{12})}{(P_4 P_7 - P_6 r_A) (P_2 P_3 - P_5 \gamma_C)}, \\
\mathcal{R}_4 &= \frac{P_4 P_5 (g P_9 - P_{13} \gamma_C)}{(P_4 P_7 - P_6 r_A) (P_2 P_3 - P_5 \gamma_C)},
\end{aligned} \tag{4.5}$$

where

$$\begin{aligned}
P_1 &= g + \mu, P_2 = k_C (1 - r_C) + g + \mu, P_3 = \gamma_C + d_C + g + \mu, \\
P_4 &= k_A (1 - r_A) + \mu, P_5 = k_C (1 - r_C), P_6 = k_A (1 - r_A), \\
P_7 &= \gamma_A + d_A + \mu, P_8 = \beta_C S_C^0, P_9 = \beta_C (1 - p_C) S_C^0, P_{10} = \beta_C p_C S_C^0, \\
P_{11} &= \beta_A S_A^0, P_{12} = \beta_A (1 - p_A) S_A^0, P_{13} = \beta_A p_A S_A^0,
\end{aligned} \tag{4.6}$$

where S_C^0 and S_A^0 represent the disease-free equilibrium (DFE).

Next, we show the stability of the system at the disease-free case, we follow the following theorem.

Theorem 4.4. For any $p_1, p_2 \in \mathbb{Z}$, such that $\gcd(p_1, p_2) = 1$. Let $\alpha = \left(\frac{p_1}{p_2}\right)$ and consider $M = p_2$, then the DFE of the system (4.1) is locally asymptotically stable (LAS) if $|\arg(\lambda)| > \frac{\pi}{2M}$, for all roots λ of the characteristic Eq. (4.7) of the matrix $J_{E_{01}}$.

$$\det(\text{diag}[\lambda^{p_1} \lambda^{p_1} \lambda^{p_1} \lambda^{p_1} \lambda^{p_1} \lambda^{p_1}] - J_{E_{01}}) = 0. \tag{4.7}$$

Proof. At the disease free equilibrium E_{01} , the Jacobian matrix is given by

$$J(E_{01}) = \begin{pmatrix} -P_1 & 0 & 0 & 0 & 0 & -P_8 \\ 0 & -P_2 & \gamma_C & 0 & 0 & P_9 \\ 0 & P_5 & -P_3 & 0 & 0 & P_{10} \\ g & 0 & 0 & -\mu & 0 & -P_{11} \\ 0 & g & 0 & 0 & -P_4 & P_{12} + \gamma_A \\ 0 & 0 & g & 0 & P_6 & P_{13} - P_7 \end{pmatrix}. \tag{4.8}$$

We obtain the characteristics equation associated to $J(E_{01})$, given by

$$(\lambda^{p_1} + \mu)(\lambda^{p_1} + \mu + g)[\lambda^{4p_1} + b_1 \lambda^{3p_1} + b_2 \lambda^{2p_1} + b_3 \lambda^{p_1} + b_4] = 0. \tag{4.9}$$

The arguments of the roots of the equation $\lambda^{p_1} + \mu = 0$, $\lambda^{p_1} + \mu + g = 0$ are as follow:

$$\arg(\Pi_k) = \frac{\pi}{p_1} + k \frac{2\pi}{p_1} > \frac{\pi}{M} > \frac{\pi}{2M}, \text{ where } k = 0, 1, \dots, (p_1 - 1). \tag{4.10}$$

In Eq. (4.9), the roots are negative, i.e., $-\mu$, $-(\mu + g)$, while the rest are computed from

$$\mathcal{Q}(\lambda) = [\lambda^{4p_1} + b_1 \lambda^{3p_1} + b_2 \lambda^{2p_1} + b_3 \lambda^{p_1} + b_4], \tag{4.11}$$

where

$$\begin{aligned}
 b_1 &= \gamma_A + d_A + k_A(1 - r_A) - \beta_A p_A S_A^0 + \gamma_C + d_C + k_C(1 - r_C) \\
 &\quad + 2g + 4\mu, \\
 b_2 &= \underbrace{(\gamma_C + d_C + g + \mu)(\gamma_A + d_A + k_A(1 - r_A) + 2\mu) - \beta_A p_A S_A^0 P_2 P_4}_{70} \\
 &\quad + (k_A(1 - r_A) + \mu)(k_C(1 - r_C) + g + \mu) + (d_A + \mu)(k_A(1 - r_A) + \mu) + \mu\gamma_A \\
 &\quad + \underbrace{P_7 P_2 - \beta_A k_A(1 - p_A)(1 - r_A) S_A^0}_{75} \\
 &\quad + \left((d_C + g + \mu)(k_C(1 - r_C) + g + \mu) + \gamma_C(g + \mu) \right) \\
 &\quad \times \left(1 - \frac{\beta_A p_A S_A^0 (\gamma_C + d_C + g + \mu) + g \beta_C P_C S_C^0}{(d_C + g + \mu)(k_C(1 - r_C) + g + \mu) + \gamma_C(g + \mu)} \right), \\
 b_3 &= (P_4 + P_7 - P_{13}) \left(d_C(k_C(1 - r_C) + g + \mu) + (g + \mu)(\gamma_C + k_C(1 - r_C) + g + \mu) \right) \\
 &\quad (P_3 + P_2) \left(d_A(k_A(1 - r_A) + \mu) + k_A(1 - r_A)(\mu - \beta_A S_A^0) + \mu(\gamma_A - \beta_A p_A S_A^0 + \mu) \right) \\
 &\quad - g \beta_C S_C^0 (k_A(1 - r_A) + p_C(g + 2\mu) + k_C(1 - r_C)), \\
 b_4 &= C_1 C_2 (1 - \mathcal{R}_0), \tag{4.12}
 \end{aligned}$$

where $C_1 = (d_A(k_A(1 - r_A) + \mu) + \mu(\gamma_A + k_A(1 - r_A) + \mu))$,
 $C_2 = (d_C(k_C(1 - r_C) + g + \mu) + (g + \mu)(\gamma_C + k_C(1 - r_C) + g + \mu))$.

The function $Q(\lambda)$ will give eigenvalues with negative real part if $b_i > 0$ for $i = 1, 2, 3, 4$ and $b_1 b_2 b_3 > b_1^2 b_4 + b_3^2$. All $b_i > 0$ when $\mathcal{R}_0 < 1$ and the conditions $b_1 b_2 b_3 > b_1^2 b_4 + b_3^2$ ensure the stability of the disease free case when $\mathcal{R}_0 < 1$. If $\mathcal{R}_0 < 1$, then the necessary condition fulfil for all the roots of characteristics equation i.e., $|\arg(\lambda)| > \frac{\pi}{2M}$. Thus the DFE is LAS for $\mathcal{R}_0 < 1$. \square

The following subsection explore the numerical simulation of the Caputo model.

4.3. Numerical simulation for Caputo model

The aim of this section is to discuss a numerical simulation of the model (4.1). The simulation is done with several fractional order values to determine population dynamics in the model. The numerical results of the Caputo derivative are obtained by the Predictor-Corrector method [46, 47]. Parameters values used in these simulations are set out in Table 2. We take the initial condition $S_C(0) = 1500$, $L_C(0) = 135$, $I_C(0) = 75$, $S_A(0) = 1000$, $L_A(0) = 103$ and $I_A(0) = 60$. Here, we use 50 years for the time horizon. The simulations are conducted with different values of the order of the fractional derivative $\alpha = 1, 0.9, 0.8, 0.7, 0.6$ and 0.5 . We have the simulation results of the TB model (4.1) in Figure 2. In this case the value of R_0 is $R_0 = 91.5021 > 1$, which indicates the disease will persist in the population. We can observe in Figure 2 that as the values of α decrease, the individuals infected with TB decreases whereas the population of susceptible individuals increases.




Figure 2. Numerical results for Caputo model.

Next, we will see the effect of chemoprophylaxis and treatment on the active TB populations in both children and adults. The numerical simulation is deployed by varying the effectiveness of chemoprophylaxis (r_C, r_A) and treatment (γ_C, γ_A). In Figures 3, 4, we display the dynamic of the active TB populations for different values of the efficacy of chemoprophylaxis and treatment. For $\alpha = 0.9$, it is shown that the population of the active TB both children and adults have decreased when the efficacy of chemoprophylaxis and treatment have increased. Conversely, when the efficacy of chemoprophylaxis and treatment falls, the population of the active TB both children and adults will rise. The likewise behavior is seen for $\alpha = 0.7$. This shows that chemoprophylaxis and treatment were given to patients with latent stage and active TB respectively can reduce the number of TB infection populations of both children and adults.




Figure 3. Numerical results for Caputo model with various r_C and r_A .




Figure 4. Numerical results for Caputo model with various γ_C and γ_A .

7
Table 2. Parameter values for simulations.

Parameter	Value	Ref.	Parameter	Value	Ref.
Λ	1000	Assumed	β_C	0.01	[48]
β_A	0.02	[48]	μ	0.0143	[49]
g	$\frac{1}{14}$	[50]	γ_C	0.5	Assumed
γ_A	0.5	Assumed	p_C	0.4	Assumed
p_A	0.3	Assumed	k_C	0.005	[49]
k_A	0.005	[49]	r_C	0.5	Assumed
r_A	0.5	Assumed	d_C	0.0575	[51]
d_A	0.05751	[51]	d	0.05751	[51]
p	0.3	Assumed	β	0.02	[48]
γ	0.5	Assumed	k	0.005	[49]
r	0.5	Assumed			

Next, we express the model of TB in Atangana-Baleanu derivative where their kernel is non-singular and non-local which best describes the dynamics of real world problem.

5. TB model in the Atangana–Baleanu sense

The newly derivative called the Atangana-Baleanu derivative where their kernel is non-local and non singular is considered here to apply on our model presented in (3.1). We follow the definition described above and have the following representation:

$$\begin{aligned}
 {}_0^{ABC}D_t^\alpha S_C &= \Lambda - \beta_C I_A S_C - (g + \mu) S_C, \\
 {}_0^{ABC}D_t^\alpha L_C &= \beta_C(1 - p_C) I_A S_C + \gamma_C I_C - k_C(1 - r_C) L_C - (g + \mu) L_C, \\
 {}_0^{ABC}D_t^\alpha I_C &= \beta_C p_C I_A S_C + k_C(1 - r_C) L_C - \gamma_C I_C - (g + \mu + d_C) I_C, \\
 {}_0^{ABC}D_t^\alpha S_A &= g S_C - \beta_A I_A S_A - \mu S_A, \\
 {}_0^{ABC}D_t^\alpha L_A &= (1 - p_A) \beta_A I_A S_A + g L_C + \gamma_A I_A - k_A(1 - r_A) L_A - \mu L_A, \\
 {}_0^{ABC}D_t^\alpha I_A &= p_A \beta_A I_A S_A + k_A(1 - r_A) L_A + g I_C - (\gamma_A + \mu + d_A) I_A,
 \end{aligned} \tag{5.1}$$

with some appropriate initial conditions. For the TB fractional model in Atangana-Baleanu form is studied further to obtain their existence and uniqueness. We have the following:

5.1. The existence and uniqueness of the solution

Here, we display the uniqueness and existence results associated to the AB model (5.1). In order to show the results for the model we follow the results of the fixed-point theory. We first write the (5.1) in the form shown below,

$$\begin{cases}
 {}_0^{ABC}D_t^\alpha u(t) = G(t, u(t)) \\
 u(0) = u_0, \quad 0 < t < T < \infty.
 \end{cases} \tag{5.2}$$

In system (5.2), $u(t) = (S_C, L_C, I_C, S_A, L_A, I_A)$ is simply a vector having the state variable and G is therefore a continuous vector function given as

$$G = \begin{pmatrix} G_1 \\ G_2 \\ G_3 \\ G_4 \\ G_5 \\ G_6 \end{pmatrix} = \begin{pmatrix} \Lambda - \beta_C I_A S_C - (g + \mu) S_C \\ \beta_C(1 - p_C) I_A S_C + \gamma_C I_C - k_C(1 - r_C) L_C - (g + \mu) L_C \\ \beta_C p_C I_A S_C + k_C(1 - r_C) L_C - \gamma_C I_C - (g + \mu + d_C) I_C \\ g S_C - \beta_A I_A S_A - \mu S_A \\ (1 - p_A) \beta_A I_A S_A + g L_C + \gamma_A I_A - k_A(1 - r_A) L_A - \mu L_A \\ p_A \beta_A I_A S_A + k_A(1 - r_A) L_A + g I_C - (\gamma_A + \mu + d_A) I_A \end{pmatrix}$$

and $u_0(t) = (S_C(0), L_C(0), I_C(0), S_A(0), L_A(0), I_A(0))$ representing the state variables initial conditions in vector form. The above function namely, G satisfies the condition of Lipschitz continuity and can be described as below:

$$\|G(t, u_1(t)) - G(t, u_2(t))\| \leq M \|u_1(t) - u_2(t)\|. \tag{5.3}$$

The following result is provided in order to show the model existence and their solution as well as uniqueness for the model (5.1).

Theorem 5.1. (Existence and uniqueness) *The model given by (5.2) has the unique solution provided that the following condition satisfy,*

$$\frac{(1-\alpha)}{ABC(\alpha)}M + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)}T_{\max}^{\alpha}M < 1. \quad (5.4)$$

Proof. To prove the above statement, we need to use the fractional integral associated to the Atangana-Baleanu model given by (2.4). The application of this integral on (5.2), leads to the following non-linear voltera integral equation:

$$u(t) = u_0 + \frac{(1-\alpha)}{ABC(\alpha)}G(t, u(t)) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G(\varepsilon, u(\varepsilon)) d\varepsilon. \quad (5.5)$$

We assure that $J = (0, T)$ and take into consideration the operator $\phi : C(J, \mathbb{R}^6) \rightarrow C(J, \mathbb{R}^6)$ defined by

$$\phi[u(t)] = u_0 + \frac{(1-\alpha)}{ABC(\alpha)}G(t, u(t)) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G(\varepsilon, u(\varepsilon)) d\varepsilon. \quad (5.6)$$

Equation (5.5) turns to

$$u(t) = \phi[u(t)]. \quad (5.7)$$

The supremum norm on J , $\|\cdot\|_J$ is $\|u(t)\|_J = \sup_{t \in J} \|u(t)\|$, $u(t) \in C$.

Obviously, $C(J, \mathbb{R}^6)$ along the norm $\|\cdot\|_J$ present a Banach space. Additionally, we can demonstrate simply the following inequality

$$\left\| \int_0^t D(t, \varepsilon) u(\varepsilon) d\varepsilon \right\| \leq T \|D(t, \varepsilon)\|_J \|u(t)\|_J. \quad (5.8)$$

with $u(t) \in C(J, \mathbb{R}^6)$, $D(t, \varepsilon) \in C(J^2, \mathbb{R})$.

In a way that

$$\|D(t, \varepsilon)\|_J = \sup_{t, \varepsilon \in J} |D(t, \varepsilon)|.$$

Applying the definition of ϕ stated in (5.7), we derive

$$\begin{aligned} \|\phi[u_1(t)] - \phi[u_2(t)]\|_J &\leq \left\| \frac{(1-\alpha)}{ABC(\alpha)}G(t, u_1(t)) - \frac{(1-\alpha)}{ABC(\alpha)}G(t, u_2(t)) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \right. \\ &\quad \left. \times \int_0^t (t-\varepsilon)^{\alpha-1} (G(\varepsilon, u_1(\varepsilon)) - G(\varepsilon, u_2(\varepsilon))) d\varepsilon \right\|. \end{aligned}$$

Furthermore, using the principles of triangular inequality and Lipschitz condition stated in (5.3) couple with the results in (5.8) we get the following after some algebraic simplification.

$$\|\phi[u_1(t)] - \phi[u_2(t)]\|_J \leq \left(\frac{(1-\alpha)}{ABC(\alpha)}M + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)}MT_{\max}^{\alpha} \right) \|u_1(t) - u_2(t)\|_J.$$

Thus, we eventually have

$$\|\phi[u_1(t)] - \phi[u_2(t)]\|_J \leq \beta \|u_1(t) - u_2(t)\|_J,$$

where

$$\beta = \frac{(1-\alpha)}{ABC(\alpha)}M + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)}MT_{\max}^{\alpha}.$$

Therefore, the operator ϕ will become a contraction if condition (5.4) holds on $C(J, \mathbb{R}^6)$. As a result of Banach fixed point theorem, system (5.2) possesses a unique solution. \square

5.2. Numerical scheme for the Atangana - Baleanu model

This section derive the numerical scheme of the TB model in the ABC form which given by Eq. (5.1) using the method in [52] by using Adams-Bashforth method. By using the same procedure in [52], we write (5.1) to the following

$$\begin{aligned} {}_0^{ABC}D_t^{\alpha}S_C &= G_1(t, S_C, L_C, I_C, S_A, L_A, I_A) \\ {}_0^{ABC}D_t^{\alpha}L_C &= G_2(t, S_C, L_C, I_C, S_A, L_A, I_A), \\ {}_0^{ABC}D_t^{\alpha}I_C &= G_3(t, S_C, L_C, I_C, S_A, L_A, I_A), \\ {}_0^{ABC}D_t^{\alpha}S_A &= G_4(t, S_C, L_C, I_C, S_A, L_A, I_A), \\ {}_0^{ABC}D_t^{\alpha}L_A &= G_5(t, S_C, L_C, I_C, S_A, L_A, I_A), \\ {}_0^{ABC}D_t^{\alpha}I_A &= G_6(t, S_C, L_C, I_C, S_A, L_A, I_A). \end{aligned} \quad (5.9)$$

By utilizing the fundamental theorem of fractional calculus, we convert (5.9) to fractional

$$\begin{aligned} S_C(t) - S_C(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_1(t, S_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G_1(\varepsilon, S_C) d\varepsilon, \\ L_C(t) - L_C(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_2(t, L_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G_2(\varepsilon, L_C) d\varepsilon, \\ I_C(t) - I_C(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_3(t, I_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G_3(\varepsilon, I_C) d\varepsilon, \\ S_A(t) - S_A(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_4(t, S_A) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G_4(\varepsilon, S_A) d\varepsilon, \\ L_A(t) - L_A(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_5(t, L_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G_5(\varepsilon, L_A) d\varepsilon, \\ I_A(t) - I_A(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_6(t, I_A) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \int_0^t (t-\varepsilon)^{\alpha-1} G_6(\varepsilon, I_A) d\varepsilon. \end{aligned} \quad (5.10)$$

At $t = t_{n+1}$, $n = 0, 1, 2, \dots$, we have

$$\begin{aligned} S_C(t_{n+1}) - S_C(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_1(t_n, S_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} (t_{n+1} - \varepsilon)^{\alpha-1} G_1(\varepsilon, S_C) d\varepsilon, \\ L_C(t_{n+1}) - L_C(0) &= \frac{(1-\alpha)}{ABC(\alpha)}G_2(t_n, L_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} (t_{n+1} - \varepsilon)^{\alpha-1} G_2(\varepsilon, L_C) d\varepsilon, \end{aligned}$$

$$\begin{aligned}
I_C(t_{n+1}) - I_C(0) &= \frac{(1-\alpha)}{ABC(\alpha)} G_3(t_n, I_C) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} (t_{n+1} - \varepsilon)^{\alpha-1} G_3(\varepsilon, I_C) d\varepsilon, \quad (5.11) \\
S_A(t_{n+1}) - S_A(0) &= \frac{(1-\alpha)}{ABC(\alpha)} G_4(t_n, S_A) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} (t_{n+1} - \varepsilon)^{\alpha-1} G_4(\varepsilon, S_A) d\varepsilon, \\
L_A(t_{n+1}) - L_A(0) &= \frac{(1-\alpha)}{ABC(\alpha)} G_5(t_n, L_A) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} (t_{n+1} - \varepsilon)^{\alpha-1} G_5(\varepsilon, L_A) d\varepsilon, \\
I_A(t_{n+1}) - I_A(0) &= \frac{(1-\alpha)}{ABC(\alpha)} G_6(t_n, I_A) + \frac{\alpha}{ABC(\alpha)\Gamma(\alpha)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} (t_{n+1} - \varepsilon)^{\alpha-1} G_6(\varepsilon, I_A) d\varepsilon.
\end{aligned}$$

The integral in Eq. (5.11) is approximated through the two-point interpolation polynomial. Hence, we have the iterative scheme for the TB model (5.1). After some calculations, we finally obtain the approximation solution as

$$\begin{aligned}
S_C(t_{n+1}) &= S_C(0) + \frac{(1-\alpha)}{ABC(\alpha)} G_1(t_n, S_C) \\
&+ \frac{\alpha}{ABC(\alpha)} \sum_{j=0}^n \left(\frac{h^\alpha G_1(t_j, S_C)}{\Gamma(\alpha+2)} ((n+1-j)^\alpha (n-j+2+\alpha) \right. \\
&\left. - (n-j)^\alpha (n-j+2+2\alpha)) - \frac{h^\alpha G_1(t_{j-1}, S_C)}{\Gamma(\alpha+2)} ((n+1-j)^{\alpha+1} - (n-j)^\alpha (n-j+1+\alpha)) \right),
\end{aligned}$$

$$\begin{aligned}
L_C(t_{n+1}) &= L_C(0) + \frac{(1-\alpha)}{ABC(\alpha)} G_2(t_n, L_C) \\
&+ \frac{\alpha}{ABC(\alpha)} \sum_{j=0}^n \left(\frac{h^\alpha G_2(t_j, L_C)}{\Gamma(\alpha+2)} ((n+1-j)^\alpha (n-j+2+\alpha) \right. \\
&\left. - (n-j)^\alpha (n-j+2+2\alpha)) - \frac{h^\alpha G_2(t_{j-1}, L_C)}{\Gamma(\alpha+2)} ((n+1-j)^{\alpha+1} - (n-j)^\alpha (n-j+1+\alpha)) \right),
\end{aligned}$$

$$\begin{aligned}
I_C(t_{n+1}) &= I_C(0) + \frac{(1-\alpha)}{ABC(\alpha)} G_3(t_n, I_C) \\
&+ \frac{\alpha}{ABC(\alpha)} \sum_{j=0}^n \left(\frac{h^\alpha G_3(t_j, I_C)}{\Gamma(\alpha+2)} ((n+1-j)^\alpha (n-j+2+\alpha) \right. \\
&\left. - (n-j)^\alpha (n-j+2+2\alpha)) - \frac{h^\alpha G_3(t_{j-1}, I_C)}{\Gamma(\alpha+2)} ((n+1-j)^{\alpha+1} - (n-j)^\alpha (n-j+1+\alpha)) \right),
\end{aligned}$$

$$\begin{aligned}
S_A(t_{n+1}) &= S_A(0) + \frac{(1-\alpha)}{ABC(\alpha)} G_4(t_n, S_A) \\
&+ \frac{\alpha}{ABC(\alpha)} \sum_{j=0}^n \left(\frac{h^\alpha G_4(t_j, S_A)}{\Gamma(\alpha+2)} ((n+1-j)^\alpha (n-j+2+\alpha) \right.
\end{aligned}$$

$$\begin{aligned}
& - (n-j)^\alpha (n-j+2+2\alpha) - \frac{h^\alpha G_4(t_{j-1}, S_A)}{\Gamma(\alpha+2)} ((n+1-j)^{\alpha+1} - (n-j)^\alpha (n-j+1+\alpha)), \\
L_A(t_{n+1}) &= L_A(0) + \frac{(1-\alpha)}{ABC(\alpha)} G_5(t_n, L_A) \\
& + \frac{\alpha}{ABC(\alpha)} \sum_{j=0}^n \left(\frac{h^\alpha G_5(t_j, L_A)}{\Gamma(\alpha+2)} ((n+1-j)^\alpha (n-j+2+\alpha) \right. \\
& \left. - (n-j)^\alpha (n-j+2+2\alpha) - \frac{h^\alpha G_5(t_{j-1}, L_A)}{\Gamma(\alpha+2)} ((n+1-j)^{\alpha+1} - (n-j)^\alpha (n-j+1+\alpha)) \right), \\
I_A(t_{n+1}) &= I_A(0) + \frac{(1-\alpha)}{ABC(\alpha)} G_6(t_n, I_A) \\
& + \frac{\alpha}{ABC(\alpha)} \sum_{j=0}^n \left(\frac{h^\alpha G_6(t_j, I_A)}{\Gamma(\alpha+2)} ((n+1-j)^\alpha (n-j+2+\alpha) \right. \\
& \left. - (n-j)^\alpha (n-j+2+2\alpha) - \frac{h^\alpha G_6(t_{j-1}, I_A)}{\Gamma(\alpha+2)} ((n+1-j)^{\alpha+1} - (n-j)^\alpha (n-j+1+\alpha)) \right).
\end{aligned}$$

Next, we present the simulation result of the TB model (5.1) using the numerical scheme.

5.3. Numerical simulation of Atangana-Baleanu model

This section provides the simulation results of Atangana-Baleanu model (5.1) using the iterative scheme obtained in (5.12). The parameter values listed in Table 2 are for obtaining the numerical results. The simulation results are shown in Figure 5 by considering different values of $\alpha = 1, 0.9, 0.8, 0.7$ and 0.6 . In Figure 5, we can see that by decreasing the values of α , the population of infected compartments decreases, while the susceptible population increases.

In Figures 6 and 7, we present the effect of the efficacy of chemoprophylaxis (r_C and r_A) and treatment (γ_C, γ_A) on the TB transmission for different values of α . From Figure 6, it can be seen that as the chemoprophylaxis strategy increases, the infected TB in both children and adults population decreases. From Figure 7, it is shown that as the treatment strategy increases, the infected TB in both children and adults population decreases significantly.




Figure 5. Numerical results for ABC model.




Figure 6. Numerical results of infected TB with various r_C and r_A , (a–b) for $\alpha = 0.9$; (c–d) for $\alpha = 0.7$.




Figure 7. Numerical results of infected TB with various γ_C and γ_A , (a–b) for $\alpha = 0.9$; (c–d) for $\alpha = 0.7$.

6. Comparison of operators

This present section demonstrates the comparison of the Caputo and Atangana-Baleanu operators graphically. The parameter values used in these simulations are given in Table 2. In order to have a comparison of these two operators we considered many values of $\alpha = 1, 0.7, 0.5, 0.3$ and the results were displayed Figures 8–11. It can be seen in Figures 8–11 that there exists a significant difference in the dynamical behavior of the TB infected population when the values of α decrease within both fractional operators. From these figures, we see that Atangana-Baleanu derivative give faster decreases in the results compare to the Caputo derivative when the fractional order parameter decrease for the reduction of TB infection. This comparison section demonstrate the effectiveness of the Atangana-Baleanu derivative over the Caputo derivative.




Figure 8. Comparison results of Caputo and ABC derivative with $\alpha = 1$.




Figure 9. Comparison results of Caputo and ABC derivative with $\alpha = 0.7$.




Figure 10. Comparison results of Caputo and ABC derivative with $\alpha = 0.5$.




Figure 11. Comparison results of Caputo and ABC derivative with $\alpha = 0.3$.

7. Conclusion

The findings reported here shed new light on the fractional-order model with Caputo and Atangana-Baleanu derivative to investigate the spread of TB with children and adults population. Brief mathematical results for the TB model are investigated. The basic reproduction number of the Caputo model and its equilibrium are obtained and presented the results. Each model is studied independently in the fractional operators of Caputo and Atangana-Baleanu. We then performed the numerical simulations of the Caputo model type for different values of the fractional-order using the predictor-corrector method. Then, we apply the fractional Atangana-Baleanu operator to the model and analyzed and obtained the associated results. We proved the uniqueness and existence of the Atangana-Baleanu model. The numerical simulation of the Atangana-Baleanu model was conducted using the Adams–Bashforth scheme. We further presented the graphical results of Atangana-Baleanu model for various fractional order parameters. A comparison of the proposed operators graphically presented. These graphical results for comparison show that the Atangana-Baleanu results are more appropriate for the better decrease in infection while the Caputo is less. We conclude from graphical results that increasing chemoprophylaxis and treatment rates then the TB infective both children and adult cases can be reduced. The graphical results show that a decrease in fractional order parameter causes a significant decrease in the infective TB population. Therefore, the present work is a novel analysis on the TB dynamics and will be more useful for the readers and public health authorities. In future work the TB age fractional model can be considered in the light of fractional impulse model and the results can be compared with non-impulse TB model, stated in [53–55]. This is important because the impulse effect is attracting the receiving considerable attention and eliciting widespread interest in epidemiological models.

Acknowledgements

Part of this research is financially supported by Universitas Airlangga 2018.

Conflict of interest

All authors declare no conflict of interest.

References

1. World Health Organization, *Anti-tuberculosis Drug Resistance in the World: Third Global Report*, No. WHO / HTM / TB / 2004. 343, World Health Organization, 2004.
2. J. Cohen, *Extensively drug-resistant TB gets foothold in South Africa*, *Science* **313** (2006), 1554.
3. L. B. Reichman, J. H. Tanne, *Timebomb: The Global Epidemic of Multi-Drug Resistant Tuberculosis*, 2002.
4. Y. Zhou, K. Khan, Z. Feng, et al. *Projection of tuberculosis incidence with increasing immigration trends*, *J. Theor. Biol.*, **254** (2008), 215–228.
5. P. Rodrigues, M. G. M. Gomes, C. Rebelo, *Drug resistance in tuberculosis: a reinfection model*, *Theor. Popul. Biol.*, **71** (2007), 196–212.

6. N. Blaser, C. Zahnd, S. Hermans, et al. *Tuberculosis in Cape Town: an age-structured transmission model*, *Epidemics*, **14** (2016), 54–61.
7. C. P. Bhunu, W. Garira, Z. Mukandavire, et al. *Tuberculosis transmission model with chemoprophylaxis and treatment*, *B. Math. Biol.*, **70** (2008), 1163–1191.
8. Centers for Disease Control and Prevention, *TB in Children in the United States*, CDC, 2014. Available from:
<https://www.cdc.gov/tb/topic/populations/tbinchildren/default.htm>.
9. S. M. Blower, P. M. Small, P. C. Hopewell, *Control strategies for tuberculosis epidemics: new models for old problems*, *Science*, **273** (1996), 497–500.
10. Fatmawati, H. Tasman, *An optimal treatment control of TB-HIV coinfection*, *International Journal of Mathematics and Mathematical Sciences*, **2016** (2016).
11. R. I. Hickson, G. N. Mercer, K. M. Lokuge, *A metapopulation model of tuberculosis transmission with a case study from high to low burden areas*, *PLoS One*, **7** (2012).
12. R. M. G. J. Houben, T. Sumner, A. D. Grant, et al. *Ability of preventive therapy to cure latent Mycobacterium tuberculosis infection in HIV-infected individuals in high-burden settings*, *P. Natl. A. Sci.*, **111** (2014), 5325–5330.
13. R. Kaplan, J. Caldwell, K. Middelkoop, et al. *Impact of ART on TB case fatality stratified by CD4 count for HIV-positive TB patients in Cape Town, South Africa (2009-2011)*, *J. Acq. Imm. Def.*, **66** (2014), 487–494.
14. Fatmawati, U. D. Purwati, F. Riyudha, et al. *Optimal control of a discrete age-structured model for tuberculosis transmission*, *Heliyon*, **6** (2020).
15. S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach Science Publishers, 1993.
16. I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, California, USA, 1999.
17. T. Sardar, S. Rana, J. Chattopadhyay, *A mathematical model of dengue transmission with memory*, *Commun. Nonlinear Sci.*, **22** (2015), 511–525.
18. J. Huo, H. Zhao, L. Zhu, *The effect on backward bifurcation in a fractional order HIV model*, *Nonlinear Analysis : Real World Applications*, **26** (2015), 289–305.
19. M. Saeedian, M. Khalighi, N. Azimi-Tafreshi, et al. *Memory effects on epidemic evolution: the susceptible-infected-recovered epidemic model*, *Phys. Rev. E.*, **95** (2017).
20. C. M. A. Pinto, A. R. M. Carvalho, *The HIV/TB coinfection severity in the presence of TB multi-drug resistant strains*, *Ecol. Complex.*, **32** (2017), 1–20.
21. Fatmawati, E. M. Shaiful, M. I. Utoyo, *A fractional order model for HIV dynamics in a two-sex population*, *International Journal of Mathematics and Mathematical Sciences*, **2018** (2018).
22. G. C. Wu, Z. G. Deng, D. Baleanu, et al. *New variable-order fractional chaotic systems for fast image encryption*, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, **29** (2019).
23. G. C. Wu, T. Abdeljawad, J. Liu, et al. *Mittag-Leffler stability analysis of fractional discrete-time neural networks via fixed point technique*, *Nonlinear Analysis: Modelling and Control*, **24** (2019), 919–936.

24. G. C. Wu, D. Q. Zeng, D. Baleanu, *Fractional impulsive differential equations: Exact solutions, integral equations and short memory case*, *Frac. Calc. Appl. Anal.*, **22** (2019), 180–192.
25. M. Itik, S. P. Banks, *Chaos in a three-dimensional cancer model*, *Int. J. Bifurcat. Chaos*, **20** (2010), 71–79.
26. M. Caputo, M. Fabrizio, *A new definition of fractional derivative without singular kernel*, *Progr. Fract. Differ. Appl.*, **1** (2015), 1–13.
27. M. Caputo, M. Fabrizio, *On the notion of fractional derivative and applications to the hysteresis phenomena*, *Meccanica*, **52** (2017), 3043–3052.
28. T. Zhang, L. Xiong, *Periodic motion for impulsive fractional functional differential equations with piecewise Caputo derivative*, *Appl. Math. Lett.*, **101** (2020), 106072.
29. A. Atangana, D. Baleanu, *New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model*, *Therm. Sci.*, **20** (2016).
30. A. Atangana, I. Koca, *Chaos in a simple nonlinear system with Atangana-Baleanu derivatives with fractional order*, *Chaos, Solitons & Fractals*, **89** (2016), 447–454.
31. A. Atangana, K. M. Owolabi, *New numerical approach for fractional differential equations*, *Math. Model. Nat. Pheno.*, **13** (2018).
32. A. Atangana, J. F. Gomez-Aguilar, *Decolonisation of fractional calculus rules: Breaking commutativity and associativity to capture more natural phenomena*, *The European Physical Journal Plus*, **133** (2018).
33. M. A. Khan, S. Ullah, M. Farooq, *A new fractional model for tuberculosis with relapse via Atangana-Baleanu derivative*, *Chaos, Solitons & Fractals*, **116** (2018), 227–238.
34. E. Bonyah, *Chaos in a 5-D hyperchaotic system with four wings in the light of non-local and non-singular fractional derivatives*, *Chaos, Solitons & Fractals*, **116** (2018), 316–331.
35. K. Muhammad Altaf, A. Atangana, *Dynamics of Ebola disease in the framework of different fractional derivatives*, *Entropy*, **21** (2019).
36. R. Jan, M. A. Khan, P. Kumam, et al, *Modeling the transmission of dengue infection through fractional derivatives*, *Chaos, Solitons & Fractals*, **127** (2019), 189–216.
37. W. Wang, M. A. Khan, P. Kumam, et al. *A comparison study of bank data in fractional calculus*, *Chaos, Solitons & Fractals*, **126** (2019), 369–384.
38. M. A. Khan, *The dynamics of a new chaotic system through the Caputo-Fabrizio and Atangana-Baleanu fractional operators*, *Adv. Mech. Eng.*, **11** (2019).
39. M. A. Khan, M. Azizah, S. Ullah, *A fractional model for the dynamics of competition between commercial and rural banks in Indonesia*, *Chaos, Solitons & Fractals*, **122** (2019), 32–46.
40. S. Ullah, M. A. Khan, M. Farooq, et al. *A fractional model for the dynamics of tuberculosis (TB) using Atangana-Baleanu derivative*, *Discrete Cont. Dyn. S*, **13** (2019).
41. A. A. Velayati, *Tuberculosis in children*, *International Journal of Mycobacteriology*, **5** (2016).
42. C. Castillo-Chaves, B. Song, *Dynamic models of tuberculosis and their applications*, *Math. Biosci. Eng.*, **1** (2004), 361–404.

43. Z. M. Odibat, N. T. Shawagfeh, *Generalized Taylors formula*, Appl. Math. Comput., **186** (2007), 286–293.
44. W. Lin, *Global existence theory and chaos control of fractional differential equations*, J. Math. Anal. Appl., **332** (2007), 709–726.
45. P. van den Driessche, J. Watmough, *Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission*, Math. Biosci., **180** (2002), 29–48.
46. K. Diethelm, N. J. Ford, A. D. Freed, *A predictor-corrector approach for the numerical solution of fractional differential equations*, Nonlinear Dynamics, **29** (2002), 3–22.
47. K. Diethelm, N. J. Ford, A. D. Freed, *Detailed error analysis for a fractional Adams method*, Numerical Algorithms, **36** (2004), 31–52.
48. C. P. Bhunu, *Mathematical analysis of a three-strain tuberculosis transmission model*, Appl. Math. Model., **35** (2011), 4647–4660.
49. S. Athithan, M. Ghosh, *Optimal control of tuberculosis with case detection and treatment*, World Journal of Modelling and Simulation, **11** (2015), 111–122.
50. World Health Organization, *Factsheet on the World Tuberculosis Report 2017*, WHO, 2017. Available from:
<https://www.who.int/en/news-room/fact-sheets/detail/tuberculosis>.
51. J. J. Tewa, S. Bowong, B. Mewoli, *Mathematical analysis of two-patch model for the dynamical transmission of tuberculosis*, Appl. Math. Model., **36** (2012), 2466–2485.
52. M. Toufik, A. Atangana, *New numerical approximation of fractional derivative with non-local and non-singular kernel: application to chaotic models*, The European Physical Journal Plus, **132** (2017), 444.
53. L. Xu, H. Hub, F. Qinc, *Ultimate boundedness of impulsive fractional differential equations*, Appl. Math. Lett., **62** (2016), 110–117.
54. L. Xu, J. Li, S. S. Ge, *Impuls ivestabilization of fractional differential systems*, ISA T., **70** (2017), 125–131.
55. L. Xu, X. Chu, H. Hu, *Exponential ultimate boundedness of non-autonomous fractional differential systems with time delay and impulses*, Appl. Math. Lett., **99** (2020), 106000.



AIMS Press

© 2020 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)

A mathematical model of tuberculosis (TB) transmission with children and adults groups: A fractional model

ORIGINALITY REPORT

19%

SIMILARITY INDEX

9%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Khan Muhammad Altaf, Abdon Atangana. "Dynamics of Ebola Disease in the Framework of Different Fractional Derivatives", Entropy, 2019
Publication 1%
- 2** Abdon Atangana, Kolade M. Owolabi. "New numerical approach for fractional differential equations", Mathematical Modelling of Natural Phenomena, 2018
Publication 1%
- 3** aip.scitation.org
Internet Source 1%
- 4** Wanting Wang, Muhammad Altaf Khan, Fatmawati, P. Kumam, P. Thounthong. "A comparison study of bank data in fractional calculus", Chaos, Solitons & Fractals, 2019
Publication 1%
- 5** www.iaeng.org
Internet Source 1%

-
- 6 "Fractional Derivatives with Mittag-Leffler Kernel", Springer Science and Business Media LLC, 2019
Publication 1 %
-
- 7 Fatmawati, Utami Dyah Purwati, Firman Riyudha, Hengki Tasman. "Optimal control of a discrete age-structured model for tuberculosis transmission", Heliyon, 2020
Publication 1 %
-
- 8 Ebraheem O. Alzahrani, M. A. Khan. "Comparison of numerical techniques for the solution of a fractional epidemic model", The European Physical Journal Plus, 2020
Publication 1 %
-
- 9 Carla M.A. Pinto, Ana R.M. Carvalho. "The HIV/TB coinfection severity in the presence of TB multi-drug resistant strains", Ecological Complexity, 2017
Publication <1 %
-
- 10 advancesindifferenceequations.springeropen.com
Internet Source <1 %
-
- 11 Fatmawati, Muhammad Altaf Khan, Muftiyatul Azizah, Windarto, Saif Ullah. "A fractional model for the dynamics of competition between commercial and rural banks in Indonesia", Chaos, Solitons & Fractals, 2019
Publication <1 %
-

12 Saif Ullah, Muhammad Altaf Khan, Muhammad Farooq. "A fractional model for the dynamics of TB virus", Chaos, Solitons & Fractals, 2018
Publication <1 %

13 Muhammad Altaf Khan. "The dynamics of a new chaotic system through the Caputo-Fabrizio and Atanagana-Baleanu fractional operators", Advances in Mechanical Engineering, 2019
Publication <1 %

14 "Mathematical Modelling in Health, Social and Applied Sciences", Springer Science and Business Media LLC, 2020
Publication <1 %

15 Rashid Jan, Muhammad Altaf Khan, Poom Kumam, Phatiphat Thounthong. "Modeling the transmission of dengue infection through fractional derivatives", Chaos, Solitons & Fractals, 2019
Publication <1 %

16 www.tandfonline.com
Internet Source <1 %

17 Bowong, S.. "Mathematical analysis of a tuberculosis model with differential infectivity", Communications in Nonlinear Science and Numerical Simulation, 200911
Publication <1 %

18 Eve McDonald-Madden. "Optimal timing for managed relocation of species faced with climate change", Nature Climate Change, 07/24/2011

Publication

<1 %

19 Muhammad Altaf Khan, Olusola Kolebaje, Ahmet Yildirim, Saif Ullah, P. Kumam, P. Thounthong. "Fractional investigations of zoonotic visceral leishmaniasis disease with singular and non-singular kernel", The European Physical Journal Plus, 2019

Publication

<1 %

20 Bowong, S.. "Global analysis of a dynamical model for transmission of tuberculosis with a general contact rate", Communications in Nonlinear Science and Numerical Simulation, 201011

Publication

<1 %

21 JATI K. SENGUPTA. "Game theory models applied to economic systems: their information, structure and stability", International Journal of Systems Science, 2007

Publication

<1 %

22 aimc46.yazd.ac.ir

Internet Source

<1 %

23 chesterrep.openrepository.com

Internet Source

<1 %

24	litis.univ-lehavre.fr Internet Source	<1 %
25	oaji.net Internet Source	<1 %
26	tel.archives-ouvertes.fr Internet Source	<1 %
27	www.m-hikari.com Internet Source	<1 %
28	Tanvi, Rajiv Aggarwal. "Dynamics of HIV-TB co-infection with detection as optimal intervention strategy", International Journal of Non-Linear Mechanics, 2020 Publication	<1 %
29	E.O. Alzahrani, M.A. Khan. "Modeling the dynamics of Hepatitis E with optimal control", Chaos, Solitons & Fractals, 2018 Publication	<1 %
30	Fathalla A. Rihan, Dumitru Baleanu, S. Lakshmanan, R. Rakkiyappan. " On Fractional SIRC Model with Bacterial Infection ", Abstract and Applied Analysis, 2014 Publication	<1 %
31	R. U. Diva Amalia, Fatmawati, Windarto, Didik Khusnul Arif. "Optimal control of predator-prey mathematical model with infection and	<1 %

harvesting on prey", Journal of Physics:
Conference Series, 2018

Publication

32

Seda İğret Araz. "A fractional optimal control problem with final observation governed by wave equation", Chaos: An Interdisciplinary Journal of Nonlinear Science, 2019

Publication

<1 %

33

Muhammad Altaf Khan, Francisco Gómez - Aguilar. "Tuberculosis model with relapse via fractional conformable derivative with power law", Mathematical Methods in the Applied Sciences, 2019

Publication

<1 %

34

Muhammad Altaf Khan, Saif Ullah, Muhammad Farooq. "A new fractional model for tuberculosis with relapse via Atangana-Baleanu derivative", Chaos, Solitons & Fractals, 2018

Publication

<1 %

35

Abdon Atangana, J.F. Gómez-Aguilar. "Fractional derivatives with no-index law property: Application to chaos and statistics", Chaos, Solitons & Fractals, 2018

Publication

<1 %

36

Coronel-Escamilla, A., J.F. Gómez-Aguilar, M.G. López-López, V.M. Alvarado-Martínez, and G.V. Guerrero-Ramírez.

<1 %

"Triple pendulum model involving fractional derivatives with different kernels", Chaos Solitons & Fractals, 2016.

Publication

37

Abdon Atangana, Sania Qureshi. "Modeling attractors of chaotic dynamical systems with fractal-fractional operators", Chaos, Solitons & Fractals, 2019

Publication

<1 %

38

Aliyu Isa Aliyu, Ali Saleh Alshomrani, Yongjin Li, Mustafa Inc, Dumitru Baleanu. "Existence theory and numerical simulation of HIV-I cure model with new fractional derivative possessing a non-singular kernel", Advances in Difference Equations, 2019

Publication

<1 %

39

ascelibrary.org

Internet Source

<1 %

40

www.mdpi.com

Internet Source

<1 %

41

Mehmet Yavuz, Ebenezer Bonyah. "New approaches to the fractional dynamics of schistosomiasis disease model", Physica A: Statistical Mechanics and its Applications, 2019

Publication

<1 %

42 Abdon Atangana, Sonal Jain. "The role of power decay, exponential decay and Mittag-Leffler function's waiting time distribution: Application of cancer spread", Physica A: Statistical Mechanics and its Applications, 2018
Publication

43 D. P. Ahokposi, Abdon Atangana, D. P. Vermeulen. "Modelling groundwater fractal flow with fractional differentiation via Mittag-Leffler law", The European Physical Journal Plus, 2017
Publication

44 Junli Liu, Zhenguo Bai, Tailei Zhang. "A periodic two-patch SIS model with time delay and transport-related infection", Journal of Theoretical Biology, 2018
Publication

45 M. Hassouna, A. Ouhadan, E.H. El Kinani. "On the solution of fractional order SIS epidemic model", Chaos, Solitons & Fractals, 2018
Publication

46 jpsr.pharmainfo.in
Internet Source

47 "Mathematics Applied to Engineering, Modelling, and Social Issues", Springer Science and Business Media LLC, 2019
Publication

48

Aliyu Isa Aliyu, Mustafa Inc, Abdullahi Yusuf, Dumitru Baleanu. "A fractional model of vertical transmission and cure of vector-borne diseases pertaining to the Atangana–Baleanu fractional derivatives", *Chaos, Solitons & Fractals*, 2018

Publication

<1 %

49

Dhage, B.C.. "Basic results on hybrid differential equations", *Nonlinear Analysis: Hybrid Systems*, 201008

Publication

<1 %

50

Ebenezer Bonyah. "Chaos in a 5-D hyperchaotic system with four wings in the light of non-local and non-singular fractional derivatives", *Chaos, Solitons & Fractals*, 2018

Publication

<1 %

51

Khaled M. Saad. "New fractional derivative with non-singular kernel for deriving Legendre spectral collocation method", *Alexandria Engineering Journal*, 2019

Publication

<1 %

52

McCluskey, C.C.. "Global stability for a class of mass action systems allowing for latency in tuberculosis", *Journal of Mathematical Analysis and Applications*, 20080201

Publication

<1 %

53

Sania Qureshi, Ebenezer Bonyah, Asif Ali Shaikh. "Classical and contemporary fractional operators for modeling diarrhea transmission dynamics under real statistical data", *Physica A: Statistical Mechanics and its Applications*, 2019

Publication

<1 %

54

citeseerx.ist.psu.edu

Internet Source

<1 %

55

internonlinearscience.org

Internet Source

<1 %

56

www.emis.de

Internet Source

<1 %

57

www.waset.org

Internet Source

<1 %

58

"HIV and Tuberculosis", Springer Science and Business Media LLC, 2019

Publication

<1 %

59

Arenas, Abraham J., Gilberto González-Parra, and Benito M. Chen-Charpentier. "Construction of nonstandard finite difference schemes for the SI and SIR epidemic models of fractional order", *Mathematics and Computers in Simulation*, 2016.

Publication

<1 %

60

Atangana, Abdon, and Ilknur Koca. "Chaos in a simple nonlinear system with Atangana-Baleanu derivatives with fractional order", *Chaos Solitons & Fractals*, 2016.

Publication

<1 %

61

E. Bonyah, A. Atangana, Mehar Chand. "Analysis of 3D IS-LM macroeconomic system model within the scope of fractional calculus", *Chaos, Solitons & Fractals: X*, 2019

Publication

<1 %

62

George Z. Voyiadjis, Wojciech Sumelka. "Brain modelling in the framework of anisotropic hyperelasticity with time fractional damage evolution governed by the Caputo-Almeida fractional derivative", *Journal of the Mechanical Behavior of Biomedical Materials*, 2019

Publication

<1 %

63

Yang, Ya Li, Jian Feng Cheng, Fang Chen, and Ya Yi Xu. "Optimal Control of a Tuberculosis Model with Chemoprophylaxis Treatment", *Advanced Materials Research*, 2013.

Publication

<1 %

64

epdf.tips
Internet Source

<1 %

65

ethesis.nitrkl.ac.in
Internet Source

<1 %

- | | | |
|----|--|------|
| 66 | repository.tudelft.nl
Internet Source | <1 % |
| 67 | repository.unair.ac.id
Internet Source | <1 % |
| 68 | www-users.math.umd.edu
Internet Source | <1 % |
| 69 | A. Al-khedhairi, A.A. Elsadany, A. Elsonbaty. "Modelling immune systems based on Atangana–Baleanu fractional derivative", Chaos, Solitons & Fractals, 2019
Publication | <1 % |
| 70 | Abdon Atangana. "Fractal-fractional differentiation and integration: Connecting fractal calculus and fractional calculus to predict complex system", Chaos, Solitons & Fractals, 2017
Publication | <1 % |
| 71 | Ana R.M. Carvalho, Carla M.A. Pinto. "Non-integer order analysis of the impact of diabetes and resistant strains in a model for TB infection", Communications in Nonlinear Science and Numerical Simulation, 2018
Publication | <1 % |
| 72 | Badr Saad T. Alkahtani. "Numerical analysis of dissipative system with noise model with the Atangana–Baleanu fractional derivative", Chaos, Solitons & Fractals, 2018 | <1 % |

73

Chuanqing Xu, Xiaoxiao Wei, Jingan Cui, Xiaojing Wang, Dashun Xu. "Mixing in regional-structure model about the influence of floating population and optimal control about TB in Guangdong province of China", *International Journal of Biomathematics*, 2017

Publication

<1 %

74

Danhua He, Yin Qing. " Boundedness theorems for non-autonomous stochastic delay differential systems driven by -Brownian motion ", *Applied Mathematics Letters*, 2019

Publication

<1 %

75

Dumitru Baleanu, Arran Fernandez. "On some new properties of fractional derivatives with Mittag-Leffler kernel", *Communications in Nonlinear Science and Numerical Simulation*, 2018

Publication

<1 %

76

Hashemi, M.S., and D. Baleanu. "Numerical approximation of higher-order time-fractional telegraph equation by using a combination of a geometric approach and method of line", *Journal of Computational Physics*, 2016.

Publication

<1 %

77

L.F. Ávalos-Ruiz, J.F. Gómez-Aguilar, A. Atangana, Kolade M. Owolabi. "On the dynamics of fractional maps with power-law,

<1 %

exponential decay and Mittag–Leffler memory", Chaos, Solitons & Fractals, 2019

Publication

78 Leonardo Martínez Jiménez, J. Juan Rosales García, Abraham Ortega Contreras, Dumitru Baleanu. "Analysis of Drude model using fractional derivatives without singular kernels", Open Physics, 2017

Publication

79 Rajnesh Kumar, Sunil Kumar. "A New Fractional Modelling on Susceptible-Infected-Recovered Equations with Constant Vaccination Rate", Nonlinear Engineering, 2014

Publication

80 Syed Azhar Ali Shah, Muhammad Altaf Khan, Muhammad Farooq, Saif Ullah, Ebraheem O. Alzahrani. "A fractional order model for Hepatitis B virus with treatment via Atangana–Baleanu derivative", Physica A: Statistical Mechanics and its Applications, 2020

Publication

81 [aims.org](https://www.aims.org)
Internet Source

82 arxiv.org
Internet Source

83	italy.usembassy.gov Internet Source	<1 %
84	journals.plos.org Internet Source	<1 %
85	msl.mit.edu Internet Source	<1 %
86	www.aims sciences.org Internet Source	<1 %
87	www.nature.com Internet Source	<1 %
88	"Mathematical Modelling, Applied Analysis and Computation", Springer Science and Business Media LLC, 2019 Publication	<1 %
89	C. Chiyaka, W. Garira, S. Dube. "Modelling Immune Response and Drug Therapy in Human Malaria Infection", Computational and Mathematical Methods in Medicine, 2008 Publication	<1 %
90	Hailay Weldegiorgis Berhe, Sania Qureshi, Asif Ali Shaikh. "Deterministic modeling of dysentery diarrhea epidemic under fractional Caputo differential operator via real statistical analysis", Chaos, Solitons & Fractals, 2020 Publication	<1 %

91 Yao Xu, Wenxue Li. "Finite-time synchronization of fractional-order complex-valued coupled systems", Physica A: Statistical Mechanics and its Applications, 2019
Publication <1 %

92 "Recent Trends in Dynamical Systems", Springer Science and Business Media LLC, 2013
Publication <1 %

93 A. Atangana, E. Bonyah. "Fractional stochastic modeling: New approach to capture more heterogeneity", Chaos: An Interdisciplinary Journal of Nonlinear Science, 2019
Publication <1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

A mathematical model of tuberculosis (TB) transmission with children and adults groups: Afractional model

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19

PAGE 20

PAGE 21

PAGE 22

PAGE 23

PAGE 24

PAGE 25

PAGE 26

PAGE 27

PAGE 28

PAGE 29

PAGE 30
