


Development of Air Polluter Model for the Carbon Monoxide (CO) Element Based on Mixed Geographically Temporal Weighted Regression (MGTWR) Kriging

Kukuh Winarso, Hari Basuki Notobroto and Fatmawati

Applied Mathematical Sciences, Vol. 8, 2014, no. 118, 5863 – 5873

HIKARI Ltd, www.m-hikari.com

<http://dx.doi.org/10.12988/ams.2014.45334>



Applied Mathematical Sciences

Indexing and Abstracting

Information for Applied Mathematical Sciences might be found in:

CrossRef (USA)
 Google Scholar (USA)
 Scilit (Switzerland)
 Microsoft Academic (USA)
 ERA-Excellence in Research for Australia (Australia)
 ROAD Directory of Open Access Scholarly Resources (EU)
 MIAR - Information Matrix for the Analysis of Journals (Spain)
 Academic Journals Database (Switzerland)
 WorldCat (USA)
 eLIBRARY.RU (Russia)
 Socolar (P.R. China)
 Research GATE (USA,EU)
 MathGuide (Germany)

Applied Mathematical Sciences ResearchGate (RG) Impact: 0.17

These values are calculated using ResearchGate (RG) data (https://www.researchgate.net/journal/1312-885X_Applied_Mathematical_Sciences) based on average citation counts from work published in the journal.

RG Journal impact history (https://www.researchgate.net/journal/1312-885X_Applied_Mathematical_Sciences)										
Year:	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
RG journal impact:	0.23	0.31	0.72	1.07	0.96	1.16	0.63	0.78	0.36	0.17

Full-text Archives and Copyright policy:

Portico (USA)
 National Library (Bulgaria)
 SHERPA/RoMEO (UK)

Recently most downloaded articles:

R. B. Paris

The expansion of the confluent hypergeometric function on the positive real axis (<ams/ams-2018/ams-1-4-2018/p/parisAMS1-4-2018.pdf>)
 Applied Mathematical Sciences, Vol. 12, 2018, no. 1, 19-26

R. Caimmi

Advanced astronomy in Bhagavata Purana: the role of the Sun and a tale from 108 (<ams/ams-2018/ams-5-8-2018/p/caimmiAMS5-8-2018.pdf>)
 Applied Mathematical Sciences, Vol. 12, 2018, no. 5, 229-254

Luca Grosset, Bruno Viscolani

A note on optimality conditions for control problems with parameters (<ams/ams-2018/ams-13-16-2018/p/grossetAMS13-16-2018.pdf>)
 Applied Mathematical Sciences, Vol. 12, 2018, no. 16, 773-782

Francisco J. Torres

Solvability for fractional boundary value problems with p-Laplacian operator (<ams/ams-2018/ams-21-24-2018/p/torresAMS21-24-2018.pdf>)
 Applied Mathematical Sciences, Vol. 12, 2018, no. 24, 1167-1178

Vol. 8, no. 89-92, 2014
doi:10.12988/ams

ISSN 1312-885X (Print)
ISSN 1314-7552 (Online)

APPLIED MATHEMATICAL SCIENCES

Journal for Theory and Applications

Editorial Board

K. S. Berenhaut (USA)	F. Hosseinzadeh Lotfi (Iran)
Nguyen Buong (Vietnam)	J. E. Macias-Diaz (Mexico)
G. Caristi (Italy)	B. J. McCartin (USA)
N. Cercone (Canada)	Michael Kwok-Po Ng (China)
Ping-Teng Chang (Taiwan)	B. Oluyede (USA)
W. Y. C. Chen (China)	J. F. Peters (Canada)
K. W. Chow (China)	V. Popov (Russia)
M. Darus (Malaysia)	Qinghua Qin (Australia)
M. Ferrara (Italy)	Z. Retchkiman (Mexico)
Ji-Huan He (China)	Cheon Seoung Ryoo (Korea)
T. C. Headrick (USA)	M. de la Sen (Spain)
Tzung-Pei Hong (Taiwan)	G. L. Silver (USA)
G. Jumarie (Canada)	F. T. Suttmeier (Germany)
S. Khardi (France)	Xue-Cheng Tai (Norway)
L. Kohaupt (Germany)	A. Vacca (Italy)
V. Kreinovich (USA)	Xin-She Yang (UK)

Managing Editor: Emil Minchev

Hikari Ltd

Applied Mathematical Sciences

Aims and scopes: The journal publishes refereed, high quality original research papers in all branches of the applied mathematical sciences.

Call for papers: Authors are cordially invited to submit papers to the editorial office by e-mail to: ams@m-hikari.com . Manuscripts submitted to this journal will be considered for publication with the understanding that the same work has not been published and is not under consideration for publication elsewhere.

Instruction for authors: The manuscript should be prepared using LaTeX or Word processing system, basic font Roman 12pt size. The papers should be in English and typed in frames 14 x 21.6 cm (margins 3.5 cm on left and right and 4 cm on top and bottom) on A4-format white paper or American format paper. On the first page leave 7 cm space on the top for the journal's headings. The papers must have abstract, as well as subject classification and keywords. The references should be in alphabetic order and must be organized as follows:

- [1] D.H. Ackley, G.E. Hinton and T.J. Sejnowski, A learning algorithm for Boltzmann machine, *Cognitive Science*, 9 (1985), 147-169.
- [2] D.O. Hebb, *Organization of Behaviour*, Wiley, New York, 1949.

Editorial office

e-mail: ams@m-hikari.com

Postal address:

HIKARI Ltd, P.O. Box 85
Ruse 7000, Bulgaria

Street address:

HIKARI Ltd, Vidin str. 40, office no. 3
Ruse 7012, Bulgaria

www.m-hikari.com

Published by Hikari Ltd



[Home](#) | [Journals](#) | [Books](#) | [Paper submission](#) | [About us and our mission](#) | [News](#) | [Contact us](#) |



APPLIED MATHEMATICAL SCIENCES

Site Search:

Online Edition, Vol. 8, 2014, no. 117-120

Mohammad Ahsanullah, Mohammad Shakil

[A note on a characterization of J-shaped distribution by truncated moment](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 117, 5801-5812

<http://dx.doi.org/10.12988/ams.2014.47556>

Stanley O. Sewe, Patrick G. O. Weke, Joseph K. Mung'atu

[Modelling dependence between the equity and foreign exchange markets using copulas](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 117, 5813-5822

<http://dx.doi.org/10.12988/ams.2014.47560>

S. Vaithyasubramanian, A. Christy, D. Saravanan

[An analysis of Markov password against Brute force attack for effective web applications](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 117, 5823-5830

<http://dx.doi.org/10.12988/ams.2014.47579>

P. Haarsa, S. Pothat

[The energy method on wave equation in a resistant medium with series expansion](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 117, 5831-5834

<http://dx.doi.org/10.12988/ams.2014.46476>

P. Haarsa, S. Pothat

[The unique solution on the Neumann B.C. of diffusion equation with energy method](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 117, 5835-5838

<http://dx.doi.org/10.12988/ams.2014.46475>

Prasanta Kumar Mohanty, Manoj Kumar Hota

[Quadrature rules for evaluation of hyper singular integrals](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 117, 5839-5845

<http://dx.doi.org/10.12988/ams.2014.47584>

Sudevi Basu, Neha, Pratibha Nair, Shreya Singh, Nisha

[Use of statistical expressions to evaluate the performance of two parameter adsorption isotherm models on dyes using natural adsorbents](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 118, 5847-5862

<http://dx.doi.org/10.12988/ams.2014.47555>

Kukuh Winarso, Hari Basuki Notobroto, Fatmawati

[Development of air polluter model for the carbon monoxide \(CO\) element based on mixed geographically temporal weighted regression \(MGTWR\) kriging](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 118, 5863-5873

<http://dx.doi.org/10.12988/ams.2014.45334>

A.V. Anufrieva, K.B. Igudesman, D.N. Tumakov

[Peculiarities of elastic wave refraction from the layer with fractal distribution of density](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 118, 5875-5886

<http://dx.doi.org/10.12988/ams.2014.46473>

Dmitrii Tumakov

[Iterative method for solving the problem of scattering of an electromagnetic wave by a partially shielded conducting sphere](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 118, 5887-5898

<http://dx.doi.org/10.12988/ams.2014.48657>

Makoto Hayashi

[An improvement for the unique existence of a limit cycle of the Duff and Levinson-type system](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 119, 5899-5905

<http://dx.doi.org/10.12988/ams.2014.48643>

Christos E. Kountzakis

[Coherent risk measures by pricing functionals](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 119, 5907-5914

<http://dx.doi.org/10.12988/ams.2014.48607>

Ka-Luen Cheung

[Exact solutions for the two-dimensional incompressible magnetohydrodynamics equations](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 119, 5915-5922

<http://dx.doi.org/10.12988/ams.2014.48641>

Zlatko Pavic

[Inequalities for convex and non-convex functions](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 119, 5923-5931

<http://dx.doi.org/10.12988/ams.2014.48601>

V. N. Dumachev

[On semideterministic finite automata games type](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 119, 5933-5941

<http://dx.doi.org/10.12988/ams.2014.48645>

Jaekeun Park

[Simpson-like type inequalities for functions whose derivatives in absolute value at certain power are s-geometrically convex](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 119, 5943-5954

<http://dx.doi.org/10.12988/ams.2014.48626>

Jaekeun Park

[Remarks on Simpson-like type inequalities for harmonically convex mappings on the co-ordinates](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 120, 5955-5969

<http://dx.doi.org/10.12988/ams.2014.48627>

Jaekeun Park

[On the Simpson and Hermite-Hadamard type inequalities for the geometrically convex functions](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 120, 5971-5983

<http://dx.doi.org/10.12988/ams.2014.48671>

Aminah Qawasmeh, Marwan Alquran

[Reliable study of some new fifth-order nonlinear equations by means of G'/G expansion method and rational sine-cosine method](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 120, 5985-5994

<http://dx.doi.org/10.12988/ams.2014.48669>

Zitian Li

[Heteroclinic breather-wave for the coupled Schrodinger Boussinesq equation](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 120, 5995-6000

<http://dx.doi.org/10.12988/ams.2014.48662>

Jong Seo Park

[On common fixed point for converse commuting maps in IFMS](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 120, 6001-6004

<http://dx.doi.org/10.12988/ams.2014.48614>

C. S. Ryoo

[A note on generalized twisted \(h; q\)-tangent numbers and polynomials](#)

Applied Mathematical Sciences, Vol. 8, 2014, no. 120, 6005-6010

<http://dx.doi.org/10.12988/ams.2014.48606>

Journals | Books | Author guidelines

Development of Air Polluter Model for the Carbon Monoxide (CO) Element Based on Mixed Geographically Temporal Weighted Regression (MGTWR) Kriging

Kukuh Winarso

Post Graduate Student of MIPA, Universitas Airlangga and Departement of
Industrial Engineering, Faculty of Engineering, Universitas Trunojoyo Madura,
JL. Raya Telang P.O. Box 2 Bangkalan, Madura, Indonesia

Hari Basuki Notobroto

Department of Biostatistics and Demography, Faculty of Public Health, Airlangga
University, JL. Mulyorejo, Kampus C, Unair Surabaya, Indonesia

Fatmawati

Department of Mathematics, Faculty of Sciences and Technology, Airlangga
University, JL. Mulyorejo, Kampus C, Unair Surabaya

Copyright © 2014 Kukuh Winarso, Hari Basuki Notobroto, Fatmawati. This is an open access
article distributed under the Creative Commons Attribution License, which permits unrestricted
use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Motor vehicle exhaust is the main source of Carbon Monoxide (CO). In urban areas, where motor vehicles are easily found, CO is commonly identified as one of the air polluter sources. CO is hazardous, and human being will suffer from serious health problems if they are exposed to this polluting agent in a prolonged period. This study aims to develop a model that will aid the effort to minimize the negative effects of CO. The model is built in which it will be able to show the cause-factors and the preventing-factors of CO generation. The Mixed Geographically Temporal Weighted Regression (MGTWR) approach is used as

the basis of the model. MTGWR is a spatial-temporal regression model, which takes into account the geographical and temporal aspects of the pollution. However, the MGTWR model cannot be used to predict the effect if it is used outside the sample location of the research, unless we predict the associated regression coefficients in the respected area beforehand. In this case, we use the estimated predictor parameter based on the Kriging method to predict the regression parameters outside the research location. As the result, the Kriging Predictor-based MTGWR model can be used to estimate the pollution level caused by CO outside the sample location of the research.

Keywords: Air Polluter, Carbon Monoxide, Spatio-Temporal, MGTWR, Kriging-predictor

1. INTRODUCTION

Previous study shows that there is a parallel relationship between the number of polluted air-generated disease sufferers and the intensity of industrialization and urbanization in countries whose level of air pollution are severely high [1]. Carbon Monoxide (CO) is one of the air pollutants, which is generated from the imperfect fuel combustion of motor vehicles and industrial machines. Motor vehicle exhaust is the main generator of Carbon Monoxide and it is common to be found in big cities. Based on data, about 60% of air pollution in major urban areas is related with public transportation [2]. Within the context of human health, Carbon Monoxide is toxic. It causes lower fetal weight, higher rate of infant death, and leads to human brain damage. The air quality standard that is related with Carbon Monoxide is 10.000 ug/Nm³ [3]

Demographically, the potential of the effect and causes of air pollution will be different one to another in different regions. Also, it is inappropriate to analyze the effect and causes of air pollution based on a global approach, as we will be unable to examine the influence of local variations if we are to use global approach [4-5]. The common spatial regression method known is the Geographically Weighted Regression (GWR). GWR is a weighted-regression technique that is based on the simple regression approach. It is a common statistical technique used for spatial heterogeneity. The term heterogeneity in this case is the measurement relationship of different variables among various locations [6]. Spatial heterogeneity occurs when an independent variable gives different responses to different locations within the same research location [7]. The crux of GWR approach is determining the regression model for each of the location nodes so that the regression models obtained are unique, in which a model for a node will be different to the others [8]. There is another model aside from GWR, which is Mixed Geographically Temporal Weighted regression (MGTWR). MTGWR is a combined modeling approach between global regression and local regression (GWR) [9]. This paper takes into consideration the element of location, as well as the time into the local variation based on MGTWR approach. The

usefulness of global regression is that it can be used to predict in each location. However, in the spatial regression models such as GWR, MGWR (Mixed Geographically Weighted Regression), and MGTWR, the global regression cannot be used for predicting outside the research sample location, unless we predict the respected location's regression coefficients beforehand [10]. In order to eliminate this obstacle, we estimate the predictor parameter using the Kriging method to predict the regression parameter outside the research location.

2. RESEARCH OBJECTIVE

This research aims to develop the air polluter model for the Carbon Monoxide (CO) element based on the Mixed Geographically Temporal Weighted Regression (MGTWR) approach, in which the developed model can be used to predict outside the research location using the Kriging predictor estimation. This way, the developed model can be used to predict the regression parameter outside the research location.

3. METHODOLOGY

3.1 Mixed Geographically Temporal Weighted Regression (MGTWR)

MGTWR is the advancement of the MGWR model [11] with the incorporation of the temporal element into the model. The temporal factor is aimed to predict the observation time, and to complement the location (location coordinate) factor. Mathematically, the temporal model can be stated as the following:

$$y_i = \sum_{j=1}^q \beta_j x_{ij} + \beta_j (u_i, v_i, t_i) x_{ij} + \varepsilon_i \quad (1)$$

Where: $i = 1, 2, 3, \dots, n$

β_i is global variable x_i

β_j is local variable x_j

y_i is Response of variables

x_i is predictors of variables

u_i is longitude at the time-i

v_i is latitude at the time-i

t_i is the time length at the time-i

3.2. Kriging Spatial Predictor Model

Kriging method is used to predict the regression parameter outside the research location [12]. The assumption that is associated with the predictor is:

$$p(b; s_0) = \sum_{i=1}^n \lambda_i b(s_i), \quad \sum_{i=1}^n \lambda_i = 1 \quad (2)$$

Where:

λ_i = the weight $b(s_i)$ for the estimation of location s , with the value of $b(s_i)$ has different weighted-coefficients for the estimation in different locations

s_i = location vector

s_0 = Location of the data to be estimated

n = number of the sample data

The Kriging spatial predictor matrix notation can be stated as the following [12]:

$$\begin{bmatrix} \gamma(s_i - s_j) & f_t(s_i) \\ f_t(s_j) & 0 \end{bmatrix} \begin{bmatrix} \lambda_j \\ m_t \end{bmatrix} = \begin{bmatrix} \gamma(s_i - s_j) \\ f_t(s_0) \end{bmatrix}, \text{ for } i = 1, 2, \dots, n \text{ and } t = 1, 2, \dots, p$$

Based on the previous matrix, we can compute the weight matrix for the Universal Kriging, which is:

$$\begin{bmatrix} \lambda_j \\ m_t \end{bmatrix} = \begin{bmatrix} \gamma(s_i - s_j) & f_t(s_i) \\ f_t(s_j) & 0 \end{bmatrix}^{-1} \begin{bmatrix} \gamma(s_i - s_j) \\ f_t(s_0) \end{bmatrix} \quad (3)$$

Notations :

$\gamma(s_i - s_j)$ = Semivariogram between sampled nodes

$\gamma(s_0 - s_i)$ = Semivariogram between sampled node and estimated node

$f_t(s_i), f_t(s_j)$ = Location coordinate of the sampled data

λ_j = Weighting value to be computed

m_t = Value of *lagrange* parameter

(s_i, s_j) = Location of the sampled data

s_0 = Location of the data to be estimated

p = Number of order in the trend equation

Semivariogram is used to determine the distance where the observation data values are not dependent one to another or have no correlation among them [12]. The equation in the Kriging model is used to model the trend of the low order polynomial, which is the first or the second order [13]. If the obtained trend has the first order at \mathbb{R}^3 , then the equation is:

$$m(s) = m(x,y,z) = a_0 + a_1x + a_2y + a_3z$$

where x,y,z are the sampled location coordinates. Thus, the equation (3) becomes as the following:

$$\sum_{j=1}^n \lambda_j \gamma(s_i - s_j) + m_0 + m_1 x_i + m_2 y_i + m_3 z_i = \gamma(s_0 - s_i), \text{ for } i = 1, 2, \dots, n$$

With assumption:

$$\sum_{j=1}^n \lambda_j x_j = x, \quad \sum_{j=1}^n \lambda_j y_j = y, \quad \sum_{j=1}^n \lambda_j z_j = z, \quad \sum_{j=1}^n \lambda_j = 1$$

If we want to transform the equation above into the Kriging matrix in the

following first order Kriging trend (KT):

$$K_{KT} = \begin{bmatrix} \gamma(s_1 - s_1) & \cdots & \gamma(s_1 - s_n) & 1 & x_1 & y_1 & z_1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma(s_n - s_1) & \cdots & \gamma(s_1 - s_n) & 1 & x_n & y_n & z_n \\ 1 & \cdots & 1 & 0 & 0 & 0 & 0 \\ x_1 & \cdots & x_n & 0 & 0 & 0 & 0 \\ y_1 & \cdots & y_n & 0 & 0 & 0 & 0 \\ z_1 & \cdots & z_n & 0 & 0 & 0 & 0 \end{bmatrix} \quad \lambda_{KT} = \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_n \\ m_0 \\ m_1 \\ m_2 \\ m_3 \end{bmatrix}$$

$$k_{KT} = \begin{bmatrix} \gamma(s_0 - s_1) \\ \vdots \\ \gamma(s_0 - s_n) \\ 1 \\ x \\ y \\ z \end{bmatrix}$$

where : $K_{KT} \lambda_{KT} = k_{KT}$

then the weighting matrices can be stated as the followings [13]:

$$\lambda_{KT} = K_{KT}^{-1} k_{KT}$$

The case study in this research uses the Carbon Monoxide (CO) element as the response variable Y, while the predictor variables are the air temperature (X₁), the wind velocity (X₂), the air humidity (X₃), the traffic velocity (X₄), the area size of the urban forest (X₅), the population density (X₆), and the business center aspect (X₇) [14]

4. RESULTS AND DISCUSSION

The first MGTWR model test is the model fitness, in which the hypothesis can be stated as the followings:

$$H_0 : \beta_k(u_i, v_i, t_i) = \beta_k(u_i, v_i) \quad k = 0, 1, 2, \dots, 4, j = \text{d a n } \dots$$

(MGTWR model is not different from the MGWR model)

$$H_1 : \text{At least one } \beta_k(u_i, v_i, t_i) \neq \beta_k(u_i, v_i)$$

(MGTWR model is significantly different from the MGWR model)

The test results are summarized in Table 1:

Table 1 Fitness Test of MGTWR Model for the CO Response

Source of Error	Sum Square	Degree of Freedom	Mean Square	F	p-value
Improvement	7,7679	15,9260	0,4877	4,2433	0,0000
MGTWR	46,1362	401,3706	0,1149		
MGWR	53,9041	417,2966			

Table 1 shows that the F test statistical value is 4,2433, and the the p-value of 0,0000. Using the significance value (α) of 5%, we must reject H_0 , and conclude that the MGTWR model is significantly different from the MGWR model. Therefore, we can further conclude that the MGTWR model is more proper to model the Air Polluter Standard Index (APSI) for Carbon Monoxide (CO). This means that the time element is influential in the APSI modeling for CO so that not only the location factor is considered, but the observation time is taken as the influential factor to the APSI model for CO as well.

Next step is the global parameter test that is conducted to identify which global predictor variables significantly influence the response. In this test, we perform the partial global parameter test using the hypothesis below:

$$H_0 : \beta_k = 0 \quad (\text{global variable } x_k \text{ is not significant})$$

$$H_1 : \beta_k \neq 0 \quad (\text{global variable } x_k \text{ is significant})$$

Table 2 shows that by using the level of significance (α) 5%, we can conclude that the global predictor variable that significantly influence the APSI for CO are the area size of the urban forest (X_5) and the business center aspect (X_7), because these variables have the p-value that are less than 0.05.

Table 2 Parsial Test of Global Variable of MGTWR Model

Global Parameter			
Variable	Beta	T	p-value
X ₅	-0,0445	-0,4171	0,0080*
X ₆	0,0088	0,3556	0,3613
X ₇	-0,0456	2,0045	0,0228*

Note: *) significant at $\alpha = 5\%$

By using $\alpha = 5\%$, the location and the observation time of APSI for CO in Surabaya city can be classified based on the significant variables for APSI values for CO. The detailed significant variable list in each location and in each observation time is presented in Table 3:

Table 3 MGTWR Model for CO at Five Observation Location Nodes

No	Observation Location	Observation Time	Significant Variables
1	SUF 1 (Location 1)	Morning	X ₃ , X ₄
		Noon	X ₃ , X ₄
		Evening	X ₁ , X ₂ , and X ₄
2	SUF 3 (Location 2)	Morning	X ₃ , X ₄
		Noon	X ₃ , X ₄
		Evening	X ₁ , X ₂ , X ₃ , X ₄
3	SUF 4 (Location 3)	Morning	X ₃ , X ₄
		Noon	X ₄
		Evening	X ₁ , X ₂ , X ₃ , X ₄
4	SUF 5 (Location 4)	Morning	X ₃ , X ₄
		Noon	X ₃ , X ₄
		Evening	X ₁ , X ₂ , X ₃ , X ₄
5	SUF 6 (Location 5)	Morning	X ₃ , X ₄
		Noon	X ₂
		Evening	X ₂ , X ₃ , and X ₄

The MGTWR model obtained in each of the location and observation time will be different one to another, depending on the parameter value of MGTWR and on which predictor variables that significantly influence the response variable of APSI (CO) with the R² value of 36.60%.

To predict the MGTWR model at a certain location node, we use the Kriging approach that is presented in the following Table 4:

Table 4 Kriging-based MGTWR Model with Local Parameter Lokal for CO Element

No	Sub-district	Time	Constant	X ₁	X ₂	X ₃	X ₄
1	Location z	Morning	4.5498	-0.0541	0.0690	-0.1437	0.0614
		Noon	4.4902	0.0164	0.0180	0.0932	0.2797
		Evening	4.5415	0.1745	0.0870	0.1284	0.2571
2	Location y	Morning	4.5523	-0.0425	0.0667	-0.1441	0.0636
		Noon	4.4903	0.0145	0.0192	0.0833	0.2773
		Evening	4.5366	0.1706	0.0840	0.1233	0.2567
3	Location x	Morning	4.5457	-0.0618	0.0665	-0.1335	0.0693
		Noon	4.4897	0.0202	0.0206	0.0923	0.2775
		Evening	4.5441	0.1686	0.0874	0.1262	0.2612

In Table 4, if we are to estimate for location x, the Kriging-MGTWR with the CO response for evening is:

$$\text{Ln (CO)} = 4.544 + 0.0874X_2 + 0.2612X_4 + 0.0445X_5 - 0.0456X_7$$

The classification map in Surabaya based on CO polluter can be seen in the following:




Figure 1. Classification of Sub-districts in Surabaya Based on the Air Polluter CO

There are eleven sub-districts that are classified as unhealthy (yellow-colored) and twenty sub-districts that are classified as fair (blue-colored).

5. CONCLUSION

Aside from the global influence, the MGTWR model of the air pollution is also influenced significantly by the location factor (geographical factor). Therefore, in this study, the mixed model is suitable to model the air polluter. The local influence of MTGWR for CO element for observation time morning and noon show that the two significantly influencing predictor variables are the air humidity (x_3) and the traffic velocity (x_4). This indicates that in the morning and noon, the influence of motor-vehicle exhaust and air humidity are very dominant in polluting the air with CO. For the night observation, almost all of the predictor variables influence the air pollution level, with the global influence factors of it are the area size of the urban forest (x_5) and the business center (x_7). To predict an

MGTWR model outside the observation nodes, we can use the Kriging estimation approach.

REFERENCES

- [1] Ebtekar, 2006. Air Pollution induced asthma and alterations in cytokine pattern. *Allergy Asthma Immunol*, 5(2):47-56. ISSN 1735-1502
- [2] Environment Minister, (1997). Air Polluter Standard Index - APSI. Jakarta, Indonesia.
- [3] Fahimi, M., Dharma, B., Fetararayani, D., Baskoro, (2012). Asosiasi antara polusi udara dengan IgE total serum dan tes faal paru pada polisi lalu lintas. *Jurnal Penyakit Dalam*, 1-9.
- [4] Robinson, D. and Lloyd, J. M. 2011. Increasing the accuracy of nitrogen dioxide (NO₂) pollution mapping using geographically weighted regression & geostatistic. *International Journal of Applied Earth Observation and Geoinformation*, In Press 1-10. ISSN: 0303-2434
- [5] Gilbert, A. and Chakraborty, J. 2011. Using geographically weighted regression for environmental justice analysis: Cumulative cancer risks from air toxics in Florida. *Social Science Research*, 40 (2011): 273-286. DOI:10.1016/j.ssresearch.2010.08.006
- [6] Mennis, J. 2006. Mapping the results of geographically weighted. *The Cartographic*, 43(2): 171-179. DOI: 10.1179/000870406X114658
- [7] Fotheringham A.S., Brunson C., Charlton (2002). *Geographically Weighted Regression. the analysis of spatially varying relationship*. John Wiley and Sons, LTD. ISBN 0-471-49616-2
- [8] Brunson, C., Fotheringham, A. and Charlton. 1999. Some notes on parametric significance tests for Geographically Weighted Regression. *Journal of Regional Science*, 39 (3): 497-524. DOI: 10.1111/0022-4146.00146
- [9] Purhadi and Hasbi, Y. 2012. Mixed geographically weighted regression model (Case study: The percentage of poor households in Mojokerto 2008). *European Journal of Scientific Research*, 69 (2): 188-196. ISSN: 1450202X
- [10] Walter J., Carsten R. and Jeremy W. Lichstein (2005) *Local and global approaches to spatial data analysis in ecology*, *Global Ecology and Biogeography* 14 (1), 97-98. DOI: 10.1111/j.1466-822X.2004.00129.x

- [11] Leung Y., Mei Chang L., and ZhangWen X. (2000a), Statistical test for spatial nonstationarity based on the geographically weighted regression model, *Environment and Planning*, 32 (5): 871–890. DOI:10.1068/a32117
- [12] Cressie, N, A, C. (1992). *Statistics For Spatial Data*. Terra nova. 4 (5): 613-617. DOI: 10.1111/j.1365-3121.1992.tb00605.x
- [13] Goovaerts, P. (1997). *Geostatistics for Natural resource Evaluation*. Oxford university Press, New York. ISBN-13 978-0-19-511538-3. ISBN 0-19-511538-4
- [14] K Winarso, H Basuki, Fatmawati, 2014, Development Of Air Polluter Model For The Carbon Monoxide (CO) Element Based On Mixed Geographically Temporal Weighted Regression (MGTWR) Kriging, *international Of Journal Academics Research*, 6 (2): 193-198, DOI: 10.7813/2075-4124.2014/6-2,

Received: May 1, 2014