

COVID-19 and Endothelial Dysfunction Biomarkers and Potential Drug Mechanisms

by Andrianto Andrianto

Submission date: 14-Mar-2022 05:07PM (UTC+0800)

Submission ID: 1783926450

File name: thelial_Dysfunction_Biomarkers_and_Potential_Drug_Mechanisms.pdf (994.52K)

Word count: 7606

Character count: 44745

Indonesian Journal of Tropical and Infectious Disease

Vol. 9 No. 2 May–August 2021

Review Article

COVID-19 and Endothelial Dysfunction: Biomarkers and Potential Drug Mechanisms

Andrianto¹, Ronaldi Rizkiawan¹, Primasitha Maharany Harsoyo¹, Syafira Yasmine²

¹ Department of Cardiology and Vascular Medicine, School of Medicine, Universitas Airlangga - Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

² School of Medicine, Universitas Airlangga, Surabaya, Indonesia

Received: 15th February 2021; Revised: 4th April 2021; Accepted: 11st April 2021

ABSTRACT

Since the first report of pneumonia outbreak in Wuhan by the end of 2019, Coronavirus Disease 2019 (COVID-19) has become a global pandemic; causing millions of deaths globally and affecting the rest of worldwide population. The disease is caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), which enters hosts by inhabiting Angiotensin-Converting Enzyme-2 (ACE-2) receptors expressed in the endothelium layer of not only the respiratory tracts, but also various organs in the body. COVID-19 has been reported to trigger multiple cardiovascular manifestations. Since endothelial dysfunction plays an important role in cardiovascular events and the endothelium is heavily involved in COVID-19 pathophysiology, it is important to investigate their associations and previously established drug potencies to improve endothelial functions as possible treatment options for COVID-19. In this review, we summarize endothelial dysfunction biomarkers involved in COVID-19 and drugs that have shown potential endothelial protective properties to better understand the incidence of endothelial dysfunction in COVID-19 and its future treatment. We searched in PubMed, Wiley Online Library, EBSCO, ScienceDirect databases for literatures containing following keywords: "Endothelial dysfunction", "COVID-19", and "biomarkers". Eligible publications were then assessed and studied to comprise our literature review. A total of 96 studies matched our criteria and provided scientific evidences for our review. Materials were then compiled into a review summarizing endothelial biomarkers involved in COVID-19 and potentially repurposed drugs targeting endothelium for COVID-19. Various endothelial dysfunction biomarkers were found to be elevated in COVID-19 and is found to be related to its severity, such as adhesion molecules, selectins, PAI-1, and von Willebrand Factors. Multiple drugs targeting the endothelium are also potential and some are under investigation for COVID-19.

Keywords: COVID-19, endothelial dysfunction, SARS-COV-2, biomarkers

ABSTRAK

Sejak pertama kali pelaporannya sebagai wabah pneumonia di Wuhan pada akhir 2019, Corona Virus disease 2019 (COVID-19) telah menjadi pandemi global yang menyebabkan jutaan kematian dan memengaruhi populasi di seluruh dunia. Penyakit ini disebabkan oleh SARS-CoV-2 yang menginvasi sel inang dan berikatan dengan reseptor angiotensin-converting enzyme-2 (ACE-2) yang diekspresikan lapisan endotel saluran pernapasan serta berbagai organ lain dalam tubuh. COVID-19 telah dilaporkan dapat menimbulkan berbagai manifestasi gangguan kardiovaskular. Mengingat peran penting disfungsi endotel yang berperan dalam kejadian kardiovaskular dan patofisiologi COVID-19, maka penting menyelidiki hubungan di antaranya. Selain itu, perlu dilakukan telaah mengenai potensi obat yang sebelumnya telah terbukti memperbaiki fungsi endotel sebagai pilihan terapi yang digunakan untuk COVID-19. Pada studi ini, kami memaparkan beberapa biomarka disfungsi endotel yang terlibat pada COVID-19 dan obat-obatan yang melindungi endotel sebagai terapi potensial COVID-19 di masa mendatang. Kami melakukan pencarian di database PubMed, Perpustakaan Online Wiley, EBSCO, dan ScienceDirect untuk literatur yang berisi kata kunci berikut: "Disfungsi endotel", "COVID-19", dan "biomarker". Publikasi yang memenuhi syarat kemudian dinilai dan ditelaah untuk menyusun tinjauan pustaka. Sebanyak 96 publikasi memenuhi kriteria dan memberikan

* Corresponding Author:
andrianto@fk.unair.ac.id

bukti ilmiah untuk tinjauan pustaka ini. Materi kemudian disusun menjadi ulasan yang merangkum biomarker-biomarker endotel yang terlibat dalam COVID-19 dan obat-obatan dengan target endotel yang potensial. Berbagai biomarker disfungsi endotel ditemukan meningkat pada COVID-19 dan terkait dengan tingkat keparahannya, seperti molekul adhesi, selektin, PAI-1, dan Faktor von Willebrand. Beberapa obat yang menargetkan endotel juga ditemukan memiliki potensi dengan beberapa tengah diinvestigasi lebih lanjut sebagai terapi COVID-19.

Kata kunci: COVID-19, disfungsi endotel, SARS-CoV-2, biomarka

How to Cite: Andrianto., Rizkiawan., Harsoyo, PM., Yasmine, S. COVID-19 and Endothelial Dysfunction: Biomarkers and Potential Drug Mechanism. Indonesian Journal of Tropical and Infectious Disease, 9(2), 108–119.

INTRODUCTION

In December 2019, a novel corona virus was isolated from the respiratory tract epithelial layer of a patient with pneumonia of unknown cause in Wuhan, China. By November 2020, the Severe Acute Respiratory Syndrome-Coronavirus-2 (SARS-CoV-2), and its consequential disease, Coronavirus Disease 2019 (COVID-19), has now infected more than 46 millions of people all over the world and caused more than 1.2 million global death.¹

Beside instigating substantial pulmonary disturbances such as pneumonia and acute respiratory distress syndromes, COVID-19 is known to trigger extrapulmonary responses, including cardiovascular manifestations. Several reported cardiovascular manifestations of COVID-19 include myocardial ischemia, arrhythmia, and even cardiogenic shock.^{1,2}

Endothelial dysfunction is a significant contributor of various cardiovascular diseases mechanism. Endothelial cells play a major role in maintaining blood tissue non-thrombogenicity, regulating thrombosis, thrombolysis, thrombocyte adherence, vascular tone, and blood flow.¹ Any disturbance of endothelial function may trigger a progression of cardiovascular problems.³

SARS-CoV-2 exploits Angiotensin-Converting-Enzyme-2 (ACE-2) as a receptor to attack human cells. ACE-2, a main target component in various cardiovascular diseases mechanism and drugs, is widely expressed in both respiratory endothelial layers and vascular endothelial layers. Therefore, it is important to learn of the endothelial involvement in COVID-19 related cardiovascular function alterations as well as the changes of biomarkers it comprises.^{1,2}

DISCUSSION

SARS-CoV-2 related endothelial changes

The innermost part of the vascular system is comprised of a single layer of endothelial cells called endothelium. A healthy endothelium maintains blood fluidity through platelet reactivity regulation, coagulation, and smooth thrombolysis by synthesizing and responding to vasoactive molecules accordingly.² The endothelium, alongside its primary immunoregulatory properties, also plays an important role in maintaining dynamic interactions between pro-coagulant and fibrinolytic factors within the vascular system. In its inactive state, the endothelium acts as a barrier that separates pro-thrombotic subendothelial layers and procoagulant factors carried within the circulation.^{3,9}

SARS-CoV-2 directly affects the endothelial cells due to the high expression of ACE-2 receptors and Transmembrane Protease, Serine 2 (TMPRSS2) enzyme. After being bound by SARS-CoV-2, ACE-2 receptors are internalized such that the diminishing number of ACE-2 in endothelial cells promotes inflammation and thrombosis, triggered by subsequent hyperactivity of local Angiotensin-II (Ang-II). The decrease of ACE-2 receptors also reduces a number of conversions it normally mediates, including Angiotensin 1-7 from Ang-II, which acts as a vasoactive ligand of the MAS receptor. This causes reduction of MAS receptor activation and induction of pro-inflammatory phenotypes through the increase of type-1 Angiotensin receptor (AT₁R) activation. Furthermore, the decrease of ACE-2 receptor prevents the degradation of des-Arg-9-

Bradykinin (DABK) into inactive peptides, which subsequently raises prothrombotic signals through bradykinin receptors (BKR) activation.^{4,5}

Endothelial activation and subsequent dysfunction is marked by imbalance of endothelium-released vasomotor factors, expressions of inflammatory cytokines and chemokines, expression and secretion of selectins and adhesive molecules as well as modulation of local thrombotic pathway.⁶ The release of cytokines and pro-inflammatory chemokines by activated macrophages augments the vicious circle of vascular integrity, coagulation, and thrombosis disturbances through reduction of endothelial glycocalyx, activation of coagulation system, and dampening of anticoagulation mechanism. Endothelial cell adhesive phenotypes induced by pro-inflammatory cytokines and chemokines promote neutrophil infiltration, producing large quantity of histotoxic components such as reactive oxygen species (ROS) and Neutrophil Extracellular Traps (NETs).

Endothelial activation is the transition of static phenotype to a specific phenotype involving responses to host immunity.⁵ Endothelial cells activation causes an increase of inflammatory cytokines and adhesion molecules, triggering releases of leukocyte, adhesion, and migration

to subendothelial chambers, constituting fundamental process of atherosclerotic lesion initiation, progress, and destabilization.⁵ Activated endothelial cells begin coagulation process by expressing P-selectin, von Willebrand factor (vWf), and fibrinogen, causing massive platelet binding, formation of fibrin and clotting of Red Blood Cells (RBC), which finally results in systemic thrombosis and Disseminated Intravascular Coagulation (DIC).⁴

Measurable biomarkers of endothelial activation and dysfunction

Activated endothelial cells express increased levels of E-selectin, P-selectin, Intercellular Adhesion Molecule 1 (ICAM-1), and Vascular Cell Adhesion Molecule 1 (VCAM-1). Upregulations of E-selectins, ICAM-1, and VCAM-1 are mediated at transcription level. E-selectin induces the rolling of circulatory leukocytes. VCAM-1 and ICAM-1 induces strong adhesion by binding Very Late Antigen 4 (VLA4) and Leucocyte Function Antigen-1 (LFA-1). After strong adhesion, leukocyte migrates through endothelial cells into its underlying tissues. Among these molecules, the adhesion molecules (ICAM-1 and VCAM-1), selectins (E-selectin and P-selectins), plasminogen activator inhibitor-1 (PAI-1), and




Figure 1. Alterations of endothelial, systemic coagulation and thrombotic functions due to SARS-CoV-2 infection.⁶

von Willebrand Factor (vWF) are easily obtained from the circulation and well measured by commercial immunoassays.⁵

Adhesion molecules

Endothelial dysfunction can be detected by the increase of circulating cellular adhesion molecules (CAM), including ICAM-1 and VCAM-1. ICAM-1 and VCAM-1 are minimally expressed in inactive endothelial cells. However, these expressions can be augmented by activation of cytokines and endotoxin liposaccharides. Thrombin and histamine stimulations selectively induce endothelial P-selectin expression, whereas cytokines and liposaccharides stimulations induce E-selectins expression.^{7,8}

Studies have shown that endothelial cellular adhesion markers, VCAM-1 and ICAM-1, are increased in COVID-19 patients compared to control group.⁹ These markers were also found significantly higher in severe COVID-19 infections, associating the severity of COVID-19 infection with increase of serum VCAM-1 and ICAM-1.⁹ Likewise, the recovery of severe COVID-19 infections was also associated with a decrease of serum VCAM-1 and ICAM-1 levels over time. Therefore, it could be inferred that according to the study, the increase of endothelial cellular adhesion molecule expression is correlated with the presence of COVID-19 and its severity, as well as indicating a contribution to patient's coagulopathy state.⁹ To further support this, a previous postmortem study had demonstrated a marked endothelial dysfunction in COVID-19 patients with significant increase of ICAM-1.¹⁰

Selectins

Besides VCAM-1 and ICAM-1, endothelial cells also express P-selectin and E-selectin. P-selectin is a cellular adhesion molecule stored inside the endothelium and thrombocytes which would be rapidly deployed into plasma membrane upon activation.¹¹ As infection progresses, P-selectin increases platelet aggregation and platelet-endothelium interaction. Soluble P-selectin (sP-selectin) is produced through enzymatic release of mobilized P-selectins during

inflammation.¹² Experiments with genetically engineered rat in pro-coagulant state revealed high level of sP-selectin expressions. On the other hand, sP-selectin was also proven to be a significant marker of several inflammatory and pro-coagulopathy disorders, including systemic inflammatory responses.^{11,14,15} Most recent study showed increase of sP-selectin in COVID-19 cases, further supporting the theory of COVID-19 being an endothelial dysfunction disorder.¹⁶

E-selectin can also provide as markers of endothelial dysfunction. E-selectin (CD62E) is a leukocyte adhesion molecule that is expressed by activated endothelial cells. Endothelial cells of normal skin and bone marrow or in the case of infantile hemangioma constitutively express E-selectin, in contrast with endothelial cells from other tissues that do not.^{18,19} However, these expressions are tightly regulated by inflammatory cytokines. Soluble E-selectins (sE-selectin) are released during inflammation and has long been proposed as a biomarker of endothelial dysfunction, especially in cases of sepsis.²⁰ Therefore, increased circulating sE-selectin alongside mRNA increase confirm the presence of endothelial dysfunction in COVID-19 as well as a linear correlation with its severity. Smadja *et al.* found in their study that there was an increase of E-Selectin level in COVID-19 patients. This strengthens the theory of endothelial dysfunction in COVID-19.²¹

Plasminogen activator inhibitor-1 (PAI-1)

Procoagulant condition formed by endothelial activation can also be measured from alterations of balance between tissue plasminogen activator (t-PA) and its endogen, plasminogen activator inhibitor-1.²² Plasminogen activator inhibitor-1, also known as endothelial PAI, is a serine protease inhibitor (serpin) which serves as main inhibitor of t-PA and urokinase type plasminogen activator (u-PA), a fibrinolytic agent with recent additional non-fibrinolytic properties reported.²³⁻²⁶ While also secreted by other tissues such as adipose tissue, PAI-1 is mainly produced by endothelium. Increase of PAI-1 is a risk factor of thrombosis and atherosclerosis.^{2,22}

One characteristic of COVID-19 is leukocyte sequestration, especially neutrophils, in the pulmonary microvasculature—which contributes to alveolar injury and infinite inflammation.²⁷ Local pro-inflammatory environment is exaggerated by the formation of NET which results in mass production of pro-inflammatory cytokines.²⁸ These cytokines trigger endothelial cells activations and possibly promotes release of t-PA and PAI-1.^{28,29} Activated endothelial cells expresses raised levels of PAI-1, inhibiting t-PA and u-PA, further instigating hemostasis balance alteration to procoagulant state.¹⁷

In a study by Zuo, PAI-1 level is found to be raised in COVID-19 patients.³⁰ This raise has a significant correlation between PAI-1 as well as circulating absolute neutrophil and calprotectin. This supports the presence of endothelial dysfunction in COVID-19.³⁰ Besides endothelial activation, there is a possibility that direct infection and endothelial cells destruction by SARS-CoV-2 cause a potential release of t-PA and PAI-1.³¹

A study by Kang *et al* revealed that in advanced cases of COVID-19 with severe respiratory dysfunction, the PAI-1 level is significantly higher than sepsis, acute respiratory distress syndrome (ARDS), or even burn cases.³² The significant increase of PAI-1 in severe cases of COVID-19, which are comparable to ARDS, had shown to induce vascular endothelium destruction. SARS-CoV-2 had also shown to directly infect the vascular endothelium, triggering endothelitis which indicates vascular endothelial destruction in patients with COVID-19.³¹ Collectively, this finding suggests that increased PAI-1 level promotes endotheliopathy and coagulopathy in severe cases of COVID-19.³²

Intensive care patients with severe conditions were reported to have significantly increased level of PAI-1 compared to non-intensive care patients. Data from a study by Nougier clearly demonstrated that balance between coagulation and fibrinolysis diminished in COVID-19 patients in significant hypercoagulability state related to hypofibrinolysis caused by high increase of PAI-1 level.³³ In a study by Blasi, plasma PAI-1 level

was found 3.7 times higher in COVID-19 patients compared to control.³⁴

Von Willebrand Factor

Von Willebrand Factor (vWF) is a major multidomain adhesive glycoprotein derived from endothelium, released into circulation by activated endothelial cells.^{35,36} vWF binds with platelet glycoprotein Ib α , α Iib β 3 and endothelial collagen, which activates the platelets and commences platelet aggregation.³⁵ As a carrier of blood clotting factor VIII, vWF also has an important role in clotting cascade.³⁷ vWF is synthesized by endothelial cells and megakaryocytes, which is then stored as ultra-large vWF multimers or multimers with large molecular weight within the endothelial Weibel-Palade bodies or platelet α -granules.

It is well established that pathological alterations of fibrinogen, D-dimer, vWF and P-selectin have important roles in abnormal coagulation and endothelial dysfunction. Both infection and inflammation can increase plasma vWF through activated endothelial cells.^{40,41} The majority of vWF originate from endothelial cells. Endothelial cells vWF (EC-vWF), instead of platelet vWF, critically promotes thrombus formation. In accordance, EC-vWF contributes in vWF-dependent atherogenesis by raising platelet adhesion and vascular inflammation.³⁹ vWF is also a biomarker that is relatively easy to measure.⁴²

In healthy individuals, ACE2 transforms angiotensin-II into angiotensin 1-7, which stimulates endothelial cells to produce nitric oxide (NO). NO aids blood vessels in vasodilation and suppressing platelet aggregation. In COVID-19, SARS-CoV-2 occupies ACE2, subsequently raising Angiotensin-II levels. This further enhance vasoconstriction and reduce blood flow. In this process, vWF stored inside Weibel Palade bodies, increasing formation of blood clots.⁴⁴

Activation of EC-vWF in relation of COVID-19 is known as acute phase protein, released from endothelial cells as inflammatory response.⁴⁵ In this case, its high level indicates a disturbance of endothelial function.⁴⁶ Several studies had been

performed to determine the role of vWF in COVID-19. COVID-19 patients have a documented higher level of vWF, as shown in a study by Blasi.³⁴ An increased activity of vWF and vWF antigen (vWF:Ag) was also found to be significantly raised in a study by Helms, which revealed a well-defined endothelial inflammation with a very high level of vWF:Ag.⁴⁷ Morici *et al.*, in their study, reported a significantly higher vWF in all COVID-19 patients, supporting previous studies.⁴⁸ Katneni *et al.* and several other previous publications similarly reported that vWF level was higher in COVID-19 patients.⁴⁹⁻⁵³ In a study by Panigada, the increase of vWF was reported up to 863 U/dL, further supporting evidences of endothelial dysfunction in COVID-19 patients and its potential use as a biomarker in COVID-19.⁵⁴

Fraser, in his study, measured three thrombosis factors and five markers of endothelial cell injury from the plasma using ELISA, showing significant higher levels in intensive care COVID-19 patients compared to healthy individuals.¹⁶ A study by Rauch has also shown evidences of associations between coagulation marker level on admission, amongst which are factor VIII and vWF, with the severity of COVID-19.⁵⁵ In other words, increased levels of vWF in COVID-19 patients is a potential biomarker associated with severity

of the disease. Previous studies have shown the ability of pulmonary virus to promote platelet-endothelium interactions through upregulation of endothelial ICAM-1, vWF, and fibronectins, culminating ongoing pulmonary injury.⁵⁶

State of hypoxia also triggers vWF expression and its following detachment from endothelial Weibel-Palade bodies.^{57,58} VWF upregulation induced by hypoxia has been associated with the presence of coronary and pulmonary vessels thrombus formation, as well as promoting leukocyte recruitment.⁵⁹ Accordingly, hypoxemia state observed in COVID-19 patients induce prothrombotic condition through upregulation of t-PA inhibitor and stimulation of procoagulant endothelial synthesis, including tissue factor and vWF.^{57,60-62}

VWF is an acute phase response protein released by activated endothelial cells as a reaction to inflammatory stimuli. This increase in activity of vWF and its antigen level contribute to platelet aggregation. In another study, cases of COVID-19 can cause up to 490% raise of vWF and vWF:Ag level. VWF is a main determinant of platelet adhesion after vascular injury and its resulting blood clot. High vWF:Ag level is an independent risk factor for ischemic stroke and myocardial infarction.⁶⁴ Thus, thrombocyte




Figure 2. Thrombus formation in COVID-19.

inhibitors can be considered in prevention of cardiovascular events in COVID-19.⁶⁵

Potential mechanisms of cardiovascular drugs in COVID-19 related endothelial function repair

Various experimental and clinical studies have shown multiple currently used of investigated drugs currently can repair endothelial function, although they may have different structures and mechanisms. Among those drugs are Angiotensin Converting Enzymes (ACE) inhibitor (ACEi), Angiotensin-II Receptor Blocker (ARB), statins and antioxidants.

ACEi and ARB

Evidences have established the role of ACE inhibitors in repairing endothelial functions in animal models with heart failure and in animal models with coronary artery disease.^{66,67} ACE inhibitors elevate endothelial functions by reducing Angiotensin-II, while regulating endothelial Nitric Oxide Synthase (eNOS) and inhibiting production of reactive oxygen species (ROS), giving the drug its protective properties for endothelium.⁶⁸⁻⁷⁰ In the TREND study, quinapril, compared to placebo, repaired endothelial dysfunction in normotensive patients with coronary artery disease.⁶⁷ Administration of ARB has shown to improve endothelial function and demonstrate an overall reduction of inflammatory biomarkers, implying its importance in the pathogenesis of atherosclerosis.⁷¹ A previous meta-analysis showed that ACEi improves endothelial function in patients with endothelial dysfunction of multiple causes.⁷²

However, despite available studies mentioned, there has not been any consensus related to COVID-19 related endothelial dysfunction and ACEi or ARB as its potential clinical approach. Several available meta-analysis regarding effectivity of ACE inhibitor or ARB administration for endothelial dysfunction in COVID-19 still have differing conclusions.^{73,74} Further investigation is necessary, especially considering that ACE-2 receptor is a binding site of SARS-CoV-2.

Statins

Beneficial effects of statin in endothelial function involves multiple mechanisms. Statins improve endothelial dysfunction due to their LDL lowering properties, considering LDL and OxLDL capability in reducing eNOS.^{75,76} Statin improves NO bioactivity by activating eNOS through PI3K/Akt signaling pathway.⁸⁰ The benefits of statin for endothelial function is also related with its anti-inflammation and antioxidant properties.⁷⁷ Statins provide direct antioxidant effects to LDL through reducing LDL electronegative forms.^{78,79}

In 1995, randomized control trial stated that lovastatin can return endothelial function of a coronary artery.⁸¹ Atorvastatin was shown to reduce pro-inflammatory cytokines (TNF- α , IL-1 and IL-6), ICAM-1 and C-reactive protein (CRP) in hypercholesterolemic patients.⁸² One study showed that simvastatin produces significant reduction in endothelial dysfunction markers, inflammation, oxidative stress and endothelial apoptosis; in this study, CRP reduction seemed to have been in relation to the lipid lowering property of simvastatin.⁸³ A meta-analysis showed that statin therapy is associated with significant improvement of both coronary and peripheral endothelial functions.⁸⁴ Otherwise, statin elevate endothelial progenitor cells circulation, which contribute in the long-term effects of statin in endothelial function.⁸⁵ The combination of ACE inhibitors and statin therapy have also been demonstrating a relaxation effects in coronary vessels, which is largely dependent on endothelial function through NO production.⁸⁶

Several studies reported benefits of statins in COVID-19. Masana *et al.* in their study revealed fewer deaths reported in statin-administered group compared to the non-statin-administered group. It has also been stated that statin therapy should not be halted in hospitalized patients due to lower death rate of SARS-CoV-2 infection patients who took statins before hospitalization.⁸⁷ An observational study by Omar *et al.* also reported that, in diabetic patients with COVID-19, statin users have a 12% lower chance of death during hospitalization compared to those who did not.⁸⁸ A retrospective cohort in Singapore

linked independent use of statin with lower ICU admission rate.⁸⁸ Evidence by Sophia *et al* suggested beneficial use of continuous statin for hospitalized COVID-19 patients, as it correlates with lower chance of invasive mechanical ventilation support.⁸⁹ However, there has not been a randomized controlled trial which further supports the use of statins in COVID-19.

Antioxidant

Several substances such as vitamin C, vitamin E and N-acetylcysteine provide an antioxidant effect through different mechanisms. Vitamin C protects the endothelium by eliminating superoxide, which in turn prevents NO decomposition, lipid peroxidation, platelet and neutrophil activation, as well as upregulation of adhesion molecules.^{90,91} Vitamin C scavenges reactive nitrogen species yielded by peroxidase and inhibits myeloperoxidase/H₂O₂.⁹² Vitamin E acts as lipid soluble antioxidant, clearing radical hydroperoxyl in lipid environment.⁹³ Meanwhile, the effects of N-acetylcysteine in endothelial dysfunction are related to inhibitions of NADPH oxidase expression, leukocyte adhesion and inflammatory cytokine secretion.⁹⁴ N-acetylcysteine also prevents platelet aggregation, which largely depends on vWF and collagen binding in human plasma, and inhibits upregulation of Caveolin-1 as well as strengthening endothelial barrier function in rats.⁹⁵ Further investigations may provide useful information regarding antioxidants in the management of endothelial dysfunction related to COVID-19.

CONCLUSION

SARS-CoV-2 infection, which has caused a global pandemic, involved various clinical manifestations and underlying mechanisms. This virus invades hosts by occupying ACE-2 receptors in endothelium, which signify the important role of endothelium in this disease. Available studies have demonstrated the increase of several endothelial biomarkers such as ICAM-1, VCAM-1, E-Selectin, P-Selectin, PAI-1 and vWF in COVID-19 patients, which

supports the presence of endothelial dysfunction in COVID-19, and could further provide helpful information for the detection of endothelial dysfunction in COVID-19. These findings may also explain the higher chance of individuals with comorbidity to contract COVID-19, with increased severity. Further studies investigating NO levels in COVID-19 is necessary to confirm COVID-19 related endothelial dysfunction.

The use of drugs that has been established to improve endothelial functions may be useful as a baseline therapy in COVID-19. It has also become a point of interest whether a cured COVID-19 patient has an increased risk of cardiovascular diseases in the future due to their endothelial impairment. This topic is potential for future research, and may provide helpful insight in prevention of cardiovascular system events after COVID-19.

ACKNOWLEDGEMENT

Authors would like to thank Department of Cardiology and Vascular Medicine, Airlangga University Faculty of Medicine for the support in conducting this study. The authors' responsibilities were as follows—AA: determined and conceptualized the framework of studied topic, provided guidance in the design of manuscript and edit manuscript content; RR: conducted literature research and wrote the manuscript; PMH: translated, edited and had primary responsibility for the final content of the manuscript; and all authors: read and approved the final manuscript.

CONFLICT OF INTEREST

None of the authors had a conflict of interest in this study.

REFERENCES

1. Galley HF, Webster NR. Physiology of the endothelium. *Br J Anaesth* [Internet]. 2004;93(1):105–13. Available from: <http://dx.doi.org/10.1093/bja/aeh163>

2. Wang M, Hao H, Leeper NJ, Zhu L. Thrombotic Regulation From the Endothelial Cell Perspectives. *Arterioscler Thromb Vasc Biol.* 2018;38(6):e90–5.
3. Siddiqi HK, Libby P, Ridker PM. COVID-19 – A vascular disease. *Trends Cardiovasc Med.* 2020;(January).
4. Perico L, Benigni A, Casiraghi F, Ng LFP, Renia L, Remuzzi G. Immunity, endothelial injury and complement-induced coagulopathy in COVID-19. *Nat Rev Nephrol* [Internet]. 2020; Available from: <http://dx.doi.org/10.1038/s41581-020-00357-4>
5. Deanfield JE, Halcox JP, Rabelink TJ. Endothelial function and dysfunction: Testing and clinical relevance. *Circulation.* 2007;115(10):1285–95.
6. Halcox JPI. Endothelial Dysfunction. *Prim Auton Nerv Syst.* 2012;319–24.
7. McLean RC, Blumenthal RS. Inflammatory markers and the risk of coronary heart disease. *Cardiol Rev.* 2005;22(6):41–2.
8. Ridker PM, Brown NJ, Vaughan DE, Harrison DG, Mehta JL. Established and emerging plasma biomarkers in the prediction of first atherothrombotic events. *Circulation.* 2004;109(25):6–19.
9. Tong M, Jiang Y, Xia D, Xiong Y, Zheng Q, Chen F, et al. Elevated Expression of Serum Endothelial Cell Adhesion Molecules in COVID-19 Patients. *J Infect Dis.* 2020;222(6):894–8.
10. Nagashima S, Mendes MC, Camargo Martins AP, Borges NH, Godoy TM, Miggiolaro AFRDS, et al. Endothelial dysfunction and thrombosis in patients with COVID-19 - Brief report. *Arterioscler Thromb Vasc Biol.* 2020;(October):2404–7.
11. Furie B. P-selectin and blood coagulation: It's not only about inflammation any more. *Arterioscler Thromb Vasc Biol.* 2005;25(5):877–8.
12. Ishiwata N, Takio K, Katayama M, Watanabe K, Titani K, Ikeda Y, et al. Alternatively spliced isoform of P-selectin is present in vivo as a soluble molecule. *J Biol Chem.* 1994;269(38):23708–15.
13. André P, Hartwell D, Hrachovinová I, Saffaripour S, Wagner DD. Pro-coagulant state resulting from high levels of soluble P-selectin in blood. *Proc Natl Acad Sci U S A.* 2000;97(25):13835–40.
14. Schrijver IT, Kemperman H, Roest M, Kesecioglu J, de Lange DW. Soluble P-selectin as a biomarker for infection and survival in patients with a systemic inflammatory response syndrome on the intensive care unit. *Biomark Insights.* 2017;12.
15. Katayama M, Handa M, Araki Y, Ambo H, Kawai Y, Watanabe K, et al. Soluble P-selectin is present in normal circulation and its plasma level is elevated in patients with thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome. *Br J Haematol.* 1993;84(4):702–10.
16. Fraser DD, Patterson EK, Slessarev M, Gill SE, Martin C, Daley M, et al. Endothelial Injury and Glycocalyx Degradation in Critically Ill Coronavirus Disease 2019 Patients: Implications for Microvascular Platelet Aggregation. *Crit Care Explor.* 2020;2(9):e0194.
17. Shapiro NI, Schuetz P, Yano K, Sorasaki M, Parikh SM, Jones AE, et al. The association of endothelial cell signaling, severity of illness, and organ dysfunction in sepsis. *Crit Care.* 2010;14(5).
18. Silva M, Videira PA, Sackstein R. E-selectin ligands in the human mononuclear phagocyte system: Implications for infection, inflammation, and immunotherapy. *Front Immunol.* 2018;8(Jan).
19. Smadja DM, Mulliken JB, Bischoff J. E-selectin mediates stem cell adhesion and formation of blood vessels in a murine model of infantile hemangioma. *Am J Pathol* [Internet]. 2012;181(6):2239–47. Available from: <http://dx.doi.org/10.1016/j.ajpath.2012.08.030>
20. Cummings CJ, Sessler CN, Beall LD, Fisher BJ, Best AM, Fowler AA. Soluble E-selectin levels in sepsis and critical illness: Correlation with infection and hemodynamic dysfunction. *Am J Respir Crit Care Med.* 1997;156(2 I):431–7.
21. Smadja DM, Guerin CL, Chocron R, Yatim N, Boussier J, Gendron N, et al. Angiopoietin-2 as a marker of endothelial activation is a good predictor factor for intensive care unit admission of COVID-19 patients. *Angiogenesis* [Internet]. 2020;23(4):611–20. Available from: <https://doi.org/10.1007/s10456-020-09730-0>
22. Vaughan DE. PAI-1 and atherothrombosis. *J Thromb Haemost.* 2005;3(8):1879–83.
23. Fay WP, Korthuis RJ. No Sweetie Pie: Newly Uncovered Role for PAI-1 in Inflammatory Responses to Ischemia/Reperfusion. *Physiol Behav.* 2019;176(3):139–48.
24. Praetner M, Zuchtriegel G, Holzer M, Uhl B, Schaubächer J, Mittmann L, et al. Plasminogen Activator Inhibitor-1 Promotes Neutrophil Infiltration and Tissue Injury on Ischemia-Reperfusion. *Arterioscler Thromb Vasc Biol.* 2018;38(4):829–42.
25. Vaughan DE, Rai R, Khan SS, Eren M, Ghosh AK. Plasminogen Activator Inhibitor-1 Is a Marker and a Mediator of Senescence. *Arterioscler Thromb Vasc Biol.* 2017;37(8):1446–52.
26. Ji Y, Weng Z, Fish P, Goyal N, Luo M, Myears SP, et al. Pharmacological Targeting of Plasminogen Activator Inhibitor-1 Decreases Vascular Smooth Muscle Cell Migration and Neointima Formation. *Physiol Behav* [Internet]. 2017;176(10):139–48. Available from: [file:///C:/Users/Carla Carolina/Desktop/Artigos para acrescentar na qualificação/The impact of birth weight on cardiovascular disease risk in the.pdf](file:///C:/Users/Carla%20Carolina/Desktop/Artigos%20para%20acrescentar%20na%20qualifica%C3%A7%C3%A3o/The%20impact%20of%20birth%20weight%20on%20cardiovascular%20disease%20risk%20in%20the.pdf)
27. Whyte CS, Morrow GB, Mitchell JL, Chowdary P, Mutch NJ. Fibrinolytic abnormalities in acute respiratory distress syndrome (ARDS) and versatility of thrombolytic drugs to treat COVID-19. *J Thromb Haemost.* 2020;18(7):1548–55.
28. Fox SE, Akmatbekov A, Harbert JL, Li G, Brown JQ, Heide RS Vander. Pulmonary and cardiac pathology in African American patients with COVID-19: an autopsy series from New Orleans. *Lancet Respir Med* 2020;(January):19–21.

29. Meltzer ME, Lisman T, De Groot PG, Meijers JCM, Le Cessie S, Doggen CJM, et al. Venous thrombosis risk associated with plasma hypofibrinolysis is explained by elevated plasma levels of TAFI and PAI-1. *Blood*. 2010;116(1):113–21.
30. Zuo Y, Wamock M, Harbaugh A, Yalavarthi S, Gockman K, Lawrence DA. Plasma tissue plasminogen activator and plasminogen activator inhibitor-1 in hospitalized COVID-19 patients. *Sci Rep*. 2020;1–13.
31. Varga Z, Flammer A, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, et al. Endothelial cell infection and endothelitis in COVID-19. *Ann Oncol*. 2020;(January).
32. Kang S, Tanaka T, Inoue H, Ono C, Hashimoto S, Kioi Y, et al. IL-6 trans-signaling induces plasminogen activator inhibitor-1 from vascular endothelial cells in cytokine release syndrome. *Proc Natl Acad Sci U S A*. 2020;117(36):22351–6.
33. Nougier C, Benoit R, Simon M, Desmurs-Clavel H, Marcotte G, Argaud L, et al. Hypofibrinolytic state and high thrombin generation may play a major role in SARS-CoV2 associated thrombosis. *J Thromb Haemost*. 2020;18(9):2215–9.
34. Blasi A, von Meijenfeldt FA, Adelmeijer J, Calvo A, Ibañez C, Perdomo J, et al. In vitro hypercoagulability and ongoing in vivo activation of coagulation and fibrinolysis in COVID-19 patients on anticoagulation. *J Thromb Haemost*. 2020;18(10):2646–53.
35. Löf A, Müller JP, Brehm MA. A biophysical view on von Willebrand factor activation. *J Cell Physiol*. 2018;233(2):799–810.
36. Da Silva ML, Cutler DF. Von Willebrand factor multimerization and the polarity of secretory pathways in endothelial cells. *Blood*. 2016;128(2):277–85.
37. Butera D, Passam F, Ju L, Cook KM, Woon H, Aponte-Santamaría C, et al. Autoregulation of von Willebrand factor function by a disulfide bond switch. *Sci Adv*. 2018;4(2).
38. Verhenne S, Denorme F, Libbrecht S, Vandenbulcke A, Pareyn I, Deckmyn H, et al. Platelet-derived VWF is not essential for normal thrombosis and hemostasis but fosters ischemic stroke injury in mice. *Blood*. 2015;126(14):1715–22.
39. Doddapattar P, Dhanesha N, Chorawala MR, Tinsman C, Jain M, Nayak MK, et al. Endothelial cell-derived von Willebrand factor, but not platelet-derived, promotes atherosclerosis in Apoe-deficient mice. 2019;38(3):319–35.
40. Kayal S, Jaïs JP, Aguiñ N, Chaudière J, Labrousse J. Elevated circulating E-selectin, intercellular adhesion molecule 1, and von Willebrand factor in patients with severe infection. *Am J Respir Crit Care Med*. 1998;157(3 PART I):776–84.
41. Meyer AA, Kundt G, Steiner M, Schuff-Werner P, Kienast W. Impaired flow-mediated vasodilation, carotid artery intima-media thickening, and elevated endothelial plasma markers in obese children: The impact of cardiovascular risk factors. *Pediatrics*. 2006;117(5):1560–7.
42. Mannucci PM. von Willebrand Factor A Marker of Endothelial Damage? Greece Rome.?? *Arterioscler Thromb Vasc Biol*. 1967;14(2):188–98.
43. Grobler C, Maphumulo SC, Grobbelaar LM, Bredenkamp JC, Laubscher GJ, Lourens PJ, et al. Covid-19: The rollercoaster of fibrin(ogen), d-dimer, von willebrand factor, p-selectin and their interactions with endothelial cells, platelets and erythrocytes. *Int J Mol Sci*. 2020;21(14):1–25.
44. Iba T, Levy JH, Connors JM, Warkentin TE, Thachil J, Levi M. The unique characteristics of COVID-19 coagulopathy. *Crit Care*. 2020;24(1):4–11.
45. Kawecki C, Lenting PJ, Denis C V. von Willebrand factor and inflammation. *J Thromb Haemost*. 2017;15(7):1285–94.
46. Breakey N, Escher R. D-dimer and mortality in COVID-19: A self-fulfilling prophecy or a pathophysiological clue? *Swiss Med Wkly*. 2020;150(21–22):1–7.
47. Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* [Internet]. 2020;46(6):1089–98. Available from: <https://doi.org/10.1007/s00134-020-06062-x>
48. Morici N, Bottioli M, Fumagalli R, Marini C, Cattaneo M. Role of von Willebrand Factor and ADAMTS-13 in the Pathogenesis of Thrombi in SARS-CoV-2 Infection: Time to Rethink. *Thromb Haemost*. 2020;120(9):1339–41.
49. Bazzan M, Montaruli B, Sciascia S, Cosseddu D, Norbiato C, Roccatello D. Low ADAMTS 13 plasma levels are predictors of mortality in COVID-19 patients. *Intern Emerg Med* [Internet]. 2020;15(5):861–3. Available from: <https://doi.org/10.1007/s11739-020-02394-0>
50. Huisman A, Beun R, Sikma M, Westerink J, Kusadasi N. Involvement of ADAMTS13 and von Willebrand factor in thromboembolic events in patients infected with SARS-CoV-2. *Int J Lab Hematol*. 2020;42(5):e211–2.
51. Adam E, Zacharowski K, Miesbach W. A comprehensive assessment of the coagulation profile in critically ill COVID-19 patients. *Thromb Res*. 2020;(January).
52. Latimer G, Corriveau C, DeBiasi RL, Jantusch B, Delaney M, Jacquot C, et al. Cardiac dysfunction and thrombocytopenia-associated multiple organ failure inflammation phenotype in a severe paediatric case of COVID-19. *Lancet Child Adolesc Health*. 2020;(January):19–21.
53. Katmeni UK, Alexaki A, Hunt RC, Schiller T, Dicuccio M, Buehler PW, et al. Coagulopathy and thrombosis as a result of severe COVID-19 infection: A microvascular focus. *Thromb Haemost*. 2020;
54. Panigada M, Bottino N, Tagliabue P, Grasselli G, Novembrino C, Chantarangkul V, et al.

- Hypercoagulability of COVID-19 patients in intensive care unit: A report of thromboelastography findings and other parameters of hemostasis. *J Thromb Haemost.* 2020;18(7):1738–42.
55. Rauch A, Labreuche J, Lassalle F, Goutay J, Caplan M, Charbonnier L, et al. Coagulation biomarkers are independent predictors of increased oxygen requirements in COVID-19. *J Thromb Haemost.* 2020;(August):1–12.
 56. Sugiyama MG, Gamage A, Zyla R, Armstrong SM, Advani S, Advani A, et al. Influenza Virus Infection Induces Platelet-Endothelial Adhesion Which Contributes to Lung Injury. *J Virol.* 2016;90(4):1812–23.
 57. Mojiri A, Nakhai-Nejad M, Phan WL, Kulak S, Radziwon-Balicka A, Jurasz P, et al. Hypoxia results in upregulation and de novo activation of von willebrand factor expression in lung endothelial cells. *Arterioscler Thromb Vasc Biol.* 2013;33(6):1329–38.
 58. Pinsky DJ, Naka Y, Liao H, Oz MC, Wagner DD, Mayadas TN, et al. Hypoxia-induced exocytosis of endothelial cell weibel-palade bodies: A mechanism for rapid neutrophil recruitment after cardiac preservation. *J Clin Invest.* 1996;97(2):493–500.
 59. Mojiri A, Alavi P, Lorenzana Carrillo MA, Nakhai-Nejad M, Sergi CM, Thebaud B, et al. Endothelial cells of different organs exhibit heterogeneity in von Willebrand factor expression in response to hypoxia. *Atherosclerosis* [Internet]. 2019;282(June 2018):1–10. Available from: <https://doi.org/10.1016/j.atherosclerosis.2019.01.002>
 60. Matsuura Y, Yamashita A, Iwakiri T, Sugita C, Okuyama N, Kitamura K, et al. Vascular wall hypoxia promotes arterial thrombus formation via augmentation of vascular thrombogenicity. *Thromb Haemost.* 2015;114(1):158–72.
 61. Ogawa S, Clauss M, Kuwabara K, Shreeniwars R, Butura C, Koga S, et al. Hypoxia induces endothelial cell synthesis of membrane-associated proteins. *Proc Natl Acad Sci U S A.* 1991;88(21):9897–901.
 62. Fearn C, Loskutoff DJ. Induction of plasminogen activator inhibitor 1 gene expression in murine liver by lipopolysaccharide: Cellular localization and role of endogenous tumor necrosis factor- α . *Am J Pathol.* 1997;150(2):579–90.
 63. Gragnano F, Sperlongano S, Golia E, Natale F, Bianchi R, Crisci M, et al. The Role of von Willebrand Factor in Vascular Inflammation: From Pathogenesis to Targeted Therapy. *Mediators Inflamm.* 2017;2017.
 64. Andersson HM, Siegerink B, Luken BM, Crawley JTB, Algra A, Lane DA, et al. High VWF, low ADAMTS13, and oral contraceptives increase the risk of ischemic stroke and myocardial infarction in young women. *Blood.* 2012;119(6):1555–60.
 65. Hoechter DJ, Becker-pennrich A, Langrehr J, Bruegel M, Zwissler B, Schaefer S. Higher procoagulatory potential but lower DIC score in COVID-19 ARDS patients compared to non-COVID-19 ARDS patients. *Thromb Res* 2020;(January).
 66. Varin R, Mulder P, Tamion F, Richard V, Henry JP, Lallemand F, et al. Improvement of endothelial function by chronic angiotensin-converting enzyme inhibition in heart failure: Role of nitric oxide, prostanoids, oxidant stress, and bradykinin. *Circulation.* 2000;102(3):351–6.
 67. Mancini GBJ, Henry GC, Macaya C, O'Neill BJ, Pucillo AL, Carere RG, et al. Angiotensin-converting enzyme inhibition with quinapril improves endothelial vasomotor dysfunction in patients with coronary artery disease: The TREND (Trial on Reversing ENdothelial Dysfunction) study. *Circulation.* 1996;94(3):258–65.
 68. Fujii M, Wada A, Tsutamoto T, Ohnishi M, Isono T, Kinoshita M. Bradykinin improves left ventricular diastolic function under long-term angiotensin-converting enzyme inhibition in heart failure. *Hypertension.* 2002;39(5):952–7.
 69. Bachetti T, Comini L, Pasini E, Cargnoni A, Curello S, Ferrari R. Ace-inhibition with quinapril modulates the nitric oxide pathway in normotensive rats. *J Mol Cell Cardiol.* 2001;33(3):395–403.
 70. Mukai Y, Shimokawa H, Higashi M, Morikawa K, Matoba T, Hiroki J, et al. Inhibition of renin-angiotensin system ameliorates endothelial dysfunction associated with aging in rats. *Arterioscler Thromb Vasc Biol.* 2002;22(9):1445–50.
 71. Sola S, Mir MQS, Cheema FA, Khan-Merchant N, Menon RG, Parthasarathy S, et al. Irbesartan and lipoic acid improve endothelial function and reduce markers of inflammation in the metabolic syndrome: Results of the Irbesartan and Lipoic Acid in Endothelial Dysfunction (ISLAND) study. *Circulation.* 2005;111(3):343–8.
 72. Shahin Y, Khan JA, Samuel N, Chetter I. Angiotensin converting enzyme inhibitors effect on endothelial dysfunction: A meta-analysis of randomised controlled trials. *Atherosclerosis* [Internet]. 2011;216(1):7–16. Available from: <http://dx.doi.org/10.1016/j.atherosclerosis.2011.02.044>
 73. Hasan SS, Kow CS, Hadi MA, Zaidi STR, Merchant HA. Mortality and Disease Severity Among COVID-19 Patients Receiving Renin-Angiotensin System Inhibitors: A Systematic Review and Meta-analysis. *Am J Cardiovasc Drugs* [Internet]. 2020;20(6):571–90. Available from: <https://doi.org/10.1007/s40256-020-00439-5>
 74. Liu X, Long C, Xiong Q, Chen C, Ma J, Su Y, et al. Association of angiotensin converting enzyme inhibitors and angiotensin II receptor blockers with risk of COVID-19, inflammation level, severity, and death in patients with COVID-19: A rapid systematic review and meta-analysis. *Clin Cardiol.* 2020;
 75. Martínez-gonzález J, Raposo B, Rodríguez C, Badimon L. Downregulation by Atherogenic Levels of Native LDLs Balance Between Transcriptional and

- Posttranscriptional Regulation. *Arterioscler Thromb Vasc Biol.* 2001;804–9.
76. Laufs U, Fata V La, Plutzky J, Liao JK. Upregulation of Endothelial Nitric Oxide Synthase by HMG CoA Reductase Inhibitors. *Circulation.* Ulrich. 1998;1129–35.
 77. Bonetti PO, Lerman LO, Napoli C, Lerman A. Statin effects beyond lipid lowering - Are they clinically relevant? *Eur Heart J.* 2003;24(3):225–48.
 78. Aviram M, Hussein O, Rosenblat M, Schlezinger S, Hayek T, Keidar S. Interactions of platelets, macrophages, and lipoproteins in hypercholesterolemia: Antiatherogenic effects of HMG-CoA reductase inhibitor therapy. *J Cardiovasc Pharmacol.* 1998;31(1):39–45.
 79. Sánchez-Quesada JL, Otal-Entraigas C, Franco M, Jorba O, González-Sastre F, Blanco-Vaca F, et al. Effect of simvastatin treatment on the electronegative low-density lipoprotein present in patients with heterozygous familial hypercholesterolemia. *Am J Cardiol.* 1999;84(6):655–9.
 80. Kureishi Y, Luo Z, Shiojima I, Bialik A, Fulton D, David J, et al. The HMG-CoA reductase inhibitor simvastatin activates the protein kinase Akt and promotes angiogenesis in normocholesterolemic animals. *2010;6(9):1004–10.*
 81. Anderson TJ, Meredith IT, Yeung AC, Frei B, Selwyn AP, Ganz P. The Effect of Cholesterol-Lowering and Antioxidant Therapy on Endothelium-Dependent Coronary Vasomotion. *J Occup Environ Med.* 1996;38(5):468.
 82. Ascer E, Bertolami MC, Venturinelli ML, Buccheri V, Souza J, Nicolau JC, et al. Atorvastatin reduces proinflammatory markers in hypercholesterolemic patients. *Atherosclerosis.* 2004;177(1):161–6.
 83. Kirmizis D, Papagianni A, Dogrammatzi F, Skoura L, Belechri AM, Alexopoulos E, et al. Effects of simvastatin on markers of inflammation, oxidative stress and endothelial cell apoptosis in patients on chronic hemodialysis. *J Atheroscler Thromb.* 2010;17(12):1256–65.
 84. Reriani MK, Dunlay SM, Gupta B, West CP, Rihal CS, Lerman LO, et al. Effects of statins on coronary and peripheral endothelial function in humans: A systematic review and metaanalysis of randomized controlled trials. *Eur J Cardiovasc Prev Rehabil.* 2011;18(5):704–16.
 85. Dimmeler S, Aicher A, Vasa M, Mildner-Rihm C, Adler K, Tiemann M, et al. Increase endothelial progenitor cells via the PI 3-kinase / Akt pathway. *J Clin Invest.* 2001;108(3):365–6.
 86. Tiefenbacher CP, Friedrich S, Bleeke T, Vahl C, Chen X, Niroomand F. ACE inhibitors and statins acutely improve endothelial dysfunction of human coronary arterioles. *Am J Physiol - Hear Circ Physiol.* 2004;286(4 55-4):1425–32.
 87. Masana L, Correig E, Borjabad CR, Anoro E, Arroyo JA, Al E. Effect of Statin Therapy on SARS-CoV-2 Infection-related Mortality in Hospitalized Patients. *Akrab Juara [Internet].* 2020;5(1):43–54. Available from: <http://www.akrabjuara.com/index.php/akrabjuara/article/view/919>
 88. Saeed O, Castagna F, Agalliu I, Xue X, Patel SR, Rochlani Y, et al. Statin Use and In-Hospital Mortality in Diabetics with COVID-19. *J Am Heart Assoc.* 2020; 15(24): 1-12
 89. Song SL, Hays SB, Panton CE, Mylona EK, Kalligeros M, Shehadeh F, et al. Statin use is associated with decreased risk of invasive mechanical ventilation in COVID-19 patients: A preliminary study. *Pathogens.* 2020;9(9):1–9.
 90. Heitzer T, Just H, Münzel T. Antioxidant Vitamin C Improves Endothelial Dysfunction in Chronic Smokers. *Circulation.* 1996;94(1):6–9.
 91. Matsumoto T, D'Uscio L V., Eguchi D, Akiyama M, Smith LA, Katusic ZS. Protective effect of chronic vitamin C treatment on endothelial function of apolipoprotein E-deficient mouse carotid artery. *J Pharmacol Exp Ther.* 2003;306(1):103–8.
 92. Carr AC, McCall MR, Frei B. Oxidation of LDL by myeloperoxidase and reactive nitrogen species: Reaction pathways and antioxidant protection. *Arterioscler Thromb Vasc Biol.* 2000;20(7):1716–23.
 93. Traber MG, Stevens JF. Vitamins C and E: Beneficial effects from a mechanistic perspective. *Free Radic Biol Med.* 2011;51(5):1000–13.
 94. Sciola MG, Bielli A, Agostinelli S, Tarquini C, Arcuri G, Ferlosio A, et al. Antioxidant treatment prevents serum deprivation- and TNF- α -induced endothelial dysfunction through the inhibition of NADPH oxidase 4 and the restoration of β -oxidation. *J Vasc Res.* 2014;51(5):327–37.
 95. Chen J, Reheman A, Gushiken FC, Nolasco L, Fu X, Moake JL, et al. N-acetylcysteine reduces the size and activity of von Willebrand factor in human plasma and mice. *J Clin Invest.* 2011;121(2):593–603.
 96. Shapiro, N.I., Schuetz, P., Yano, K. et al. The association of endothelial cell signaling, severity of illness, and organ dysfunction in sepsis. *Crit Care* 14, R182 (2010). <https://doi.org/10.1186/cc9290>

COVID-19 and Endothelial Dysfunction Biomarkers and Potential Drug Mechanisms

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nektarios Barabutis, Vikramjit Khangoora, Paul E. Marik, John D. Catravas. "Hydrocortisone and Ascorbic Acid Synergistically Prevent and Repair Lipopolysaccharide-Induced Pulmonary Endothelial Barrier Dysfunction", Chest, 2017
Publication 1%
- 2 H. F. Galley. "Physiology of the endothelium", British Journal of Anaesthesia, 07/01/2004
Publication 1%
- 3 TERRY LEE, A CAU, MP CHENG, A LEVIN et al. "ANGIOTENSIN RECEPTOR BLOCKERS AND ANGIOTENSIN CONVERTING ENZYME INHIBITORS IN COVID-19 –META-ANALYSIS/META-REGRESSION ADJUSTED FOR CONFOUNDING FACTORS", CJC Open, 2021
Publication <1%
- 4 D.J. Hoechter, A. Becker-Pennrich, J. Langrehr, M. Bruegel, B. Zwissler, S. Schaefer, M. Spannagl, L.C. Hinske, M. Zoller. "Higher <1%

procoagulatory potential but lower DIC score in COVID-19 ARDS patients compared to non-COVID-19 ARDS patients", Thrombosis Research, 2020

Publication

5

Danielle Brain, Alex Plant-Hatley, Bethany Heaton, Usman Arshad et al. "Drug delivery systems as immunomodulators for therapy of infectious disease: Relevance to COVID-19", Advanced Drug Delivery Reviews, 2021

Publication

<1 %

6

Hasan K. Siddiqi, Peter Libby, Paul M Ridker. "COVID-19 – A vascular disease", Trends in Cardiovascular Medicine, 2021

Publication

<1 %

7

Zhangjing Ma, Kevin Y. Yang, Yu Huang, Kathy O. Lui. "Endothelial contribution to COVID-19: an update on mechanisms and therapeutic implications", Journal of Molecular and Cellular Cardiology, 2021

Publication

<1 %

8

Nirav Dhanesha, Prem Prakash, Prakash Doddapattar, Ira Khanna et al. "Endothelial Cell-Derived von Willebrand Factor Is the Major Determinant That Mediates von Willebrand Factor-Dependent Acute Ischemic Stroke by Promoting Postischemic Thrombo-

<1 %

Inflammation", Arteriosclerosis, Thrombosis, and Vascular Biology, 2016

Publication

9	surgicalcasereports.springeropen.com Internet Source	<1 %
10	Minerva-access.unimelb.edu.au Internet Source	<1 %
11	coek.info Internet Source	<1 %
12	plus.mcmaster.ca Internet Source	<1 %
13	rxivist.org Internet Source	<1 %
14	www.scielo.br Internet Source	<1 %
15	Prakash Doddapattar, Nirav Dhanesha, Mehul R. Chorawala, Chandler Tinsman et al. "Endothelial Cell-Derived Von Willebrand Factor, But Not Platelet-Derived, Promotes Atherosclerosis in Apolipoprotein E-Deficient Mice", Arteriosclerosis, Thrombosis, and Vascular Biology, 2018 Publication	<1 %
16	atm.amegroups.com Internet Source	<1 %

repository.upnvj.ac.id

17	Internet Source	<1 %
18	turkishjournalofurology.com Internet Source	<1 %
19	uokerbala.edu.iq Internet Source	<1 %
20	waspada.id Internet Source	<1 %
21	Annabel Blasi, Fien A. Meijenfeldt, Jelle Adelmeijer, Andrea Calvo et al. " hypercoagulability and ongoing activation of coagulation and fibrinolysis in COVID - 19 patients on anticoagulation ", Journal of Thrombosis and Haemostasis, 2020 Publication	<1 %
22	Dohke, T.. "Proteomic Analysis Reveals Significant Alternations of Cardiac Small Heat Shock Protein Expression in Congestive Heart Failure", Journal of Cardiac Failure, 200602 Publication	<1 %
23	circ.ahajournals.org Internet Source	<1 %
24	www.kompasiana.com Internet Source	<1 %
25	www.thieme-connect.de Internet Source	<1 %

26

Abdollah Jafarzadeh, Maryam Nemati, Sara Jafarzadeh. "Contribution of STAT3 to the pathogenesis of COVID-19", *Microbial Pathogenesis*, 2021

Publication

<1 %

27

Hong Yu, Yajun Liu, Meifang Wang, Ricardo J. Restrepo et al. "Myeloperoxidase instigates proinflammatory responses in a cecal ligation and puncture rat model of sepsis", *American Journal of Physiology-Heart and Circulatory Physiology*, 2020

Publication

<1 %

28

Irina Portier, Robert A. Campbell, Frederik Denorme. "Mechanisms of immunothrombosis in COVID-19", *Current Opinion in Hematology*, 2021

Publication

<1 %

29

Isabelle Bernard, Daniel Limonta, Lara K. Mahal, Tom C. Hobman. "Endothelium Infection and Dysregulation by SARS-CoV-2: Evidence and Caveats in COVID-19", *Viruses*, 2020

Publication

<1 %

30

Justin Marley, Nisha Marley. "Characterising COVID-19 as a Viral Clotting Fever: A Mixed Methods Scoping Review", *Cold Spring Harbor Laboratory*, 2020

Publication

<1 %

31

Luca Perico, Ariela Benigni, Giuseppe Remuzzi. "Angiotensin-converting enzyme 2: from a vasoactive peptide to the gatekeeper of a global pandemic", *Current Opinion in Nephrology & Hypertension*, 2021

Publication

<1 %

32

Roberto F. Nicosia, Giovanni Ligresti, Nunzia Caporarello, Shreeram Akilesh, Domenico Ribatti. "COVID-19 Vasculopathy: Mounting Evidence for an Indirect Mechanism of Endothelial Injury", *The American Journal of Pathology*, 2021

Publication

<1 %

33

Sean X. Gu, Tarun Tyagi, Kanika Jain, Vivian W. Gu et al. "Thrombocytopeny and endotheliopathy: crucial contributors to COVID-19 thromboinflammation", *Nature Reviews Cardiology*, 2020

Publication

<1 %

34

Sergey Brankovich Bolevich, Peter Frantzevich Litvitsky, Sergei Vitalievich Grachev, Sergey Ivanovich Vorobyev et al. "Fundamental Basis of COVID-19 Pathogenesis", *Serbian Journal of Experimental and Clinical Research*, 2020

Publication

<1 %

35

Yilmaz, Banu, and Fusun Toruner. "277", *Endothelium Molecular Aspects of Metabolic Disorders*, 2013.

<1 %

36	digitalcollections.ohsu.edu Internet Source	<1 %
37	docero.net Internet Source	<1 %
38	epdf.tips Internet Source	<1 %
39	hrcak.srce.hr Internet Source	<1 %
40	pubag.nal.usda.gov Internet Source	<1 %
41	sites.kowsarpub.com Internet Source	<1 %
42	www.dovepress.com Internet Source	<1 %
43	www.gahmj.com Internet Source	<1 %
44	www.qscience.com Internet Source	<1 %
45	www.thieme-connect.com Internet Source	<1 %
46	Alpo Vuorio, Petri T. Kovanen. "Statins as Adjuvant Therapy for COVID-19 to Calm the	<1 %

Stormy Immunothrombosis and Beyond", Frontiers in Pharmacology, 2021

Publication

47

Anders G. Kjaergaard, Anders Dige, Jeppe S. Nielsen, Else Tønnesen, Jan Krog. "The use of the soluble adhesion molecules sE-selectin, sICAM-1, sVCAM-1, sPECAM-1 and their ligands CD11a and CD49d as diagnostic and prognostic biomarkers in septic and critically ill non-septic ICU patients", APMIS, 2016

Publication

<1 %

48

Chin-Chou Huang, Wan-Leong Chan, Yu-Chun Chen, Tzeng-Ji Chen et al. "The beneficial effects of statins in patients undergoing hemodialysis", International Journal of Cardiology, 2013

Publication

<1 %

49

Cléa Melenotte, Aymeric Silvin, Anne-Gaëlle Goubet, Imran Lahmar et al. "Immune responses during COVID-19 infection", OncoImmunology, 2020

Publication

<1 %

50

Fabio Mangiacapra, Marco Giuseppe Del Buono, Antonio Abbate, Tommaso Gori et al. "Role of endothelial dysfunction in determining angina after percutaneous coronary intervention: Learning from

<1 %

pathophysiology to optimize treatment",
Progress in Cardiovascular Diseases, 2020

Publication

51

Hidesaku Asakura, Haruhiko Ogawa.
"Perspective on fibrinolytic therapy in COVID-19: the potential of inhalation therapy against suppressed-fibrinolytic-type DIC", Journal of Intensive Care, 2020

Publication

<1 %

52

Malay S. Sarkar, Irappa V. Madabhavi, Pham Nguyen Quy, Manjunath B. Govindagoudar.
"COVID - 19 AND COAGULOPATHY", The Clinical Respiratory Journal, 2021

Publication

<1 %

53

Masoumeh Farahani, Zahra Niknam, Leila Mohammadi Amirabad, Nasrin Amiri-Dashatan et al. "Molecular pathways involved in COVID-19 and potential pathway-based therapeutic targets", Biomedicine & Pharmacotherapy, 2022

Publication

<1 %

54

Priya Veluswamy, Max Wacker, Dimitrios Stavridis, Thomas Reichel et al. "The SARS-CoV-2/Receptor Axis in Heart and Blood Vessels: A Crisp Update on COVID-19 Disease with Cardiovascular Complications", Viruses, 2021

Publication

<1 %

55 Radha Gopal, Michael A. Marinelli, John F. Alcorn. "Immune Mechanisms in Cardiovascular Diseases Associated With Viral Infection", *Frontiers in Immunology*, 2020

Publication

<1 %

56 Romeo Gabriel Mihaila, Marius Dragos Mihaila. "Coagulation disorders in SARS-CoV-2 infection", *Biomedical Papers*, 2020

Publication

<1 %

57 archive-ouverte.unige.ch

Internet Source

<1 %

58 d.researchbib.com

Internet Source

<1 %

59 encyclopedia.pub

Internet Source

<1 %

60 eprints.gla.ac.uk

Internet Source

<1 %

61 investigadores.uandes.cl

Internet Source

<1 %

62 journals.eco-vector.com

Internet Source

<1 %

63 lpi.oregonstate.edu

Internet Source

<1 %

64 openaccess.leidenuniv.nl

Internet Source

<1 %

65	pure.rug.nl Internet Source	<1 %
66	qjmed.oxfordjournals.org Internet Source	<1 %
67	smw.ch Internet Source	<1 %
68	translational-medicine.biomedcentral.com Internet Source	<1 %
69	web.archive.org Internet Source	<1 %
70	www.ecdc.europa.eu Internet Source	<1 %
71	www.endotext.org Internet Source	<1 %
72	www.spandidos-publications.com Internet Source	<1 %
73	Li, Shuang, Yan Wu, Ge Yu, Qing Xia, and Yawei Xu. "Angiotensin II Receptor Blockers Improve Peripheral Endothelial Function: A Meta-Analysis of Randomized Controlled Trials", PLoS ONE, 2014. Publication	<1 %
74	Suowen Xu, Iqra Ilyas, Peter J. Little, Hong Li et al. "Endothelial Dysfunction in Atherosclerotic Cardiovascular Diseases and Beyond: From	<1 %

Mechanism to Pharmacotherapies",
Pharmacological Reviews, 2021

Publication

75

Toshiaki Iba, Jean Marie Connors, Jerrold H. Levy. "The coagulopathy, endotheliopathy, and vasculitis of COVID-19", Inflammation Research, 2020

Publication

<1 %

76

"Adhesion Molecules: Function and Inhibition", Springer Nature, 2007

Publication

<1 %

77

Christina D. Camell, Matthew J. Yousefzadeh, Yi Zhu, Larissa G. P. Langhi Prata et al. "Senolytics reduce coronavirus-related mortality in old mice", Science, 2021

Publication

<1 %

78

Daniella M. Mizurini, Eugenio D. Hottz, Patrícia T. Bozza, Robson Q. Monteiro. "Fundamentals in Covid-19-Associated Thrombosis: Molecular and Cellular Aspects", Frontiers in Cardiovascular Medicine, 2021

Publication

<1 %

79

Geoffrey D. Wool, Jonathan L. Miller. "The Impact of COVID-19 Disease on Platelets and Coagulation", Pathobiology, 2021

Publication

<1 %

80

Jawahar L. Mehta, Giuseppe Calcaterra, Pier P. Bassareo. " - 19, thromboembolic risk, and

<1 %

Virchow's triad: Lesson from the past ", Clinical Cardiology, 2020

Publication

81

Juniastuti, Dedy Wahyuddin,
Nihayatussa'adah, Mochamad Amin et al.
"Analysis of genetic and serology of hepatitis
A virus infection during and after outbreak in
two junior high schools in Surabaya,
Indonesia", Journal of Medical Virology, 2019

Publication

<1 %

82

Kirmizis, Dimitrios, Aikaterini Papagianni, Fani
Dogrammatzi, Lemonia Skoura, Anna-Maria
Belechri, Efstathios Alexopoulos, Georgios
Efstratiadis, and Dimitrios Memmos. "Effects
of simvastatin on markers of inflammation,
oxidative stress and endothelial cell apoptosis
in patients on chronic hemodialysis", Journal
of Atherosclerosis and Thrombosis, 2010.

Publication

<1 %

83

Scioli, Maria Giovanna, Pietro Lo Giudice,
Alessandra Bielli, Valeria Tarallo, Alfonso De
Rosa, Sandro De Falco, and Augusto Orlandi.
"Propionyl-L-Carnitine Enhances Wound
Healing and Counteracts Microvascular
Endothelial Cell Dysfunction", PLoS ONE,
2015.

Publication

<1 %

84

U. M. VISCHER. "von Willebrand factor, endothelial dysfunction, and cardiovascular disease", Journal of Thrombosis and Haemostasis, 6/2006

Publication

<1 %

85

Widlansky, M.E.. "The clinical implications of endothelial dysfunction", Journal of the American College of Cardiology, 20031001

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

COVID-19 and Endothelial Dysfunction Biomarkers and Potential Drug Mechanisms

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12
