

SAINS MALAYSIANA



www.ukm.my/jsm



Utama

Sidang Editor

Panduan untuk Penulis

Kandungan, Abstrak dan Rujukan

English



KANDUNGAN, ABSTRAK DAN RUJUKAN
sains malaysia

❖ Jilid 50 ♦ Bilangan 3 ♦ Mac 2021

Muka Surat

Abstract and References Full Text PDF (692KB)	
Effect of Aging Heat Treatment on Corrosion Behavior and Corrosion Kinetics of 17-4PH Stainless Steel in Artificial Saliva <i>Onthida Kosasang, Manutchaya Wongkaewmoon & Somwan Chumphongphan</i>	849 - 858
Abstract and References Full Text PDF (1.98MB)	
Simple and Fast Generalized-M (GM) Estimator and Its Application to Real Data Set <i>Habshah Midi, Shelan Saied Ismael, Jayanthi Arasan & Mohammed A Mohammed</i>	859 - 867
Abstract and References Full Text PDF (1.12MB)	
Analysis of Environmental Status during the Movement Control Order Period due to COVID-19 Pandemic in Malaysia (Analisis Status Alam Sekitar semasa Tempoh Perintah Kawalan Pergerakan Akibat Pandemik COVID-19 di Malaysia) <i>Jeffrey Lawrence D'Silva, Haliza Abdul Rahman, Dzuhalimi Dahalan</i>	869 - 878
Abstract and References Full Text PDF (324KB)	
Association between Climatic Conditions, Population Density and Covid-19 in Indonesia <i>R. Azizah, Santi Martini, Lilis Sulistyorini, Mahmudah, Aditya Sukma Pawitra, Didik Budijanto, Stefanny Surya Nagari, Cendana Fitrihanjani, Fairuz Haniyah Ramadhani & Mohd. Talib Latif</i>	879 - 887
Abstract and References Full Text PDF (400KB)	



Home

Editorial Board

Guide for Authors

Contents, Abstract and References

Malay



EDITORIAL BOARD

sains malaysiana

❖ Editor-In-Chief

Rusli Daik

❖ Editors

Biology

**Choong Chee Yen/ Wee Suk Ling/ Kalaivani
a/p K. Nadarajah/ Nurul Yuziana Mohd
Yusof**

Bioscience and Biotechnology

**Roohaida Othman/ Mohd Firdaus Mohd
Raih/ Noor Liyana Sukiran/ Noraziyah Abd
Aziz Shamsudin**

Chemistry

**Siti Aishah Hasbullah/ Nadhratun Naiim
Mobarak**

Engineering Geology

Goh Thian Lai

Environmental Science and Natural Resources

**Mohd Talib Latif/ Marlia Mohd Hanafiah/
Mohammad Rozaimi Jamaludin**

Earth Science

Norbert Simon

Food Science and Nutrition

Wan Aida Wan Mustapha/ Lim Seng Joe

Health Science

**Siti Balkis Budin/ Nur Azlina Mohd Fahami/
Dalia Abdullah/ Haliza Katas/ Dayang
Fredalina Basri/ Chua Kien Hui/ Leong Xin
Fang/ Noraziah Mohamad Zin**

Materials Science

Chia Chin Hua/ Sahrim Hj. Ahmad

Mathematics

Anuar Mohd Ishak/ Ishak Hashim

❖ International Advisory Board

Biodiversity

Abdul Latiff Mohamad

Universiti Kebangsaan Malaysia, Malaysia

Environmental Science

P. Brimblecombe

University of East Anglia, UK

Materials Science

M. P. Laborie

University of Freiburg, Germany

Materials Science

D. Hui

University of New Orleans, USA

Materials Chemistry

M. Oyama

Kyoto University, Japan

Mathematics

I. Pop

University of Cluj, Romania

Physics

P. Quentin

France

Health Sciences

K.L. Tucker

Northeastern University, USA

Food Science and Technology

L. Wicker

USA

Physics
Yap Chi Chin

Statistics
Noriszura Ismail/ Zamira Hasanah Zamzuri

[Home](#)

[Editorial Board](#)

[Guide for Authors](#)

[Contents, Abstract and References](#)

[Malay](#)

Copyright Reserved © 2007 Sains Malaysiana



SURAT KETERANGAN

Nomor : 3113/UN3.1.10/KP/2023

Yang bertanda tangan di bawah ini :

Nama : Dr. M Atoillah Isfandiari, dr., M.Kes
NIP : 197603252003121002
Pangkat / Golongan : Penata/Gol. III/c
Jabatan : Wakil Dekan II

Dengan ini menerangkan bahwa :

Nama : Dr. Santi Martini, dr., M.Kes
NIP : 196609271997022001
Pangkat / Golongan : Pembina/Gol. IV/a / TMT 01-04-2021
Jabatan : Lektor Kepala / TMT 1 Desember 2018

Telah melaksanakan penelitian dengan judul sebagai berikut :

No	Judul Karya Ilmiah	Tahun Pelaksanaan
1	Association between percentage of smokers and prevalence of smoking attributable morbidity in Indonesia: one decade after implementation of smoke-free area regulation (https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-022-14435-8)	2021
2	Association between climatic conditions, population density and covid-19 in indonesia (https://www.ukm.my/jsm/pdf_files/SM-PDF-50-3-2021/28.pdf)	2020

Adapun penelitian tersebut layak dilakukan, meskipun belum ada *Uji Etical Clearence* karena menggunakan data sekunder yang telah di publikasikan dan menghasilkan output yang sangat baik.

Demikian surat keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional Guru Besar.

Surabaya, 17 April 2023

A.n. Dekan
Wakil Dekan II,



Dr. M. Atoillah Isfandiari, dr., M.Kes
NIP 197603252003121002

Association between Climatic Conditions, Population Density and COVID-19 in Indonesia

(Perkaitan antara Keadaan Iklim, Kepadatan Penduduk dan COVID-19 di Indonesia)

R. AZIZAH*, SANTI MARTINI, LILIS SULISTYORINI, MAHMUDAH, ADITYA SUKMA PAWITRA, DIDIK BUDIJANTO, STEFANNY SURYA NAGARI, CENDANA FITRAHANJANI, FAIRUZ HANIYAH RAMADHANI & MOHD. TALIB LATIF

ABSTRACT

The first emergence of Corona Virus Disease 2019 (COVID-19) confirmed cases found in Wuhan, China, has become a global crisis. At least 177 countries have been affected over 43,000,000 confirmed cases of corona positive and more than one million deaths until October 27th, 2020. Recent research has analyzed any possible factors causing the COVID-19 spreads were climate factors and population density. Indonesia was a tropical region known as the high-populated country in the World, with a 52.9% area with a high mean air temperature and over 267.7 million populations. Our study aims to analyze the correlation between climate, population density, and COVID-19 in Indonesia. We used the K-means cluster method and Fisher's exact test to determine climatic conditions, population density, and COVID-19 clusters and study the correlation. Our research found that there is a correlation between climatic conditions and population density with COVID-19 ($p: 0,034$; $p:0,004$). Warmer climate conditions and densely populated regions contributed to the risen COVID-19 transmission in Indonesia. These are highlighted by the evidence of the top six provinces with highest COVID-19 cases are province classified in warmer climatic conditions (high air temperature, low rainfall, and humidity) and a fairly-dense to densely populated region.

Keywords: Climatic conditions; COVID-19; population density

ABSTRAK

Kemunculan pertama Penyakit Virus Corona 2019 (COVID-19) mengesahkan kes yang dijumpai di Wuhan, China, telah menjadi krisis global. Sekurang-kurangnya 177 negara telah dijangkiti dengan lebih 43,000,000 kes yang disahkan positif korona dan lebih daripada satu juta kematian sehingga 27 Oktober 2020. Penyelidikan terkini telah menganalisis faktor kemungkinan yang menyebabkan penyebaran COVID-19 iaitu faktor iklim dan kepadatan penduduk. Indonesia adalah wilayah tropik yang terkenal sebagai negara berpenduduk tinggi di dunia dengan luas 52.9% dan min suhu udara tinggi dengan lebih 267.7 juta penduduk. Kajian kami bertujuan untuk menganalisis hubungan antara iklim, kepadatan penduduk dan COVID-19 di Indonesia. Kami menggunakan kaedah kluster K-means dan ujian tepat Fisher untuk menentukan keadaan iklim, kepadatan penduduk dan kelompok COVID-19 dan mengkaji korelasi tersebut. Penyelidikan kami mendapati bahawa terdapat hubungan antara keadaan iklim dan kepadatan penduduk dengan COVID-19 ($p: 0,034$; $p: 0,004$). Keadaan iklim yang lebih panas dan kawasan berpenduduk padat menyumbang kepada peningkatan jangkitan COVID-19 di Indonesia. Ini disoroti dengan bukti enam wilayah teratas dengan kes COVID-19 tertinggi adalah wilayah yang dikelaskan dalam keadaan iklim yang lebih panas (suhu udara tinggi, hujan yang rendah dan kelembapan) dan wilayah yang hampir padat hingga berpenduduk padat.

Kata kunci: COVID-19; keadaan iklim; kepadatan penduduk

INTRODUCTION

Starting at the end of 2019, a new respiratory disease, which was later named COVID-19, were found in Wuhan, China. This disease is caused by a novel coronavirus (SARS-CoV-2). The virus has spread throughout China within two months, which then develops into various countries. The World Health Organization (WHO) has declared the epidemic of public health emergencies since January 30th, 2020 (WHO 2020a). This pandemic has caused panic around the World since the beginning of 2020. There

were 177 countries with over 43,000,000 confirmed cases and more than 1,000,000 deaths until October 27th, 2020 (WHO 2020b).

Elderly and people with congenital diseases are considered to be a high-risk group for COVID-19. The disease is spread through direct and indirect contact (Dangi 2020). Through direct contacts, droplets can spread up to 1-2 m from the contact source. Many studies predict that the virus can spread through the air (Wang & Du 2020), looking at the rapid spread of COVID-19. Some research was done

to prove the spread of the SARS-CoV-2 virus in addition to through droplets. In October, The Center for Disease Control and Prevention (CDC) announced that the virus could spread through the air (CDC 2020). The CDC's statement further confirmed that the spread of the virus is very frightening because the potential spread of the virus is more comfortable and faster when it is through the air. The theory states that crowds can be the most vulnerable source for COVID-19 spread.

To date, the increase in daily COVID-19 cases in Indonesia is still an alarming number. Apart from manageable factors that can control the COVID-19 confirmed cases increase, there are some uncontrollable factors that we cannot handle, such as climate and population density and many other pre-existing conditions which increase the risk of death from COVID-19. Scientists in the United States government estimate that COVID-19 could kill 10 out of 1000 Americans. Crowd increases the potential for human contact and is a primary source of human-to-human transmissions. COVID-19 growth significantly in denser areas (Therese 2020). Another study on climate and COVID-19 in Brazil, with an annual average temperature, ranges from 16.8 °C to 27.4 °C, found a negative linear association between temperature and the number of COVID-19 cases. They also found that the increase of 1 °C is associated with a decrease of -4.8951% ($t = -2.29$, $p = 0.0226$) in the cumulative daily number of confirmed cases of COVID-19 (Prata et al. 2020). Research on population density, wind speed and COVID 19 has also been conducted in Turkey by Coskun et al. (2020). The results showed that population density and wind, accounting for 94% of the virus spread variance, had a significant impact on the spread of the virus or the number of cases. Coskun et al. (2020) stated that the virus is spreading more in windy weather, indicating that the air threatens humans with wind speeds. A cluster and regression model was carried out on the group of cities with the highest COVID-19 cases, and the highest population density in Algeria shows a strong correlation between population density and the number of COVID-19 infections.

The results showed that population density positively affected the spread of COVID-19 (Kadi et al. 2020).

Indonesia is a highly-populated country; Indonesia's population deploys across 34 provinces, so does the climatic conditions. Although it is a tropical area, climatic conditions in various provinces in Indonesia are diverse. Some regions may have a lower air temperature than others. Given these diverse conditions in Indonesia, it is necessary to study the climate conditions and population density to see whether these two factors are related to the enormous COVID-19 event in Indonesia. This study analyzes the correlation between climatic conditions (temperature, rainfall, and humidity), population density, and COVID-19 in 34 provinces in Indonesia.

MATERIALS AND METHODS

This research was conducted in Indonesia using climatic conditions, population density, and COVID-19 data from secondary resources. Climatic conditions data are collected from Indonesia's Environmental Statistics 2019 report (BPS 2019). Due to the limitation of data in Indonesia, we used the 2019 climate conditions in this study. Population density data is collected from the Indonesia Statistics 2020 report (BPS 2019). COVID-19 data used in this study is a cumulative data of total, recovered, and death cases from early March until July 22, 2020. The data collected from COVID-19 Daily Media Report, Indonesian Health Ministry. There were 34 Province in Indonesia, which illustrated in Figure 1.

This study used computer-based data processing with a statistical approach. There were two analyses conducted in this study: K-means cluster and Fisher's exact test. K-means cluster was used to determine the clusters of climatic conditions, population density, and COVID-19. Fisher's exact test is used to determine the correlation between the climatic condition and COVID-19 and the correlation between population density and COVID-19.

RESULTS AND DISCUSSION




FIGURE 1. Indonesia Map

Source: <https://peta-hd.com/peta-34-provinsi-di-indonesia/>

COVID-19 CLUSTER SPREADS IN INDONESIA
 K-means cluster is used to determine the COVID-19 clusters. There are three indicators used to determine the

COVID-19 clusters; total, recovered, and death cases. Table 1 shows three COVID-19 clusters; low, medium, and high.

TABLE 1. COVID-19 cluster cases positive, recovered, and

deaths

Measure	COVID-19 Clusters								
	Low			Medium			High		
	Total cases	Recovered	Deaths	Total cases	Recovered	Deaths	Total cases	Recovered	Deaths
Min	136	75	1	3149	1472	144	17621	10460	751
Max	8407	5034	287	7726	3720	444	19093	11139	1496
Mean ± SD	1185,54 ± 1677,51	674,61 ± 981,39	41,39 ± 62,29	5458 ± 1880,95	2441,75 ± 941,11	263,25 ± 129,77	18357 ± 1040,86	10799,5 ± 480,12	1123,5 ± 526,79

From Table 1, low clusters are cluster with lower total COVID-19 cases than the medium and high cluster. The recovery rate means recovered cases per mean total cases in each cluster in the low cluster are 56.9%. It is higher than the medium cluster with a 44.73% recovery rate and lower than the high cluster is 58.83%. The case fatality rate means death cases per mean total cases in each cluster; in the low cluster, they are lower with a 3.49% than the medium cluster 4.82% and high cluster 6.12%.

From the cluster analysis, 28 provinces classified in low clusters are Aceh; Bali; Banten; Bangka Belitung; Bengkulu; DI Yogyakarta; Jambi; West Kalimantan; East Kalimantan; Central Kalimantan; North Kalimantan;

Riau Islands; West Nusa Tenggara; South Sumatra; West Sumatra; North Sulawesi; North Sumatra; Southeast Sulawesi; Central Sulawesi; Lampung; Riau; North Maluku; Maluku; West Papua; Papua; West Sulawesi; East Nusa Tenggara; and Gorontalo. Four provinces classified in medium clusters are West Java Province; Central Java; Kalimantan Selatan; and South Sumatra. Two clusters are classified in high clusters are East Java and Jakarta.

Figure 2 shows the six provinces in Indonesia with the highest number of total cases: East Java 19,093; Jakarta 17,621; South Sulawesi 8,407; Central Java 7,726; West Java 5,741; and South Kalimantan 5,216.




FIGURE 2. Number of COVID-19 cases spread across 34 provinces in Indonesia

CORRELATION BETWEEN CLIMATIC CONDITIONS
(TEMPERATURE, HUMIDITY, AND RAINFALL) AND
COVID-19 IN INDONESIA

*Climatic Conditions (Temperature, Humidity, and Rainfall)
In Indonesia*

There are three climate indicators used to determine the Climatic Conditions cluster; air temperature, rainfall, and

humidity. The three indicators used different units, so a value standardization is performed. There are two Climatic Conditions clusters; Climatic Conditions I and Climatic Conditions II (Table 2).

TABLE 2. Climate clusters (temperature, rainfall, and humidity) in climatic conditions i and climatic conditions ii of 34 provinces in Indonesia

Measure	Climate clusters					
	Climatic conditions I			Climatic conditions II		
	Temperatures (°C)	Rainfall (mm/year)	Humidity (%)	Temperatures (°C)	Rainfall (mm/year)	Humidity (%)
Min	26,7	1585	80	25,9	490	67
Max	28,3	3823	87	29,5	3048	82
Mean ± SD	27,3 ± 0,44	2693,5 ± 597,88	83,19 ± 2,07	28,22 ± 0,91	1840,5 ± 578,81	76,72 ± 3,94

The mean air temperature in Climatic Conditions I is lower than Climatic Conditions II, while the mean rainfall and humidity in Climatic Conditions I are higher than the Climatic Conditions II. These show that Climatic Conditions I is a colder area than Climatic Conditions II. From the cluster analysis, 16 provinces within Climatic conditions I are Aceh; West Sumatra; Riau; Jambi; Bangka Belitung; Riau Islands; West Kalimantan; Central Kalimantan; North Kalimantan; North Sulawesi; South Sulawesi; Southeast Sulawesi; Maluku; North Maluku;

West Papua; Papua; and 18 provinces within Climatic conditions II are North Sumatra; South Sumatra; Bengkulu; Lampung; Jakarta; West Java; Central Java; Yogyakarta; East Java; Banten; Bali; West Nusa Tenggara; East Nusa Tenggara; South Kalimantan; East Kalimantan; Central Sulawesi; Gorontalo; West Sulawesi. This finding shows that 52.9% of Indonesia's area indicates having a high mean air temperature, low rainfall and humidity, included in Climatic conditions II (Table 3).

TABLE 3. The difference of cluster COVID-19 based on climate climatic conditions

Climatic conditions	Cluster COVID-19						Total	P _{value}	Correlation coefficient	
	Low		Medium		High					
	n	%	n	%	n	%				
Climatic conditions I	16	100	0	0	0	0	16	100	0,034	0,400
Climatic conditions II	12	66,7	4	22,2	2	11,1	18	100		
Total	28	82,4	4	11,8	2	5,9	34	100		

Our study found a correlation between Climatic Conditions and COVID-19 clusters $p=0.034$ with a positive correlation coefficient of 40%, indicates a strong

correlation between climatic conditions and COVID-19. It shows a linear correlation, that higher climate conditions led to higher COVID-19 cases. All provinces in Climatic

Conditions I have low COVID-19 cases. We predict that the coronavirus can live under high temperatures. Indonesia's mean air temperature ranges from 26.7-29.5 °C and COVID-19 spread in Indonesia is also high. The condition is not according to the previous studies state that coronavirus can survive with lower air temperatures 5-11 °C (Sajadi et al. 2020). However, Indonesia's humidity level is relatively high, ranging from 67-87% to the range of humidity, which is considered following the coronavirus environment (44-88%) (Sajadi et al. 2020). The humidity level in Indonesia is suitable for the survival of the coronavirus. We predict humidity is an essential climate factor in the survival of the coronavirus.

A study conducted in Brazil showed that warm temperature (above 25 °C) were not associated with the declined of the COVID-19 cases (Prata et al. 2020). Pan et al. (2020) suggest that warmer weather unlikely to reduce the COVID-19 transmissions. Although a surge in COVID-19 cases in the hot and humid region is not as rapid as in the cold and dry region, COVID-19 cases in the hot and humid region keep increasing consistently. Warm weather alone may not be able to stop the COVID-19 transmissions and several other factors other than meteorological factors should be taken into account (Bukhari et al. 2020). The spread of COVID-19 appears to be lower in warm and wet climates, but the certainty of the resulting evidence is rated low. Temperature and humidity alone do not explain most of the variability of the COVID-19 outbreak. The spread of COVID-19 may be influenced by other factors such as public isolation policy, herd immunity, migration patterns, population density, and cultural aspects (Macenas et al. 2020).

CORRELATION BETWEEN POPULATION DENSITY (POPULATION AND AREA OF PROVINCE) AND COVID-19 IN INDONESIA

Population Density (Population and Area of Province) In Indonesia

Various studies on population density correlation and COVID-19 spread are conducted in the world. Some studies found that population density correlates with COVID-19 distribution rates, while others found the opposite. According to research conducted in China, population density is not related to COVID-19 events. However, Sun et al. (2020) stated that territorial restriction is a useful measure in reducing the chance of infection in the community. Territorial restrictions, commonly referred to as regional quarantine, are traditional solutions used in antiquity to control the spread of infectious diseases (Wilder-Smith & Freedman 2020). This step is useful to limit communities' interaction within the region so that the virus does not increasingly spread mainly in densely populated areas.

High-density regions can cause the spread of the SARS-Cov-2 virus faster than lower density regions, based on research conducted in Japan (Sajadi et al. 2020). High-density levels increase the likelihood of physical contact with others due to increasingly limited wiggle room (Ramadhani 2020). Besides, population in densely populated areas will have less chance of treating properly if the health service in the region is inadequate inpatient surges. Population density is related to the number of COVID-19 cases, the case fatality rate from COVID-19, and the willingness of health services in the region (Amoo et al. 2020) (Table 4).

TABLE 4. Population density of 34 provinces in Indonesia

Measure	Population density		
	Low	Medium	High
Min	9	750	15900
Max	273	1394	15900
Mean ± SD	101,15 ± 77,79	1099,67 ± 266,77	15900 ± .*

*Standard Deviation unable to calculate, as there is only 1 Province

We divided population density into three categories; low, medium, and high. Population density is the amount that represents how dense the population in one particular area. Population density is calculated by dividing the

population by the area. There is a large discrepancy in high population density compared to the low and medium population.

There are 27 low density provinces, Aceh; North Sumatra; West Sumatra; Riau; Jambi; South Sumatra; Bengkulu; Lampung; Bangka Belitung; Riau Islands; West Nusa Tenggara; East Nusa Tenggara; West Kalimantan; Central Kalimantan; South Kalimantan; East Kalimantan; North Kalimantan; North Sulawesi; Central Sulawesi;

South Sulawesi; North Sulawesi; Gorontalo; West Sulawesi; Maluku; North Maluku; West Papua; and Papua. The six medium density provinces are West Java; Central Java; Yogyakarta; East Java; Banten; Bali. The high-density province is Jakarta. The province in Java Island has a medium to high population density (Figure 3).




FIGURE 3. Population density of 34 provinces in Indonesia

Jakarta province has a high population of 15,900 people/km² with population is 10,557,800 and 664.01 km². Jakarta is one of the metropolitan areas with a dense population. In addition to having a dense population, Jakarta is one of the areas with a high positive case. Most metropolitan areas in different countries have dense populations. New York is one of the states of the United States with high population density and positive

cases. In research conducted in the United States, New York accounted for 37% of positive cases in the United States. In addition to New York, other metropolitan areas such as Chicago, Washington, and Los Angeles are also among the top ten densely populated metropolitan areas contributing to positive cases in the United States (Zhang & Schwartz 2020) (Table 5).

TABLE 5. The difference of Cluster COVID-19 Based on Population Density

Population density	Cluster COVID-19						Total	P _{value}	Correlation coefficient	
	Low		Medium		High					
	n	%	n	%	n	%				
Low	25	92,6	2	0	0	0	27	100	0,004	0,633
Medium	3	50	2	33,3	1	16,7	6	100		
High	0	0	0	0	1	100	1	100		
Total	28	82,4	4	11,8	2	5,9	34	100		

Our study found a correlation between population density and COVID-19 with $p=0.004$ and a positive correlation coefficient of 63.3%. Based on cross-tabulation results, 92.6% of low-density provinces have low COVID-19 cases. As well as 100% high-density have high COVID-19. This research is in line with research (Sarmadi et al. 2020) that population density in the United States affects deaths from COVID-19 in the United States. The greater the population density, the greater the mortality rate in the United States. Similar research was conducted in India, which ranks in the top ten of the world's top population as Indonesia. The research shows an association between population density and COVID-19 spread (Bhadra et al. 2020). This result is similar to this study and the United States study, which shows that population density has a role in the spread of COVID-19.

SIX PROVINCES WITH THE HIGHEST COVID-19 CASES, CLIMATE CONDITIONS AND POPULATION DENSITY

The Top Six Provinces with the Highest Cases, Climatic Conditions and Population Density

Our study found that the six provinces with the highest COVID-19 cases were in Climatic conditions II (high mean air temperature with low rainfall and humidity) (Table 6). The findings are in stark contrast, as research analyzing the effects of climate and weather on COVID-19 cases found that countries with high mean air temperatures and low humidity have the effect of lowering the number of deaths caused by COVID-19. A study found that tropical regions with high mean temperature and high relative humidity, i.e., India, Pakistan, Saudi Arabia, Malaysia,

Thailand, and Australia, have low death cases and high recovery rates (Pavani & Sunalini 2020). Muhammad Mazhar et al. (2020) also discovered that tropical regions, i.e. Malaysia, Thailand, and Indonesia, are less affected by the coronavirus. However, Muhammad Mazhar et al. (2020) study the regional climatic condition's effect by June 2020. Some researchers also conducted a study in the subtropical climate, i.e. China. They also found that higher average air temperatures will lower the risk of COVID-19 (Ma et al. 2020; Wang et al. 2020). However, Indonesia's high mean temperature did not follow the same conditions related to COVID-19 in other regions. Coronavirus will still last inside the human body and infect other humans during the incubation period. The more frequent human activities outside the house, the more the virus will likely spread and infect other humans.

Based on our results, the top 6 provinces with the highest COVID-19 cases are located in a fairly dense and dense region. As a high-density region, Jakarta is the region with the second-highest COVID-19 cases, while East Java is in the first position is a fairly dense region. The rest following a sequence order are West Java, Central Java, South Kalimantan, and South Sumatra. This region is fairly dense. Our result found a similar study conducted by Jawad (2020), which states that the results of an analysis conducted in countries vulnerable to COVID-19 spread show that population density affects the peak period of COVID-19 spread. However, population density is not an essential factor in the spread of COVID-19 under strict lockdown policies. China's lockdown policy could effectively limit the speed at which COVID-19 spreads (Sun 2020).

TABLE 6. Climatic conditions and population density in six provinces with the highest COVID-19 cases

No.	Province	Cluster		
		Covid 19	Climatic conditions	Population density
1.	East Java	High Cases	II	Medium
2.	Jakarta	High Cases	II	High
3.	West Java	Medium Cases	II	Medium
4.	Central Java	Medium Cases	II	Medium
5.	South Kalimantan	Medium Cases	II	Medium
6.	South Sumatra	Medium Cases	II	Medium

CONCLUSION

Indonesia region is identified in high air temperature with 26.7-29.5 °C and high humidity (67-87%), due to its tropical climate. However, the majority of the province in Indonesia are classified in Climatic Conditions II in 52,9%. Climatic conditions II is a Climatic condition with higher mean air temperatures, lower rainfall and humidity than the rest. Indonesia is also known as a highly-dense country. Jakarta is the most populated region in Indonesia (population density: 15,900). Our results found a correlation between the climatic conditions and the COVID-19 cluster ($p:0,034$), and there is a correlation between population density and COVID-19 clusters ($p:0,004$). Population density is more related to the high number of COVID-19 cases in Indonesia, which has a positive correlation of 63.3%. The top six provinces that have the highest COVID-19 cases in Indonesia are classified in Climatic Conditions II and a fairly-dense to densely populated region. There need to be adaptation and mitigation measures in COVID-19 control related to climate conditions in Climatic conditions II and high population density so that policies in their respective provinces in crowd control, social distancing, and efforts to reduce COVID-19 cases through increased community-based awareness.

ACKNOWLEDGEMENTS

We would like to thank the Department of Environmental Health and Environmental Health Master Program, Faculty of Public Health, Universitas Airlangga which has facilitated the activities of writing scientific articles in the framework of international journal publication. We would also thank Prof. Mohd. Talib Latif, Universiti Kebangsaan Malaysia who guided and corrected this research until this article was compiled.

REFERENCES

- Amoo, E.O., Adekeye, O., Olawole-Isaac, A., Fasina, F., Adekola, P.O., Samuel, G.W., Akanbi, M.A., Oladosun, M. & Azuh, D.E. 2020. Nigeria and Italy divergences in coronavirus experience: Impact of population density. *Scientific World Journal* 2020: 8923036. DOI: 10.1155/2020/8923036.
- Bhadra, A., Mukherjee, A. & Sarkar, K. 2020. Impact of population density on Covid-19 infected and mortality rate in India. *Modeling Earth Systems and Environment* 7: 623-629.
- BPS. 2020. *Statistics Indonesia Statistical Year Book of Indonesia 2020*. Accessed by https://drive.google.com/file/d/18EfhKxnIARMh0h_4qqWsEszYsk_X0W8G/view.
- BPS. 2019. *Population by Age and Gender Group 2019*. Indonesian Central Bureau of Statistics. https://www.bps.go.id/indikator/indikator/view_data_pub/0000/api_pub/58/da_03/1.
- BPS. 2019. *Indonesia Environmental Statistics 2019*. <https://www.bps.go.id/publication/2019/12/13/e11bfc8ff8392e5e13a8cff3/statistik-lingkungan-hidup-indonesia-2019.html>.
- Bukhari, Q., Massaro, J.M., D'Agostino Sr., R.B. & Sheraz Khan. 2020. Effects of weather on coronavirus pandemic. *International Journal of Environmental Research and Public Health* 17(15): 5399. <https://doi:10.3390/ijerph17155399>.
- CDC. SARS-CoV-2 & Potential Airborne Transmission [Internet]. 2020. *Center for Disease Control and Prevention*. <https://www.cdc.gov/coronavirus/2019-ncov/more/scientific-brief-sars-cov-2.html>.
- Coşkun, H., Yıldırım, N. & Gündüz, S. 2020. The spread of COVID-19 virus through population density and wind in Turkey cities. *Science of The Total Environment* 751(10): 141663.
- Dangi, R.R. & George, M.A. 2020. Review on theories and models of disease causation for COVID-19. *Journal SSRN Electron* 2020 <https://doi.org/10.2139/ssrn.3584080>.

- Jawad, A.J. 2020. Effectiveness of population density as natural social distancing in COVID-19 spreading. *Ethics Medicine and Public Health* 15(C): 100556.
- Kadi, N. & Khelifaoui, M. 2020. Population density, a factor in the spread of COVID-19 in Algeria: Statistic study. *Bulletin of the National Research Centre* 44: 138. <https://doi.org/10.1186/s42269-020-00393-x>.
- Ma, Y., Zhao, Y., Liu, J., He, X., Wang, B., Fu, S., Yan, J., Niu, J., Zhou, J. & Luo, B. 2020. Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Science of The Total Environment* 724: 138226.
- Mecenas, P., da Rosa Moreira Bastos, R.T., Rosário Vallinoto, A.C. & Normando, D. 2020. Effects of temperature and humidity on the spread of COVID-19: A systematic review. *PLoS ONE* 15(9): e0238339.
- Ministry of Health of Indonesia. 2020. *Daily Media Report Covid-19* dated July 22, 2020 at 12.00 WIB.
- Ministry of Health of Indonesia. 2020. *COVID-19 Transmission Prevention Guide*. <https://promkes.kemkes.go.id/panduan-pencegahan-penularan-covid-19-untuk-masyarakat>.
- Muhammad Mazhar Iqbal, Irfan Abid, Saddam Hussain, Naem Shahzad, Muhammad Sohail Waqas & Muhammad Jawed Iqbal. 2020. The effects of regional climatic condition on the spread of COVID-19 at global scale. *Sci. Total Environ.* 15(739): 140101.
- National Health Commission of People's Republic of China. 2020. *Pneumonia Diagnosis and Treatment of 2019-nCoV Infection from Chinese NHC and CDC 2020*.
- Pan, J., Yao, Y., Liu, Z., Meng, X., John, S. Ji., Qiu, Y., Wang, W., Zhang, L., Wang, W. & Kan, H. 2020. Warmer weather unlikely to reduce the COVID-19 transmission: An ecological study in 202 locations in 8 countries. *Science of The Total Environment* 753: 142272.
- Pavani, A. & Sunalini, K.K. 2020. Effects of temperature and relative humidity on Covid-19. *Journal of Critical Reviews* 7(9): 3177-3182.
- Prata, D.N., Rodrigues, W. & Bermejo, P.H. 2020. Temperature significantly changes COVID-19 transmission in (sub) tropical cities of Brazil. *Science of The Total Environment* 729: 138862.
- Ramadhani, F.H. 2020. Literature review: Healthy home as the new normal for Covid-19 prevention. *Jurnal Kesehatan Lingkungan* doi: 10.20473/jkl.v12i1si.2020.1-10.
- Sajadi, M.M., Habibzadeh, P., Vintzileos, A., Shokouhi, S., Miralles-Wilhelm, F. & Amoroso, A. 2020. Temperature, humidity, and latitude analysis to estimate potential spread and seasonality of Coronavirus Disease 2019 (COVID-19). *JAMA Network Open* 3(6): e2011834.
- Sarmadi, M., Moghandam, V.K., Dickerson, A.S. & Martelletti, L. 2020. Association of COVID-19 distribution with air quality, sociodemographic factors, and comorbidities: An ecological study of US states. *Air Quality, Atmosphere & Health* <https://doi.org/10.1007/s11869-020-00949-w>.
- Sun, Z., Zhang, H., Yang, Y., Wan, H. & Wang, Y. 2020. Impacts of geographic factors and population density on the COVID-19 spreading under the lockdown policies of China. *Science of The Total Environment* 746(666): 141347.
- Therese, K.L.S., White, L.F. & Nichols, B. 2020. Population density and basic reproductive number of COVID-19 across United States counties. *National Library of Medicine* <https://doi.org/10.1101/2020.06.12.20130021>.
- Wang, H., Chen, C. & Lin, Q. 2020. Short-term effects of ambient temperature and relative humidity on the risk of COVID-19 and SARS in Guangzhou, China: A time-series analysis. *Research Square* doi: 10.21203/rs.3.rs-50445/v1.
- Wang, J. & Du, G. 2020. COVID-19 may transmit through aerosol. *Irish Journal of Medical Science* 89: 1143-1144.
- WHO. 2020a. Infection prevention and control guidance for long-term care facilities in the context of COVID-19. *Interim Guidance of World Health Organization*. <https://apps.who.int/iris/handle/10665/331508>.
- WHO. 2020b. WHO Corona Disease (COVID-19) Dashboard. <https://covid19.who.int/>.
- Wilder-Smith, A. & Freedman, D.O. 2020. Isolation, quarantine, social distancing and community containment: Pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak. *Journal of Travel Medicine* 27(2): 1-4. <https://doi.org/10.1093/jtm/taaa020>.
- Zhang, C.H. & Schwartz, G.G. 2020. Spatial disparities in coronavirus incidence and mortality in the United States: An ecological analysis as of May 2020. *Journal of Rural Health* 36(3): 433-445.
- R. Azizah*, Santi Martini, Lilis Sulistyorini, Mahmudah, Aditya Sukma Pawitra, Stefanny Surya Nagari, Cendana Fitrahanjani, Fairuz Haniyah Ramadhani & Mohd. Talib Latif
Faculty of Public Health
Universitas Airlangga
Surabaya
Indonesia
- Didik Budijanto
The Ministry of Health of the Republic of Indonesia
Jakarta Selatan
12950 Jakarta
Indonesia
- Mohd. Talib Latif
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
43600 UKM Bangi, Selangor Darul Ehsan
Malaysia

*Corresponding author; email: azizah@fkm.unair.ac.id

Received: 5 November 2020

Accepted: 17 December 2020