Association between Climatic Conditions, Population Density and COVID-19 in Indonesia

by R. Azizah

Submission date: 01-Nov-2021 11:39AM (UTC+0800) Submission ID: 1689586008 File name: tic_Conditions,_Population_Density_and_COVID-19_in_Indonesia.pdf (396.92K) Word count: 4800 Character count: 26385 Sains Malaysiana 50(3)(2021): 879-887 http://dx.doi.org/10.17576/jsm-2021-5003-28

Association between Climatic Conditions, Population Density and COVID-19 in

Indonesia

(Perkaitan antara Keadaan Iklim, Kepadatan Penduduk dan COVID-19 di Indonesia)

R. Azizah*, Santi Martini, Lilis Sulistyorini, Mahmudah, Aditya Sukma Pawitra, Didik Budijanto, Stefanny Surya Nagari, Cendana Fitrahanjani, Fairuz Haniyah Ramadhani & Mohd. Talib Latif

ABSTRACT

The first emergence of Corona Virus Disease 2019 (COVID-19) confirmed cases found in Wuhan, China, has become a global crisis. At least 177 countries have been affected over 43,000,000 confirmed cases of corona positive and more than one million deaths until October 27th, 2020. Recent research has analyzed any possible factors causing the COVID-19 spreads were climate factors and population density. Indonesia was a tropical region known as the high-populated country in the World, with a 52.9% area with a high mean air temperature and over 267.7 million populations. Our study aims to analyze the correlation between climate, population density, and COVID-19 in Indonesia. We used the K-means cluster method and Fisher's exact test to determine climatic conditions, population density, and COVID-19 clusters and study the COVID-19 (p: 0,034; p:0,004). Warmer climate conditions and densely populated regions contributed to the risen COVID-19 transmission in Indonesia. These are highlighted by the evidence of the top six provinces with highest COVID-19 cases are province classified in warmer climatic conditions (high air temperature, low rainfall, and humidity) and a fairhy-dense to densely populated region.

Keywords: Climatic conditions; COVID-19; population density

ABSTRAK

Kemunculan pertama Penyakit Virus Corona 2019 (COVID-19) mengesahkan kes yang dijumpai di Wuhan, China, telah menjadi krisis global. Sekurang-kurangnya 177 negara telah dijangkiti dengan lebih 43,000,000 kes yang disahkan positif korona dan lebih daripada satu juta kematian sehingga 27 Oktober 2020. Penyelidikan terkini telah menganalisis faktor kemungkinan yang menyebabkan penyebaran COVID-19 iaitu faktor iklim dan kepadatan penduduk. Indonesia adalah wilayah tropik yang terkenal sebagai negara berpopulasi tinggi di dunia dengan luas 52.9% dan min suhu udara tinggi dengan lebih 267.7 juta penduduk. Kajian kami bertujuan untuk menganalisis hubungan antara iklim, kepadatan penduduk dan COVID-19 di Indonesia. Kami menggunakan kaedah kluster K-means dan ujian tepat Fisher untuk menentukan keadaan iklim, kepadatan penduduk dan kelompok COVID-19 dan mengkaji korelasi tersebut. Penyelidikan kami mendapati bahawa terdapat hubungan antara kaedaan iklim dan kepadatan penduduk dengan COVID-19 (p: 0,034; p: 0,004). Keadaan iklim yang lebih panas dan kawasan berpenduduk padat menyumbang kepada peningkatan jangkitan COVID-19 di Indonesia. Ini disoroti dengan bukti enam wilayah teratas dengan kes COVID-19 tertinggi adalah wilayah yang dikelaskan dalam keadaan iklim yang lebih panas (suhu udara tinggi, hujan yang rendah dan kelembapan) dan wilayah yang hampir padat hingga berpenduduk padat.

Kata kunci: COVID-19; keadaan iklim; kepadatan penduduk

INTRODUCTION

Starting at the end of 2019, a new respiratory disease, which was later named COVID-19, were found in Wuhan, China. This disease is caused by a novel coronavirus (SARS-CoV-2). The virus has spread throughout China within two months, which then develops into various countries. The World Health Organization (WHO) has declared the epidemic of public health emergencies since January 30th, 2020 (WHO 2020a). This pandemic has caused panic around the World since the beginning of 2020. There were 177 countries with over 43,000,000 confirmed cases and more than 1,000,000 deaths until October 27th, 2020 (WHO 2020b).

Elderly and people with congenital diseases are considered to be a high-risk group for COVID-19. The disease is spread through direct and indirect contact (Dangi 2020). Through direct contacts, droplets can spread up to 1-2 m from the contact source. Many studies predict that the virus can spread through the air (Wang & Du 2020), looking at the rapid spread of COVID-19. Some research was done 880

to prove the spread of the SARS-CoV-2 virus in addition to through droplets. In October, The Center for Disease Control and Prevention (CDC) announced that the virus could spread through the air (CDC 2020). The CDC's statement further confirmed that the spread of the virus is very frightening because the potential spread of the virus is more comfortable and faster when it is through the air. The theory states that crowds can be the most vulnerable source for COVID-19 spread.

To date, the increase in daily COVID-19 cases in Indonesia is still an alarming number. Apart from manageable factors that can control the COVID-19 confirmed cases increase, there are some uncontrollable factors that we cannot handle, such as climate and population density and many other pre-existing conditions which increase the risk of death from COVID-19. Scientists in the United States government estimate that COVID-19 could kill 10 out of 1000 Americans. Crowd increases the potential for human contact and is a primary source of human-to-human transmissions. COVID-19 growth significantly in denser areas (Therese 2020). Another study on climate and COVID-19 in Brazil, with an annual average temperature, ranges from 16.8 °C to 27.4 °C, found a negative linear association between temperature and the number of COVID-19 cases. They also found that the increase of 1 °C is associated with a decrease of -4.8951% (t = -2.29, p = 0.0226) in the cumulative daily number of confirmed cases of COVID-19 (Prata et al. 2020). Research on population density, wind speed and COVID 19 has also been conducted in Turkey by Coskun et al. (2020). The results showed that population density and wind, accounting for 94% of the virus spread variance, had a significant impact on the spread of the virus or the number of cases. Coskun et al. (2020) stated that the virus is spreading more in windy weather, indicating that the air threatens humans with wind speeds. A cluster and regression model was carried out on the group of cities with the highest COVID-19 cases, and the highest population density in Algeria shows a strong correlation between population density and the number of COVID-19 infections.

The results showed that population density positively affected the spread of COVID-19 (Kadi et al. 2020).

Indonesia is a highly-populated country; Indonesia's population deploys across 34 provinces, so does the climatic conditions. Although it is a tropical area, climatic conditions in various provinces in Indonesia are diverse. Some regions may have a lower air temperature than others. Given these diverse conditions in Indonesia, it is necessary to study the climate conditions and population density to see whether these two factors are related to the enormous COVID-19 event in Indonesia. This study analyzes the correlation between climatic conditions (temperature, rainfall, and humidity), population density, and COVID-19 in 34 provinces in Indonesia.

MATERIALS AND METHODS

This research was conducted in Indonesia using climatic conditions, population density, and COVID-19 data from secondary resources. Climatic conditions data are collected from Indonesia's Environmental Statistics 2019 report (BPS 2019). Due to the limitation of data in Indonesia, we used the 2019 climate conditions in this study. Population density data is collected from the Indonesia Statistics 2020 report (BPS 2019). COVID-19 data used in this study is a cumulative data of total, recovered, and death cases from early March until July 22, 2020. The data collected from COVID-19 Daily Media Report, Indonesian Health Ministry. There were 34 Province in Indonesia, which illustrated in Figure 1.

This study used computer-based data processing with a statistical approach. There were two analyses conducted in this study: K-means cluster and Fisher's exact test. K-means cluster was used to determine the clusters of climatic conditions, population density, and COVID-19. Fisher's exact test is used to determine the correlation between the climatic condition and COVID-19 and the correlation between population density and COVID-19.

RESULTS AND DISCUSSION



44 FIGURE 1. Indonesia Map Source: https://peta-hd.com/peta-34-provinsi-di-indonesia/

COVID-19 CLUSTER SPREADS IN INDONESIA K-means cluster is used to determine the COVID-19 clusters. There are three indicators used to determine the COVID-19 clusters; total, recovered, and death cases. Table 1 shows three COVID-19 clusters; low, medium, and high.

TABLE 1. COVID-19 cluster cases positive, recovered, and

	COVID-19 Clusters										
Measure		Low		Medium			High				
	Total cases	Recovered	Deaths	Total cases	Recovered	Deaths	Total cases	Recovered	Deaths		
Min	136	75	1	3149	1472	144	17621	10460	751		
Max	8407	5034	287	7726	3720	444	19093	11139	1496		
Mean \pm SD	$^{1185,54\pm}_{1677,51}$	674,61 ± 981,39	$^{41,39\pm}_{62,29}$	$5458 \pm 1880,95$	2441,75± 941,11	263,25 ± 129,77	$18357 \pm 1040,86$	$^{10799,5\pm}_{480,12}$	1123,5 ± 526,79		

deaths

From Table 1, low clusters are cluster with lower total COVID-19 cases than the medium and high cluster. The recovery rate means recovered cases per mean total cases in each cluster in the low cluster are 56.9%. It is higher than the medium cluster with a 44.73% recovery rate and lower than the high cluster is 58.83%. The case fatality rate means death cases per mean total cases in each cluster; in the low cluster, they are lower with a 3.49% than the medium cluster 4.82% and high cluster 6.12%.

From the cluster analysis, 28 provinces classified in low clusters are Aceh; Bali; Banten; Bangka Belitung; Bengkulu; DI Yogyakarta; Jambi; West Kalimantan; East Kalimantan; Central Kalimantan; North Kalimantan; Riau Islands; West Nusa Tenggara; South Sumatra; West Sumatra; North Sulawesi; North Sumatra; Southeast Sulawesi; Central Sulawesi; Lampung; Riau; North Maluku; Maluku; West Papua; Papua; West Sulawesi; East Nusa Tenggara; and Gorontalo. Four provinces classified in medium clusters are West Java Province; Central Java; Kalimantan Selatan; and South Sumatra. Two clusters are classified in high clusters are East Java and Jakarta.

Figure 2 shows the six provinces in Indonesia with the highest number of total cases: East Java 19,093; Jakarta 17,621; South Sulawesi 8,407; Central Java 7,726; West Java 5,741; and South Kalimantan 5,216.




FIGURE 2. Number of COVID-19 cases spread across 34 provinces in Indonesia

CORRELATION BETWEEN CLIMATIC CONDITIONS (TEMPERATURE, HUMIDITY, AND RAINFALL) AND COVID-19 IN INDONESIA

humidity. The three indicators used different units, so a value standardization is performed. There are two Climatic Conditions clusters; Climatic Conditions I and Climatic Conditions II (Table 2).

Climatic Conditions (Temperature, Humidity, and Rainfall) In Indonesia

There are three climate indicators used to determine the Climatic Conditions cluster; air temperature, rainfall, and

TABLE 2. Climate clusters (temperature, rainfall, and hu	midity) in climatic conditions	s i and climatic conditions ii of 34 provinces
	in Indonesia	

Measure			Cli	mate clusters				
	C	limatic conditions	I	Climatic conditions II				
	Temperatures (°C)	Rainfall (mm/year)	Humidity (%)	Temperatures (°C)	Rainfall (mm/year)	Humidity (%)		
Min	26,7	1585	80	25,9	490	67		
Max	28,3	3823	87	29,5	3048	82		
Mean \pm SD	$27{,}3\pm0{,}44$	$2693,5 \pm 597,88$	$83,\!19\pm2,\!07$	$28,22 \pm 0,91$	$1840,5\pm 578,81$	$76,\!72\pm3,\!94$		

The mean air temperature in Climatic Conditions I is lower than Climatic Conditions II, while the mean rainfall and humidity in Climatic Conditions I are higher than the Climatic Conditions II. These show that Climatic Conditions I is a colder area than Climatic Conditions II. From the cluster analysis, 16 provinces within Climatic conditions I are Aceh; West Sumatra; Riau; Jambi; Bangka Belitung; Riau Islands; West Kalimantan; Central Kalimantan; North Kalimantan; North Sulawesi; South Sulawesi; Southeast Sulawesi; Maluku; North Maluku;

West Papua; Papua; and 18 provinces within Climatic conditions II are North Sumatra; South Sumatra; Bengkulu; Lampung; Jakarta; West Java; Central Java; Yogyakarta; East Java; Banten; Bali; West Nusa Tenggara; East Nusa Tenggara; South Kalimantan; East Kalimantan; Central Sulawesi; Gorontalo; West Sulawesi. This finding shows that 52.9% of Indonesia's area indicates having a high mean air temperature, low rainfall and humidity, included in Climatic conditions II (Table 3).

TABLE 3. The difference of cluster COVID-19 based on climate climatic conditions

	Cluster COVID-19						T- (-1		P	Correlation
Climatic conditions	Low		Medium		High		- Total		P_{value}	coefficient
	n	%	n	%	n	%	n	%		
Climatic conditions I	16	100	0	0	0	0	16	100		
Climatic conditions II	12	66,7	4	22,2	2	11,1	18	100	0,034	0,400
Total	28	82,4	4	11,8	2	5,9	34	100		

Our study found a correlation between Climatic Conditions and COVID-19 clusters p=0.034 with a positive correlation coefficient of 40%, indicates a strong correlation between climatic conditions and COVID-19. It shows a linear correlation, that higher climate conditions led to higher COVID-19 cases. All provinces in Climatic

Conditions I have low COVID-19 cases. We predict that the coronavirus can live under high temperatures. Indonesia's mean air temperature ranges from 26.7-29.5 °C and COVID-19 spread in Indonesia is also high. The condition is not according to the previous studies state that coronavirus can survive with lower air temperatures 5-11 °C (Sajadi et al. 2020). However, Indonesia's humidity level is relatively high, ranging from 67-87% to the range of humidity, which is considered following the coronavirus environment (44-88%) (Sajadi et al. 2020). The humidity level in Indonesia is suitable for the survival of the coronavirus. We predict humidity is an essential climate factor in the survival of the coronavirus.

A study conducted in Brazil showed that warm temperature (above 25 °C) were not associated with the declined of the COVID-19 cases (Prata et al. 2020). Pan et al. (2020) suggest that warmer weather unlikely to reduce the COVID-19 transmissions. Although a surge in COVID-19 cases in the hot and humid region is not as rapid as in the cold and dry region, COVID-19 cases in the hot and humid region keep increasing consistently. Warm weather alone may not be able to stop the COVID-19 transmissions and several other factors other than meteorological factors should be taken into account (Bukhari et al. 2020). The spread of COVID-19 appears to be lower in warm and wet climates, but the certainty of the resulting evidence is rated low. Temperature and humidity alone do not explain most of the variability of the COVID-19 outbreak. The spread of COVID-19 may be influenced by other factors such as public isolation policy, herd immunity, migration patterns, population density, and cultural aspects (Macenas et al. 2020).

CORRELATION BETWEEN POPULATION DENSITY (POPULATION AND AREA OF PROVINCE) AND COVID-19 IN INDONESIA

Population Density (Population and Area of Province) In Indonesia

Various studies on population density correlation and COVID-19 spread are conducted in the world. Some studies found that population density correlates with COVID-19 distribution rates, while others found the opposite. According to research conducted in China, population density is not related to COVID-19 events. However, Sun et al. (2020) stated that territorial restriction is a useful measure in reducing the chance of infection in the community. Territorial restrictions, commonly referred to as regional quarantine, are traditional solutions used in antiquity to control the spread of infectious diseases (Wilder-Smith & Freedman 2020). This step is useful to limit communities' interaction within the region so that the virus does not increasingly spread mainly in densely populated areas.

High-density regions can cause the spread of the SARS-Cov-2 virus faster than lower density regions, based on research conducted in Japan (Sajadi et al. 2020). High-density levels increase the likelihood of physical contact with others due to increasingly limited wiggle room (Ramadhani 2020). Besides, population in densely populated areas will have less chance of treating properly if the health service in the region is inadequate inpatient surges. Population density is related to the number of COVID-19 cases, the case fatality rate from COVID-19, and the willingness of health services in the region (Amoo et al. 2020) (Table 4).

TABLE 4. Population density of 34 provinces in Indonesia

Measure				
Measure	Low	Medium	High	
Min	9	750	15900	
Max	273	1394	15900	
Mean \pm SD	$101,15 \pm 77,79$	1099,67 ± 266,77	$15900 \pm .*$	

Standard Deviation unable to calculate, as there is only 1 Province

16

We divided population density into three categories; low, medium, and high. Population density is the amount that represents how dense the population in one particular area. Population density is calculated by dividing the population by the area. There is a large discrepancy in high population density compared to the low and medium population.

There are 27 low density provinces, Aceh; North Sumatra; West Sumatra; Riau; Jambi; South Sumatra; Bengkulu; Lampung; Bangka Belitung; Riau Islands; West Nusa Tenggara; East Nusa Tenggara; West Kalimantan; Central Kalimantan; South Kalimantan; East Kalimantan; North Kalimantan; North Sulawesi; Central Sulawesi; South Sulawesi; North Sulawesi; Gorontalo; West Sulawesi; Maluku; North Maluku; West Papua; and Papua. The six medium density provinces are West Java; Central Java; Yogyakarta; East Java; Banten; Bali. The highdensity province is Jakarta. The province in Java Island has a medium to high population density (Figure 3).




FIGURE 3. Population density of 34 provinces in Indonesia

Jakarta province has a high population of 15,900 people/km² with population is 10,557,800 and 664.01 km². Jakarta is one of the metropolitan areas with a dense population. In addition to having a dense population, Jakarta is one of the areas with a high positive case. Most metropolitan areas in different countries have dense populations. New York is one of the states of the United States with high population density and positive cases. In research conducted in the United States, New York accounted for 37% of positive cases in the United States. In addition to New York, other metropolitan areas such as Chicago, Washington, and Los Angles are also among the top ten densely populated metropolitan areas contributing to positive cases in the United States (Zhang & Schwartz 2020) (Table 5).

		Cluster COVID-19								Correlation
Population density	Low		Medium		High		Total		\mathbf{P}_{value}	coefficient
	n	%	n	%	n	%	n	%		
Low	25	92,6	2	0	0	0	27	100		0,633
Medium	3	50	2	33,3	1	16,7	6	100		
High	0	0	0	0	1	100	1	100	0,004	
Total	28	82,4	4	11,8	2	5,9	34	100		

TABLE 5. The difference of Cluster COVID-19 Based on Population Density

Our study found a correlation between population density and COVID-19 with p=0.004 and a positive correlation coefficient of 63.3%. Based on cross-tabulation results, 92.6% of low-density provinces have low COVID-19 cases. As well as 100% high-density have high COVID-19. This research is in line with research (Sarmadi et al. 2020) that population density in the United States affects deaths from COVID-19 in the United States. The greater the population density, the greater the mortality rate in the United States. Similar research was conducted in India, which ranks in the top ten of the world's top population as Indonesia. The research shows an association between population density and COVID-19 spread (Bhadra et al. 2020). This result is similar to this study and the United States study, which shows that population density has a role in the spread of COVID-19.

SIX PROVINCES WITH THE HIGHEST COVID-19 CASES, CLIMATE CONDITIONS AND POPULATION DENSITY

The Top Six Provinces with the Highest Cases, Climatic Conditions and Population Density

Our study found that the six provinces with the highest COVID-19 cases were in Climatic conditions II (high mean air temperature with low rainfall and humidity) (Table 6). The findings are in stark contrast, as research analyzing the effects of climate and weather on COVID-19 cases found that countries with high mean air temperatures and low humidity have the effect of lowering the number of deaths caused by COVID-19. A study found that tropical regions with high mean temperature and high relative humidity, i.e., India, Pakistan, Saudi Arabia, Malaysia, Thailand, and Australia, have low death cases and high recovery rates (Pavani & Sunalini 2020). Muhammad Mazhar et al. (2020) also discovered that tropical regions, i.e. Malaysia, Thailand, and Indonesia, are less affected by the coronavirus. However, Muhammad Mazhar et al. (2020) study the regional climatic condition's effect by June 2020. Some researchers also conducted a study in the subtropical climate, i.e. China, They also found that higher average air temperatures will lower the risk of COVID-19 (Ma et al. 2020; Wang et al. 2020). However, Indonesia's high mean temperature did not follow the same conditions related to COVID-19 in other regions. Coronavirus will still last inside the human body and infect other humans during the incubation period. The more frequent human activities outside the house, the more the virus will likely spread and infect other humans.

Based on our results, the top 6 provinces with the highest COVID-19 cases are located in a fairly dense and dense region. As a high-density region, Jakarta is the region with the second-highest COVID-19 cases, while East Java is in the first position is a fairly dense region. The rest following a sequence order are West Java, Central Java, South Kalimantan, and South Sumatra. This region is fairly dense. Our result found a similar study conducted by Jawad (2020), which states that the results of an analysis conducted in countries vulnerable to COVID-19 spread show that population density affects the peak period of COVID-19 spread. However, population density is not an essential factor in the spread of COVID-19 under strict lockdown policies. China's lockdown policy could effectively limit the speed at which COVID-19 spreads (Sun 2020).

TABLE 6. Climatic conditions and population density in six provinces with the highest COVID-19 cases

No.			Cluster					
INO.	Province	Covid 19	Climatic conditions	Population density				
1.	East Java	High Cases	П	Medium				
2.	Jakarta	High Cases	П	High				
3.	West Java	Medium Cases	П	Medium				
4.	Central Java	Medium Cases	П	Medium				
5.	South Kalimantan	Medium Cases	П	Medium				
6.	South Sumatra	Medium Cases	П	Medium				

CONCLUSION

Indonesia region is identified in high air temperature with 26.7-29.5 °C and high humidity (67-87%), due to its tropical climate. However, the majority of the province in Indonesia are classified in Climatic Conditions II in 52,9%. Climatic conditions II is a Climatic condition with higher mean air temperatures, lower rainfall and humidity than the rest. Indonesia is also known as a highly-dense country. Jakarta is the most populated region in Indonesia (population density: 15,900). Our results found a correlation between the climatic conditions and the COVID-19 cluster (p:0,034), and there is a correlation between population density and COVID-19 clusters (p:0,004). Population density is more related to the high number of COVID-19 cases in Indonesia, which has a positive correlation of 63.3%. The top six provinces that have the highest COVID-19 cases in Indonesia are classified in Climatic Conditions II and a fairly-dense to densely populated region. There need to be adaptation and mitigation measures in COVID-19 control related to climate conditions in Climatic conditions II and high population density so that policies in their respective provinces in crowd control, social distancing, and efforts to reduce COVID-19 cases through increased community-based awareness.

ACKNOWLEDGEMENTS

We would like to thank the Department of Environmental Health and Environmental Health Master Program, Faculty of Public Health, Universitas Airlangga which has facilitated the activities of writing scientific articles in the framework of international journal publication. We would also thank Prof. Mohd. Talib Latif, Universiti Kebangsaan Malaysia who guided and corrected this research until this article was compiled.

REFERENCES

- Amoo, E.O., Adekeye, O., Olawole-Isaac, A., Fasina, F., Adekola, P.O., Samuel, G.W., Akanbi, M.A., Oladosun, M. & Azuh, D.E. 2020. Nigeria and Italy divergences in coronavirus experience: Impact of population density. *Scientific World Journal* 2020: 8923036. DOI: 10.1155/2020/8923036.
- Bhadra, A., Mukherjee, A. & Sarkar, K. 2020. Impact of population density on Covid-19 infected and mortality rate in India. *Modeling Earth Systems and Environment* 7: 623-629.
- BPS. 2020. Statistics Indonesia Statistical Year Book of Indonesia 2020. Accessed by https://drive.google.com/ file/d/18EfhKxnIARMh0h_4qqWsEszYsk_X0W8G/view.
- BPS. 2019. Population by Age and Gender Group 2019. Indonesian Central Bureau of Statistics. https://www.bps. go.id/indikator/indikator/view_data_pub/0000/api_pub/58/ da_03/1.
- BPS. 2019. Indonesia Environmental Statistics 2019. https://www.bps.go.id/publication/2019/12/13/ e11bfc8ff8392e5e13a8cff3/statistik-lingkungan-hidupindonesia-2019.html.
- Bukhari, Q., Massaro, J.M., D'Agostino Sr., R.B. & Sheraz Khan. 2020. Effects of weather on coronavirus pandemic. *International Journal of Environmental Research* and Public Health 17(15): 5399. https://doi:10.3390/ ijerph17155399.
- CDC. SARS-CoV-2 & Potential Airborne Transmission [Internet]. 2020. Center for Disaese Control and Prevention. https://www.cdc.gov/coronavirus/2019-ncov/more/scientificbrief-sars-cov-2.html.
- Coşkun, H., Yıldırım, N. & Gündüz, S. 2020. The spread of COVID-19 virus through population density and wind in Turkey cities. *Science of The Total Environment* 751(10): 141663.
- Dangi, R.R. & George, M.A. 2020. Review on theories and models of disease causation for COVID-19. *Journal SSRN Electron 2020* https://doi.org/10.2139/ssm.3584080.

- Jawad, A.J. 2020. Effectiveness of population density as natural social distancing in COVID19 spreading. *Ethics Medicine and Public Health* 15(C): 100556.
- Kadi, N. & Khelfaoui, M. 2020. Population density, a factor in the spread of COVID-19 in Algeria: Statistic study. *Bulletin of the National Research Centre* 44: 138. https://doi.org/10.1186/ s42269-020-00393-x.
- Ma, Y., Zhao, Y., Liu, J., He, X., Wang, B., Fu, S., Yan, J., Niu, J., Zhou, J. & Luo, B. 2020. Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Science of The Total Environment* 724: 138226.
- Mecenas, P., da Rosa Moreira Bastos, R.T., Rosário Vallinoto, A.C. & Normando, D. 2020. Effects of temperature and humidity on the spread of COVID-19: A systematic review. *PLoS ONE* 15(9): e0238339.
- Ministry of Health of Indonesia. 2020. Daily Media Report Covid-19 dated July 22, 2020 at 12.00 WIB.
- Ministry of Health of Indonesia. 2020. COVID-19 Transmission Prevention Guide. https://promkes.kemkes.go.id/panduanpencegahan-penularan-covid-19-untuk-masyarakat.
- Muhammad Mazhar Iqbal, Irfan Abid, Saddam Hussain, Naeem Shahzad, Muhammad Sohail Waqas & Muhammad Jawed Iqbal. 2020. The effects of regional climatic condition on the spread of COVID-19 at global scale. *Sci. Total Environ.* 15(739): 140101.
- National Health Commission of People's Republic of China. 2020. Pneumonia Diagnosis and Treatment of 2019-nCoV Infection from Chinese NHC and CDC 2020.
- Pan, J., Yao, Y., Liu, Z., Meng, X., John, S. Ji., Qiu, Y., Wang, W., Zhang, L., Wang, W. & Kan, H. 2020. Warmer weather unlikely to reduce the COVID-19 transmission: An ecological study in 202 locations in 8 countries. *Science of The Total Environment* 753: 142272.
- Pavani, A. & Sunalini, K.K. 2020. Effects of temperature and relative humidity on Covid-19. *Journal of Critical Reviews* 7(9): 3177-3182.
- Prata, D.N., Rodrigues, W. & Bermejo, P.H. 2020. Temperature significantly changes COVID-19 transmission in (sub) tropical cities of Brazil. *Science of The Total Environment* 729: 138862.
- Ramadhani, F.H. 2020. Literature review: Healthy home as the new normal for Covid-19 prevention. Jurnal Kesehatan Lingkungan doi: 10.20473/jkl.v12i1si.2020.1-10.
- Sajadi, M.M., Habibzadeh, P., Vintzileos, A., Shokouhi, S., Miralles-Wilhelm, F. & Amoroso, A. 2020. Temperature, humidity, and latitude analysis to estimate potential spread and seasonality of Coronavirus Disease 2019 (COVID-19). JAMA Network Open 3(6): e2011834.
- Sarmadi, M., Moghanddam, V.K., Dickerson, A.S. & Martelletti, L. 2020. Association of COVID-19 distribution with air quality, sociodemographic factors, and comorbidities: An ecological study of US states. *Air Quality, Atmosphere & Health* https://doi.org/10.1007/s11869-020-00949-w.

- Sun, Z., Zhang, H., Yang, Y., Wan, H. & Wang, Y. 2020. Impacts of geographic factors and population density on the COVID-19 spreading under the lockdown policies of China. *Science of The Total Environment* 746(666): 141347.
- Therese, K.L.S., White, L.F. & Nichols, B. 2020. Population density and basic reproductive number of COVID-19 across United States counties. *National Library of Medicine* https:// doi.org/10.1101/2020.06.12.20130021.
- Wang, H., Chen, C. & Lin, Q. 2020. Short-term effects of ambient temperature and relative humidity on the risk of COVID-19 and SARS in Guangzhou, China: A time-series analysis. *Research Square* doi: 10.21203/rs.3.rs-50445/v1.
- Wang, J. & Du, G. 2020. COVID-19 may transmit through aerosol. Irish Journal of Medical Science 89: 1143-1144.
- WHO. 2020a. Infection prevention and control guidance for long-term care facilities in the context of COVID-19. *Interim Guidance of World Health Organization*. https://apps.who. int/iris/handle/10665/331508.
- WHO. 2020b. WHO Corona Disease (COVID-19) Dashboard. https://covid19.who.int/.
- Wilder-Smith, A. & Freedman, D.O. 2020. Isolation, quarantine, social distancing and community containment: Pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak. *Journal of Travel Medicine* 27(2): 1-4. https://doi.org/10.1093/jtm/taaa020.
- Zhang, C.H. & Schwartz, G.G. 2020. Spatial disparities in coronavirus incidence and mortality in the United States: An ecological analysis as of May 2020. *Journal of Rural Health* 36(3): 433-445.

R. Azizah*, Santi Martini, Lilis Sulistyorini, Mahmudah, Aditya Sukma Pawitra, Stefanny Surya Nagari, Cendana Fitrahanjani, Fairuz Haniyah Ramadhani & Mohd. Talib Latif Faculty of Public Health

Universitas Airlangga Surabaya Indonesia

Didik Budijanto

The Ministry of Health of the Republic of Indonesia Jakarta Selatan 12950 Jakarta Indonesia

Mohd. Talib Latif Faculty of Science and Technology

Universiti Kebangsaan Malaysia 43600 UKM Bangi, Selangor Darul Ehsan Malaysia

*Corresponding author; email: azizah@fkm.unair.ac.id

Received: 5 November 2020 Accepted: 17 December 2020

Association between Climatic Conditions, Population Density and COVID-19 in Indonesia

ORIGINALITY REPORT

	0% ARITY INDEX Y SOURCES	13% INTERNET SOURCES	18% PUBLICATIONS	0% STUDENT P/	APERS
1	Gündüz. through Turkey ci	oşkun, Nazmiye "The spread of population den ties", Science o hent, 2020	COVID-19 vir sity and wind	us	1 %
2	Angioi, G Napodar respirato solid was and effec settings"	Bellizzi, Paola M ualtiero Canu, no, Antonio Mai ory syndrome co ste treatment: A tive strategies , Waste Manag nal for a Sustain , 2021	Catello M Par nca. "Severe a oronavirus-2 i A need for effi in low-resour ement & Rese	nu acute medical icient iced arch:	1 %
3	agupubs	.onlinelibrary.w	viley.com		1%
4		srullah, Muhan dayani, Lilis Roh		Retno	1%

"Psychological impact among health workers in effort to facing the COVID-19 in Indonesia", International Journal of Public Health Science (IJPHS), 2021 Publication

1%

1%


1%

1%

philpapers.org Internet Source
ukmsarjana.ukm.my Internet Source
"Analyze K-Value Selected Method of K-Means Clustering Algorithm to Clustering Province Based on Disease Case", International Journal of Innovative Technology and Exploring

Engineering, 2020

- Ramadhan Tosepu, Joko Gunawan, Devi
 Savitri Effendy, La Ode Ali Imran Ahmad,
 Hariati Lestari, Hartati Bahar, Pitrah Asfian.
 "Correlation between weather and Covid-19
 pandemic in Jakarta, Indonesia", Science of
 The Total Environment, 2020
 Publication
- 9 Singgih Tri Sulistiyono. "The Expulsion of KPM and its Impact on the Inter-island Shipping and Trade in Indonesia, 1957–1964", Itinerario, 2010 Publication



16 Nurjaeni Nurjaeni, Yothin Sawangdee, Umaporn Pattaravanich, Charamporn

Holumyong, Aphichat Chamratrithirong. "The Role of Structural and Process Quality of Family Planning Care in Modern Contraceptive Use in Indonesia: A Multilevel Analysis", Research Square Platform LLC, 2021 Publication

17	archive.org Internet Source	<1%
18	Siew Bee Aw, Bor Tsong Teh, Gabriel Hoh Teck Ling, Pau Chung Leng, Weng Howe Chan, Mohd Hamdan Ahmad. "The COVID-19 Pandemic Situation in Malaysia: Lessons Learned from the Perspective of Population Density", International Journal of Environmental Research and Public Health, 2021 Publication	<1%
19	www.thelandmark.com	<1%
20	www.ukm.my Internet Source	<1%

21 pubmed.ncbi.nlm.nih.gov

www.cambridge.org

22

23

yourcountrychoice.blogspot.com

<1%

24	Arpad A. Vass, Rob R. Smith, Cyril V. Thompson, Michael N. Burnett et al. "Decompositional Odor Analysis Database", Journal of Forensic Sciences, 2004 Publication	<1%
25	Ketut Prasetyo. "Measuring the Achievement of the Unemployment and Poverty Target in the Sustainable Development Goals Program in Pandemi Covid-19", Walter de Gruyter GmbH, 2020 Publication	<1%
26	Te Faye Yap, Colter J. Decker, Daniel J. Preston. "Effect of daily temperature fluctuations on virus lifetime", Science of The Total Environment, 2021 Publication	<1%
27	Zhaomin Dong, Xiarui Fan, Jiao Wang, Yixin Mao, Yueyun Luo, Song Tang. "Data-related and methodological obstacles to determining associations between temperature and COVID-19 transmission", Environmental Research Letters, 2021 Publication	<1%
28	environmentportal.in Internet Source	<1%
29	www.ncbi.nlm.nih.gov Internet Source	<1%

- Jie Yuan, Yu Wu, Wenzhan Jing, Jue Liu, Min Du, Yaping Wang, Min Liu. "Association between meteorological factors and daily new cases of COVID-19 in 188 countries: A time series analysis", Science of The Total Environment, 2021 Publication
- 31 Keerthi Sasikumar, Debashis Nath, Reshmita Nath, Wen Chen. "Impact of extreme hot climate on COVID - 19 outbreak in India", GeoHealth, 2020

Publication

32

mdpi-res.com

- Abderrazek Ben Maatoug, Mohamed Bilel Triki, Hesham Fazel. "How do air pollution and meteorological parameters contribute to the spread of COVID-19 in Saudi Arabia?", Environmental Science and Pollution Research, 2021 Publication
- Amir Reza Khavarian-Garmsir, Ayyoob Sharifi, Nabi Moradpour. "Are high-density districts more vulnerable to the COVID-19 pandemic?", Sustainable Cities and Society, 2021 Publication



Declan C. Murphy, Sergio Razzano, Ryckie G. Wade, Richard M. Haywood, Andrea Figus. "Inferior Gluteal Artery Perforator (IGAP) flap versus Profunda Artery Perforator (PAP) flap as an alternative option for free autologous breast reconstruction", Journal of Plastic, Reconstructive & Aesthetic Surgery, 2021 Publication

<1 %

- Jayanthi Devaraj, Rajvikram Madurai Elavarasan, Rishi Pugazhendhi, G.M. Shafiullah et al. "Forecasting of COVID-19 cases using Deep learning models: Is it reliable and practically significant?", Results in Physics, 2021 Publication
- Meiting Tu, Wenxiang Li, Olivier Orfila, Ye Li, Dominique Gruyer. "Exploring nonlinear effects of the built environment on ridesplitting: Evidence from Chengdu", Transportation Research Part D: Transport and Environment, 2021 Publication
- Najaf Iqbal, Zeeshan Fareed, Farrukh Shahzad, Xin He, Umer Shahzad, Ma Lina. "Nexus between COVID-19, temperature and exchange rate in Wuhan City: New findings from Partial and Multiple Wavelet Coherence", Science of The Total Environment, 2020

39	Nir Barak, Udi Sommer, Nir Mualam. "Urban attributes and the spread of COVID-19: The effects of density, compliance and socio- political factors in Israel", Science of The Total Environment, 2021 Publication	<1%
40	Sukanta Malakar. "Geospatial modelling of COVID-19 vulnerability using an integrated fuzzy MCDM approach: a case study of West Bengal, India", Modeling Earth Systems and Environment, 2021 Publication	<1%
41	Tamer Oraby, Michael G Tyshenko, Jose Campo Maldonado, Kristina Vatcheva et al. "Modeling the Effect of Lockdown Timing as a COVID-19 Control Measure in Countries with Differing Social Contacts", Cold Spring Harbor Laboratory, 2020 Publication	<1%
42	depot.gdnet.org Internet Source	<1%
43	jurnal.undhirabali.ac.id	<1%
44	peta-hd.com Internet Source	<1%

Charlie H. Zhang, Gary G. Schwartz. "Spatial Disparities in Coronavirus Incidence and Mortality in the United States: An Ecological Analysis as of May 2020", The Journal of Rural Health, 2020 Publication

<1%

<1 %

- Dimitrios Kapetanios, Maria Petropoulou, Dimitrios Chatzelas, Georgios Pitoulias et al. "The Effect of Transfusion of Two Units of Fresh Frozen Plasma on the Perioperative Fibrinogen Levels and the Outcome of Patients Undergoing Elective Endovascular Repair for Abdominal Aortic Aneurysm", Annals of Vascular Surgery, 2021 Publication
- Jennifer Viezzer, Daniela Biondi. "The influence of urban, socio-economic, and ecoenvironmental aspects on COVID-19 cases, deaths and mortality: A multi-city case in the Atlantic Forest, Brazil", Sustainable Cities and Society, 2021 Publication
- Juan Qiu, Rendong Li, Dongfeng Han, Qihui Shao, Yifei Han, Xiyue Luo, Yanlin Wu. "A multiplicity of environmental, economic and social factor analyses to understand COVID-19 diffusion", One Health, 2021 Publication

49	Mengyang Liu, Mengmeng Liu, Zhiwei Li, Yingxuan Zhu, Yue Liu, Xiaonan Wang, Lixin Tao, Xiuhua Guo. "The spatial clustering analysis of COVID-19 and its associated factors in mainland China at the prefecture level", Science of The Total Environment, 2021 Publication	<1%
50	Supari Supari, Danang Eko Nuryanto, Amsari Mudzakir Setiawan, Furqon Alfahmi et al. "The association between initial COVID-19 spread and meteorological factors in Indonesia", Environmental Sustainability, 2021 Publication	<1%
51	covid.yale.edu Internet Source	<1%
52	dugi-doc.udg.edu Internet Source	<1%
53	koreascience.or.kr Internet Source	<1%
54	m.scirp.org Internet Source	<1%
55	peerj.com Internet Source	<1%
56	photography-now.com Internet Source	<1%

repub.eur.nl



⁵⁸ Álvaro Briz-Redón, Ángel Serrano-Aroca. "The effect of climate on the spread of the COVID-19 pandemic: A review of findings, and statistical and modelling techniques", Progress in Physical Geography: Earth and Environment, 2020 Publication

59 Mahesh Jayaweera, Hasini Perera, Buddhika Gunawardana, Jagath Manatunge. "Transmission of COVID-19 virus by droplets and aerosols: A critical review on the unresolved dichotomy", Environmental Research, 2020 Publication

- 60 Matthew Groves, Alexander Domling, Angel Jonathan Ruiz Moreno, Atilio Reyes Romero et al. "Gliptin Repurposing for COVID-19", American Chemical Society (ACS), 2020 Publication
- 61 Álvaro Briz-Redón, Ángel Serrano-Aroca. "A spatio-temporal analysis for exploring the effect of temperature on COVID-19 early evolution in Spain", Science of The Total Environment, 2020 Publication

62	"COVID in the Islands: A comparative perspective on the Caribbean and the Pacific", Springer Science and Business Media LLC, 2021 Publication	<1 %
63	"Impact of AI and Data Science in Response to Coronavirus Pandemic", Springer Science and Business Media LLC, 2021 Publication	<1%
64	Alessandro Camerotto, Andrea Sartorio, Anna Mazzetto, Milena Gusella et al. "Early Phase Management of the SARS-CoV-2 Pandemic in the Geographic Area of the Veneto Region, in One of the World's Oldest Populations", International Journal of Environmental Research and Public Health, 2020 Publication	<1%

Exclude	quotes	Off
Exclude	bibliography	On

Exclude matches Off

Association between Climatic Conditions, Population Density and COVID-19 in Indonesia

GRADEMARK REPORT		
FINAL GRADE	GENERAL COMMENTS	
/0	Instructor	
PAGE 1		
PAGE 2		
PAGE 3		
PAGE 4		
PAGE 5		
PAGE 6		
PAGE 7		
PAGE 8		
PAGE 9		