

MALAYSIAN JOURNAL OF
Medicine and Health Sciences
Vol. 17 No. 2, April 2021

MALAYSIAN JOURNAL OF

Medicine and Health Sciences

Vol. 17 No. 2 April 2021

Malaysian Journal of Medicine and Health Sciences Vol. 17 No. 2, April 2021



Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://medic.upm.edu.my/>

UPM Press
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://penerbit.upm.edu.my>

PENERBIT
UPM
UNIVERSITI PUTRA MALAYSIA
P R E S S



A scientific journal published by Universiti Putra Malaysia Press

About the Journal

The Malaysian Journal of Medicine and Health Sciences (MJMHS) is published by the Faculty of Medicine and Health Sciences, Universiti Putra Malaysia. The main aim of the MJMHS is to be a premier journal on all aspects of medicine and health sciences in Malaysia and internationally. The focus of the MJMHS will be on results of original scientific research and development, emerging issues and policy analyses pertaining to medical, biomedical and clinical sciences. The Malaysian Journal of Medicine and Health Sciences is now indexed in the following data bases: Scopus, EBSCOhost, ISC, and Rubriq.

Editorial Board

Editor in Chief:

Prof. Dr. Normala Ibrahim

Members:

Prof. Dr. Rukman Awang Hamat

Prof. Dr. Christopher Lim Thiam Seong

Prof. Dr. Ching Siew Mooi

Prof. Dr. Sharmili Vidyadaran

Assoc. Prof. Dr. Rajesh Ramasamy

Assoc. Prof. Dr. Sethu Thakachy Subha

Assoc. Prof. Dr. Loh Su Peng

Assoc. Prof. Dr. Nor Afiah Mohd Zulkefli

Assoc. Prof. Dr. Wan Aliaa Wan Sulaiman

Assoc. Prof. Dr. Ling King Hwa

Assoc. Prof. Dr. Habibah Abdul Hamid

Assoc. Prof. Dr. Zubaidah Nor Hanipah

Assoc. Prof. Dr. Intan Hakimah Ismail

Assoc. Prof. Dr. Ho Yu Bin

Assoc. Prof. Dr. Karmegam Karuppiah

Assoc. Prof. Dr. Subapriya Suppiah

Technical Coordinator:

Dr. Tan Kit Aun

International Advisory Board

Prof. Dr. Pierce Anthony Grace

Prof. Dr. David Isaacs

Prof. Dr. P. Ronan O'Connell

Prof. Dr. Graham Alexander Mc Gregor

Prof. Dr. Tan Ser Kiat

Prof. Dr. Gregory Y.H. Lip

Prof. Dr. Roger Pepperell

Malaysian Journal of Medicine and Health Sciences Vol. 17(2), April 2021

Contents

EDITORIAL

The Faculty of Medicine and Health Sciences - Charting the Future 1
Zamberi Sekawi

ORIGINAL ARTICLES

Quantification of Muscle Metabolites Using Proton Magnetic Resonance Spectroscopy (¹H-MRS) for Incomplete Spinal Cord Injury Patients: Preliminary Study 4
Nur Syahfinaz Hidayah Rusli, Faikah Zakaria, Hairil Rashmizal Abdul Razak

The Proportion of Non-retention and Its Associated Factors Among Clients Receiving Methadone Maintenance Therapy in Kelantan, Malaysia 11
Mohd Hazwan Baharuddin, Mohd Azhar Mohd Yasin, Mohd Nazri Shafei

The Outcomes of Team-based Learning on Test Performance, Cognitive Engagement and Motivation in Preclinical Anatomy Practical Classes 18
Shamsi Amalina Shamsuddin, Fazlina Kasim, Jamilah Al-Muhammady Mohammad, Muhamad Saiful Bahri Yusoff, Siti Nurma Hanim Hadie

Survey on Knowledge and Practices in Erosive Tooth Wear Among Dental Practitioners in Kuantan, Pahang: A Preliminary Study 28
Noorhazayti Ab. Halim, Nur Atiyah Fakhri, Nur Izzati Aisyah Azhar, Md Mustafa Md-Muziman-Syah

Barriers to Physical Activity and Exercise Amongst Persons With Physical Disability in a Government-funded Teaching Hospital, Kuala Lumpur 34
Haidzir Manaf, Aishah Shaïd, Maria Justine, Hafifi Hisham, Nazirah Hasnan, Ummi Mohlisi Mohd Asmawi, Amirah Mustapa

Correlation Between Blood Glucose and Salivary Glucose in Type 2 Diabetes Mellitus Patients 40
Kaushalya Indunil Dharmakeerthi, Malaka Priyashan Ponweera, Evindu Hasanjana Moragoda, Lahiru Sandaruwan Galgamuwa, Kithsiri Jayasekara, Vidumini Kaluarachchi, Uditha Bulugahapitiya

The Prevalence of Candidiasis Oris, Tuberculosis, and Anemia in Hospitalized HIV Patients Admitted in Dr. Soetomo General Hospital, Surabaya, Indonesia 46
Erwin Astha Triyono, Feriawan Tan, Manik Retno Wahyunitisari

Factors Correlated With Internet Gaming Disorder Among Malaysian University Students 54
Nik Ruzyanei Nik Jaafar, Azlin Baharudin, Izzat Tajjudin, Ling Shiao Ling, Muhammad Amirul Safarudin, Don Shakir Sufia, Tan Ying Hui, Nur Hasya Zulkifli, Kit-Aun Tan

Tumour Growth Inhibition and Systemic Responses of Δ sopB Δ sopD Δ pipD Disrupted *Salmonella* Agona and *Salmonella* Typhimurium in Mice 63
Ubaidah Naim Taraq Naem Zia, Emy Sarah Ng Amar Ng, Mohd Amirudin Sidik, Mohamad Fauzi Mohd Idris, Khoo Chai Hoon, Sabrina Sukardi, Yeap Swee Keong, Cheah Yoke Kqueen

Dosimetric Study of Custom-made Pelvic Perspex Phantom using Single-Energy Mode (SECT) and Dual-Energy Mode (DECT) Computed Tomography 72
Siti Hajar Zuber, Nursyatina Abdul Raof, Rafidah Zainon, Abd Aziz Tajuddin

Association of Temperature and Rainfall with *Aedes* Mosquito Population in 17th College of Universiti Putra Malaysia 78
Banulata Gopalsamy, Latifah Saiful Yazan, Nissa Nabila Abdul Razak, Mustafa Man

Identify Prognostic Risk Factor of Breast Cancer in North West of Iran 85
Davoud Adham, Somaieh Matin, Payam Amini, Malek Abazari

Positive Predictive Value of Sonographic BI-RADS Final Assessment Categories for Breast Lesions 91
Nursyahrah @ Sheila Salinah Md Bohari, Norzailin Abu Bakar, Sharifah Majedah Idrus Alhabshi, Nur Yazmin Yaacob

Knowledge, Attitude and Practice on Vegetables Intake among Adolescents in Rural Terengganu, Malaysia 98
Mohd Adzim Khalili Rohin, Norhayati Abd Hadi, Sahirah Sariff, Siti Syarma Mohd Shariff, Norhaslinda Ridzwan, Mimie Noratiqah Jumli

Tuberculosis Infection and Incidence of Loss to Follow Up among HIV Patients at Saiful Anwar General Hospital, Indonesia: A Retrospective Study 106
Tika Dwi Tama, Rosita Dewi Ambarwati, Hartati Eko Wardani

Successful Academic Remediation of Undergraduate Medical Students for Exit Examination: Lessons Learned 112
Alam Sher Malik, Rukhsana Hussain Malik, Muhammad Najib Alwi

Psychological Impact of Covid-19 Pandemic on Mental Health Among Medical Students in Malaysia <i>Md Mizanur Rahman, Ai Ling Ang, Nithiya Lakshmi, Kishwani Hiram Chakraverty, Dyg Shafiqah, Krithiga Selvarajoo</i>	119
The Effect of Balance Exercise Training on Balance Status, and Quality of Life in Elderly Women: A Randomized Controlled Trial <i>Mohammad Hossein Kaveh, Farzaneh Bahadori, Afsaneh Doosti, Abdolrahim Asadollahi</i>	129
Work-related Quality of Life and Its Associated Factors Among House Officers Working at Two Hospitals in Selangor <i>Nadia Abd Gaffar, Zaliha Ismail, Hasidah Abdul-Hamid, Suraya Abdul-Razak</i>	137
Analysis of Heavy Metals (Cadmium, Chromium, Lead, Manganese, and Zinc) in Well Water in East Java Province, Indonesia <i>Mochammad Sholehuddin, R. Azizah, Arif Sumantri, Shaharuddin Mohd Sham, Zainul Amiruddin Zakaria, Mohd Talib Latif</i>	146
Isolation and Characterization of <i>Chromobacterium</i> sp. from Lake Water at Manipal International University <i>Pratheep Sandrasagar, Pavitraa Rajandrai, Mak Woei Loon, Hanan Hasan</i>	154
Effects of Pulsed Ultrasound with Low-Intensity on Recovery of Physical Impairments After Total Knee Arthroplasty: A Preliminary Quasi-Experimental Study <i>Munayati Munajat, Nor Azlin Mohd Nordin, Nor Hamdan Mohamad Yahaya, Ahmad Hafiz Zulkifly</i>	162
Inhibition Effect of Garlic (<i>Allium sativum</i>) Extract on <i>Streptococcus sanguinis</i> Biofilm Formation Involving Bacterial Motility Mechanism <i>Suryani Hutomo, Denise Utami Putri, Beatric Chindy Welviyanda, Heni Susilowati</i>	169
The Effect of Submaximal Incremental Running Test on Heart Rate Variability in University-level Male Football Athletes <i>Nurul Syafiqah Ahmad Naspi, Marilyn Li Yin Ong</i>	175
Development of a Patient Safety-training Program for Health Workers in Indonesia: Perspectives of Health Workers and Hospital Stakeholders <i>Inge Dhamanti, Taufik Rachman, Muhammad Ardian C.L, Nanda Aulya Ramadhan, Elida Zairina, Rista Fauziningtyas</i>	183
Food Craving and Symptoms of Premenstrual Syndrome among University Students <i>Nor Fatimah Abdullah, Aliza Haslinda Hamirudin, Suriati Sidek, Nur Aainna Amira Mat Hassan</i>	189
Mesopic Pupillometry in Pre-LASIK Patients by a Placido-Disc Topographer and Hartmann-Shack Aberrometer <i>Md Mustafa Md-Muziman-Syah, Muhammad Aiman Suhaimi, Umar Hakimi Sulaiman, Noorhazayti Ab. Halim, Ahmad Tajudin Liza-Sharmini, Khairidzan Mohd Kamal</i>	197
Features of an Ergonomic Chair Designed for Surgeons Performing Microscopic Surgeries <i>Shahrzad Riaei, Hadi Daneshmandi, Mohsen Razeghi, Bahram Kouhnavard, Zahra Zamanian</i>	203
Diabetic Knowledge of T2DM Patients in Hospital Pakar Sultanah Fatimah, Muar, Johor, Malaysia <i>C.H. Chua, Y.M. Chua, J.L. Cheong</i>	210
Types of Nutrition Resource Kit Needed by Community Living Elderly in Malaysian Health Clinic Setting: A Qualitative Study <i>Iman Nabihah Noor Azam, Aliza Haslinda Hamirudin, Sakinah Harith, Mohd Aznan Md Aris, Karimah Hanim Abd Aziz, Nurul Syaireen A Rashid</i>	217
Assessment of Bangladeshi People's Knowledge, Awareness, Precautionary Behavior and Preventive Actions During the COVID-19 Pandemic: A Cross-sectional Survey <i>Ashfia Tasnim Munia, K. M. Yasif Kayes Sikdar, Md. Anamul Haque, Mohammad Ashraful Islam, Md Raihan Sarkar</i>	225
Effects of Low Birth Weight Babies on Cochlear Function In Newborns at Dr Soetomo Hospital Surabaya <i>Wahyu Bagus Handoko, Risa Etikaa, Agus Hariantoa, Nyilo Purnami</i>	234
Fibronectin-binding Protein F1 (<i>prtF1</i>) Gene Is Highly Distributed in Tetracycline-resistant, Erythromycin-sensitive <i>Streptococcus pyogenes</i> Isolates <i>Simon Onyema Azi, Suresh Kumar, Leslie Than Thian Lung, Malina Osman, Rukman Awang Hamat</i>	239
REVIEW ARTICLES	
The Potential of Gallic Acid as a Radiosensitizer on Human Prostate Cancer: A Systematic Review of Preclinical Studies <i>Agung Tri Cahyono, Melva Louisa, Tiara Bunga Mayang Permata, Handoko, Endang Nuryadi, Henry Kodrat, Heri Wibowo, Agus Rizal Ardy Hariandy Hamid, Sri Mutya Sekarutami, Soehartati Argadikoesoema Gondhowiardjo</i>	246
Knowledge, Attitude and Practices (KAP) On Aedes Mosquito-Borne Diseases Amongst Community Members in Malaysia: A Review <i>Neevaarthana Subramaniam, Siti Marwanis Anua, Nor Fazila Che Mat</i>	255

Review on the In Vitro Cytotoxicity Assessment in Accordance to the International Organization for Standardization (ISO) <i>Muhammad Aminuddin Mohd Shafiee, Mohd Ashraf Muhamad Asri, Sharifah Sakinah Syed Alwi</i>	261
Investigating the Effect of Saffron on Anxiety, Fatigue, Labor Pain and Vaginal Childbirth: Review Article <i>Dina Abadibavil, Sareh Dashti</i>	270
Thyrotoxic Heart Failure: A Narrative Review of Pathophysiology and Principle Management <i>Yoga Alfian Noor, Eka Prasetya Budi Mulia, Jongky Hendro Prajitno</i>	276
COMMENTARY	
Empathetic Dentistry: Noble Opportunity to Restore Broken Hearts of Malaysian Homes <i>Ramasamy Chidambaram</i>	283
Transforming Fragile Perfects into Dynamic Perfects to tide over the 2020 Pandemic Crises <i>Shivali Shamsheer, Thiagarajan Praba, KR Sethuraman</i>	286
SHORT COMMUNICATION	
Inconsistency of Lesion Quantitative Assessment in 2D SUV and 3D SUV Quantification Techniques for [18F]-FDG PET/CT: A Phantom Study <i>Muhammad Hafiz Hanafi, Noramaliza Mohd Noor, Muhammad Hishar Hassan</i>	289
CASE SERIES	
Case Series on Drug Reaction with Eosinophilia and Systemic Syndrome (DRESS) with Different Ocular Manifestation <i>Josephine Lee En Hui, Kursiah Mohd Razali, Adil Hussein</i>	294
CASE REPORT	
A Novel Approach in Treating Diabetic Foot Ulcer With Hyperbaric Oxygen Therapy: A Case Report <i>Mohd Syazwan Zainal, Khairul Azmi Ghazali, Masdiamin Mohamad Nor, Navin Kumar Devaraj, Rampal S</i>	298
Return to Work Assessment in a Radiographer Assistant Suffering a Hemianopsia due to an Ischemic Stroke <i>Iqbal Mochtar, Nuri Purwito Adi, Astrid Widayati Hardjono, Muchtaruddin Mansyur</i>	301
An Experience of Embolization of Multiple Recurrence Pheochromocytoma Nodules <i>Chong Eng Ngen, Zuria Amaleen binti Zulkifli, Ridzuan Abdul Rahim, Ezamin Abdul Rahim</i>	305
Successful Pregnancy in a Patient With Recurrent Pregnancy Loss Due to Afibrinogenemia Managed With Cryoprecipitate Prophylaxis in a Resource-limited Setting <i>Athira Sasidharan, Aboobacker Mohamed Rafi, Ramesh Bhaskaran, Nithya M Baiju, Susheela Jacob Innah</i>	308
Severe Pulmonary Tuberculosis With Organizing Pneumonia: A Diagnostic Ambiguity <i>Nurashikin Mohammad, Mohd Jazman Che Rahim, Wan Aireene Wan Ahmed, Wan Syamimee Wan Ghazali</i>	311
Case of Acute ST-elevation Myocardial Infarction (STEMI) Secondary to Anaphylactic Reaction: Type 2 Kounis Syndrome <i>Toh Chong Rui, Huzairi Sani, Nada Syazana Zulkufli</i>	314
Exceptionally Rare Krukenberg Tumour in a Patient With Dextrocardia and Situs Inversus Totalis <i>Vickneswaren Thever Ramasamy, Yong Chee Meng</i>	317

ORIGINAL ARTICLE

Analysis of Heavy Metals (Cadmium, Chromium, Lead, Manganese, and Zinc) in Well Water in East Java Province, Indonesia

Mochammad Sholehhudin¹, R. Azizah¹, Arif Sumantri² Shahrudin Mohd Sham^{1,3}, Zainul Amiruddin Zakaria^{1,4}, Mohd Talib Latif^{1,5}

¹ Department of Environmental Health, Faculty of Public Health, Universitas Airlangga, 60115 Mulyorejo, Surabaya, Indonesia

² Public Health Study Program, Faculty of Health Science, State Islamic University (UIN) Syarif Hidayatullah Jakarta, Indonesia

³ Department of Environmental & Occupational Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁴ Department of Biomedical Science, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁵ Department of Earth Sciences and Environment, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

ABSTRACT

Introduction: Indonesia is one of the countries whose inhabitants use well water for drinking and cooking purposes. In East Java, 48.90% of the population uses well water for their daily needs. Well water contained heavy metals had bad effects on health such as cancer, damage of liver, kidneys, and others. The objective of this study was to evaluate the concentration of heavy metals in well water and relate them to a potential health outcomes. **Methods:** The method used in this study was analytical descriptive. Data used was secondary from East Java Environmental Office. A total of 101 samples were collected from 33 locations. There were 5 heavy metals analyzed, namely cadmium, chromium, lead, manganese, and zinc. Equipment using ICPMS and AAS. Data analyzed with descriptive statistics by SPSS. Data obtained were compared to the WHO Standard for Drinking Water Quality. **Results:** Concentration for cadmium was 0.002 mg/l, followed by manganese at 1.80 mg/l and zinc at 0.020 mg/l. Besides, all water samples had levels of chromium and lead below the detection limit. **Conclusion:** All heavy metals had concentrations below the maximum allowable standard, except for five water samples from three locations with levels of manganese which was above the maximum standard. Long term effects of manganese include neurological problems, intelligence, and low birth weight. Further studies need to be done to determine the source of manganese contamination. It is recommended that bottled water is used for drinking purposes in an area where heavy metal concentration is above the allowable limit.

Keywords: Heavy metal, Well water, Manganese, Drinking water, Health risk

Corresponding Author:

R. Azizah, PhD

Email: azizah@fkm.unair.ac.id

Tel: +6285851885999

INTRODUCTION

Water is important to human life that is always needed every day. Water consumed by humans is needed by all organs of the body so that health maintained. The availability of water on earth is tremendous, but only a small portion can be used for drinking. Experts mentioned that the surface of the earth is covered by water by 71% but only 2.5% is freshwater that can be consumed by humans (1). Freshwater can be obtained from underground water such as wells. Many countries use well water as their main source of drinking water, in study Indonesia. As many as 27.04% of households in Indonesia use well water for drinking and cooking

purposes. That number is the second-highest after bottled drinking water sources, at 31.30% of households (2). One province that uses well water as a source of drinking water is East Java. As many as 48.90% of households in East Java use well water, which is the highest compared to other sources such as bottled water 29% and 8.70% pipe water (3).

Well water quality consumed by the community for their daily needs is the most important. Currently, the quality of well water is threatened by a large number of water pollutants and it mainly occurs in urban areas and areas that have become centers of intensive agriculture (1). Contamination that occurs in well water may be caused by natural processes and the consequences of human activities. Some contaminants that have become important to comply with guidelines of the United States Environmental Protection Agency (USEPA) are microorganisms (bacteria, viruses, and parasites), nitrates

and nitrites through chemical fertilizers, human sewage, and animal waste; heavy metals like copper, chromium, arsenic, lead, antimony, cadmium, selenium, organic chemical; radionuclides; and fluoride (4,5).

Heavy metals are a group of metal elements that have a density of more than 5 gr/cm³ and a high relative density. Heavy metals have toxic properties down to ppb levels. Examples are Cd, Zn, Fe, Pb, Mg, As, Pt, and Cu. The sources of these metals from natural and human or anthropogenic activities such as mining, transportation, and industrial activities. Heavy metals cannot be lost naturally because they continue to accumulate so that they can cause harm to human health if exposed continuously for a long time. Some research has been conducted to examine the quality of well water in several regions in Indonesia. The results show that the concentration of manganese (Mn) in some areas has exceeded the maximum drinking water standard. For example well water in Jakarta indicated that Mn levels in Jakarta residents have exceeded the maximum standard of 0.970 - 1.022 mg/l (6). Also, the research in Sukoharjo shows water is at 1.43 mg/l, exceeded drinking water standards (7). Iron (Fe) concentration, also showed exceeded standards such as in areas around the Kaliyasa river, Cilacap City where the level was 2.3 mg/l (8). Similarly, well water in Sukoharjo has a high Fe level of 1.45 mg/l (7). In addition to these two elements, lead (Pb) level also recorded high reading from the maximum standard (0.01 mg/l), such as in Palembang, where it was known that Pb in well water around landfills as 0.03 – 0.05 mg/l (9). And research in Pekalongan also showed that Pb levels in well water was high 0.04 mg/l (10). Cadmium (Cd) levels were also reported to be above the maximum standard (0.003 mg/l). In a Yogyakarta study, cadmium levels in well water reached 0.0178 mg/l (11). Similarly, a research report in Jember, East Java, showed the level of Cd in well water around landfills were 80% higher than the maximum standard (12). This shows a little picture of the condition of well water pollution in Indonesia. Besides, other studies discussed the quality of well water in East Java, especially heavy metals contamination.

Contaminated well water may cause a variety of public health problems. Many studies have reported that pollutants entering the human body through drinking water will cause negative health impacts on the consumers. Some health problems that may arise include microorganism infections, reduction of the blood's ability to carry oxygen, heavy metal risks such as acute and chronic toxicity on some organs including the liver, kidneys, and intestines, and medical conditions like anemia, cancer, and hormonal disorders (4,13). A Studied in northeast Iran on the content of heavy metals in drinking water showed that Cr levels exceed the safe level of US EPA risk. This condition is stated from the results of the Hazard Index that will be carcinogenic in children and adults through the consumption of water

and skin pathways (14).

As many people rely on well water for their daily needs and the challenges of health problems that may arise due to contamination, a study is needed that will illustrate well water quality in East Java. This province is the third-largest population in Indonesia. Various kinds of population activities such as very dense transportation, many large-scale industries to accommodate workers, home industries in each district area, as well as household activities in densely populated housing carryout negative impacts on the environment including the occurrence of well water pollution. This situation is very appropriate to be the basis for evaluating heavy metals in well water because a study in Port Klang, Malaysia also states that heavy metal pollution in water and sediment will increase along with the intention industrialization and urbanization (15). This is also an effort to improve the 6th sustainable development goals indicator, which is about safe drinking water. A managed safe drinking water services are defined as one definite location, always available for daily needs, and safe from contamination (16). The aims of this study, therefore, were to identify and analyze the quality of well water and the health impacts caused by heavy metal contaminants in East Java Province, Indonesia.

MATERIALS AND METHODS

Study Design, Location, and Time

This was an analytical descriptive study using secondary data. The data was sourced from the Information on Environmental Management Performance Report in 2016 - 2017 by the Environmental Office, East Java Province, Indonesia.

Population dan Sample

Water samples were obtained from 101 groundwater sources at 33 district locations and were collected from February 2016 - August 2017. The sample selection in these areas was based on community reports to the environmental office in each region. The community provided information on the differences in water quality based on their observations by the physical quality of the water. Therefore it needs proper monitoring with the appropriate equipment to ensure the quality of this water. Well water sampling technique refers to the guidelines from the Ministry of Health of The Republic of Indonesia. To ensure that quality of sampling and inspection, was carried out by a research team who had received training and a certificate from the environmental office training center agency.

Variable Data

Variables in this study were 5 heavy metals, namely Lead (Pb), Cadmium (Cd), Manganese (Mn), Chromium (Cr), and Zinc (Zn) in well water.

Processing, Analysis, and Presentation of Data

The level of each heavy metals were determined at the Laboratory of Technical Center of Environmental Health and Disease Control in Surabaya. The equipment used were Inductively Coupled Plasma Mass Spectrometry (ICP-MS) and Atomic Absorption Spectrophotometer (AAS). ICP-MS was an analytical technique that can be used to measure elements at trace levels in biological fluids. The advantages of this technique were multi-element technique, large analytical range, low detection limit, high sample throughput, low sample volume, simple sample preparation, high-resolution and tandem mass spectrometry (triple-quadrupole) instruments offer a very high level of interference control.

Then data was analyzed with descriptive statistics by SPSS. While water quality assessment was compared to The Drinking Water Quality Standard by the World Health Organization (2017). According to the regulation, there were the limit standards of heavy metals, 0.003 mg/l for cadmium, 0.05 mg/l for chromium, 0.01 mg/l for lead, 0.4 mg/l for manganese, and 3 mg/l for zinc (17). A review of potential health effects was also discussed by literature research. Recommendations with regards to health impacts were presented at the conclusion.

RESULTS

The results of heavy metals measurement in this study were presented in Figure 1. It showed that only manganese (Mn) had 5 readings which exceeded than standard (0.4 mg/l) detected in samples from location 55 (1.8 mg/l) in Malang District, location 87 (1.78 mg/l), 100 (0.68 mg/l) and 101 (0.43 mg/l) in Surabaya City, and location 89 in Mojokerto District (1.20 mg/l). While the other metals had a low level in all of the locations.




Figure 1: The measurement result of heavy metal samples (cadmium, chromium, lead, manganese, and zinc)

Table I showed a comparison between measurement results with the maximum standard for heavy metals in drinking water. The levels of cadmium (Cd), chromium (Cr), lead (Pb), and zinc (Zn) were below the allowable standard, even the level of chromium and lead were below limit detection in all samples. While the level of manganese (Mn) exceeded the WHO standard for drinking water.

Table I: The results of descriptive analyze

Desc. Analyze	Cd (mg/l)	Cr (mg/l)	Pb (mg/l)	Mn (mg/l)	Zn (mg/l)
Standard	0,003	0,05	0,01	0,4	3
Limit Detection	0,001	0,003	0,004	0,05	0,008
Minimun	< LD	< LD	< LD	< LD	< LD
Maxium	0,002	< LD	< LD	1,800	0,020
Mean	0,000	0,000	0,000	0,578	0,016
St. Deviation	0,000	0,000	0,000	0,652	0,004

< LD = Less then limit detection

According to figure 2, all water samples did not exceed the drinking water quality standard for Cd (0.003 mg/l) but there are 2 locations (location 55 and 56 in Malang) with the highest level of Cd ie 0.0024 mg/l. While Mn levels (showed in figure 4) in 95% of the samples were under the detection limit of 0.05 mg/l, but 5 samples were above the standard (0.4 mg/l) and the points of locations were mentioned above (figure 3). Zn level in figure 4 showed that all levels below the standard in drinking water (3 mg/l), but there were 2 locations with the highest level of Zn ie 0.020 mg/l (location 26




Figure 2: The result of cadmium measurement in 101 samples location




Figure 3: The result of manganese measurement in 101 samples location




Figure 4: The result of zinc measurement in 101 samples location

in Bangkalan and location 28 in Sampang). Very good results were showed by Cr and Pb which had all samples (100%) in figure 1 had levels under the detection limit. It was below the measurement ability of the equipment.

This figure (Fig. 5) was the result of mapping from the measurement of heavy metals level in samples area (East Java Province). This area was one of the 34 provinces in Indonesia country. This map showed 28 district locations with the blue color had below standard for all heavy metals, 2 district locations with the green color had the highest level of Zn, 1 district location with the yellow color had the highest level of Cd, and 3 district locations with the red color had Mn level above the standard. While the black color was not sampled locations.




Figure 5: The map of the result of sample locations and the level of heavy metals

DISCUSSION

Cadmium (Cd)

Cd sources in natural waters are more commonly found in basic sediments and suspended particles. Besides, that Cd will be affected by the degree of acidity of the water. While, the anthropogenic sources of Cd are from agriculture, landfill, sewage sludge, and a lot of activities in urban areas such as high emissions, industry, and mining. However, identification can be difficult because one pollution route can be from multiple sources simultaneously leading to groundwater sources (18). The results of the data in this study showed similarities with research on assessment of well water quality in the city of Kakamega, Kenya which stated that the levels of cadmium were lower than the WHO standard (19). However, this level is different from research in East Delhi, India. It is stated in the study that the sample of groundwater had a level up to 0.28 mg/l, it was above the safe level standard (20). The research was conducted in the Krishna Vigar industrial area. The presence of cadmium in groundwater or well water is dangerous if it is consumed by humans. As happened in Namo Bintang, Deli Serdang, North Sumatera Province, Indonesia, it was stated that 65 respondents (65.7%) were exposed by drinking water which contains

cadmium with a concentration level above the safe standard. Apart from the research also showed the urine test 99% of the sample showed a Cd level exceeded the normal limit, so the recommendation of researchers was residents do not use well water as the main source of drinking water (21). Cd that has levels exceeding the WHO standard will potentially damage human health. In general, organs that are targeted by Cd are kidneys and bones. Canada Health Authority explained that “Cadmium exposure is well known to result in damage to the nephron’s proximal tubule, causing impaired reabsorption of low molecular weight proteins and enzymes by the kidneys” (19 p. 21). Besides, “cadmium exposure has long been associated with reduced bone mineral density, osteoporosis, and fractures” (19 p. 23). Another long-term effect is on the onset of cancers based on animal testing, making cadmium a carcinogen to humans (23). Various ways can be taken to reduce the level of Cd in water, one of which is the Bioreduction Adsorbent with the bacteria *Bacillus* sp. and durian leather. The biosorbent is carried out in several steps, starting with the manufacture of activated charcoal from durian leather and continuously by administering the isolated bacteria with medium Cd. Both of them can be done sequentially to get the optimum results (24).

Chromium (Cr)

This study is supported by research conducted at Agbor and Owa Community of Nigeria, which stated that chromium levels in all groundwater samples were below the detection limit (25). In contrast to chromium detection in the Unnao district of India, Cr levels were higher than the safe limit of 0.141 mg/l in the water hand pump samples at Gupta Gate, Shivnagar (26). The high level of chromium at this location is probably due to the study location being close to the Ganga River, whereas the river holds nearly 70% of the untreated waste from the 50 leather industries. In addition, research in Bangladesh also has different results from this study. It is stated that the Cr pollutant in groundwater located close to the Meghna Ghat industrial area has a Cr level that exceeds the safe standard because it reaches 0.07 mg/l (27). Reviewing from these studies, the value of Cr is always high in areas that are located close to industrial areas. This indicates that the potential for pollution occurs due to the lack of monitoring of waste management from the factory. Chromium in the environment is usually divided into 2 groups namely Cr3+ and Cr6+ but for drinking water quality, total chromium. In general, the concentration of chromium in groundwater is low (< 1 µg/liter). Because of that, it takes a long time to find out the effects of Cr exposure on the human body. One way is through an examination of Cr levels in blood and hair. As mentioned in previous studies, personals exposure dose of chromium for drinking water during lifetime showed associations with levels of chromium in the blood and hair, furthermore in other parameters such as hematological and biochemical also (28). Another way is urine examination by the previous study, which showed

that the method is effective to determine the chromium level in the body. It's mentioned that 53% of workers sampled detected chromium levels are exceeded the normal standard (29).

Lead (Pb)

This research had similar data with the study conducted in Surabaya. It was stated that the Pb contamination in groundwater was below the detection limit (30). The research was conducted in the coastal area of Surabaya City. Compared to research in Algeria where Pb levels (0.072 up to 0.458 mg/l) had exceeded the WHO standard (0.01 mg/l). It has a potential health risk especially impact for the population who consumed the water (31). It similar to a study in East of Algeria also said that the concentration of the lead is above the maximum permissible standard, the level of sample ranges from 0.017 to 0.292 mg/l (32). In addition, the case study in Surulere also showed that 36.73% of well water samples had a Pb concentration level above the maximum contaminant level (33). Children and infants are two groups who are highly sensitive towards health problems caused by Pb. Estimated from a 5 µg/l Pb concentration in drinking water, the Pb intake for infants is assumed to be around 3.8 µg/day, whereas for adults up to 10 µg/day (34). Health problems caused by Pb are very diverse, including neurological effects in children and adults, and also renal, cardiovascular, hematological, immunological, and reproductive effects (35). But, this study had no worries about health risk potential to the population because the level of Pb was safe.

Manganese (Mn)

This study showed similarities with results of a study on heavy metal concentrations in well water in Tamil Nadu, India, that mentioned manganese (1.276 mg/l) in the water exceeded the maximum standard (36). One study which analyzed well water quality in the Sidoarjo mudflow area, stated that Mn level exceeded the WHO standard as the mud is the source of Mn contamination of groundwater (37). The other research in Western Amazonia, Peru also had similarities with this study. Which is the contamination of Mn in groundwater was very high from maximum standard, which was 4 mg/l (38). However, it is different from research on groundwater at Kilvelur Taluk, Nagapattinan District, Tamil Nadu, India which showed that the Mn levels detected in 3 phases (pre-monsoon, monsoon, post-monsoon seasons) from 20 locations had a safe condition of concentration levels (0.02 – 0.26 mg/l) below WHO maximum limit (39).

The high concentration of Mn in the three areas mentioned can be caused by several sources, the first of which is the natural content of groundwater. This statement is supported by the theory that Mn in water can come from natural sources in bedrock, specifically, water from deep wells (40) and the United States National Water Quality Assessment Program showed

that the levels of manganese in groundwater is about 99% is commonly higher than in surface waters. The natural presence of Mn that enters groundwater and well water can be influenced by several factors such as TDS, fluctuation of groundwater level, and time of residence of water in the ground (41). Other sources of Mn that can pollute well water are agricultural practices and leachate disposal. This may be the reason behind the high Mn level in sample 55 where the well is located near a landfill area. This is evidenced by research that discussed the content of Mn in well water and leachate in Banyuurip landfill, Magelang, Indonesia. It stated that Mn is the main substance in leachate that originates from metal wastes that have accumulated at the landfill. This researcher also explained that "there is a very strong and significant correlation between landfill-to-well distance and total Mn concentration" (32 p. 4). Besides, contamination may occur by industrial activities as evident through samples obtained in Surabaya City and Mojokerto District which are centers of industries. WHO, in Concise International Chemical Assessment Document 63 about manganese, mentioned that Mn pollution which originates from industry has been happening for a long time. Mn can enter water bodies through industrial waste disposal facilities. Since 1983, it has been estimated that Mn that enter the waters from anthropogenic sources worldwide amounted to 109,000 to 414,000 tons. Domestic wastewater and sewage sludge are the most important sources. Reports from the USA stated that Mn does not only pollute surface water, but is also able to penetrate groundwater up to 0.114 tons (43). Although smaller than surface water pollution (up to 17.2 tons), this fact is a real threat to health in the future. Therefore, it can be said that these 3 sources are eligible to be potential causes of Mn pollution.

One health disorder that is often linked to Mn contamination is neurologically-based, such as abnormal gait, ataxia, muscle hypotonicity, and a face without permanent emotions. Besides, liver dysfunction has also been reported (44). Initially, research conducted in Canada involving 259 children between 2012 - 2014 stated that there was nothing to show about the relationship between manganese and the cognitive development of school children, thus allowing for gender-based associations (45). Later, in 2018, another study on Mn in drinking water was conducted in Canada involving 630 children (aged 5.9 - 13.7 years) which discussed drinking water with manganese content and were associated with decreased performance IQ scores up to 5 %. It showed that Mn level of 78 µg/L (0.078 mg/l) contributed to a decrease of 1% Performance IQ, followed by a decrease of 2% for levels of 156 µg/L (0.156 mg/l), and a decrease of 5% for levels of 406 µg/L (0.406 mg/l). Therefore, it is very important to re-checking the maximum concentration of Mn in drinking water, with the aims are to protect and prevent problems in children's health (46). Potential health problem is also shown for adults, such as risks to

pregnant women who consume water with Mn. Cohort studies were conducted in three cities in South Korea, with 331 mother-infant respondents from July 2007 to December 2009. The results of the study mentioned that any relationship of Mn levels in the blood (high and low) with low birth weight of infants (less than 3000 g). It is, therefore, clear that high Mn (more than 36 µg/l) in the blood has an association with the presence of low birth weight of infants (47). Some of these studies showed that health problems caused by Mn actually occur. Because of that, it is not recommended for people especially children and pregnant women to consume water with high level of Mn during their lifetime.

Zinc (Zn)

This concurs with results from research in Nigeria which stated that Zn concentration of 0.02 mg/l did not exceed the maximum standard by WHO (25). In contrast with a study in Ghana showed that Zn levels in shallow groundwater were exceeded the recommended value of WHO regulation in 2008 (48). Zn is naturally present in groundwater and it can be influenced by the acidity of the water. According to the acidity theory, the higher the acidity of the water, the higher Zn concentration will become. Some types of Zn found in the soil can seep into groundwater so that it is possible if waste pollution occurs at a location, it will affect Zn concentration in the water (49). In general, it is stated that the natural presence of Zn is indeed smaller than other heavy metals. One study that discussed the assessment of heavy metals mobility states that the mobility and bioavailability of Zn were the lowest when compared to Mn and Cu (Mn > Cu > Zn) (50). Even though, Zn is one of the minerals needed by the body for cell and tissue growth. While, recommendation of WHO showed if consuming too much, zinc will cause damages to organs such as prostate, bone, muscle, liver, and gastrointestinal system. The recommended zinc consumption limits based on Recommended Dietary Allowances (RDAs) is 11 mg/day for male and 8 mg/l for female (49).

CONCLUSION

Nearly all the heavy metals studied such as Chromium, Cadmium, Zinc, and Lead had levels below the maximum allowable standard by the World Health Organization, except for manganese (Mn) where 5 samples had a concentration which exceeded the standard ie 0.43 mg/l until 1.80 mg/l. The source of Mn in the water can occur by earth's layer content, agriculture practice, leachate disposal, industrial activities, and domestic sewage. Therefore, further studies need to be done to determine the source of Mn contamination at the three locations which have high concentration of Mn. Besides, to determine the long term effects of Mn on the general population. More importantly, to vulnerable populations such as women pregnant and children. Based on previous research Mn that enters the body will cause important problems for the body,

including neurological health problems, a decrease in children's intelligence, and affect the low weight of baby birth. Therefore the research team suggests that the government also needs to frequently control the level of Mn in well water. Then examined in-depth the causes and consequences together with researchers from the university. Collaboration between the government and academia is also needed in an effort to improve the quality of well water, more specifically to reduce the level of concentration of heavy metal, manganese. The hope is to realize a technology that is easily applied in the community as a solution to the high levels of manganese in well water. The authors, therefore, would strongly suggest the use of bottled water for drinking purposes.

ACKNOWLEDGMENTS

This research was supported by the Department of Environmental and Occupational Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia and Department of Environmental Health, Faculty of Public Health, Universitas Airlangga, Surabaya.

REFERENCES

1. World Health Organization. Water Services for Health. 2019.
2. Statistics Center Office. Percentage of Households by Province and Drinking Water Source 2000-2016. Indonesia; 2017.
3. Environmental Office. Environmental Management Performance Information Document East Java Province 2017. Surabaya; 2018.
4. Environmental Protection Agency USA. Potential Well Water Contaminants and Their Impacts. 2019.
5. Al-bahry S, Mahmoud IY. Microbial and chemical pollution of water-wells relative to sewage effluents in Oman. 2014;(January).
6. Lestari; Thoriq. The Quality of The Population's Well Water In The East Jakarta Pulogadung Village. J Petro. 2017;VI(2):59-65.
7. Kurniawan TP. Chemical Quality of Well Water in Perum Pondok Baru Permai Bulak Rejo Village Sukoharjo District Sukoharjo Regency, Year 2015. 2017;9(1):26-30.
8. Sasongko EB, Widyastuti E, Priyono RE. Water Quality Assessment and Use Of Well Duck by Communities Around Kaliyasa River, Cilacap District. J Ilmu Lingkung. 2014;12(2):72-82.
9. Maksuk, Suzanna. Study of Lead Containing in Dug Well Water at Sukawinatan Dumping Site in Palembang City. J Ilmu Kesehat Masy. 2018;9(2):107-14.
10. Budiyanto S, Purnaweni H, Sunoko HR. Environmental Analysis of The Impacts of Batik Waste Water Pollution on The Quality of Dug Well Water in The Batik Industrial Center of Jenggot

- Pekalongan City. Proc ICENIS 2017. 2018;9008.
11. Putrinta I, Dwantari S, Indonesia UI, Wiyantoko B, Indonesia UI. Analysis of Total Hardness, Lead Metal (Pb), and Cadmium (Cd) in Well Water Using the Complexometric Titration and Atomic Absorption Spectrophotometry Method. *Indones J Chem Anal.* 2019;2(1):11–9.
 12. Qadriyah L, Moelyaningrum AD, Ningrum PT. The Cadmium Levels in Well Water Around Landfill (Study in Landfill, Jember Distric, Indonesia). *J Biol Lingkungan, Ind dan Kesehat.* 2019;6(1):41–9.
 13. Adeogun AO, Ibor OR, Adeduntan SD, Arukwe A. Science of the Total Environment Intersex and alterations in reproductive development of a cichlid, *Tilapia guineensis*, from a municipal domestic water supply lake (Eleyele) in Southwestern Nigeria. *Sci Total Environ.* 2016;541:372–82.
 14. Alidadi H, Tavakoly Sany SB, Zarif Garaati Oftadeh B, Mohamad T, Shamszade H, Fakhari M. Health risk assessments of arsenic and toxic heavy metal exposure in drinking water in northeast Iran. *Environ Health Prev Med.* 2019;24(1):1–17.
 15. Sany SBT, Salleh A, Sulaiman AH, Sasekumar A, Rezayi M, Tehrani GM. Heavy metal contamination in water and sediment of the Port Klang coastal area, Selangor, Malaysia. *Environ Earth Sci.* 2013;69(6):2013–25.
 16. SDG-Tracker.org. Sustainable Development Goal 6. 2018.
 17. World Health Organization. Guidelines for Drinking-water Quality. 4th ed. Geneva: WHO Press; 2017.
 18. Kubier A, Wilkin RT, Pichler T. Cadmium in soils and groundwater: A review. *Appl Geochemistry.* 2019;108(August).
 19. Christine AA, Kibet JK, Kiprop AK, Were ML. The assessment of bore-hole water quality of Kakamega County, Kenya. *Appl Water Sci.* 2018;8(1):1–8.
 20. Panwar RM, Ahmed S. Assessment of contamination of soil and groundwater due to e-waste handling. *Curr Sci.* 2018;114(1):166–73.
 21. Ashar YK, .Z, Wulandari RA, Susana D. Well-Water Consumption of High Cadmium and the Resulting Urinary Cadmium Levels in a Community near a Dumping Site. *KnE Life Sci.* 2018;4(1):129.
 22. Canada Health Office. Cadmium in Drinking Water. Ontario: Canada Health; 2019.
 23. IARC. IARC Monographs on The Evaluation of Carcinogenic Risks to Humans. France: International Agency for Research on Cancer; 2012.
 24. Sueb, Diartika EIA, Sripalupi K, Irmawati A. Bioreduction Adsorbent (Biosorbent): Recovery Technology Of Heavy Metal Pollution (Cadmium / Cd) In Polluted Lapindo Water Sources Using Bacteria And Durian Leather. *Proceeding 1st IBSC Towar Ext Use Basic Sci Enhancing Heal Environ Energy Biotechnol.* 2016;7–9.
 25. Oyem HH, Oyem IM, Usese AI. Iron, Manganese, Cadmium, Chromium, Zinc and Arsenic Groundwater Contents of Agbor and Owa communities of Nigeria. *Springer Plus.* 2015;4(104).
 26. Mani Tripathi S, Chaurasia SR. Detection of Chromium in surface and groundwater and its bio-absorption using bio-wastes and vermiculite. *Eng Sci Technol an Int J.* 2020;23(5):1153–61.
 27. Rahman MATMT, Paul M, Bhoumik N, Hassan M, Alam MK, Aktar Z. Heavy metal pollution assessment in the groundwater of the Meghna Ghat industrial area, Bangladesh, by using water pollution indices approach. *Appl Water Sci.* 2020;10(8):1–15.
 28. Sazakli E, Villanueva CM, Kogevinas M, Maltezi K. Chromium in Drinking Water : Association with Biomarkers of Exposure and Effect. *Int J Environ Res Public Health.* 2014;11:10125–45.
 29. Ningrum NF., Azizah R. KM. Correlation Between Characteristic with Heavy Metal Chromium (Cr) Levels in Urine of Metal Coating Workers in Sidoarjo, East Java, Indonesia. *Psychosoc J.* 2020;24(4):1158–73.
 30. Rochaddi B, Atmodjo W, Satriadi A, Suryono CA, Irwani I, Widada S. The Heavy Metal Contamination in Shallow Groundwater at Coastal Areas of Surabaya East Java Indonesia. *J Kelaut Trop.* 2019;22(1):69.
 31. Balli N, Leghouchi E. Assessment of Lead and Cadmium in Groundwater Sources Used for Drinking Purposes in Jijel (Northeastern Algeria). *Glob NEST J.* 2018;20(2):417–23.
 32. Belkhir L, Tiri A, Mouni L. Assessment of Heavy Metals Contamination in Groundwater: A Case Study of the South of Setif Area, East Algeria. *IntechOpen.* 2018;17–31.
 33. Momodu MA, Anyakora CA. Heavy metal contamination of ground water: the Surulere case study. *Res J Environ Earth Sci.* 2010;2(1):39–43.
 34. World Health Organization. Lead in Drinking-water. Geneva: WHO Press; 2011.
 35. ATSDR. Toxicological Profile for Lead. Georgia: Agency for Toxic Substances and Disease Registry; 2019.
 36. Sridhar SGD, Sakthivel AM, Sangunathan U, Balasubramanian M, Jenefer S, Rafik MM, et al. Heavy Metal Concentration in Groundwater from Besant Nagar to Sathankuppam, South Chennai, Tamil Nadu, India. *Appl Water Sci.* 2017;7(8):4651–62.
 37. Sholehuddin M, Susanto BH, Adriyani R, Laksono D. Heavy Metals Contamination in Groundwater Around Sidoarjo Mud Vulcano Area, East Java Indonesia. *Malaysian J Med Heal Sci.* 2019;15(Supplement 3):86–9.
 38. de Meyer CMC, Rodriguez JM, Carpio EA, Garcia PA, Stengel C, Berg M. Arsenic, manganese and aluminum contamination in groundwater resources of Western Amazonia (Peru). *Sci Total*

- Environ. 2017;607–608:1437–50.
39. Punitha S, Selvarajan G. Analysis of Heavy Metals Concentration in Ground Water From Kilvelur Taluk, Nagapattinam District, Tamil Nadu, India. *J Chem Chem Sci.* 2018;8(3):538–47.
 40. Wendel B Van, Jooode D, Barbeau B, Bouchard MF, Марна А, Quesada R, et al. Manganese Concentrations In Drinking Water from Villages Near Banana Plantations With Aerial Mancozeb Spraying in Costa Rica : Results from the Infants Environmental Health Study (ISA). *Environ Pollut.* 2016;215:247–57.
 41. Zhang Z, Xiao C, Adeyeye O, Yang W, Liang X. Source and mobilization mechanism of iron, manganese and arsenic in groundwater of Shuangliao City, Northeast China. *Water (Switzerland).* 2020;12(2).
 42. Budihardjo MA, Nugraheni AS. Analysis of Groundwater Quality Surrounding Municipal Solid Waste Landfill : Banyuurip Landfill , Magelang , Indonesia. *IJCAET ISAMPE 2017.* 2018;1031.
 43. World Health Organization. Concise International Chemical Assessment Document 63 Manganese and Its Compounds : Environmental Aspects. Geneva: WHO Press; 2004.
 44. ATSDR. Toxicological Profile for Manganese. Georgia: Agency for Toxic Substances and Disease Registry; 2012.
 45. Bouchard MF, Surette C, Cormier P, Foucher D. Neurotoxicology Low Level Exposure to Manganese From Drinking Water And Cognition in School-Age Children. *Neurotoxicology.* 2018;64:110–7.
 46. Kullar SS, Shao K, Surette C, Foucher D, Mergler D, Cormier P, et al. A Benchmark Concentration Analysis for Manganese in Drinking Water And IQ Deficits in Children. *Environ Int.* 2019;130(February):104889.
 47. Eum J, Cheong H, Ha E, Ha M, Kim Y, Hong Y, et al. Maternal Blood Manganese Level and Birth Weight : A MOCEH Birth Cohort Study. *Environ Heal.* 2014;13(31):1–7.
 48. Anim-Gyampo M, Anornu GK, Appiah-Adjei EK, Agodzo SK. Quality and health risk assessment of shallow groundwater aquifers within the Atankwidi basin of Ghana. *Groundw Sustain Dev.* 2019;9(December 2018):100217.
 49. ATSDR. Public Health Statement Zinc. Georgia: Agency for Toxic Substances and Disease Registry; 2005.
 50. Borah P, Gujre N, Rene ER, Rangan L, Paul RK, Karak T, et al. Assessment of mobility and environmental risks associated with copper, manganese and zinc in soils of a dumping site around a Ramsar site. *Chemosphere.* 2020;254:126852.