

Risk Factors for Neonatal Asphyxia Occurrence at General Hospital Dr. M. Soewandhie, Surabaya

by Domunicus Husada

Submission date: 21-Aug-2020 05:00PM (UTC+0800)

Submission ID: 1372173695

File name: LK_-_TA_Sekarwati_-_7757-Article_Text-14260-1-10-20200709.pdf (427.41K)

Word count: 3471

Character count: 17870

Risk Factors for Neonatal Asphyxia Occurrence at General Hospital Dr. M. Soewandhie, Surabaya

Tri Adharia Sekarwati¹, Ninik Darsini², Dominicus Husada³

¹Midwifery Programme, ²Biomedical Department, ³Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya

Abstract

Background: Neonatal mortality rates in Indonesia are still high at 15 per 1000 live births. Surabaya is one of the regions with the highest number of neonatal deaths in East Java. Neonatal asphyxia is the highest cause of neonatal death after preterm birth. Some factors that cause neonatal asphyxia include maternal factors, labor and fetal factors. This study aims to identify the risk factors that influence the incidence of neonatal asphyxia.

Method: This medical-based study using a case-control approach used neonatal medical record data that was born in the period of January 1, 2018, to December 31, 2018, in the General Hospital of dr. M. Soewandhie, Surabaya. Samples were selected based on inclusion and exclusion criteria. Sampling was done by total sampling in the case group (93 neonates) and simple random sampling in the control group (93 neonates) and matching was done based on sex and birth month. Data analysis was performed by univariate, chi-square and fisher's exact test and multivariate with multiple logistic regression.

Results: Neonates with diagnosed neonatal asphyxia were born to mothers who had low education (63.4%) and did not work (68.8%). Risk factors that significantly increased the incidence of neonatal asphyxia were: non-spontaneous labor with OR=5.56 (95%CI: 2.50-12.34), preeclampsia with OR=2.52 (95%CI: 1.15-5.54), meconium-stained amniotic fluid with OR=2.51 (95%CI: 1.17-5.38) and primiparous parity with OR=2.15 (95%CI: 1.06-4.39).

Conclusion: Non-spontaneous labor, preeclampsia, meconium-stained amniotic fluid and primiparous parity affect the incidence of neonatal asphyxia.

Keywords: Neonatal asphyxia, non-spontaneous labor, preeclampsia, meconium-stained amniotic fluid, primiparous.

Introduction

The number of child deaths under five years is as many as 5.4 million children with an under-five mortality

rate of 39 per 1,000 live births, as much as 47% of these deaths occur in neonatal, which is 18 per 1,000 live births^[1]. The neonatal mortality rate in Indonesia is 15 per 1000 live births. This figure is still quite far from the Sustainable Development Goals (SDGs) target, which is to reduce neonatal mortality to a minimum of 12 per 1,000 live births^[2]. Neonatal mortality rates in East Java experienced an insignificant decrease from 2015 to 2017^[3] with neonatal asphyxia as the highest cause of neonatal death after preterm birth^[4].

Neonatal asphyxia is a result of intrapartum hypoxia/ischemia of the fetus and resuscitation measures which

Corresponding Author:

Dominicus Husada

Faculty of Medicine, Universitas Airlangga Surabaya,
Jawa Timur, Indonesia

Phone: +62818337734

e-mail: dominicushusada@yahoo.com

can immediately restore the condition of the newborn^[5]. This condition results in malfunctioning of vital organs, brain damage to death^[6]. Causes of neonatal asphyxia include hypertension in pregnancy, post-term pregnancy, maternal narcotics during labor, uterine contractions (hypertonic or tetania uterine), disorders of the umbilical cord and hypovolemic shock^[7]. Risk factors for neonatal asphyxia include maternal age, maternal education, parity, anemia during pregnancy, prolonged labor, premature rupture of membranes, low birth weight^[8] and very low birth weight in neonates^[9]. Preeclampsia in pregnancy can increase the risk of neonatal asphyxia^[10]. Obesity in pregnant women also increases the incidence of severe asphyxia in neonates^[11].

This study aims to identify risk factors for the incidence of neonatal asphyxia so that it is expected to be prevented and reduced.

Material and Method

This was medical-based research with a case-control approach. The population in this study were neonates born at the General Hospital of dr. M. Soewandhie, Surabaya in the period January 1, to December 31, 2018, obtained from medical records. The inclusion criteria for case samples were neonates born with neonatal asphyxia diagnoses and had complete medical record data and could be read by the author. The inclusion criteria for the control sample were neonates who were born without neonatal asphyxia and had complete medical record data and could be read by the author. The exclusion criteria in this study were neonates born with congenital abnormalities such as congenital heart defects and diaphragmatic hernias. The sampling technique was total sampling in the case group and simple random sampling in the control group and matching was done based on sex and birth month. Samples obtained were 186 neonates divided into 93 cases and 93 controls. The variables in the study consisted of the dependent variables namely neonatal asphyxia and independent variables namely parity, maternal age, prematurity, birth weight, obesity, anemia during pregnancy, preeclampsia, type of labor, premature rupture of membranes (PROM), prolonged labor and *meconium*-stained *amniotic* fluid (MSAF). Data were analyzed by univariate, Chi-Square and Fisher's Exact Test and multivariate (Backward Stepwise: Likelihood Ratio) using a statistical program (SPSS version 16).

General Hospital of dr. M. Soewandhie, Surabaya

was one of the largest secondary hospitals in Surabaya. This hospital had 356 patient beds with as many patients (inpatient = 26,590, outpatient = 275,628) per year. There were 104 medical personnel and 527 paramedics.

Results

The percentage of neonatal asphyxia births was 4.4% (142/3,267). The total sample of cases was 142, as many as 93 samples stated included inclusion criteria. Case and control samples were matched based on sex and birth month of newborns. Table 1 showed that the sex in each case and control group were 52 samples (55.9%) for male and 41 samples (44.1%) for female.

Table 1: Comparability by sex in the case and control groups

Sex of newborns	Neonates	
	Cases n = 93 (%)	Controls n = 93 (%)
Male	52 (55,9%)	52 (55,9%)
Female	41 (44,1%)	41 (44,1%)

Table 2 showed that as many as 59 (63.4%) case samples and 55 (59.1%) control samples were born to mothers with low education. A total of 64 (68.8%) case samples and 57 (61.3%) control samples were born to mothers who did not work.

Table 2: Characteristics of mothers based on education and occupation in cases and controls

Characteristics	Neonates	
	Cases n = 93 (%)	Controls n = 93 (%)
Education		
Low (primary school/middle school)	59 (63,4%)	55 (59,1%)
High (high school/diploma/ bachelor)	34 (36,6%)	38 (40,9%)
Occupation		
Not working (housewife)	64 (68,8%)	57 (61,3%)
Working	29 (31,2%)	36 (38,7%)

Table 3 showed the results on each of the independent variables that are significantly related to neonatal asphyxia were parity, obesity, preeclampsia, type of labor, prolonged labor and MSAF. The highest crude odds ratio was found in non-spontaneous labor, which was 3.71 (95% CI: 1.81-7.64). As for the variables of maternal age, prematurity, birth weight, anemia and PROM were not related (p-value > α) with the incidence of neonatal asphyxia.

Table 3 the relationship of risk factors for the incidence of neonatal asphyxia

Variables	Neonates		Crude OR	95% CI	P-value*
	Cases n = 93 (%)	Controls n = 93 (%)			
Parity					
Primiparous	40 (43,0%)	26 (28,0%)	1,95	1,06-3,58	0,032
Multiparous	53 (57,0%)	67 (72,0%)			
Maternal Age					
<20 and>35	20 (21,5%)	25 (26,9%)	0,75	0,38-1,46	0,392
20-35	73 (78,5%)	68 (73,1%)			
Prematurity (weeks)					
Premature (<37)	16 (17,2%)	10 (10,8%)	1,73	0,74-4,03	0,205
Not premature (≥37)	77 (82,8%)	83 (89,2%)			
Birth weight (gram)					
<2500	14 (15,1%)	9 (9,7%)	1,65	0,68-4,04	0,265
≥2500	79 (84,9%)	84 (90,3%)			
Obesity					
Yes	14 (15,1%)	5 (5,4%)	3,12	1,08-9,05	0,029
No	79 (84,9%)	88 (94,6%)			
Anemia					
Yes	6 (6,5%)	1 (1,1%)	6,35	0,75-53,78	0,118
No	88 (93,5%)	92 (98,9%)			
Preeclampsia					
Yes	28 (30,1%)	15 (16,1%)	2,24	1,10-4,55	0,024
No	65 (69,9%)	78 (83,9%)			
Type of labor					
Non-spontaneous (CS/VE/Manual Aid)	35 (37,6%)	13 (14,0%)	3,71	1,81-7,64	< 0,001
Spontaneous	58 (62,4%)	80 (86,0%)			
PROM					
Yes	11 (11,8%)	7 (7,5%)	1,65	0,61-4,46	0,321
No	82 (88,2%)	86 (92,5%)			
Prolonged labor					
Yes	9 (9,7%)	0 (0%)	2,12	1,81-2,46	0,003
No	84 (90,3%)	93 (100%)			
MSAF					
Yes	30 (32,3%)	16 (17,2%)	2,29	1,15-4,58	0,017
No	63 (67,7%)	77 (82,8%)			

*Chi Square/Fisher's Exact Test

Table 4 showed the results of multivariate analysis with the Backward Stepwise method in order to obtain the final model of multiple logistic regression. Variables with p values <0.25 in the Chi-Square/Fisher's Exact Test were included in the multivariate analysis, which included eight variables. Parity, preeclampsia, type of labor and MSAF were risk factors for neonatal asphyxia.

The analysis showed that the type of labor with non-spontaneous (CS/VE/manual aid) had a risk of 5.6 times to deliver babies with neonatal asphyxia (OR = 5.56; 95% CI: 2.50-12.34) compared to spontaneous labor. The risk of neonatal asphyxia in women with preeclampsia was 2.5 times higher than in women without preeclampsia (OR = 2.52; 95% CI: 1.15-5.54).

Similar to preeclampsia, MSAF also had a 2.5-fold risk of increasing the incidence of neonatal asphyxia (OR = 2.51; 95% CI: 1.17-5.38). Compared to multiparous

mothers, mothers with primiparous parity had a 2.1 risk of giving birth to infants with neonatal asphyxia (OR = 2.15; 95% CI: 1.06-4.39).

Table 4: Results of multivariate regression

Variables	B	P-value	Odds Ratio	95.0% C.I. for OR	
				Lower	Upper
Parity (1)	0,767	0,035	2,152	1,055	4,392
Preeclampsia (1)	0,924	0,022	2,519	1,146	5,540
Type of labor (1)	1,715	< 0,001	5,555	2,500	12,343
Prolonged labor (1)	20,970	0,999	1,280E9	< 0,001	.
MSAF (1)	0,919	0,018	2,507	1,170	5,376
Constant	-1,223	< 0,001	0,294		

Discussion

This study aimed to identify risk factors for neonatal asphyxia, which was by including several determinants that might be risk factors for neonatal asphyxia. Therefore, neonatal asphyxia could be prevented and inherited.

This study showed that neonatal asphyxia mostly occurred in male neonates. This study was in accordance with a study in the United States, which states that 56.2% of neonates with neonatal asphyxia were male. There was no clear theory yet, but this might be related to male vulnerability (XY chromosome) to cerebral anoxia compared to women because there was an additional X chromosome^[12].

Maternal characteristics showed that the majority of neonatal asphyxia incidents were born to mothers with low education and no work. This result was in line with a study conducted in Jakarta^[13], neonates with neonatal asphyxia were born to mothers with low education (90.6%) and from mothers who did not work by 77.8%. In Northern Ethiopia, the majority of neonatal asphyxia were born to mothers with low education (84.1%) and from mothers who did not work (housewives) by 53.4%^[6].

Parity had a significant relationship with the incidence of neonatal asphyxia. Mothers with primiparous parity had a 2.2 times higher risk of giving birth to babies with neonatal asphyxia (OR = 2.15; 95% CI: 1.06-4.39) compared to multiparous mothers. The results of this study differed from studies conducted

in Ethiopia, which showed that parity did not affect the incidence of asphyxia^[14]. However, another study showed that primiparous mothers had 3.1 times higher risk of giving birth to babies with neonatal asphyxia. This might be related to primiparous mothers who often did not know their pregnancy needs and even ignored antenatal care visits^[6]. In the first pregnancy, cervical muscles tended to be stiff so that labor took a long time, which could increase the risk of neonatal asphyxia^[15].

Mothers with preeclampsia during pregnancy had 2.5 times the risk of giving birth to neonatal asphyxia babies (OR = 2.52; 95% CI: 1.15-5.54) compared to women who did not experience preeclampsia during pregnancy. This was in line with a study conducted in Ethiopia^[6],^[14], which showed that preeclampsia increases the risk of the incidence of neonatal asphyxia by 4.1 times and 2.6 times, respectively. The result might be related to changes in the cardiovascular system due to an increase in blood pressure and vascular spasm^[10], resulting in placental insufficiency and fetal hypoxia^[16].

The type of delivery had a significant relationship with the incidence of neonatal asphyxia. Labor with non-spontaneous (CS/VE/manual aid) had a 5.6 times higher risk of delivering a baby with neonatal asphyxia (OR = 5.56; 95% CI: 2.50-12.34) compared to spontaneous labor. These results were not consistent with a study conducted in Bali^[8] but were consistent with studies in Ethiopia which stated that labor with cesarean section had a 7-fold risk of increasing the risk of neonatal asphyxia^[14]. This was related to impaired perfusion and vascular resistance due to anesthesia during the

cesarean section and this action also resulted in no chest compressions in infants such as in vaginal delivery^[17]. In addition, the used of medical devices such as forceps and or a vacuum extractor during labor could result in trauma and intracranial hemorrhage, which hindered the baby's circulation^[16].

Meconium-stained amniotic fluid at delivery have a 2.5 times higher risk for neonatal asphyxia in infants (OR = 2.51; 95% CI: 1.17-5.38) compared with clear membranes. These results were consistent with studies in Ethiopia^[14], which showed that MSAF had a risk of 8.6 times to occurred neonatal asphyxia. The presence of MSAF at delivery indicated that fetal distress had occurred^[18]. This results in meconium aspiration syndrome, which resulted in the obstruction of the baby's respiratory tract^[15].

The limitation of this study was that it used secondary data, namely medical record data, so that much of the information needed was not available. In addition, this study was only conducted in one hospital in the city of Surabaya, so the study could not be generalized to a wider population.

Conclusion

Non-spontaneous labor, preeclampsia, primiparous and MSAF are risk factors for neonatal asphyxia. Midwives as a companion for pregnant women and maternity mothers are expected to improve the quality of education and early detection of risk factors. In addition, midwives are expected to be able to carry out close monitoring during labor and identify any complications during labor. Having the right diagnosis and treatment can improve the quality of fetal output.

It is expected that further authors will conduct a more specific study on the incidence of neonatal asphyxia with the factors that influence it. In addition, it is hoped that the community can increase awareness in recognizing high risks in pregnancy and childbirth so that it can reduce morbidity and mortality rates for mothers and infants.

Conflict of Interest: This study was conducted without conflict of interest from person or organization or commercial interests such as financial relationship, personal relationship, academic competition or intellectual passion.

Source of Funding: Self-funding.

Ethical Clearance: This study has obtained an ethical clearance certificate from the Health Research Ethics Committee, Faculty of Medicine, Airlangga University, Surabaya No. 96/EC/KEPK/FKUA/2019, obtained on April 8, 2019 and from the Medical Research Ethics Committee of General Hospital of Dr. M Soewandhie, Surabaya No. 006/KE/KEPK/2019, obtained on April 24, 2019.

Acknowledgments: The authors would like to thank General Hospital of dr. M. Soewandhie, Surabaya for making this study possible by allowing us to access patient medical record data.

And the kind guidance from Mrs. Atika as a lecturer in study methodology and Mrs. Ilya Krisnana as a lecturer examiner that enables this study to be carried out well.

References

1. United Nations Inter-Agency Group for Child Mortality (UN IGME). Level and trends in child mortality: report 2018 estimated development by the UN Inter-Agency Group for Child Mortality. New York: United Nation Children's Fund (UNICEF);2018.
2. Kementerian Kesehatan Republik Indonesia. Profil kesehatan Indonesia tahun 2017 (Indonesia health profile in 2017). Jakarta: Kementerian Kesehatan Republik Indonesia;2018.
3. Dinas Kesehatan Provinsi Jawa Timur. Profil kesehatan Provinsi Jawa Timur Tahun 2017 (East Java Province Health Profile 2017). Surabaya: Dinas Kesehatan Provinsi Jawa Timur; 2018
4. World Health Organization. Newborn: reducing mortality [internet]; 2018[cited 2019 July 12]. Available from: <https://www.who.int/en/news-room/fact-sheets/detail/newborns-reducing-mortality>
5. LissauerT, FanaroffAA, Miall L,Fanaroff J. Neonatology at a glance. 3rd ed. UK: John Wiley and Sons Ltd; 2016. p. 26-30.
6. TasewH, Zemicheal M, TeklayG, Mariye T, Ayele E. Risk factors of birth asphyxia among newborns in public hospitals of Central Zone, Tigray, Ethiopia 2018. BMC Res Notes [internet]. 2018 [cited 2019 Feb 5];11(496):[about 7 pp.]. Available from: <https://doi.org/10.1186/s13104-018-3611-3>
7. DwiendaO, Maita L, Saputri EM, Yulviana R.

- 22 Asuhan kebidanan neonatus, bayi/balita dan anak prasekolah untuk para bidan (Neonatal midwifery care, infants/toddlers and preschoolers for midwives). Yogyakarta: Deepublish;2014. p. 15-17.
8. Widiani NNA, Kurniati DPY, Windiani IGAT. Faktor risiko ibu dan bayi terhadap kejadian asfiksia neonatorum di Bali : penelitian case control (Mother and infant risk factors for the incidence of neonatal asphyxia in Bali: a case-control study). Pub Health and Prev Med Archive. 2016 Des; 4(2):120–126.
 9. De AlmeidaMFB, MoreiraLMO, Vaz Dos SantosRM, Kawakami MD, AnchietaLM, Guinsburg R. Early neonatal deaths with perinatal asphyxia in very low birth weight Brazilian infants. J of Perinatology [internet]. 2015 Sep [cited 2018 Des 2]; 35(11):954–957. Available from: <https://doi.org/10.1038/jp.2015.114>
 10. Indah SN, Apriliana E. Hubungan antara preeklamsia dalam kehamilan dengan kejadian asfiksia pada bayi baru lahir (The relationship between preeclampsia in pregnancy with the incidence of asphyxia in newborns). Majority [internet]. 2016Des [cited 2018 Des 2];5(5):55–60. Available from: <http://juka.kedokteran.unila.ac.id/index.php/majority/article/view/924>
 11. Persson M, Johansson S, CnattingiusS. Interpregnancy weight change and risks of severe birth-asphyxia-related outcomes in Singleton infants born at term: a Nationwide Swedish cohort study. PLoS Medicine [internet]. 2016 Jun [cited 2019 Jan 25];13(6):[about 15 pp.]. Available from: <https://doi.org/10.1371/journal.pmed.1002033>
 12. Mohamed MA, Aly H. Impact of race on male predisposition to birth asphyxia. J of Perinatology [internet]. 2014 Feb[cited 2018 Des 7]; 34(6):449-452. Available from: <https://doi.org/10.1038/jp.2014.27>.
 13. Opitasari C. Maternal education, prematurity and the risk of birth asphyxia in selected hospitals in Jakarta. Health Sci J of Indonesia [internet]. 2015 Des [cited 2019 Jan 25]; 6(2):111-115. Available from: <https://doi.org/10.22435/hsji.v6i2.4769.111-115>.
 14. Gebreheat G, Abraha D, Teame H, Tsegay T, Welu G, Etsay N, et al. Prevalence and associated factors of perinatal asphyxia among neonates in general hospitals of Tigray, Ethiopia, 2018. Bio Med Research International [internet].2018Nov [cited 2019 Feb 20]; [about 7 pp.]. Available from: <https://doi.org/10.1155/2018/5351010>
 15. Aisyiah, LubisR, Kurnia S. Hubungan paritas, anemia dan ketuban pecah dini dengan kejadian asfiksia neonatorum pada bayi baru lahir di RSUD Cilegon Provinsi Banten (Relationship between parity, anemia and premature rupture of membranes with the incidence of neonatal asphyxia in newborns in Cilegon District Hospital, Banten Province). PHI Health Journal. 2016 Aug; 2(7):1-9.
 16. Gomella T, Cunningham M, Eyal F. Common multiple congenital anomaly syndromes. neonatology: management, procedures, on call problems, disease and drugs.7th ed. New York: Lange Medical Books/McGraw-Hill; 2013. p. 530.
 17. FannyF. Sectio caesarea sebagai faktor risiko kejadian asfiksia neonatorum (the cesarean section as the risk factors of neonatal asphyxia).Majority [internet]. 2015 Nov [cited 2019 Feb 20]; 4(8):57–62. Available from:<http://juka.kedokteran.unila.ac.id/index.php/majority/article/view/1474>.
 18. Rahmatullah, Kurniawan. 9 bulan dibuat penuh cinta dibuai penuh harap menjalani kehamilan & persalinan (9 months made full of love cradled hopefully through pregnancy & childbirth). Jakarta: Gramedia; 2019. p. 126-128.

Risk Factors for Neonatal Asphyxia Occurrence at General Hospital Dr. M. Soewandhie, Surabaya

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

10%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	bmcpublichealth.biomedcentral.com Internet Source	1%
2	journals.lww.com Internet Source	1%
3	eprints.undip.ac.id Internet Source	1%
4	Submitted to University of Utah Student Paper	1%
5	protocolexchange.researchsquare.com Internet Source	1%
6	ejurnal.undana.ac.id Internet Source	1%
7	bmcpediatr.biomedcentral.com Internet Source	1%
8	Submitted to Universitas Brawijaya Student Paper	1%
9	jurnal.stikesphi.ac.id	

Internet Source

1%

10

Estin Gita Maringga, Nunik Ike Yuniasari.
"Hubungan Usia Gestasi terhadap Kejadian
Asfiksia Neonatorum di RSUD Kabupaten Kediri
Periode Januari s/d Juni 2017", Jurnal
Kebidanan Midwiferia, 2017

Publication

1%

11

www.signavitae.com

Internet Source

1%

12

repositorio.unb.br

Internet Source

<1%

13

www.frontiersin.org

Internet Source

<1%

14

sinta3.ristekdikti.go.id

Internet Source

<1%

15

Submitted to University of Newcastle

Student Paper

<1%

16

ejournal.helvetia.ac.id

Internet Source

<1%

17

proquest.com

Internet Source

<1%

18

Submitted to International Medical University

Student Paper

<1%

19	www.medscape.com Internet Source	<1%
20	demikarina.blogspot.com Internet Source	<1%
21	Submitted to Nazarbayev University Student Paper	<1%
22	www.providefreebookedition.com Internet Source	<1%
23	www.ajol.info Internet Source	<1%
24	sites.kowsarpub.com Internet Source	<1%
25	cityofdekalb.com Internet Source	<1%
26	Persson, Martina, Stefan Johansson, Eduardo Villamor, and Sven Cnattingius. "Maternal Overweight and Obesity and Risks of Severe Birth-Asphyxia-Related Complications in Term Infants: A Population-Based Cohort Study in Sweden", PLoS Medicine, 2014. Publication	<1%
27	ejournal.litbang.kemkes.go.id Internet Source	<1%
28	liebertpub.com Internet Source	<1%

29 Martina Persson, Stefan Johansson, Sven Cnattingius. "Inter-pregnancy Weight Change and Risks of Severe Birth-Asphyxia-Related Outcomes in Singleton Infants Born at Term: A Nationwide Swedish Cohort Study", PLOS Medicine, 2016

Publication

<1%

30 www.pacificjournals.com

Internet Source

<1%

31 "Nutrition and Diet in Maternal Diabetes", Springer Science and Business Media LLC, 2018

Publication

<1%

32 file.scirp.org

Internet Source

<1%

33 Maharani Sulistiyo Putri, Ira Titisari, Arika Indah Setyarini. "Hubungan Usia Kehamilan Dengan Komplikasi Pada Bayi Baru Lahir Di Rs Aura Syifa Kabupaten Kediri Tahun 2017", Jurnal Ilmu Kesehatan, 2017

Publication

<1%

34 Abdulai Abubakari, Francis Zinenuba Taabia, Zakari Ali. "Maternal determinants of low birth weight and neonatal asphyxia in the Upper West region of Ghana", Midwifery, 2019

Publication

<1%

35

Endah Purwanti, Ichroom Septa Preswari, Ernawati Ernawati. "Early Risk Detection of Pre-eclampsia for Pregnant Women Using Artificial Neural Network", International Journal of Online and Biomedical Engineering (iJOE), 2019

Publication

<1%

36

Biselele, T., G. Naulaers, P. Bunga Muntu, E. Nkidiaka, M. Kapepela, L. Mavinga, and B. Tady. "A Descriptive Study of Perinatal Asphyxia at the University Hospital of Kinshasa (Democratic Republic Of Congo)", Journal of Tropical Pediatrics, 2013.

Publication

<1%

37

Gdiom Gebreheat, Tesfay Tsegay, Dessalegn Kiros, Hirut Teame, Natnael Etsay, Guesh Welu, Desta Abraha. "Prevalence and Associated Factors of Perinatal Asphyxia among Neonates in General Hospitals of Tigray, Ethiopia, 2018", BioMed Research International, 2018

Publication

<1%

38

Junrong Wang, Enfu Tao, Miaojun Mo, Weimin Ding, Junhui Yuan, Min Wang, Changhua Zheng, Haixiao Zheng. "Perinatal Risk Factors Influencing Neonatal Hypoxic Ischemic Encephalopathy in Southern China: A Case–Control Study", American Journal of Perinatology, 2020

Publication

<1%

39

Yunarsih Yunarsih. Jurnal Ilmu Kesehatan, 2019

Publication

<1%

40

Cholifah Cholifah, Navyati Asrita Putri.
"FAKTOR-FAKTOR YANG BERPENGARUH
TERHADAP PENCAPAIAN K4 DI DESA
SUMBEREJO WONOAYU SIDOARJO",
Midwiferia, 2016

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Risk Factors for Neonatal Asphyxia Occurrence at General Hospital Dr. M. Soewandhie, Surabaya

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
