

Validity of Modified Whisper Test as Hearing Screening Method

by Rizki Najoa

Submission date: 06-Apr-2022 06:00AM (UTC+0800)

Submission ID: 1802796804

File name: alidity_of_Modified_Whisper_Test_as_Hearing_Screening_Method.pdf (425.3K)

Word count: 2286

Character count: 11816

Validity of Modified Whisper Test as Hearing Screening Method in Presbycusis Patients

Rizki Najohan, Nyilo Purnami

Department of Otorhinolaryngology-Head and Neck Surgery, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Hospital, Surabaya, Indonesia

Abstract

Background: Presbycusis is a hearing loss due to the degeneration process which is found at the age of more than 65 years old, with decreased hearing sensitivity in both ears, asymmetrical bilateral sensorineural detection type. **Objective:** To identify the validity of modified whisper test by detecting the hearing level in presbycusis patients. **Methods:** The design of this study was comparative, cross-sectional, prospective study. Subjects were elderly undergoing hearing monitoring at URJ Geriatric and URJ Audiology Dr. Soetomo Hospital, in August–September 2018 and examined using whisper test and audiometry. The statistical analysis used 2×2 tables, sensitivity, specificity, positive predictive value, and negative predictive value. **Results:** The lowest sensitivity at frequencies >41–55 dB was at 100%, the lowest specificity at 100%, lowest sensitivity at frequencies >56–70 dB was at 23.07% with specificity at 100%. The positive predictive value (NRP) at frequencies >25–40 dB was 88.46% while the negative predictive value (NPV) was of 100%. **Conclusion:** The modified whisper test can be used as early detection of hearing impairment.

Keywords: Audiogram, elderly, presbycusis, whisper test

INTRODUCTION

Presbycusis is a hearing loss due to the degeneration process, which is characterized by decreased hearing sensitivity in both ears resulting in an increase of the hearing threshold with the symmetrical bilateral type of sensorineural hearing loss.^[1,2] A striking change in the cochlea is the atrophy and degeneration of the supporting hair cells in the organ of Corti.^[3,4] This degeneration process causes high-frequency sensorineural hearing loss with bilateral and symmetrical nature.^[4,5] The degree of hearing loss is determined based on the average value of the hearing threshold or pure tone averages (PTA).^[6,7] Due to the limitations of audiometry, a simpler modification of the whisper test is performed.^[8,9]

This study intends to prove the validity of the modified whisper test by detecting the degree of hearing loss based on the hearing threshold value in presbycusis patients

METHODS

This study was cross-sectional comparative study with a prospective design. The subjects of this study were elderly

patients with complaints of hearing loss who came to the Geriatric Outpatient Unit (URJ) and Audiology URJ Dr. Soetomo Hospital, Surabaya, Indonesia. The study was performed between August and September 2018 with inclusion criteria were patients aged 65 years or older with complaints of symmetrical and bilateral sensorineural hearing loss, never used hearing aids, and no history of surgery and ear infections, while exclusion criteria were inability to concentrate or remember the words spoken due to dementia, disturbance of consciousness, or disturbance of balance. The number of samples obtained was 45 elderly or 90 ears. Ethical approval for this study (Ethical Committee number 0482/KEPK/VIII/2018) was provided by the Ethical Committee of Dr. Soetomo General Hospital, Surabaya, on 11 August 2018.

Address for correspondence: Dr. Nyilo Purnami, Department of Otorhinolaryngology-Head and Neck Surgery, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Hospital, Jalan Mayjen Prof. Dr. Moestopo 47, Surabaya 60286, Indonesia. E-mail: nyilo@fk.unair.ac.id

Submitted: 15-Oct-2019 Accepted: 27-May-2020 Published: 26-Oct-2021

Access this article online

Quick Response Code:



Website:
www.indianjotol.org

DOI:
10.4103/indianjotol.INDIANJOTOL_114_19

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Najohan R, Purnami N. Validity of modified whisper test as hearing screening method in presbycusis patients. *Indian J Otol* 2021;27:11-3.

The examination was performed as follows : the examiner sat in front of the patient, and the examiner’s left and right index fingers were pressed on to the tragus of the unexamined ear of the patient and the ear was covered. The fingers were moved so as to produce sound as a masking. The examiner whispered 10 words then the patient has to repeat the words with light and loud sound. If the patient correctly repeated 80% of the whispered words, the ear was regarded as having pass outcome, while correct repetition of <80% was declared not to have pass outcome.

The hearing threshold value (PTA) is the average value of the airborne hearing threshold at frequencies of 500, 1000, 2000, and 4000 Hz. The hearing threshold is the lowest pure tone that can be heard by someone at a certain frequency. The degree of hearing loss based on hearing threshold values in this study refers to International Standard Organization (ISO) criteria or the ISO: normal degree ≤25 dB, mild degree >26–40 dB, moderate degree >41–55 dB, moderately severe degree >56–70 dB, severe degree >71–90 dB, and profound degree >90 dB.^[10]

Statistical analysis in this study examined the first performance or the ability of the modified voice whisper test to predict

hearing loss using receiver operating characteristic (ROC) curve with an output in the form of area under the curve (AUC). [Table 3]. We also measured the sensitivity and specificity of the modified whispered test. The subsequent analysis used a 2 × 2 table, with the output of sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) [Table 4].^[11]

RESULTS

In this study, the results of modified whisper test was compared to the results of hearing threshold measurement using audiometry as the as the gold standard and basic examination.

From a total of 90 ears of elderly patients who performed the modified whisper test, 65 ears (72.22%) had pass outcome and 25 ears (27.77%) had refer outcome [Table 1].

Audiometry results were calculated on an average of all frequencies (250 Hz–4000 Hz); there were 6 normal ears (6.66%), mild impairment in 55 (61.11%), moderate in 7 ears (7.77%), moderately severe in 13 (14.44%), severe in 4 ears (4.44%), and profound in as many as 5 ears (5.55%) [Table 2].

The ROC curve which shows that the modified whisper test had a good diagnostic value because the curve was far from the 50% line and close to 100%.

The AUC value obtained from the ROC method was 94.2% (95% confidence interval [CI], 91.65%–98.72%). Statistically, the AUC value of 94.2% is relatively strong.

Table 1: Result of the modified whisper test

Modified whisper test	n (%)
Pass	65 (72.2)
Refer	25 (27.7)
Total	90 (100)

Table 2: Distribution of hearing loss degree

Hearing loss degree	n (%)
Normal	6 (6.66)
Mild	55 (61.11)
Moderate	7 (7.77)
Moderately severe	13 (14.44)
Severe	4 (4.44)
Profound	5 (5.55)
Total	90 (100)

Table 3: Area under the curve value in the modified whispered test of hearing loss

Obs	ROC area	SE	Asymptotic interval	95% confidence
90	0.9420	0.0161	0.91653	0.98726

SE: Standard error, ROC: Receiver operating characteristic




Figure 1: The receiver operating characteristic curve in the modified whispered test hearing loss

Table 4: Sensitivity and specificity of the modified whispered test

Modified whispered test	Audiogram result (%)				
	>25-40 dB (mild)	>41-55 dB (moderate)	>56-70 dB (moderately severe)	>71-90 dB (severe)	>90 dB (profound)
Sensitivity	83	100	23.07	25	40
Specificity	100	100	100	100	100
PPV	88.46	53.84	33.33	14.28	25
NPV	100	0	100	100	100

PPV: Positive predicitive value, NPV: Negative predicitive value

AUC value of 94.2% means that a modified whisper test score [Figure 1] is reliable for use as a diagnostic tool. The AUC percentage of 94.2% means that if a modified whisper test is applied to determine the presence or absence of hearing impairment in 90 normal ears, the accurate outcome may be obtained from 90 ears of the patients. In this study, the resulted confidence interval of the modified whisper test in the population ranged between 91.65% to 98.72%.

Audiogram examination results at all frequencies showed that the highest sensitivity was at the frequencies of >41–55 dB (100%) with specificity at 100%, while the lowest sensitivity 23.07% was at frequencies of >56–70 dB with specificity at 100%. The positive predictive value (PPV) was at the frequencies >25–40 dB of 88.46% while the negative predictive value (NPV) was 100%.

DISCUSSION

The results of audiometry examination at all frequencies were as many as 6 normal ears (6.66%) while those with hearing impaired were 84 ears (93.33%). These results are in accordance with a study by Lee *et al.* who reported that there is a correlation between age and sex on hearing loss in the elderly. The average hearing threshold increases by 1 dB every year at the age of 60 years and over and there is a significant decrease in the hearing threshold at frequencies of 4 and 8 kHz between men and women.^[12] Another study in South Carolina, USA, found that presbycusis was found mostly in those aged 60–70 years.^[13] In contrast, a study in Qatar found that the highest prevalence of presbycusis in the middle age group of 50–59 years.^[14]

In this study, modification of whispered voice test in 45 elderly revealed that as many as 65 ears (72.22%) had pass outcome and the remaining 25 ears (27.77%) did not pass. The whisper voice test is a simple method; it does not require sophisticated equipment or provide manipulation to the patient, so this procedure is safe and is not dangerous to elderly.

The results of the audiogram as the gold standard for hearing loss are calculated from the average hearing threshold of all frequencies. The 3-meter whisper test study conducted by Purnami, obtained quite reliable results. The sensitivity of the examination was 80% and the specificity of 95.2%.^[9] These results are consistent with Pirozzo's research, which found that the whisper test is more specific in adults (sensitivity 80%–96%, specificity 90%–98%).^[15] The results of the examination at the age of the elderly are considered adequately effective enough to diagnose hearing loss by considering the practical implementation procedures, its simplicity, and it can

be performed anywhere so that it meets the required screening method criteria.

CONCLUSION

Modified whispered test can be used as an early detection of hearing loss.

Acknowledgment

The researchers would like to thank all participants who took part in this study.

Financial support and sponsorship

This study was supported by the authors.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Busis SN. Presbycusis. In: Calhoun KH, Eibling DE, editors. *Geriatric Otolaryngology*. New York: Taylor and Francis; 2006. p. 77-90.
2. Fatmawati R, Dewi Y. Karakteristik penderita presbikusis di bagian ilmu kesehatan THT-KL RSUP Dr. Hasan Sadikin Bandung periode januari 2012-Desember 2014. *JSK* 2016;2:201-5.
3. Roland PS, Kutz Jr JW, Issacson B. Aging and the auditory and vestibular system. In Bailey BJ, Johnson JT, Newlands SP, editors. *Head and Neck Surgery-Otolaryngology*. 5th ed., Vol. 2. Philadelphia: Lippincott Williams and Wilkins; 2014. p. 2615-23.
4. Binstock R. *Hearing Impairment and Elderly People*. Washington: BC Decker; 2008. p. 35-42.
5. Pumami N. THT komunitas. In: Dalam Prajitno S, Djuari L, Husein H, editors. *Buku Ajar Kedokteran Komunitas*. Surabaya: Airlangga University Press; 2013. p. 263-72.
6. Gates GA, Murphy M, Rees TS, Fraher A. Screening for handicapping hearing loss in the elderly. *J Fam Pract* 2003;52:56-62.
7. Servidoni AB, Conterno LO. Hearing loss in the elderly: Is the hearing handicap inventory for the elderly – Screening version effective in diagnosis when compared to the audiometric test? *Int Arch Otorhinolaryngol* 2017;12:89-94.
8. Wiyadi MS. Tes Pendengaran suara bisik. *Media Perhati* 1997;3:180-4.
9. Pumami N. The modified whispered test for screening of hearing impairment in children at the elementary school. *J Physics* 2018;Conf. Ser.1075012066.
10. Soetirto I, Hendarmin H, Bashiruddin J. Hearing Loss (Deafness). In: Soepardi EA, Iskandar N, Editor. *Science Textbook Ear Nose Throat Head Neck Health*. Edition 6. Jakarta: Publishing house FK UI : 2011, p. 10-22.
11. Akobeng AK. Understanding diagnostic tests 1: Sensitivity, specificity and predictive values. *Acta Paediatr* 2007;96:338-41.
12. Lee FS, Matthews LJ, Dubno JR, Mills JH. Longitudinal study of pure-tone thresholds in older persons. *Ear Hear* 2005;26:1-11.
13. Gates GA, Mills JH. Presbycusis. *Lancet* 2005;366:1111-20.
14. Bener A, Salahudin A, Darwish S, Al-Hamaq A, Gansan L. Association between hearing loss and type 2 diabetes mellitus in elderly people in a newly developed society. *Biomed Res* 2008;19:187-95.
15. Pirozzo S, Papinczak T, Glasziou P. Whispered voice test for screening for hearing impairment in adults and children: Systematic review. *BMJ* 2003;327:967.

Validity of Modified Whisper Test as Hearing Screening Method

ORIGINALITY REPORT

16%

SIMILARITY INDEX

7%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	kjorl.org Internet Source	2%
2	"Phoniatics I", Springer Science and Business Media LLC, 2020 Publication	2%
3	mafiadoc.com Internet Source	1%
4	Yuyun Yueniwati, Evelyn Christina. "The challenges in differentiating tuberculous from pyogenic spondylitis using magnetic resonance imaging", Reports in Medical Imaging, 2017 Publication	1%
5	ijop.net Internet Source	1%
6	link.springer.com Internet Source	1%
7	Waleed B. Alshuaib, Jasem M. Al-Kandari, Sonia M. Hasan. "Chapter 2 Classification of Hearing Loss", IntechOpen, 2015	1%

8

Obi L Griffith, Connie G Chiu, Allen M Gown, Steven JM Jones, Sam M Wiseman.

"Biomarker panel diagnosis of thyroid cancer: a critical review", Expert Review of Anticancer Therapy, 2014

Publication

1 %

9

Purnami Nyilo, Manyakori Serafika Permoni Putri. "The Association of Reactive Oxygen Species Levels on Noise Induced Hearing Loss of High Risk Workers in Dr. Soetomo General Hospital Surabaya, Indonesia", Indian Journal of Otolaryngology and Head & Neck Surgery, 2018

Publication

1 %

10

cyberleninka.org

Internet Source

1 %

11

Anthony K Akobeng. "Understanding diagnostic tests 3: receiver operating characteristic curves", Acta Paediatrica, 5/2007

Publication

1 %

12

El Kady, Heba Mahmoud. "Prevalence of hearing impairment and its correlates among a group of hospitalized chronically ill elderly patients in Alexandria, Egypt :", Journal of the Egyptian Public Health Association, 2012.

Publication

1 %

13 Jui-Kun Chiang, Yen-Chang Lin, Hsiao-Chen Yu, Chih-Ming Lu, Yee-Hsin Kao. "Long-term benefits of a new oral appliance on adult snoring: a trend analysis", Multidisciplinary Respiratory Medicine
Internet Source 1 %

14 S. Pirozzo. "Whispered voice test for screening for hearing impairment in adults and children: systematic review", BMJ, 2003
Publication <1 %

15 Haihong Liu, Hua Zhang, Ruth A. Bentler, Lingyan Mo, Demin Han, Luo Zhang. "Audiometric Records Analysis in a Clinical Population in China", ORL, 2011
Publication <1 %

16 N Purnami. "The modified whispered test for screening of hearing impairment in children at the elementary school", Journal of Physics: Conference Series, 2018
Publication <1 %

17 Sennaroglu, Gonca, and Levent Sennaroglu. "Hearing Loss in Inner Ear Malformations", Encyclopedia of Otolaryngology Head and Neck Surgery, 2013.
Publication <1 %

18 scielosp.org
Internet Source <1 %

19

Elida Mustikaningtyas, Sri Herawati Juniati, A. C. Romdhoni. "Intracell Heat Shock Protein 70 Expression and Nasopharyngeal Carcinoma Stage", Indian Journal of Otolaryngology and Head & Neck Surgery, 2018

Publication

<1 %

20

Kelvin Yuwanda, I Putu Eka Widyadharma, Dewa Putu Gde Purwa Samatra, I Made Oka Adnyana, Anna Marita Gelgel, I Komang Arimbawa. "High leukotriene B4 serum levels increase risk of painful diabetic neuropathy among type 2 diabetes mellitus patients", The Egyptian Journal of Neurology, Psychiatry and Neurosurgery, 2021

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Validity of Modified Whisper Test as Hearing Screening Method

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
