

Tropical Animal Science Journal

Artificial Insemination in Sheep with Fresh Diluted Semen: Comparison Between Two Different Semen Extenders and Management Protocols

A. Madrigali, A. Rota, D. Panzani, S. Castellani, M. Shawahina, A. Hassan, F. Di Iacovo, C. Rossignoli, & F. Camillo

The Protective Impacts of α -tocopherol Supplementation on the Semen Quality of Sapera Goat Preserved at 4°C

R. A. Prastiya, Rimayanti, M. M. Munir, & A. P. Nugroho

Detection of Tetracycline Resistance Genes among *Escherichia coli* Isolated from Layer and Broiler Breeders in West Java, Indonesia

A. Indrawati, K. Khoirani, S. Setyaningsih, U. Affif, Safika, & S. G. Ningrum

Growth Performance of Male Line of Superior Tropical Rabbit

B. Brahmantiyo, N. Pratiwi, F. Saputra, Y. C. Raharjo, & L. H. Prasetyo

Genetic Diversity of Rabbit (*Oryctolagus cuniculus*) Population in South Eastern Nigeria Using Microsatellite Markers

A. I. Adeolu, M. Wheto, V. U. Oloforuh-Okoleh, R. N. Nwose, A. S. Adenaike, A. Yakubu, E. M. Abiola, & B. G. Mohammed

Feed Intake and Nutrient Digestibility, Rumen Fermentation Profiles, Milk Yield and Compositions of Lactating Dairy Cows Supplemented by *Flemingia macrophylla* Pellet

B. Phesatcha, K. Phesatcha, B. Viennaxay, N. T. Thao, & M. Wanapat

Nutrients, Fiber Fraction, and *In Vitro* Fiber Digestibility of Brown-Midrib Sorghum Mutant Lines Affected by The Maturity Stages

R. Sriagtula, P. D. M. H. Karti, L. Abdullah, Supriyanto, D. A. Astuti, & Zurmiati

Feed Intake, Nutrient Digestibility, Antioxidant Activity in Plasma, and Growth Performance of Male Dairy Cattle Fed Black Rice and Purple Corn Extracted Residue

R. Prommachart, J. Uriyapongson, A. Cherdthong, & S. Uriyapongson

Performances of Crossbred Chickens Fed Fermented Papaya Leaf and Seed Powder at High Stocking Density

S. Sugiharto, T. Yudiarti, E. Widiastuti, H. Wahyuni, T. Sartono, & A. R. Pratama

Comparison of Different Herbal Additives on Immune Response and Growth Performance of Broiler Chickens

S. Masoud-Moghaddam, J. Mehrzad, A. H. Alizadeh-Ghamsari, R. Bahari Kashani, & J. Saeidi

Environmental Monitoring of Zoonotic Fungal Infection in Broiler Chickens: Novel Approach to Control using Nanofungicide Composite

A. N. Mohammed & G. K. Abdel-Latef

Composition and Yield of Colostrum and Milk from Murrah and "Murrah x Carabao" Crosses in the Philippines

O. L. Bondoc, T. Almendral-Saludes, A. G. Tandang, A. R. Bustos, A. R. Ramos, & A. O. Ebron

Comparison between the Brix Refractometer and ELISA for Assessment of Colostrum IgG of Montbeliard x Holstein Cross Dairy Cows

N. H. Fahim & T. A. Imbabi

Detection of Class 1 Integron Encoding Gene in Multidrug Resistance (MDR) *Citrobacter freundii* Isolated from Healthy Broiler Chicken

A. M. Witaraningrum, F. J. Wibisono, D. A. Permatasari, & M. H. Eftendi

Characterization and Antimicrobial Activity of Whey Edible Film Composite Enriched with Clove Essential Oil

Fahrullah, M. Ervandi, & D. Rosyidi

Scopus[®]

Clarivate
Analytics
EMERGING SOURCES
CITATION INDEX

SJR

DOAJ

DIRECTORY OF
OPEN ACCESS
JOURNALS

ASEAN
CITATION
INDEX

Crossref

CAB Abstracts

EBSCO
PUBLISHING

Google
scholar

Sinta

Tropical Animal Science *Journal*

Accredited by Directorate General of Strengthening for Research and Development,
Ministry of Research, Technology and Higher Education of the Republic of Indonesia
No: 30/E/KPT/2018

Vol. 44 No. 3, September 2021

PUBLISHER

Faculty of Animal Science, IPB University (Bogor Agricultural University) associated with Animal Scientist's Society of Indonesia

CHIEF EDITOR

Komang G. Wiryawan

ASSOCIATE EDITORS

Tuti Suryati
Sri Suharti
Jakaria

EDITORIAL BOARDS

Abdul Razak Alimon (Malaysia) ~2023
Anjas Asmara B. Samsudin (Malaysia) ~2021
Anuraga Jayanegara (Indonesia) ~2021
Arief Boediono (Indonesia) ~2024
Armagan Hayirli (Turkey) ~2023
Asep Gunawan (Indonesia) ~2023
Budi Santoso (Indonesia) ~2021
Burhanudin Sundu (Indonesia) ~2023
Cahyo Budiman (Indonesia) ~2021

Cece Sumantri (Indonesia) ~2024
Elizabeth Wina (Indonesia) ~ 2023
Epi Taufik (Indonesia) ~2021
Irma Isnafia Arief (Indonesia) ~2024
Jong K. Ha (Korea) ~2021
Junichi Takahashi (Japan) ~2024
Loh Teck Chwen (Malaysia) ~2022
Lucia Cyrilla (Indonesia) ~2022
Luki Abdullah (Indonesia) ~2021

Metha Wanapat (Thailand) ~2024
Mulyoto Pangestu (Australia) ~2022
Myunggi Baik (Korea) ~2021
Nahrowi (Indonesia) ~2024
Rita Mutia (Indonesia) ~2022
Sri Mulatsih (Indonesia) ~2021
Wasmen Manalu (Indonesia) ~2024

TECHNICAL EDITORS

Irma Nuranthi Purnama
Yuni Cahya Endrawati

SCOPE OF JOURNAL

Tropical Animal Science Journal receives original manuscripts encompass a broad range of research topics in tropical animal sciences: breeding and genetics, reproduction and physiology, nutrition, feed sciences, agrostology, animal products, biotechnology, behaviour, welfare, health and veterinary, livestock farming system, socio-economic, animal waste management, and policy.

Authors have to prepare the manuscripts according to Tropical Animal Science Journal's guidelines. Manuscript and statement of originality and copyright release form (Form A) should be submitted via online submission through journal website: <http://journal.ipb.ac.id/index.php/tasj>.

ABOUT JOURNAL

Tropical Animal Science Journal (Trop. Anim. Sci. J.) previously Media Peternakan is a scientific journal covering broad aspects of tropical animal sciences. Started from 2018, the title is changed from Media Peternakan in order to develop and expand the distribution as well as increase the visibility of the journal. The journal is published FOUR times a year in March, June, September, and December by Faculty of Animal Science, IPB University (Bogor Agricultural University), associated with Animal Scientist's Society of Indonesia. Tropical Animal Science Journal is a peer reviewed journal that has been indexed and abstracted in Elsevier products (Scopus, Reaxys), Clarivate Analytics product (Emerging Sources Citation Index), Scimago Journal Rank, ASEAN Citation Index, DOAJ, Dimensions-Digital Science, CABI, EBSCO, Science and Technology Index (SINTA), Google Scholar, and other scientific databases. The journal also uses Similarity Check to prevent any suspected plagiarism in the manuscripts.

SECRETARIAT OF TROPICAL ANIMAL SCIENCE JOURNAL

Faculty of Animal Science Building, IPB University (Bogor Agricultural University)
Jalan Agatis, Kampus IPB Dramaga, Bogor 16680, Indonesia
Phone/Fax: +62-251-8421692
e-mail: mediapeternakan@apps.ipb.ac.id; mediapeternakan@yahoo.co.id
website: <http://journal.ipb.ac.id/index.php/tasj>

Subscription Rate (Not including shipping and handling):

IDR 100.000/ edition and IDR 350.000/ year (Indonesian subscribers); 50 US\$ (Foreign subscribers)

Bank account: Bank Rakyat Indonesia KCP KAMPUS IPB No. 0595-01-005208-53-3, o.b. Media Peternakan, SWIFT CODE: BRINIDJA

Tropical Animal Science *Journal*

Accredited by Directorate General of Strengthening for Research and Development,
Ministry of Research, Technology and Higher Education of the Republic of Indonesia
No: 30/E/KPT/2018

HISTORY

Tropical Animal Science Journal (Trop. Anim. Sci. J.) (p-ISSN 2615-787X and e-ISSN 2615-790X) previously Media Peternakan (published from 1967-2017 with p-ISSN 0126-0476 and e-ISSN 2087-4634) is a scientific journal covering broad aspects of tropical animal sciences. Started from 2018, the title is changed from Media Peternakan in order to develop and expand the distribution as well as increase the visibility of the journal. The journal is published FOUR times a year in March, June, September, and December started from the year 2020 by Faculty of Animal Science, IPB University (Bogor Agricultural University), associated with Animal Scientist's Society of Indonesia.

The first edition with the new title was published in April 2018 edition (Vol 41 No 1 2018), while the previous edition (up to 2017 edition) still use Media Peternakan as the title. The online version of Tropical Animal Science Journal could be accessed in the new website (<http://journal.ipb.ac.id/index.php/tasj>) and the previous editions are available in the old website (<http://medpet.journal.ipb.ac.id/>).

Tropical Animal Science Journal has been indexed and abstracted in Elsevier products (Scopus, Reaxys), Clarivate Analytics products (Emerging Sources Citation Index), Scimago Journal Rank, ASEAN Citation Index, DOAJ, Dimensions-Digital Science, CABI, EBSCO, Science and Technology Index (SINTA), Google Scholar, and other scientific databases. The journal also uses Similarity Check to prevent any suspected plagiarism in the manuscripts.

Tropical Animal Science Journal

Accredited by Directorate General of Strengthening for Research and Development,
Ministry of Research, Technology and Higher Education of the Republic of Indonesia
No: 30/E/KPT/2018

LIST OF CONTENT

Artificial Insemination in Sheep with Fresh Diluted Semen: Comparison Between Two Different Semen Extenders and Management Protocols	255
A. Madrigali, A. Rota, D. Panzani, S. Castellani, M. Shawahina, A. Hassan, F. Di Iacovo, C. Rossignoli, & F. Camillo	
The Protective Impacts of α-tocopherol Supplementation on the Semen Quality of Sopera Goat Preserved at 4°C	261
R. A. Prastiya, Rimayanti, M. M. Munir, & A. P. Nugroho	
Detection of Tetracycline Resistance Genes among <i>Escherichia coli</i> Isolated from Layer and Broiler Breeders in West Java, Indonesia	267
A. Indrawati, K. Khoirani, S. Setyaningsih, U. Affif, Safika, & S. G. Ningrum	
Growth Performance of Male Line of Superior Tropical Rabbit	273
B. Brahmantiyo, N. Pratiwi, F. Saputra, Y. C. Raharjo, & L. H. Prasetyo	
Genetic Diversity of Rabbit (<i>Oryctolagus cuniculus</i>) Population in South Eastern Nigeria Using Microsatellite Markers	280
A. I. Adeolu, M. Wheto, V. U. Oleforuh-Okoleh, R. N. Nwose, A. S. Adenaike, A. Yakubu, E. M. Abiola, & B. G. Mohammed	
Feed Intake and Nutrient Digestibility, Rumen Fermentation Profiles, Milk Yield and Compositions of Lactating Dairy Cows Supplemented by <i>Flemingia macrophylla</i> Pellet	288
B. Phesatcha, K. Phesatcha, B. Viennaxay, N. T. Thao, & M. Wanapat	
Nutrients, Fiber Fraction, and <i>In Vitro</i> Fiber Digestibility of Brown-Midrib Sorghum Mutant Lines Affected by The Maturity Stages	297
R. Sriagtula, P. D. M. H. Karti, L. Abdullah, Supriyanto, D. A. Astuti, & Zurmiati	
Feed Intake, Nutrient Digestibility, Antioxidant Activity in Plasma, and Growth Performance of Male Dairy Cattle Fed Black Rice and Purple Corn Extracted Residue	307
R. Prommachart, J. Uriyapongson, A. Cherdthong, & S. Uriyapongson	
Performances of Crossbred Chickens Fed Fermented Papaya Leaf and Seed Powder at High Stocking Density	316
S. Sugiharto, T. Yudiarti, E. Widiastuti, H. Wahyuni, T. Sartono, & A. R. Pratama	
Comparison of Different Herbal Additives on Immune Response and Growth Performance of Broiler Chickens	327
S. Masoud-Moghaddam, J. Mehrzad, A. H. Alizadeh-Ghamsari, R. Bahari Kashani, & J. Saeidi	
Environmental Monitoring of Zoonotic Fungal Infection in Broiler Chickens: Novel Approach to Control using Nano-fungicide Composite	336
A. N. Mohammed & G. K. Abdel-Latef	
Composition and Yield of Colostrum and Milk from Murrah and "Murrah x Carabao" Crosses in the Philippines	347
O. L. Bondoc, T. Almendral-Saludes, A. G. Tandang, A. R. Bustos, A. R. Ramos, & A. O. Ebron	
Comparison between the Brix Refractometer and ELISA for Assessment of Colostrum IgG of Montbeliard x Holstein Cross Dairy Cows	356
N. H. Fahim & T. A. Imbabi	
Detection of Class 1 Integron Encoding Gene in Multidrug Resistance (MDR) <i>Citrobacter freundii</i> Isolated from Healthy Broiler Chicken	363
A. M. Witaningrum, F. J. Wibisono, D. A. Permatasari, & M. H. Effendi	
Characterization and Antimicrobial Activity of Whey Edible Film Composite Enriched with Clove Essential Oil	369
Fahrullah, M. Ervandi, & D. Rosyidi	

The Protective Impacts of α -tocopherol Supplementation on the Semen Quality of Sopera Goat Preserved at 4°C

R. A. Prastiya^{a*}, Rimayanti^a, M. M. Munir^b, & A. P. Nugroho^c

^aDepartment of Veterinary Reproduction, Faculty of Veterinary Medicine, Universitas Airlangga

^bStudent, Faculty of Veterinary Medicine, Universitas Airlangga

Jalan Mulyorejo Kampus C Universitas Airlangga 60115, Indonesia

^cFaculty of Animal Husbandry, Universitas Jenderal Soedirman

Jalan Dr. Soeparno Karangwangkal, Purwokerto 53122, Indonesia

*Corresponding author: ragilap@fkh.unair.ac.id

(Received 04-06-2020; Revised 13-11-2020; Accepted 30-11-2020)

ABSTRACT

Supplementation of vitamin E (α -tocopherol) as an antioxidant into an extender is useful to maintain sperm quality during the process of cryopreservation by protecting the sperms from reactive oxygen species (ROS). This study aims to evaluate the protective impacts of vitamin E supplementation on the motility, viability, and reducing the abnormality of sperms in Sopera goats. The sample of semen used in the present study was obtained from fresh semen and good-quality ejaculates of the Sopera goats collected by an artificial vagina. The groups involve one controlled treatment group, which contains semen and tris egg yolk (TEY) extender without vitamin E (P0), and three treatment groups that contain semen, TEY extender, and different doses of vitamin E, i.e., 1 mg/mL (P1), 2 mg/mL (P2), and 3 mg/mL (P3), respectively. Data obtained from the experiments were analyzed using Repeated Measures Multivariate Analysis (MANOVA). The results showed that TEY extenders added with vitamin E at a dose of 1 mg/mL of vitamin E (P1) produced the highest motility and viability with the lowest abnormalities of sperm after 24th hour of cryopreservation. The lowest sperm motility and viability among the treatment groups was observed in the TEY extender added with vitamin E at a dose of 3 mg/mL (P3) at 96th hour after cryopreservation, which was almost the same as the control group without vitamin E (P0) that had the lowest motility and viability. In conclusion, the current study found that supplementation of vitamin E as an antioxidant at a dose of 1 mg/mL in TEY extender can preserve semen of the Sopera goats in refrigerator up to 96 h at 4°C and provides higher sperm motility and viability percentage, with a lower percentage of sperm abnormality.

Keywords: vitamin E (α -tocopherol); extender; Sopera goat; sperm assessment; animal medicine

INTRODUCTION

The Sopera goat is one of the dairy goats in Indonesia, which is a crossbreed between the Saanen goat from Switzerland and the Etawah crossbreed goat (Peranakan Etawa). The crossbreed between the two goats is intended to boost both milk and meat productions with a greater carcass than the Saanen goats. The Sopera goats can produce 2 to 4 liters of milk per day during the first lactation, which is higher than the Etawah goats and the Saanen goats (Praharani *et al.*, 2016). In order to increase the national milk production of dairy goats, it is necessary to increase the population of dairy goats, such as the Sopera goats.

One of the efforts to increase the population of the Sopera goats is through the application of artificial insemination (AI), which allows a higher ratio of does to bucks compared to the natural breeding system. AI in goats has become a powerful method for breeders or farmers to obtain genetically-high quality offsprings by using superior bucks. Moreover, AI can also minimize the transmission of reproductive diseases and provide a

more accurate birth of the offsprings (Cseh *et al.*, 2012; Faigl *et al.*, 2012). The success of AI using fresh semen relies on the techniques of sperm cryopreservation that semen inseminated to the female reproductive tract can reach the ovum cells (Wang *et al.*, 2015). Various cellular and molecular damages of sperm during the cryopreservation process will affect the sperm function and fertilization potential. Decreased sperm fertilization potential during semen cryopreservation can be observed and calculated from the excessive production of reactive oxygen species (ROS) that induce cryo-injury during oxidative stress (Johnston *et al.*, 2012; Kim *et al.*, 2012; Liu *et al.*, 2019). To overcome this problem, the addition of antioxidants such as vitamin E in the extenders during cryopreservation of semen is necessary (Sharafi *et al.*, 2015; Forouzanfar *et al.*, 2013a; Forouzanfar *et al.*, 2013b; Widjaya, 2017).

Antioxidants can maintain the qualities of sperms by protecting the biological membrane from damages caused by free radicals, neutralizing hydroxyl groups, superoxide, and hydrogen peroxide radicals, as well as preventing sperm agglutination

(Agarwal & Prabakaran, 2005; Breininger *et al.*, 2005). Supplementation of vitamin E into the extender will inhibit the process of lipid peroxidation reaction that helps to manage the oxidation process of phosphorylation, which is responsible for the increasing levels of Reactive Oxygen Species (ROS) in the sperm. High levels of ROS in the cells can oxidize lipids, proteins, and DNA so which can cause DNA damage, increased membrane permeability, and even reduce reproductive capacity (Noori, 2012; Paul *et al.*, 2018; Aitken, 2017). Vitamin E can function as an intracellular antioxidant that plays an important role in preventing lipid peroxidation of unsaturated fatty acids inside the cells and on the cell membrane so that it can prevent peroxidative damage that affects spermatogenesis (Spirlandeli *et al.*, 2014).

Information about the effect of antioxidant supplementation on Sapera buck's semen is limited from the previous studies. This gap in the literature drives us to study *in vitro* supplementation of Vitamin E in the semen extender to maintain the quality of Sapera buck semen.

MATERIALS AND METHODS

Animals

All experiments were approved by the Airlangga Health Research Ethical Clearance Committee number 011/HRECC.FODM/I. The study was conducted with three Sapera goats (2-3 years of age) with good body conditions were used and placed in hygienic and properly-ventilated sheds. The temperature inside the shed was maintained at 33-41°C (average 37°C). The semen collection and semen analysis were carried out at the PSDKU teaching farm, Airlangga University, during the period extended from January to May 2019. The animals were supplied with clean water that was available *ad libitum* and were fed twice a day consisting of calliandra hay and Napier grass silage as forage, tofu (soybean curd) by-product, and commercial concentrate as feed supplement according to the nutritional requirements.

Preparations of Vitamin E and Extender

Pure Vitamin E (DL- α -Tocopherol acetate solution (10 mM), MDL Number: MFCD00006848) used in this study was obtained from Sigma (St. Louis, MO, USA) and stored in a refrigerator at 5°C. Prior to the addition into the semen extender, α -tocopherol was dissolved using ethanol 0.05%. Vitamin E was divided into 3 different concentrations (1 mg/mL, 2 mg/mL, and 3 mg/mL), and each concentration of vitamin E was mixed with the Tris-Egg-Yolk (TEY) extender. The extender material was made of Tris-Egg-Yolk obtained from the Singosari Artificial Insemination Center. Extender contains 1.6% Tris Aminomethane, 0.9% citric acid, 1.4% lactose, 80% distilled water, 20% egg yolk, penicillin 1000 IU/mL, streptomycin 1 mg/mL, and vitamin E (α -tocopherol) according to the aforementioned doses. Antibiotics such as penicillin and streptomycin were added to the extender to suppress bacterial growth. The equipment used for

this process were cleaned and sterilized. The ratio between semen and tris-egg-yolk extender was 1:10.

Semen Collection

Semen collections from 3 Sapera bucks were started with a physical examination of the reproductive organs of the bucks. The organs were healthy with a high libido. Semen collection was performed using an artificial vagina once a week in the morning with a total of one ejaculate. Fresh semen that was collected was evaluated macroscopically (volume, consistency, odor, color, and pH) and microscopically (sperm motility, viability, and concentration). The standard of sperm progressive motility in this study was above or equal to 70%. The sperm concentration was evaluated using a spectrophotometer.

Experimental Design

The samples used in this study were 36 semen ejaculates collected from three Sapera bucks having good fresh semen and high libido at the ages of 2-3 years old. Tris-egg yolk extender was added to the semen, with a ratio of 1:10 (semen volume to extender). The semen mixed with TEY extender was assigned into four different groups based on the different doses of vitamin E supplementation, namely P0 (0 mg/mL), P1 (1 mg/mL), P2 (2 mg/mL), and P3 (3 mg/mL). All samples were preserved in liquid storage at 4°C for 96 hours and assessed for their parameters, namely, sperm progressive motility, percent of viability, and percent of abnormalities at 24, 48, 72, and 96 hours of storage.

Assessment of Sperm Motility

To evaluate the motion characteristics of rewarmed sperm, computer-assisted sperm analysis (CASA, Version 5.1; Microptic, Barcelona, Spain) was used. The evaluation was performed at 37°C to facilitate the optimal movement of sperms. Calculation of sperm motility was based on the speed and direction of movement. As for the direction of movement, sperms were classified into 4 categories, i.e., P for progressive, R for reverse, O for oscillatory, and N for Necrospermia or no movement, while for the speed of movement, they were categorized with a number, i.e., 0 for no movement, 1 for slow movement, 2 for moderate movement, 3 for rapid movement, and 4 for very fast movement.

Assessment of Sperm Viability

Sperm viability was assessed by using the Eosin-Nigrosine staining method (Najafi *et al.*, 2013). The glass slide was prepared, and then a small drop of semen (2 μ L) was put on the glass slide, and eosin-nigrosin (10 μ L) solution was put next to the semen drop. Subsequently, a thin preparation was made by fixation, and the preparation was dried for about 15 seconds. The sperm viability was assessed after fixation and drying of the preparation by calculating 200 sperm cells under bright field microscopy (CKX41; Olympus, Tokyo,

Japan) at 400X magnifications. Live sperms remained unstained, while dead sperms were stained, turning into a red-purple color.

Assessment of Sperm Abnormality

Evaluation for abnormalities of sperm was conducted by using preparation used for viability testing (Felipe-Perez *et al.*, 2008). At least 200 sperms were examined under a microscope with 400x magnification. Abnormal sperm were determined based on head and tail abnormalities.

Statistical Analysis

Data about motility, viability, and abnormal morphology of sperm were subsequently arranged in a table presented as average ± SD. Multivariate (MANOVA) on SPSS version 20.0 (SPSS Inc., Chicago, IL, USA) was employed to analyze the data. If the results indicated $p < 0.05$, the subsequent analysis using Duncan's multiple range test was performed.

RESULTS

The quality of fresh semen of Sapera bucks is presented in Table 1. The average volume of semen was 1.00 ± 0.20 mL from the macroscopic examination, with a creamy color, a goat-smelly odor, and thick consistency that were observed by tilting the tubes. The pH obtained in this study was 6.60 ± 0.54 . The standard volume of semen in goats is an average of 1 ml (0.5-2 mL), the color of the semen is creamy white with a thick consistency, and pH 6.4-6.8. The initial macroscopic examination results listed in Table 1 showed the normal standards for Sapera goat semen and met the criteria for further examination.

The results of a microscopic evaluation in this study showed that the average concentration of sperms was 3876 ± 167.57 (million/mL). The average mass motility was (+++) marked by a large-wave formation of sperms. Furthermore, the sperms had progressive motility (forward movements) with an average of $82.80 \pm 2.16\%$. The average result was 82%, with a speed of 3-4. The average

sperms viability was $89.40 \pm 4.33\%$. The results found in this study are within the normal standard of goat sperm concentrations. Sperm abnormalities in Sapera bucks sperm ranged from 3%-4.5%, with an average of 3.20 ± 0.83 .

The effects of doses of vitamins E addition in the TEY extender on the sperm characteristics (progressive sperm motility, viability, and abnormal sperm (mean ± SD) of Sapera bucks following the preservation at 4 °C for 96 hours are presented in Table 2. Means of sperm motility and viability were significantly affected by the treatment of doses of tocopherol (Vitamin E) addition in the TEY extender. No significant difference in sperm abnormality was observed among the doses of tocopherol addition in the extender. However, semen diluted in TEY extender and added with vitamin E at a dose of 1 mg/ml tended to have a lower sperm abnormality. The evaluation of sperm motility showed that doses of vitamin E in the extender (0 mg/mL, 1 mg/mL, 2 mg/mL, and 3 mg/mL) ($p < 0.05$) significantly affected the sperm motility. However, looking at the more detailed comparison among the doses of vitamin E, there was no significant difference in sperm motility between semen preserved in TEY extender contained vitamin E at doses of 3 and 0 mg/mL, and the highest motility was found in the semen preserved in TEY extender contained 1 mg/mL vitamin E (P1). In the semen preserved in the TEY extender added with vitamin E, the sperm motilities declined gradually when preserved at 4°C from 24 to 96 hours to $61.30 \pm 3.88\%$ and $28.78 \pm 3.57\%$, respectively, compared to semen preserved in the TEY extender without vitamins (control) which was declined markedly to $47.99 \pm 2.85\%$ and $33.75 \pm 3.57\%$, respectively.

In terms of sperm viability, there were significant differences in the sperm viability among the doses of vitamin E and time of preservation ($p < 0.05$). In addition, there were significant decline patterns in treatments P1, P2, P3, and P0 ($p < 0.05$). The highest sperm viability was found in semen preserved in TEY extender added with vitamin E at a dose of 1 mg/mL (P1) ($p < 0.05$). The results revealed that there were significant differences between P1 and the other treatments ($p < 0.05$). Semen, preserved in TEY extender added with vitamin E at a dose of 1 mg/mL, showed the lowest sperm abnormal-

Table 1. Ejaculates characteristics of Sapera buck semen (mean ± standard deviation)

Evaluation	Repetition					Mean±SD
	1	2	3	4	5	
Macroscopy						
Volume (mL)	1.20	0.80	1.20	1.00	0.80	1.00 ± 0.20
pH	7	7	6	7	6	6.60 ± 0.54
Consistency	Thick	Thick	Thick	Thick	Thick	Thick
Microscopy						
Sperm motility (%)	80	82	85	82	85	82.80 ± 2.16
Sperm concentration (million/mL)	3720	3840	4020	3720	4080	3876 ± 167.57
Sperm viability (%)	85	86	90	96	90	89.40 ± 4.33
Mass movement	+++	+++	++	+++	+++	+++
Sperm abnormalities (%)	4	3	2	4	3	3.20 ± 0.83

Note: SD= Standard deviation. ++ = Frequent and seldom wave motion of mass sperm motility; +++ = Fast and thick wave motion of mass sperm motility

Table 2. Sperm motility, viability, and abnormality of Sapera buck treated by different vitamin E doses and time

Vitamin E doses	Variables (%) (Mean ± SD)			
	Time	Motility	Viability	Abnormality
(P0) 0 mg/mL	24 hours	47.99 ± 2.85 ^{cd}	71.22 ± 6.19 ^b	5.41 ± 1.22 ^b
	48 hours	47.60 ± 5.80 ^{cd}	62.20 ± 6.19 ^c	5.45 ± 1.04 ^b
	72 hours	41.22 ± 4.61 ^d	60.32 ± 5.49 ^c	5.33 ± 1.88 ^b
	96 hours	33.75 ± 3.57 ^e	49.56 ± 6.28 ^e	6.09 ± 1.19 ^b
(P1) 1 mg/mL	24 hours	61.30 ± 3.88 ^a	77.85 ± 6.37 ^a	3.41 ± 1.22 ^a
	48 hours	55.55 ± 6.17 ^b	70.99 ± 8.05 ^b	3.54 ± 1.66 ^a
	72 hours	50.40 ± 3.93 ^c	60.52 ± 5.49 ^c	3.75 ± 1.01 ^a
	96 hours	42.78 ± 3.57 ^d	54.27 ± 6.33 ^d	4.29 ± 0.16 ^{ab}
(P2) 2 mg/mL	24 hours	49.30 ± 3.80 ^c	70.33 ± 4.33 ^b	5.01 ± 1.22 ^b
	48 hours	42.40 ± 6.64 ^d	60.17 ± 9.01 ^c	5.95 ± 1.66 ^b
	72 hours	39.40 ± 4.02 ^{de}	57.50 ± 5.22 ^{cd}	5.33 ± 1.09 ^b
	96 hours	30.78 ± 2.99 ^e	48.20 ± 5.09 ^e	5.29 ± 1.04 ^b
(P3) 3 mg/mL	24 hours	48.56 ± 2.58 ^c	71.85 ± 6.29 ^b	4.48 ± 1.22 ^{ab}
	48 hours	49.28 ± 5.91 ^c	60.42 ± 8.67 ^c	5.22 ± 1.53 ^b
	72 hours	39.40 ± 4.33 ^{de}	59.50 ± 6.49 ^c	5.65 ± 1.11 ^b
	96 hours	28.78 ± 3.57 ^e	47.24 ± 6.00 ^e	6.22 ± 0.99 ^{bc}

Note: Means in the same column with different superscripts differ significantly (p<0.05). Control (P0)= No addition of vitamin E; P1= addition of 1mg/mL vitamin E to the extender; P2= addition of 2mg/mL vitamin E to the extender; P3= addition of 3mg/mL vitamin E to the extender, respectively.

ity. Subsequent analysis was conducted to evaluate the effect of time, i.e., at 4°C preservation for 24, 48, 72, and 96 hours, on the motility, viability, and abnormalities of spermatozoa. The results showed a significant effect of time on motility, as was shown by the significant difference among the 24th, 48th, 72nd, and 96th hours after freezing (p<0.05). Furthermore, the results of the viability analysis showed a significant difference between the 24th and 48th hours, and between the 72nd and 96th hours (p<0.05), and a significant decrease in patterns between the 24th, 48th, 72th, and 96th hours (p<0.05).

Finally, from the two initial parameters, the motility and viability, vitamin E's addition at a dose of 1 g/mL (P1) showed the best results to Sapera bucks semen diluted with TEY on sperm quality.

DISCUSSION

Sperm motility is powered by cellular energy production, which is resulted from oxidative phosphorylation and is transferred into the microtubules in the tail to stimulate and support tail movements (O'Connell *et al.*, 2002). There are two main pathways of energy production in the sperm, namely glycolysis and oxidative phosphorylation (Storey, 2008). The previous research shows that oxidative phosphorylation is more effective in energy production for mammalian sperm (Tafif, 2019). The process of oxidative phosphorylation occurs in the mitochondria through the electron transport chain. Sperm mitochondria are located in the central part, and energy production takes place in the gyre (Gibb & Aitken, 2016). ROS are formed during the process of oxygen reduction in oxidative phosphorylation. Although the ROS binding system exists in the electron transport chain, electron leakage by previous components can still occur (Halliwell & Gutteridge, 2015).

The cooling process reduces membrane fluidity, sperm motility, and speed (Fang *et al.*, 2017). Cryopreservation also decelerates the metabolic rate in the sperm. However, a decrease in metabolic rate cannot stop cells from producing ROS (Gibb & Aitken, 2016; Prastiya *et al.*, 2019). The previous results of a study suggested that decreased sperm motility and mitochondrial function were found in refrigerated semen and mitochondrial dysfunction is a major cause of decreased sperm motility (O'Connell *et al.*, 2002; Tafif, 2019).

The positive outcomes of vitamin E supplementation in domestic animals have been widely studied, including in a sample of goats (Azawi & Hussein, 2013). Adding vitamin E to the semen extender also results in improving the male's fertility parameters (Hadi, 2016). Sperm membranes are mostly composed of Polyunsaturated Fatty Acids (PUFA) that makes the sperm is very susceptible to lipid peroxidation. Vitamin E supplementation helps in maintaining the sperm motility of cooled semen of bulls (Ratnani *et al.*, 2017). This helps in finding free radicals created from lipid peroxidation by creating two new stable radicals from one vitamin E molecule (Hajibabaei, 2016). Vitamin E has lipophilic properties (Azzi, 2007). Therefore, the percentage of lipids in the structure of sperm and lipid compounds from semen extender can influence the amount of vitamin E required for supplementation. In this study, high vitamin E concentrations showed a detrimental effect in maintaining motility, presumably because the formation of free radicals from metal reduction is facilitated by the high doses of vitamin E (Khorramabadi *et al.*, 2017). These metals are Fe and Cu (Halliwell & Gutteridge, 2015), which are contained in large quantities in the seminal plasma (Hardijanto *et al.*, 2010). In the previous studies, the effect of supplementation of vitamin E with concentrations of 1.50 mg

and 1 mg/mL in bull semen increases the percentage of sperm motility (Ullah *et al.*, 2019). These findings are similar to data reported in various species of mammals, including in bulls, Murrah buffaloes, and cattle families (Ullah *et al.*, 2019). These results indicate that vitamin E supplementation can maintain sperm motility during cryopreservation. The low antioxidant content of seminal plasma also decreases sperm motility, viability, and normal morphology (Am-in *et al.*, 2011). Vitamin E, which presents in the mammalian cell membranes and seminal plasma, cleanses ROS and consequently relieves oxidative stress (Sikka, 2004). The implementation of synthetic analogs of vitamin E also substantially protects membrane ultrastructure and improves intact plasma and mitochondrial membranes (Soares *et al.*, 2015).

Reactive oxygen is generated by leakage of the electron transport chain of the mitochondria and can also be resulted from damages in the mitochondrial structure (Halliwell & Gutteridge, 2015). ROS also impairs the mitochondria of sperm, creating multiple amounts of ROS production. They bind PUFA hydrogen and produce lipid peroxyl radicals that will bind the adjacent PUFA molecules, producing lipid peroxide and an alkyl radical. This cycle is known as a chain reaction (Hajibabaei, 2016).

An antioxidant supplement that can supply hydrogen atoms to ROS is necessary to prevent chain reactions. OH α -tocopherol actively supplies hydrogen atoms (Herdis *et al.*, 2013). Therefore, such antioxidants can be promising to prevent oxidative damage such as lipid peroxidation (Maya *et al.*, 2012). Vitamin E is also known as a chain-breaking antioxidant because it provides its hydrogen atom and forms alpha-tocopheroxyl radicals that will react with the second free radical. Therefore, a single molecule of α -tocopherol will consume two free radicals in the total (Hajibabaei, 2016). However, the percentage of lipids in the environment affects the solubility of vitamin E, a lipophilic antioxidant that directly affects the lipid domain. The amount of lipid content differs between semen extender and sperm architecture. Therefore, the effect of the dose of vitamin E supplementation can differ depending on the content of the extender and animal species (Azzi, 2007). According to Khorramabadi *et al.* (2019), vitamin E shows a pro-oxidant activity, so that vitamin E may be useful for eliminating complications resulting from oxidative stress. In this study, the evaluation of results produced by the addition of vitamin E at a dose of 3 mg/mL (P3) suggests that high vitamin E supplementation concentrations can bring a detrimental effect. This is likely due to the ability of vitamin E to reduce Fe and Cu (Halliwell & Gutteridge, 2015).

CONCLUSION

In conclusion, the current study found that supplementation of vitamin E as an antioxidant at a dose of 1 mg/mL in TEY extender can preserve semen of the Sapera goats in a refrigerator up to 96 h at 4°C and provides higher sperm motility and viability percentage, and a lower sperm abnormality percentage.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

ACKNOWLEDGEMENT

The authors are thankful to the Dean of Faculty of Veterinary Medicine and the Coordinator PSDKU Airlangga University (Universitas Airlangga) for approval of research work with SK no.1185/UN3/2018.

REFERENCES

- Agarwal, A. & S. A. Prabakaran. 2005. Oxidative stress and antioxidants in male infertility: A difficult balance. *Iran. J. Reprod. Med.* 3:1-8.
- Aitken, R. J. 2017. Reactive oxygen species as mediators of sperm capacitation and pathological damage. *Mol. Reprod. Dev.* 84:1039-1052. <https://doi.org/10.1002/mrd.22871>
- Am-In, N., R. Kirkwood, M. Techakumphu, & W. Tantasuparuk. 2011. Lipid profiles of sperm and seminal plasma from boars having normal or low sperm motility. *Theriogenology* 75:897-903. <https://doi.org/10.1016/j.theriogenology.2010.10.032>
- Am-In, N., R. Kirkwood, M. Techakumphu, & W. Tantasuparuk. 2010. Effect of storage for 24 h at 18°C on sperm quality and a comparison of two assays for sperm membrane lipid peroxidation. *Can. J. Anim. Sci.* 90:389-392. <https://doi.org/10.4141/CJAS09122>
- Azzawi, O. I. and E. K. Hussein. 2013. Effect of vitamins C or E supplementation to tris diluent on the semen quality of Awassi rams preserved at 5°C. *Vet. Res. Forum.* 4:157-160.
- Azzi, A. 2007. Molecular mechanism of alpha-tocopherol action. *Free. Radic. Biol. Med.* 43:16-21. <https://doi.org/10.1016/j.freeradbiomed.2007.03.013>
- Breiner, E., N. B. Beorlegui, C. M. O'Flaherty, & M. T. Beconi. 2005. Alpha-tocopherol improves biochemical and dynamic parameters in cryopreserved boar semen. *Theriogenology* 63:2126-2135. <https://doi.org/10.1016/j.theriogenology.2004.08.016>
- Cseh, S., V. Faigl, & G. S. Amiridis. 2012. Amiridis, semen processing and artificial insemination in health management of small ruminants. *Anim. Reprod. Sci.* 130:187-192. <https://doi.org/10.1016/j.anireprosci.2012.01.014>
- Faigl, V., N. Vass, A. Javor, M. Kulcsar, L. Solti, G. Amiridis, & S. Cseh. 2012. Artificial insemination of small ruminants: A review. *Acta Vet. Hung.* 60: 115-129. <https://doi.org/10.1556/AVet.2012.010>
- Fang, Q., J. Wang, Y. Y. Hao, H. Li, J. X. Hu, & G. S. Yang. 2017. Effects of iodine methionine on boar sperm quality during liquid storage at 17°C. *Reprod. Domest. Anim.* 52: 1061-1066. <https://doi.org/10.1111/rda.13024>
- Felipe-Perez, Y. E., M. L. Juarez-Mosqueda, E. O. Hernandez-Gonzalez, & J. J. Valencia. 2008. Viability of fresh and frozen bull sperm compared by two staining techniques. *Acta Vet. Bras.* 2:123-130.
- Forouzanfar, M., A. Abid, S. M. Hosseini, M. Hajian, & M. H. N. Esfahani. 2013a. Supplementation of sperm cryopreservation media with cell permeable superoxide dismutase mimetic agent (MnTE) improves goat blastocyst formation. *Cryobiology.* 67:394-397. <https://doi.org/10.1016/j.cryobiol.2013.08.004>
- Forouzanfar, M., S. F. Ershad, S. M. Hosseini, M. Hajian, S. Ostad-Hosseini, A. Abid, M. Tavalae, A. Shahverdi, A. V. Dizaji, & M. H. N. Esfahani. 2013b. Can permeable super oxide dismutase mimetic agents improve the quality of frozen-thawed ram semen?. *Cryobiology* 66:126-130.

- <https://doi.org/10.1016/j.cryobiol.2012.12.006>
- Gibb Z. & R. J. Aitken. 2016. The impact of sperm metabolism during in vitro storage: the stallion as a model. *Biomed Res. Int.* 2016:1-8. <https://doi.org/10.1155/2016/9380609>
- Hadi, I. H. 2016. Effect of vitamin E on sperm motility and survival in chilled-stored semen. *Iraqi Journal of Embryo and Infertility Researches.* 6:1-7.
- Hajibabaei, K. 2016. Antioxidant properties of vitamin E. *Ann. Res. Antioxid.* 1:e22.
- Halliwell, B. & J. M. C. Gutteridge. 2015. *Free radicals in biology and medicine*, 5th ed. Oxford, Oxford University Press.
- Hardijanto, S. Susilowati, T. Hernawati, T. Sardjito, T. W. Suprayogi. 2010. *Buku Ajar Inseminasi Buatan*. 1st ed. Pusat Penerbitan dan Percetakan Unair, Surabaya.
- Herdis, H., I. Kusuma, & D. I. W. Angga. 2013. Pengaruh penambahan a-tokoferol pada media pengencer tris kuning telur terhadap kualitas semen cair domba garut (Indonesian). *Jurnal Sains dan Teknologi Indonesia.* 11:175-180. <https://doi.org/10.29122/jsti.v11i3.836>
- Johnston, S. D., N. Satake, Y. Zee, C. Lopez-Fernandez, W.V. Holt, & J. Gosalvez. 2012. Osmotic stress and cryoinjury of koala sperm: An integrative study of the plasma membrane, chromatin stability and mitochondrial function. *Reproduction.* 143:787-797. <https://doi.org/10.1530/REP-11-0436>
- Khorramabadi, K. M., A. A. Sarcheshmeh, A. R. Talebi, & A. Mirjalili. 2019. Protective effect of vitamin E on oxidative stress and sperm apoptosis in diabetic mice. *Int. J. Reprod. Biomed.* 17:127-134. <https://doi.org/10.18502/ijrm.v17i2.3990>
- Kim, S., C. Agca, & Y. Agca. 2012. Changes in rat sperm function after cooling, cryopreservation and centrifugation processes. *Cryobiology.* 65:215-223. <https://doi.org/10.1016/j.cryobiol.2012.06.006>
- Liu T., Y. Han, T. Zhou, R. Zhang, H. Chen, S. Chen, & H. Zhao. 2019. Mechanisms of ROS-induced mitochondria dependent apoptosis underlying liquid storage of goat spermatozoa. *Aging.* 18:7880-7898. <https://doi.org/10.18632/aging.102295>
- Maya, W., K. Mayur, & S. Ashar. 2012. Pharmaceutical profile of alpha-tocopherol: A brief review. *Int. J. Pharm. Chem. Sci.* 1:1023-1039
- Najafi, A., M. Zhandi, A. Towhidi, M. Sharafi, A. A. Sharif, M. K. Motlagh, & F. Martinez-Pastor. 2013. Trehalose and glycerol have a dose-dependent synergistic effect on the post-thawing quality of ram semen cryopreserved in a soybean lecithin-based extender. *Cryobiology* 66:275-282. <https://doi.org/10.1016/j.cryobiol.2013.03.002>
- Noori, S. 2012. An overview of oxidative stress and antioxidant defensive system. *J. Clin. Cell. Immunol.* 1:1-9. <https://doi.org/10.4172/scientificreports.413>
- O'Connell, M., N. McClure, & S. E. Lewis. 2002. The effects of cryopreservation on sperm morphology, motility and mitochondrial function. *Hum. Reprod.* 17:704-709. <https://doi.org/10.1093/humrep/17.3.704>
- Paul, R. K., K. Balaganur, S. V. Bahire, D. Kumar, & R. Singh. 2018. Supplementation of cauda epididymal plasma improves sperm characteristics following liquid preservation of ram semen at 3-5°C. *Reprod. Fertil. Dev.* 30:1389-1401. <https://doi.org/10.1071/RD18063>
- Praharani, L., S. Supriyati, & R. Krisnan. 2016. A preliminary study on some reproductive traits and heterosis effects of Anglo Nubian and Etawah grade crossbred does. Indonesian Center for Animal Research and Development (ICARD), Indonesia. pp. 252-260. <https://doi.org/10.14334/Proc.Intsem.LPVT-2016-p.252-260>
- Prastiya, R. A., A. Samik, M. T. E. Purnama, & A. L. Saputro. 2019. Protective effect of mycotoxin binders on ovarian gestation mice exposed by zearalenone. *Indian Vet. J.* 96:44-46.
- Ratnani, H., M. N. Ihsan, G. Ciptadi, & S. Suyadi. 2017. Effect of alpha-tocopherol supplementation in the extender on the sperm quality of Maduran bull before and after quick freezing. *Int. J. Adv. Res.* 5:1378-1389. <https://doi.org/10.21474/IJAR01/4849>
- Sharafi, M., M. Zhandi, A. Shahverdi, & M. Shakeri. 2015. Beneficial effects of nitric oxide induced mild oxidative stress on post-thawed bull semen quality. *Int. J. Fertil. Steril.* 29:230-237. <https://doi.org/10.22074/ijfs.2015.4244>
- Sikka, S. 2004. Role of oxidative stress and antioxidants in andrology and assisted reproductive technology. *J. Androl.* 25:5-18. <https://doi.org/10.1002/j.1939-4640.2004.tb02751.x>
- Soares, A. T., S. V. Silva, A. M. Batista, F. C. Almeida, J. F. Nunes, C. A. Peixoto, & M. M. P. Guerra. 2015. Ultrastructure evaluation of goat sperm after freezing in a skim milkbased extender with Trolox supplementation. *Andrologia* 47:470-476. <https://doi.org/10.1111/and.12279>
- Spirlandeli, A. L., R. Deminice, & A. A. Jordao. 2014. Plasma malondialdehyde as biomarker of lipid peroxidation: effects of acute exercise. *Int. J. Sports Med.* 35:14-18. <https://doi.org/10.1055/s-0033-1345132>
- Storey, B. T. 2008. Mammalian sperm metabolism: Oxygen and sugar, friend and foe. *Int. J. Dev. Biol.* 52:427-437. <https://doi.org/10.1387/ijdb.072522bs>
- Tafif, K. 2019. The effect of alpha-tocopherol supplementation in skim milk extender of sapudi ramon on sperm motility, viability, plasma membrane integrity, and cromatin damage. Dissertation, Airlangga University.
- Ullah, Z., H. Khan, S. M. Hussain, M. T. Tunio, S. M. R. Dilshad, A. Gohar, H. Zahid, & A. Ali. 2019. Enhancement of extender excellence of frozen bull semen using α -Tocopherol as an antioxidant. *Acta Sci. Vet.* 47:1641. <https://doi.org/10.22456/1679-9216.90447>
- Wang, W., J. Luo, S. Sun, L. Xi, Q. Gao, A. B. Haile, H. Shi, W. Zhang, & H. Shi. 2015. The effect of season on sperm motility, plasma membrane and acrosome integrity in fresh and frozen-thawed Semen from Xinong Saanen Bucks. *Reprod. Domest. Anim.* 250:23-28. <https://doi.org/10.1111/rda.12444>
- Widjaya, N. 2017. Pengaruh pemberian susu skim dengan pengencer tris kuning telur terhadap daya tahan hidup sperm sapi pada suhu penyimpanan 5°C (Indonesian). *Sains Peternakan: Jurnal Penelitian Ilmu Peternakan.* 9:72-76. <https://doi.org/10.20961/sainspet.9.2.72-76>