


**HETEROCYCLES**  
An  
International  
Journal  
for  
Reviews  
and  
Communications  
in  
Heterocyclic  
Chemistry

Volume 81  
Number 9  
September 1  
2010

WILEY  
ISSN 0361-0316



The Japan Institute of Heterocyclic Chemistry Publication

# HETEROCYCLES

An International Journal for Reviews and Communications in Heterocyclic Chemistry  
Web Edition ISSN: 1881-0942

Published online by The Japan Institute of Heterocyclic Chemistry



[Login / Registration](#)

[Home / Prepress](#) | [e-Journal](#) | [Natural Products Database](#) | [Brush Up](#) | [Authors & Referees](#) | [about Heterocycles](#)

[Login for Authors & Referees](#)  
[Login for Librarians](#)

## about Heterocycles Aims & Scope / Editors

### Aims & Scope

#### Aims & Scope

Since its inception in 1973 HETEROCYCLES has provided a platform for the rapid exchange of research in the areas of organic, pharmaceutical, analytical, and medicinal chemistry of heterocyclic compounds in addition to communications, papers, reviews, a special section of the journal presents newly-discovered natural products whose structure has recently been established. Another section is devoted to the total synthesis of previously documented natural products with heterocyclic ring systems. Due to the fact that the journal is able to publish articles within two months of receipt of the manuscripts, researchers in this field can obtain up-to-date information on heterocyclic research by reading Heterocycles regularly.

#### Audience

Organic and Physical Organic Chemists, Biochemists, Pharmacologists and Scientists studying heterocyclic compounds

#### Indexed / Abstracted

Chemical Abstracts, Chemic Index, Current Contents, PASCAL/CNRS etc.

### Editors

#### Founded by

[Tetsuji Kametani](#)

#### Editor

Kunio Ogasawara

#### Executive Editor

Toshio Honda

#### Honorary Editorial Associates

M. Ihara (Japan)  
[Y. Kanaoka](#) (Japan)  
[Y. Kishi](#) (U.S.A.)  
[J. P. Kutney](#) (Canada)  
[R. Noyori](#) (Japan)  
[A. Suzuki](#) (Japan)  
[E. C. Taylor](#) (U.S.A.)

#### Honorary Advisors to the Editorial Board

M. Banwell (Australia)  
M. A. Ciufolini (Canada)  
J. Cossy (France)  
N. De Kimpe (Belgium)  
T. C. Gallagher (U. K.)  
S. Hashimoto (Japan)  
H. Heimgarther (Switzerland)  
S. H. Kang (Korea)  
Y. Kita (Japan)  
G.-J. Koomen (The Netherlands)  
F. Piozzi (Italy)  
V. H. Rawal (U.S.A.)  
M. D. Rozwadowska (Poland)  
S. Ruchirawat (Thailand)  
M. Shibasaki (Japan)  
M. Shimizu (Japan)  
V. Šimánek (Czech Republic)  
Cs. Szántay (Hungary)  
L. F. Tietze (Germany)  
K. Tomioka (Japan)  
H. N. C. Wong (Hong Kong SAR, P. R. China)  
M. Yamaguchi (Japan)  
L. Zhang (Peoples' Republic of China)

#### Assistant Editors

Masayoshi Tsubuki

#### Editorial Assistants

Sayaka Kametani  
Seiya Kametani  
Saki Kida  
Kaori Tsuji

#### Publisher


Koichi Kametani

#### Published by

The Japan Institute of Heterocyclic Chemistry  
503 1-1-7 Motoakasaka, Minato-ku, Tokyo 107-0051, Japan  
Tel: +81 3 3404 5019  
Fax: +81 3 3497 9370  
E-mail: [editorial@heterocycles.com](mailto:editorial@heterocycles.com)

**1107 Indole Alkaloids from the Leaves of *Alstonia scholaris***


Yusuke Hirasawa, Saori Miyama, Nobuo Kawahara,  
Yukihiro Goda, Abdul Rahman, Wiwied Ekasari,  
Aty Widyawaruyanti, Gunawan Indrayanto,  
Noor Cholies Zaini, and Hiroshi Morita\*



*Alstonia scholaris* Akuammidine-*N*-oxide Akuammidine Antiplasmodial Activity

**1113 Organic Chemistry Using Weakly Electrophilic Salts:  
The Reaction with Nitrogen Nucleophiles**


Hironichi Fujioka,\* Ozora Kubo, Kento Senami,  
Kazuhisa Okamoto, Takashi Okitsu, and  
Yasuyuki Kita\*



*N,O*-Acetal Hydrazone Chemoselective Reaction Weakly Basic Condition Acid-Labile Functional Group

**1121 Artopeden A, a New Antiplasmodial Isoprenylated Flavone  
from *Artocarpus champeden***

Tutik Sri Wahyuni, Wiwied Ekasari, Aty Widyawaruyanti,  
Yusuke Hirasawa, Hiroshi Morita,\* and Noor Cholies Zaini\*



artopeden A

*Artocarpus champeden* Artopeden A Isoprenylated Flavone Antiplasmodial Activity

■ INDEXES

- 1127 Author Index  
1141 Subject Index

HETEROCYCLES, Vol. 79, 2009, pp. 1121 - 1126. © The Japan Institute of Heterocyclic Chemistry  
Received, 17th October, 2008, Accepted, 18th December, 2008, Published online, 19th December, 2008.  
DOI: 10.3987/COM-08-S(D)72

## ARTOPEDEN A, A NEW ANTIPLASMODIAL ISOPRENYLATED FLAVONE FROM *ARTOCARPUS CHAMPEDEN*

Tutik Sri Wahyuni,<sup>a</sup> Wiwied Ekasari,<sup>a</sup> Aty Widyawaruyanti,<sup>a</sup> Yusuke  
Hirasawa,<sup>b</sup> Hiroshi Morita,<sup>b,\*</sup> and Noor Cholies Zaini<sup>a,\*</sup>

<sup>a</sup> Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia

<sup>b</sup> Faculty of Pharmaceutical Sciences, Hoshi University, Shinagawa-ku, Tokyo  
142-8501, Japan

**Abstract** – A new isoprenylated flavone, artopeden A (**1**) was isolated from the barks of *Artocarpus champeden* (Moraceae), and the structure was elucidated by NMR spectral analysis. Artopeden A (**1**) showed a potent antiplasmodial activity with an IC<sub>50</sub> of 0.045 µg/mL against *Plasmodium falciparum* 3D7.

### INTRODUCTION


Malaria, which is caused by parasites of the genus *Plasmodium*, is a most serious infections in many tropical and subtropical regions.<sup>1</sup> Since the number of patients continues to rise mainly due to the emergence of widespread chloroquine-resistant and multiple-drug-resistant strains of malaria parasites, the discovery of new antimalarial drugs is urgently needed.<sup>2</sup>

*Artocarpus champeden* Spreng., belonging to the family Moraceae, is widely distributed throughout the tropical and subtropical regions of Southeast Asia. In Indonesia, this plant is commonly known as ‘‘Cempedak’’ and is traditionally used for treatment of fever, diarrhoea, and malaria.<sup>3</sup> Several isoprenyl flavonoids with cytotoxicity,<sup>4</sup> antimalarial activity,<sup>5</sup> and interaction with IgE and IgM,<sup>6</sup> have been isolated from *A. champeden*.<sup>7</sup>

As a part of our collaborative study, recent investigation of extracts from *Cassia siamea*,<sup>8</sup> *Peganum harmala*,<sup>9</sup> and *Chisocheton ceramicus*<sup>10</sup> resulted in the isolation of some new alkaloids and limonoids with antiplasmodial activity. In this paper, we report the isolation and structure elucidation of a new isoprenylated flavone, artopeden A (**1**) with an antiplasmodial activity from the barks of *Artocarpus champeden*.

---

<sup>†</sup>Dedicated to Dr. John Daly, National Institutes of Health scientist emeritus.



## RESULTS AND DISCUSSION

The dried stem bark of *A. champeden* was extracted successively with hexane,  $\text{CH}_2\text{Cl}_2$ , and MeOH.  $\text{CH}_2\text{Cl}_2$ -soluble materials were subjected to a vacuum silica gel column ( $\text{CHCl}_3$ -EtOAc-MeOH mixture as gradient mobile phase). Fractionation was continued by an open silica gel column chromatography with  $\text{CHCl}_3$ -EtOAc-MeOH mixture as gradient mobile phase. The active fraction was purified by preparative thin layer chromatography ( $\text{CH}_3\text{CN}$ -MeOH- $\text{H}_2\text{O}$ , 4:1:1 as mobile phase) followed by preparative thin layer chromatography ( $\text{CHCl}_3$ -MeOH, 19:1 as mobile phase) and an ODS column (MeOH/ $\text{H}_2\text{O}$  7:3  $\rightarrow$  1:0) to afford artopeden A (**1**, 0.000016%) together with artoindonesianin S (**2**, 0.00004%),<sup>11</sup> artoindonesianin E (12.0 mg: 0.00048%),<sup>5b</sup> artonin A (12.0 mg: 0.00048%),<sup>12</sup> and cycloheterophyllin (19.3 mg: 0.00077%).<sup>13</sup> MeOH-soluble materials were subjected to a silica gel column ( $\text{CHCl}_3$ -MeOH mixtures as mobile phase), followed by an ODS column (MeOH/ $\text{H}_2\text{O}$  3:2  $\rightarrow$  4:1) to give morachalcone A<sup>14</sup> (**3**, 0.00002%).

Artopeden A (**1**,  $[\alpha]_{\text{D}}^0$ ) showed the pseudomolecular ion peak at  $m/z$  397 ( $\text{M}+\text{H}$ )<sup>+</sup> in ESIMS, and the molecular formula,  $\text{C}_{22}\text{H}_{20}\text{O}_7$ , was established by HRTOFMS [ $m/z$  397.1268, ( $\text{M}+\text{H}$ )<sup>+</sup>  $\Delta$  -1.9 mDa]. IR spectrum suggested the presence of OH ( $3450\text{ cm}^{-1}$ ) and conjugated carbonyl ( $1640\text{ cm}^{-1}$ ) groups. The  $^{13}\text{C}$  NMR (Table 1) spectrum of **1** disclosed twenty-two carbon signals due to one conjugated carbonyl ( $\delta_{\text{C}}$  179.7), eleven  $sp^2$  quaternary carbons ( $\delta_{\text{C}}$  104.4, 104.5, 112.0, 133.4, 135.3, 148.3, 150.6, 156.2, 160.2, 161.3, and 164.6), one  $sp^3$  quaternary carbon ( $\delta_{\text{C}}$  93.7), three  $sp^2$  methines ( $\delta_{\text{C}}$  93.1, 98.8, and 102.8), one  $sp^3$  methine ( $\delta_{\text{C}}$  47.2), one  $sp^3$  methylene ( $\delta_{\text{C}}$  20.4), two methyls ( $\delta_{\text{C}}$  22.8 and 28.3), and two methoxy groups ( $\delta_{\text{C}}$  56.3 and 56.6).  $^1\text{H}$  and  $^{13}\text{C}$  signals for **1** were assigned by detailed analysis of the 2D NMR spectra.

The  $^1\text{H}$  NMR spectrum of **1** showed the signals for the following protons: three aromatic protons [ $\delta_{\text{H}}$  6.33 (1H, d, 2.2), 6.72 (1H, d, 2.2), and 6.45 (1H, s)], two methyl protons [ $\delta_{\text{H}}$  1.36 (3H, s) and 1.68 (3H, s)], ABX type protons [ $\delta_{\text{H}}$  2.40 (1H, dd, 15.2, 15.2), 3.24 (1H, dd, 15.2, 7.1), and 3.44 (1H, dd, 15.2, 7.1)], and two methoxy protons [ $\delta_{\text{H}}$  3.89 (3H, s) and 3.92 (3H, s)]. The chemical shifts and coupling patterns of **1** were similar to those of artonin L,<sup>15</sup> which has already been isolated from *Artocarpus heterophyllus*.<sup>13</sup> The chemical shifts of all the carbons were similar to those of the relevant carbons of artonin L with the exception of the shifts of the four carbons (C-2', 3', 4', and 5') (Table 1). In the case

of artonin L, the presence of methoxy group at C-2' caused higher field shift ( $\Delta +0.15$  ppm) of the proton at the C-8 position when compared to that of hydroxy group (artoinin L: 6.51 ppm at C-8; artopeden A: 6.72 ppm at C-8).<sup>15</sup> Thus, the gross structure of artopeden A was assigned as **1** with a methoxy group at C-4', which was an isomer of artonin L. Since artopeden A (**1**) did not show the optical rotation, **1** might be biogenetically derived from artoindonesianin S (**2**) through the oxidative coupling, which was the same case as in artonins A and B.<sup>12</sup>

Malaria is the most prevalent parasitic disease. According to World Health Organization (WHO), as many as 300–500 million people suffer from malaria. *Artocarpus champeden* has been traditionally used for treatment of malaria,<sup>3</sup> and some isoprenylated flavonoids and stilbenes from *Artocarpus* species have already been reported on inhibitory activity against some parasites.<sup>5</sup> Artopeden A (**1**) showed a potent *in vitro* antiplasmodial activity against *Plasmodium falciparum* 3D7 (artopeden A: IC<sub>50</sub> 0.045  $\mu\text{g/ml}$ ; chloroquine: IC<sub>50</sub> 0.0061  $\mu\text{g/mL}$ ).

Table 1. <sup>1</sup>H and <sup>13</sup>C NMR Data [ $\delta_{\text{H}}$  (J, Hz) and  $\delta_{\text{C}}$ ] of artopeden A (**1**) in acetone-*d*<sub>6</sub> at 300K

Position	$\delta_{\text{H}}$	$\delta_{\text{C}}$
2		161.3
3		112.0
4		179.7
4a		104.5
5		160.2
6	6.33 (1H, d, 2.2)	98.8
7		164.6
8	6.72 (1H, d, 2.2)	93.1
8a		156.2
1'		104.4
2'		150.6
3'	6.45 (1H, s)	102.8
4'		148.3
5'		135.3
6'		133.4
9a	2.40 (1H, dd, 15.2, 15.2)	20.4
9b	3.24 (1H, dd, 15.2, 7.1)	
10	3.44 (1H, dd, 15.2, 7.1)	47.2
11		93.7
12	1.68 (3H, s)	28.3
13	1.36 (3H, s)	22.8
14	3.89 (3H, s)	56.3
15	3.92 (3H, s)	56.6

## EXPERIMENTAL

**General Experimental Procedures.**  $^1\text{H}$  and 2D NMR spectra were recorded on a Bruker AV 400 spectrometer and chemical shifts were reported using residual acetone- $d_6$  ( $\delta_{\text{H}}$  2.06 and  $\delta_{\text{C}}$  29.8) as internal standards. Standard pulse sequences were employed for the 2D NMR experiments. Mass spectra were obtained with a Micromass LCT spectrometer.

**Plant Material.** Dried stem bark of *A. champeden* was collected from "Mak Balim" village, Sorong, Irian Jaya, Indonesia, in August 2004. A voucher specimen was identified by Dr. Eko Baroto Walujo at the Herbarium Bogoriense, LIPI - Research center For Biology, Bogor, Indonesia, and it was deposited at the herbarium of Bogor Botanical Garden.

**Extraction and Isolation.** Dried powder of *A. champeden* stem bark (2.5 kg) was macerated successively in hexane,  $\text{CH}_2\text{Cl}_2$ , and MeOH. The  $\text{CH}_2\text{Cl}_2$  extract (32.0 g) was fractionated by a vacuum silica gel column chromatography with  $\text{CHCl}_3$ -EtOAc-MeOH mixtures as mobile phases. Fractionation was continued by an open silica gel column chromatography with  $\text{CHCl}_3$ -EtOAc-MeOH mixture as gradient mobile phase. The active fraction was purified by preparative thin layer chromatography ( $\text{CH}_3\text{CN}$ -MeOH- $\text{H}_2\text{O}$ , 4:1:1 as mobile phase) followed by preparative thin layer chromatography ( $\text{CHCl}_3$ -MeOH, 19:1 as mobile phase) and an ODS column (MeOH/ $\text{H}_2\text{O}$  7:3  $\rightarrow$  1:0) to afford artopeden A (**1**, 0.4 mg: 0.000016%) together with artoindonesianin S (**2**, 1.0 mg: 0.00004%), artoindonesianin E (12.0 mg: 0.00048%),<sup>5b</sup> artonin A (12.0 mg: 0.00048%),<sup>12</sup> and cycloheterophyllin (19.3 mg: 0.00077%).<sup>13</sup> The MeOH fraction was further chromatographed on silica gel column chromatography, with  $\text{CHCl}_3$ -MeOH mixtures as mobile phases, followed by repeated purification by reversed-phase column chromatography with MeOH- $\text{H}_2\text{O}$  mixture as mobile phases to yield morachalcone A<sup>14</sup> (**3**, 0.5 mg: 0.00002%).

**Artopeden A (1):** yellow powder;  $[\alpha]_{\text{D}}^{23}$  0 ( $c$  1.0, MeOH); IR (KBr)  $\nu_{\text{max}}$  3450, 2940, and 1640  $\text{cm}^{-1}$ ; UV (MeOH)  $\lambda_{\text{max}}$  370 ( $\log \epsilon$  4.3) and 205 ( $\log \epsilon$  4.3) nm;  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (Table 1); ESIMS  $m/z$  397 ( $\text{M}+\text{H}$ )<sup>+</sup>; HRTOFMS  $m/z$  397.1268 [ $\text{M}+\text{H}$ ]<sup>+</sup>, calcd for  $\text{C}_{22}\text{H}_{21}\text{O}_7$ , 397.1287].

**Antiplasmodial Assay.** Human malaria parasites were cultured according to the method by W. Trager *et al.*<sup>16</sup> The antimalarial activity of the isolated compounds was determined by the procedure described by Budimulja *et al.*<sup>17</sup> In brief, Stock solution of the samples were prepared in DMSO (final DMSO concentrations of < 0.5%) and were diluted to the required concentration with complete medium (RPMI 1640 supplemented with 10% human plasma, 25 mM HEPES and 25 mM  $\text{NaHCO}_3$ ) until the final concentration of samples at well culture plate were : 10; 1; 0.1; 0.01; 0.001  $\mu\text{g}/\text{mL}$ . The malarial parasite *P. falciparum* 3D7 clone was propagated in a 24-well culture plate in the presence of a wide range of concentrations of each compound. The growth of the parasite was monitored by making a blood smear fixed with MeOH and stained with Geimsa solution. The antimalarial activity of each

compound was expressed as an IC<sub>50</sub> value, defined as the concentration of the compound causing 50% inhibition of parasite growth relative to an untreated control.

The percentage of growth inhibition was expressed according to following equation : Growth inhibition % = 100 - [(test parasitaemia/control parasitemia) × 100. Chloroquine: IC<sub>50</sub>= 0.0061 µg/mL.

## ACKNOWLEDGMENTS

We acknowledge the financial support provided by Assessment Service Unit, Faculty of Pharmacy, Airlangga University. This work was also supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and The Open Research Center Project.

## REFERENCES

1. J. Wiesner, R. Ortmann, H. Jomaa, and M. Schlitzer, *Angew. Chem. Int. Ed.*, 2003, **42**, 5274.
2. (a) M. H. Gelb and W. G. Hol, *Science*, 2002, **297**, 343; (b) P. A. G. N. De Smet, *Drugs*, 1997, **54**, 801.
3. S. A. Achmad, *Bull. Soc. Nat. Prod. Chem. (Indonesia)*, 2004, **4**, 35.
4. E. H. Hakim, A. Fahriyati, M. S. Kau, S. A. Achmad, L. Makmur, E. L. Ghisalberti, and T. Nomura, *J. Nat. Prod.*, 1999, **62**, 613.
5. (a) C. Boonlaksiri, W. Oonant, P. Kongsaree, P. Kittakoop, M. Tanticharoen, and Y. Thebtharanonth, *Phytochemistry*, 2000, **54**, 415; (b) A. Widyawaruyanti, Subehan, S. K. Kalauni, S. Awale, M. Nindatu, N. C. Zaini, D. Syafruddin, P. B. S. Asih, Y. Tezuka, and S. Kadota, *J. Nat. Med.*, 2007, **61**, 410.
6. S. A. Achmad, E. H. Hakim, L. D. Juliawaty, L. Makmur, Suyatno, N. Aimi, and E. L. Ghisalberti, *J. Nat. Prod.*, 1996, **59**, 878.
7. Y. M. Syah, L. D. Juliawaty, S. A. Achmad, E. H. Hakim, and E. L. Ghisalberti, *J. Nat. Med.*, 2007, **60**, 308.
8. (a) H. Morita, S. Oshimi, Y. Hirasawa, K. Koyama, T. Honda, W. Ekasari, G. Indrayanto, and N. C. Zaini, *Org. Lett.*, 2007, **9**, 3691; (b) S. Oshimi, Y. Tomizawa, Y. Hirasawa, T. Honda, W. Ekasari, A. Widyawaruyanti, M. Rudyanto, G. Indrayanto, N. C. Zaini, and H. Morita, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 3761.
9. A. Astulla, K. Zaima, Y. Matsuno, Y. Hirasawa, W. Ekasari, A. Widyawaruyanti, N. C. Zaini, and H. Morita, *J. Nat. Med.*, 2008, **62**, 470.
10. K. Mohamad, Y. Hirasawa, M. Litaudon, K. Awang, A. H. A. Hadi, K. Takeya, W. Ekasari, A. Widyawaruyanti, N. C. Zaini, and H. Morita, *Bioorg. Med. Chem.*, in press.



11. Y. M. Syah, S. A. Achmad, E. L. Ghisalberti, E. H. Hakim, L. Makmur, and D. Mujahidin, *Phytochemistry*, 2002, **61**, 949.
12. Y. Hano, M. Aida, M. Shiina, T. Nomura, T. Kawai, H. Ohe, and K. Kagei, *Heterocycles*, 1989, **29**, 1447.
13. B. L. Wei, J. R. Weng, P. H. Chiu, C. F. Hung, J. P. Wang, and C. N. Lin, *J. Agric. Food Chem.*, 2005, **53**, 3867.
14. Y. Hano, S. Suzuki, T. Nomura, and S. Ueda, *Heterocycles*, 1989, **29**, 807.
15. M. Aida, K. Shinomiya, Y. Hano, and T. Nomura, *Heterocycles*, 1993, **36**, 575.
16. W. Trager and J. B. Jensen, *Science*, 1976, **193**, 673.
17. A. S. Budimulja, Syafruddin, P. Tapchaisri, P. Wilairat, and S. Marzuki, *Mol. Biochem. Parasitol.*, 1997, **84**, 137.

**Heterocycles**

**Q4** Analytical Chemistry  
best quartile

**SJR 2021**  
0.19

powered by scimagojr.com

also developed by scimago: **SCIMAGO INSTITUTIONS RANKINGS**

**SJR** Scimago Journal & Country Rank

Home Journal Rankings Country Rankings Viz Tools Help About Us

Enter Journal Title, ISSN or Publisher Name

## Heterocycles

<b>COUNTRY</b> Japan Universities and research institutions in Japan	<b>SUBJECT AREA AND CATEGORY</b> Chemistry Analytical Chemistry Organic Chemistry Pharmacology, Toxicology and Pharmaceutics Pharmacology	<b>PUBLISHER</b> Japan Institute of Heterocyclic Chemistry	<b>H-INDEX</b> <b>66</b>
<b>PUBLICATION TYPE</b> Journals	<b>ISSN</b> 03855414	<b>COVERAGE</b> 1983-2021	<b>INFORMATION</b> Homepage How to publish in this journal editorial@heterocycles.com

Scopus Preview

Author Search Sources Create account Sign in

## Source details

Feedback > Compare sources >

### Heterocycles

Scopus coverage years: from 1983 to Present  
 Publisher: Elsevier  
 ISSN: 0385-5414  
 Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmacology Chemistry: Analytical Chemistry Chemistry: Organic Chemistry  
 Source type: Journal

View all documents > Set document alert Save to source list Source Homepage

CiteScore 2021 **1.2**  
 SJR 2021 **0.189**  
 SNIP 2021 **0.283**

CiteScore CiteScore rank & trend Scopus content coverage

**i** Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

**CiteScore 2021**

$$1.2 = \frac{979 \text{ Citations } 2018 - 2021}{789 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

**CiteScoreTracker 2022**

$$1.0 = \frac{644 \text{ Citations to date}}{629 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly

**CiteScore rank 2021**

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics Pharmacology	#229/303	24th
Chemistry Analytical Chemistry	#99/130	24th
Chemistry Organic Chemistry	#162/192	15th