

# Ceramicines B-D, new antiplasmodial limonoids from Chisocheton ceramicus

*by* Wiwied Ekasari

---

**Submission date:** 17-May-2021 12:32PM (UTC+0800)

**Submission ID:** 1587645421

**File name:** C-14 - Bioorg Med Chem\_naskah.pdf (349.1K)

**Word count:** 3776

**Character count:** 15652



## Ceramicines B–D, new antiplasmodial limonoids from *Chisocheton ceramicus*

Khalit Mohamad<sup>a,b</sup>, Yusuke Hirasawa<sup>a</sup>, Marc Litaudon<sup>c</sup>, Khalijah Awang<sup>d</sup>, A. Hamid A. Hadi<sup>d</sup>, Koichi Takeya<sup>e</sup>, Wiwied Ekasari<sup>f</sup>, Aty Widayawaruyanti<sup>f</sup>, Noor Cholies Zaini<sup>f</sup>, Hiroshi Morita<sup>a,\*</sup>

<sup>a</sup> Faculty of Pharmaceutical Sciences, Hoshi University, Shinagawa, Tokyo 142-8501, Japan

<sup>b</sup> Department of Pharmacy, Faculty of Medicine, University Malaya, 50603 Kuala Lumpur, Malaysia

<sup>c</sup> Institut de chimie des Substances Naturelles, CNRS, 91198 Gif-sur-Yvette, France

<sup>d</sup> Department of Chemistry, Faculty of Science, University Malaya, 50603 Kuala Lumpur, Malaysia

<sup>e</sup> Tokyo University of Pharmaceutical and Life Science, 1432-1 Horinouchi, Hachioji, Tokyo 192-0392, Japan

<sup>f</sup> Faculty of Pharmacy, Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia

### ARTICLE INFO

#### Article history:

Received 10 September 2008

Revised 15 November 2008

Accepted 18 November 2008

Available online 24 November 2008

#### Keywords:

Limonoids

*Chisocheton ceramicus*

Ceramicines B–D

Antiplasmodial activity

### ABSTRACT


Three new limonoids, ceramicines B–D (**1–3**), have been isolated from the bark of *Chisocheton ceramicus*. Structures and stereochemistry of **1–3** were fully elucidated and characterized by 2D NMR analysis. Ceramicines exhibited a moderate antiplasmodial activity.

© 2008 Elsevier Ltd. All rights reserved.

## 1. Introduction

Limonoids, highly oxidative unique secondary metabolites obtained from Meliaceae are produced by a unique biosynthetic route through tetranortriterpenoid nucleus.<sup>1,2</sup> Insecticidal, insect anti-feedant, antibacterial, antifungal, antimalarial, anticancer, and antiviral activities have been reported for many limonoids.<sup>3</sup> Especially, malaria caused by parasites of the genus *Plasmodium* is one of the leading infectious diseases in many tropical and some temperate regions.<sup>4</sup> The emergence of widespread chloroquine-resistant and multiple-drug-resistant strains of malaria parasites leads to the need for the development of new therapeutic agents against malaria.<sup>5</sup>

Recently, we have isolated cassiarin A with an unprecedented tricyclic skeleton and a potent antiplasmodial activity from the leaves of *Cassia siamea*.<sup>6</sup> Previous investigations on limonoids from Meliaceae have led to the isolation of several unique tetranortriterpenoids.<sup>7</sup> In continuation of our antiplasmodial research on *Chisocheton ceramicus* belonging to Meliaceae family, we have isolated three new limonoids, ceramicines B–D (**1–3**) together with ceramicine A (**4**),<sup>7b</sup> which showed a moderate antiplasmodial activity. Herein, we report the structure elucidation and antiplasmodial activity of ceramicines B–D (**1–3**) from *C. ceramicus*.



### 1.1. Structures of ceramicines B–D (**1–3**)

The crude ethanol extract of bark was partitioned between  $\text{CHCl}_3$ , *n*-BuOH, and water. Chromatographic purification of the chloroform soluble fraction by a silica gel column (hexane/EtOAc and toluene/EtOAc solvent system) led to isolation of three new limonoids, ceramicines B (**1**, 147.4 mg, 0.074% yield), C (**2**,

\* Corresponding author. Tel./fax: +81 3 5498 5778.

E-mail address: moritah@hoshi.ac.jp (H. Morita).

44.4 mg, 0.022% yield), and D (**3**, 15.0 mg, 0.0075% yield), together with ceramidine A.<sup>7b</sup>

Ceramidine B (**1**,  $[\alpha]_D^{20} +30$  (c 1.0, CHCl<sub>3</sub>)) was obtained as a colorless solid and was revealed to have the molecular formula C<sub>26</sub>H<sub>32</sub>O<sub>4</sub>, by HRESITOFMS  $[m/z 409.2370$  (M+H)<sup>+</sup>,  $\Delta -0.9$  mmu]. IR absorptions implied the presence of an  $\alpha,\beta$ -unsaturated ketone (1669 cm<sup>-1</sup>) and a hydroxyl (3535 cm<sup>-1</sup>) groups. UV spectrum (219 nm) indicated the presence of an unsaturated carbonyl group. <sup>1</sup>H and <sup>13</sup>C NMR data (Tables 1 and 2, respectively) revealed 26 carbon resonances due to one carbonyl, two sp<sup>2</sup> quaternary carbons, four sp<sup>3</sup> quaternary carbons, six sp<sup>2</sup> methines, five sp<sup>3</sup> methines, four sp<sup>3</sup> methylenes, and four methyls. Among them, three sp<sup>3</sup> carbons ( $\delta_C$  72.5, 73.8, and 79.8) and two sp<sup>2</sup> methines ( $\delta_C$  139.7 and 142.6) were ascribed to those bearing an oxygen atom.

Five partial structures **a** (from C-2 to C-3), **b** (from C-5 to C-7), **c** (from C-9, C-11 to C-12), **d** (from C-15 to C-17), and **e** (from C-22 to C-23) were deduced from <sup>1</sup>H–<sup>1</sup>H COSY analysis of **1** in CDCl<sub>3</sub> (Fig. 1). The presence of an enone group in ring A was supported by HMBC correlations as shown in Figure 1. HMBC correlations for H-3 and H<sub>3</sub>-19 of C-1 ( $\delta_C$  202.9), for H<sub>3</sub>-29 of C-5 ( $\delta_C$  47.7), and for H-7, H<sub>3</sub>-19, and H<sub>3</sub>-30 of C-9 ( $\delta_C$  35.9) gave rise to the connectivity of partial structures **a**, **b**, and **c** through C-4, C-8, and C-10 atoms. Connection between partial structures **c** and **d** could be assigned by HMBC correlations for H<sub>3</sub>-18 of C-12 ( $\delta_C$  33.1), C-13 ( $\delta_C$  47.0), C-14 ( $\delta_C$  159.8), and C-17 ( $\delta_C$  51.9). The presence of a  $\beta$ -furyl ring at C-17 was also assigned by the HMBC correlations as shown in Figure 1. In addition, HMBC correlations for H-7 and H<sub>2</sub>-28 of C-6 ( $\delta_C$  73.8) indicated the presence of a tetrahydrofuran ring at C-4–C-6 and C-28. Thus, ceramidine B (**1**) was concluded to be a new limonoid possessing cyclopenta[*a*]phenanthren ring system with a  $\beta$ -furyl ring at C-17 and a tetrahydrofuran ring.


The relative stereochemistry of **1** was elucidated by ROESY correlations as shown in computer-generated 3D drawing (Fig. 2). ROESY correlations of H-6/H<sub>2</sub>-28, H<sub>3</sub>-19, and H<sub>3</sub>-30, H-7/H-15, and H-12/H-17 together with the <sup>3</sup>J proton coupling constants (<sup>3</sup>J<sub>H-5/H-6</sub> = 12.4 Hz and <sup>3</sup>J<sub>H-6/H-7</sub> = 3.8 Hz) suggested that each of H-6, H-7, and H-17 adopts a  $\beta$ -configuration. Furthermore, the  $\alpha$  configurations of H-5 and H-9 was indicated by ROESY correlation of H-5/H-9.

**Table 1**  
<sup>1</sup>H NMR data [ $\delta_H$  (J, Hz)] of ceramidines B–D (**1–3**) in CDCl<sub>3</sub> at 300 K


	<b>1</b>	<b>2</b>	<b>3</b>
2	5.83 (d, 9.6 Hz)	5.82 (d, 9.6 Hz)	5.73 (d, 9.6 Hz)
3	6.95 (d, 9.6 Hz)	6.95 (d, 9.6 Hz)	6.91 (d, 9.6 Hz)
5	2.73 (d, 12.4 Hz)	2.69 (d, 12.4 Hz)	2.59 (d, 12.4 Hz)
6	4.28 (dd, 12.4, 3.8 Hz)	4.28 (dd, 12.4, 3.8 Hz)	4.19 (dd, 12.4, 3.8 Hz)
7	4.23 (d, 3.8 Hz)	4.25 (d, 3.8 Hz)	4.10 (d, 3.8 Hz)
9	2.40 m	2.61 m	2.26 m
11a	2.52 m	2.48 m	2.40 m
11b	1.80 m	2.29 m	1.69 m
12a	1.89 m	5.13 m	1.81 m
12b	1.60 m		1.60 m
15	5.59 br s	5.69 m	5.47 br s
16a	2.55 m	2.57 m	2.53 m
16b	2.40 m	2.43 m	2.25 m
17	2.85 m	3.04 m	2.76 m
18	0.82 s	1.08 s	0.79 s
19	1.17 s	1.16 s	1.07 s
21	7.25 s	7.14 s	
22	6.29 (d, 1.6 Hz)	6.23 (d, 1.6 Hz)	7.21 (dd, 1.8, 1.8 Hz)
23	7.37 (d, 1.6 Hz)	7.28 (d, 1.6 Hz)	4.77 (br s)
28a	3.63 (d, 7.3 Hz)	3.63 (d, 7.3 Hz)	3.54 (d, 7.3 Hz)
28b	3.79 (d, 7.3 Hz)	3.79 (d, 7.3 Hz)	3.69 (d, 7.3 Hz)
29	1.34 s	1.33 s	1.25 s
30	1.13 s	1.13 s	1.03 s
3'		1.76 s	
4'		5.86, 5.47 (s)	

**Table 2**  
<sup>13</sup>C NMR data ( $\delta_C$ ) of ceramidines B–D (**1–3**) in CDCl<sub>3</sub> at 300 K

	<b>1</b>	<b>2</b>	<b>3</b>
1	202.9	202.4	203.3
2	130.2	130.0	129.8
3	151.0	151.9	151.2
4	41.9	41.8	41.7
5	47.7	47.6	47.4
6	73.8	73.7	73.7
7	72.5	72.0	72.1
8	47.2	46.5	46.9
9	35.9	36.5	35.3
10	47.2	47.1	47.1
11	17.7	26.8	17.4
12	33.1	77.8	32.7
13	47.0	51.6	47.1
14	159.8	156.5	158.9
15	120.4	122.7	120.0
16	34.3	36.5	33.8
17	51.9	50.6	50.6
18	21.5	16.0	21.3
19	14.3	14.1	14.0
20	124.5	124.3	133.5
21	139.7	140.3	174.8
22	111.0	111.7	147.1
23	142.6	141.2	70.3
28	79.8	79.8	79.5
29	20.1	20.1	19.8
30	26.0	26.5	25.9
1'		167.1	
2'		136.4	
3'		18.0	
4'		125.9	



**Figure 1.** Selected 2D NMR correlations for ceramidine B (**1**).



**Figure 2.** Selected ROESY correlations for ceramidine B (**1**).

HRESITOFMS data  $[m/z 515.2397$  (M+Na)<sup>+</sup>,  $\Delta -1.3$  mmu] of ceramidine C (**2**,  $[\alpha]_D^{20} +47$  (c 1.0, CHCl<sub>3</sub>)) established the molecular




Figure 3. Selected 2D NMR correlations for ceramicine C (2).

formula to be  $C_{30}H_{36}O_6$ , which was larger than that of ceramicine B (1) by a  $C_4H_4O_2$  unit.  $^1H$  and  $^{13}C$  NMR data (Tables 1 and 2) of 2 were analogous to those of 1, although H-12 ( $\delta_H$  5.13) and C-12 ( $\delta_C$  77.8) bearing an oxygen atom were observed for 2. The presence of methacrylic acid ( $\delta_H$  1.76, 5.47, and 5.86;  $\delta_C$  18.0, 125.9, 136.4, and 167.1) at C-12 was confirmed by HMBC correlations for H<sub>3</sub>-3' and H-12 of C-1' ( $\delta_C$  167.1) and H<sub>2</sub>-4' of C-1' and C-3' ( $\delta_C$  18.0).

The gross structure of 2 was elucidated by 2D NMR ( $^1H$ - $^1H$  COSY, HMQC, and HMBC) data shown in Figure 3. Analysis of ROESY spectrum (Fig. 4), suggested that the relative stereochemistry of H-12 and H-17 to be  $\beta$  through correlations among H-12, H-17, and H<sub>3</sub>-30.

Ceramicine D (3), colorless amorphous solid,  $[\alpha]_D^{20} +38$  (c 1.0,  $CHCl_3$ ), was shown to have the molecular formula of  $C_{26}H_{32}O_5$  by HRESITOFMS [ $m/z$  425.2306, (M+H)<sup>+</sup>,  $\Delta$  -2.2 mmu], which was lar-




Figure 4. Selected ROESY correlations for ceramicine C (2).




Figure 5. Selected 2D NMR correlations for ceramicine D (3).




Figure 6. Selected ROESY correlations for ceramicine D (3).

ger than that of ceramicine B by 16 mmu.  $^1H$  and  $^{13}C$  NMR data of 3 were analogous to those of ceramicine B, although the  $\beta$ -furyl ring signals for ceramicine B were lacking for 3. The presence of an  $\alpha,\beta$ -unsaturated- $\gamma$ -lactone ring instead of the  $\beta$ -furyl ring at C-17 was deduced by the  $^1H$  [ $\delta_H$  4.77 (br s, H-23) and 7.21 (dd,  $J$  = 1.8, 1.8 Hz, H-22)] and  $^{13}C$  [ $\delta_C$  133.5 (C-20), 174.8 (C-21), 147.1 (C-22), and 70.3 (C-23)] signals.<sup>18</sup> HMBC correlations for H-17 of C-20, C-21, and C-22 indicated the presence of  $\beta$ -substituted- $\gamma$ -lactone ring (Fig. 5). The gross structure of 3 was elucidated by 2D NMR ( $^1H$ - $^1H$  COSY, HMQC, and HMBC) data, and the relative stereochemistry of 3 was assigned as the same as that of ceramicine B by ROESY correlations shown in Figure 6.

## 1.2. Antiplasmodial activity

Malaria caused by parasites of the genus *Plasmodium* is one of the leading infectious diseases in many tropical and some temperate regions.<sup>4</sup> The emergence of widespread chloroquine-resistant and multiple-drug-resistant strains of malaria parasites leads to the need for the development of new therapeutic agents against malaria.<sup>5</sup> Ceramicine B (1) showed a potent in vitro antiplasmodial activity against *Plasmodium falciparum* 3D7 (IC<sub>50</sub> 1: 0.23  $\mu$ g/ml),<sup>9</sup> whereas ceramicines C (2) and D (3) did a moderate activity and ceramicine A (4) did a weak activity (IC<sub>50</sub> 2, 2.38  $\mu$ g/ml; 3, 2.15  $\mu$ g/ml; 4, 44.22  $\mu$ g/ml). Ceramicines B–D (1–3) showed a weak cytotoxicity on P388 cells (IC<sub>50</sub> 15  $\mu$ g/ml for 1; 5.5  $\mu$ g/ml for 2; 27  $\mu$ g/ml for 3). These compounds belong to two groups with a tetrahydrofuran ring at C-4–C-6 and C-28 and without this function. In comparison of antiplasmodial activity among these compounds, ceramicines B (1)–D (3) with a tetrahydrofuran ring showed a potent activity, whereas ceramicine A (4) without this function exhibited a relatively weak activity.

## 2. Experimental

### 2.1. General methods

$^1H$  and 2D NMR spectra were recorded on a 400 MHz spectrometers at 300K, while  $^{13}C$  NMR spectra were measured on a 100 MHz spectrometer. Each NMR sample of ceramicines B–D (1–3) were prepared by dissolving 3.0 mg in 250  $\mu$ l of  $CDCl_3$  in 2.5 mm microcells (Shigemi Co., Ltd) and chemical shifts were reported using residual  $CDCl_3$  ( $\delta_H$  7.21 and  $\delta_C$  77.0) as an internal standard. Standard pulse sequences were employed for the 2D NMR experiments.  $^1H$ - $^1H$  COSY, HOHAHA, and ROESY spectra were measured with spectral widths of both dimensions of 4800 Hz, and 32 scans with two dummy scans were accumulated into 1 K data points for each of 256  $t_1$  increments. ROESY and HOHAHA spectra in the phase sensitive mode were measured with a mixing time of 800 and

30 ms, respectively. For HMQC spectra in the phase sensitive mode and HMBC spectra, a total of 256 increments of 1 K data points were collected. For HMBC spectra with Z-axis PFG, a 50 ms delay time was used for long-range C–H coupling. Zero-filling to 1K for  $F_1$  and multiplication with squared cosine-bell windows shifted in both dimensions were performed prior to 2D Fourier transformation.

## 2.2. Material

The barks of *C. ceramicus* were collected at Pahang, Malaysia in 1996. The botanical identification was made by Mr. Teo Leong Eng, Faculty of Science, University of Malaya. Voucher specimens are deposited in the Herbarium of Chemistry Department, University of Malaya.

## 2.3. Extraction and isolation

The dried ground barks of *C. ceramicus* (200 g) were extracted successively with ethanol and the extract (10.5 g) was partitioned with 10% aq MeOH and  $\text{CHCl}_3$ . The  $\text{CHCl}_3$ -soluble materials were subjected to a silica gel column (hexane/EtOAc, 1:0 → 0:1), in which a fraction eluted with hexane/EtOAc (7:3) was further purified on a silica gel column with toluene/EtOAc (4:1) to afford ceramicine B (**1**, 147.4 mg, 0.074% yield) as colorless solids. The fraction eluted with hexane/EtOAc (3:2) was purified on a silica gel column with toluene/EtOAc (4:1) to obtain ceramicine C (**2**, 44.4 mg, 0.022% yield). The fraction eluted with hexane/EtOAc (2:3) was purified on a silica gel column with toluene/EtOAc (3:2) to give ceramicine D (**3**, 15 mg, 0.0075% yield).

### 2.3.1. Ceramicine B (1)

Colorless solid;  $[\alpha]_D^{20} +30$  (c 1.0,  $\text{CHCl}_3$ ); IR (KBr)  $\nu_{\text{max}}$  3535, 3457, 2969, 2927, 2862, 1720, 1669, 1457, 1387, and 1247  $\text{cm}^{-1}$ ; UV (MeOH)  $\lambda_{\text{max}}$  219 ( $\epsilon$  9300) nm; CD (MeOH)  $\lambda_{\text{max}}$  340 ( $\theta$  –3300) and 220 nm ( $\theta$  +30,800);  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (Tables 1 and 2); HRESITOFMS  $m/z$  409.2370 (M+H; calcd for  $\text{C}_{26}\text{H}_{32}\text{O}_4$ , 409.2379).

### 2.3.2. Ceramicine C (2)

Colorless solid;  $[\alpha]_D^{20} +47$  (c 1.0,  $\text{CHCl}_3$ ); IR (KBr)  $\nu_{\text{max}}$  3448, 2932, 1711, 1677, 1451, 1389, 1248, and 1160  $\text{cm}^{-1}$ ; UV (MeOH)  $\lambda_{\text{max}}$  219 ( $\epsilon$  9400) nm; CD (MeOH)  $\lambda_{\text{max}}$  334 ( $\theta$  –2700) and 220 nm ( $\theta$  +29600);  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (Tables 1 and 2); HRESITOFMS  $m/z$  515.2397 (M+Na; calcd for  $\text{C}_{30}\text{H}_{36}\text{O}_6\text{Na}$ , 515.2410).

### 2.3.3. Ceramicine D (3)

Colorless solid;  $[\alpha]_D^{20} +38$  (c 1.0,  $\text{CHCl}_3$ ); IR (KBr)  $\nu_{\text{max}}$  3435, 2929, 1750, 1677, 1459, 1388, and 1249  $\text{cm}^{-1}$ ; UV (MeOH)  $\lambda_{\text{max}}$  222 ( $\epsilon$  10,000) nm; CD (MeOH)  $\lambda_{\text{max}}$  338 ( $\theta$  –2100) and 223 nm ( $\theta$  +17,900);  $^1\text{H}$  and  $^{13}\text{C}$  NMR data (Tables 1 and 2); HRESITOFMS  $m/z$  425.2306 (M+H; calcd for  $\text{C}_{26}\text{H}_{32}\text{O}_5$ , 425.2328).

## 2.4. Antiplasmodial activity

Human malaria parasites were cultured according to the method by Trager et al.<sup>9</sup> The antimalarial activity of the isolated compounds was determined by the procedure described by Budimulja et al.<sup>10</sup> In brief, Stock solution of the samples were prepared in DMSO (final DMSO concentrations of <0.5%) and were diluted to the required concentration with complete medium (RPMI-1640 supplemented with 10% human plasma, 25 mM Hepes and 25 mM  $\text{NaHCO}_3$ ) until the final concentration of samples at well culture plate are: 10, 1, 0.1, 0.01, 0.001  $\mu\text{g}/\text{ml}$ . The malarial parasite

*P. falciparum* 3D7 clone was propagated in a 24-well culture plate in the presence of a wide range of concentrations of each compound. The growth of the parasite was monitored by making a blood smear fixed with MeOH and stained with Geimsa stain. The antimalarial activity of each compound was expressed as an  $\text{IC}_{50}$  value, defined as the concentration of the compound causing 50% inhibition of parasite growth relative to an untreated control.

The percentage of growth inhibition was expressed according to following equation: Growth inhibition% =  $100 - [(\text{test parasitaemia}/\text{control parasitemia}) \times 100]$ . Chloroquine:  $\text{IC}_{50} = 0.0061 \mu\text{g}/\text{ml}$ .

## 2.5. Cytotoxicity

P-388 murine leukemia cells were maintained in RPMI-1640 medium supplemented with 5% fetal calf serum and kanamycin (100  $\mu\text{g}/\text{ml}$ ). The cells ( $3 \times 10^3$  cells/well) were cultured in Corning disposable 96-well plates containing 100  $\mu\text{l}$  of growth medium per well and were incubated at 37 °C in a humidified atmosphere of 5%  $\text{CO}_2$ . Various drug concentrations (10  $\mu\text{l}$ ) were added to the cultures at day one after the transplantation. At day three, 20  $\mu\text{l}$  MTT solution (5 mg/ml) per well was added to each cultured medium. After a further 4 h of incubation, 100  $\mu\text{l}$  of 10% SDS–0.01 N HCl solution was added to each well and the formazan crystals in each well were dissolved by stirring with a pipette. The optical density measurements were made using a microplate reader (Tohso MPR-A4i) with a two wavelength system (550 and 700 nm). In all experiments, three replicate wells were used to determine each point.

## Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan and grants from Tokyo Biochemical Foundation, The Open Research Center Project in Hoshi university, ScienceFund 12-02-03-2034, Ministry of Science and Technology, Malaysia, Scientific grant from Institut de chimie des Substances Naturelles, CNRS, France, and Assessment Service Unit, Faculty of Pharmacy, Airlangga University.

## References and notes

- (a) Taylor, D. A. H. In Herz, W., Grisebach, H., Kirby, G. W., Eds.; Progress in the Chemistry of Organic Natural Products; Springer: New York, 1984; Vol. 45. (b) Mulholland, D.; Parel, A.; Coombes, P. H. *Curr. Org. Chem.* **2000**, *4*, 1011.
- (a) Yin, S.; Wang, X. N.; Fan, C. Q.; Liao, S. G.; Yue, J. M. *Org. Lett.* **2007**, *9*, 2353; (b) Zhang, C. R.; Yang, S. P.; Liao, S. G.; Fan, C. Q.; Wu, Y.; Yue, J. M. *Org. Lett.* **2007**, *9*, 3383; (c) Di, Y. T.; He, H. P.; Liu, H. Y.; Yi, P.; Zhang, Z.; Ren, Y. L.; Wang, J. S.; Sun, Q. Y.; Yang, F. M.; Fang, X.; Li, S. L.; Zhu, H. J.; Hao, X. J. *J. Nat. Prod.* **2007**, *70*, 1352.
- (a) Zhang, H.; Wang, X.; Chen, F.; Androulakis, X. M.; Wargovich, M. J. *Phytother. Res.* **2007**, *21*, 731; (b) Roy, A.; Saraf, S. *Biol. Pharm. Bull.* **2006**, *29*, 191; (c) Bray, D. H.; Warhurst, D. C.; Connolly, J. D.; O'Neill, M. J.; Phillipson, J. D. *Phytother. Res.* **1990**, *4*, 29.
- Wiesner, J.; Ortmann, R.; Jomaa, H.; Schlitzer, M. *Angew. Chem. Int. Ed.* **2003**, *42*, 5274.
- (a) Gelb, M. H.; Hol, W. G. *Science* **2002**, *297*, 343; (b) De Smet, P. A. G. N. *Drugs* **1997**, *54*, 801.
- Morita, H.; Oshimi, S.; Hirasawa, Y.; Koyama, K.; Honda, T.; Ekasari, W.; Indrayanto, G.; Zaini, N. C. *Org. Lett.* **2007**, *9*, 3691.
- (a) Awang, K.; Lim, C. S.; Mohamad, K.; Morita, H.; Hirasawa, Y.; Takeya, K.; Thoison, O.; Hadi, A. H. A. *Bioorg. Med. Chem.* **2007**, *15*, 5997; (b) Mohamad, K.; Hirasawa, Y.; Lim, C. S.; Awang, K.; Hadi, A. H. A.; Takeya, K.; Morita, H. *Tetrahedron Lett.* **2008**, *49*, 4276.
- Krief, S.; Martin, M.-T.; Grelhier, P.; Kasenene, J.; Sevenet, T. *Antimicrob. Agents Chemother.* **2004**, 3196.
- Trager, W.; Jensen, J. B. *Science* **1976**, *193*, 673.
- Budimulja, A. S.; Syafruddin, T. P.; Wilairat, P.; Marzuki, S. *Mol. Biochem. Parasitol.* **1997**, *84*, 137.

# Ceramicines B–D, new antiplasmodial limonoids from *Chisocheton ceramicus*

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Chee-Yan Choo. "Cytotoxic activity of *Typhonium flagelliforme* (Araceae)", *Phytotherapy Research*, 05/2001  
Publication 2%
- 2 Mio Takagi, Yosuke Tachi, Jie Zhang, Takuro Shinozaki et al. " Cytotoxic and Melanogenesis-Inhibitory Activities of Limonoids from the Leaves of (Neem) ", *Chemistry & Biodiversity*, 2014  
Publication 2%
- 3 [www.jstage.jst.go.jp](http://www.jstage.jst.go.jp)  
Internet Source 2%
- 4 Chee Yan Choo, Kit Lam Chan, Teng Wah Sam, Yukio Hitotsuyanagi, Koichi Takeya. "The cytotoxicity and chemical constituents of the hexane fraction of *Typhonium flagelliforme* (Araceace)", *Journal of Ethnopharmacology*, 2001  
Publication 2%

5	Ishak Zakaria, Norizan Ahmat, Faridahanim M. Jaafar, Aty Widyawaruyanti. "Flavonoids with antiplasmodial and cytotoxic activities of <i>Macaranga triloba</i> ", <i>Fitoterapia</i> , 2012 Publication	1 %
6	<a href="http://dashboardffup.info">dashboardffup.info</a> Internet Source	1 %
7	<a href="http://www.thieme-connect.com">www.thieme-connect.com</a> Internet Source	1 %
8	<a href="http://protocolexchange.researchsquare.com">protocolexchange.researchsquare.com</a> Internet Source	1 %
9	Liu, J.Q.. "Limonoids from the leaves of <i>Toona ciliata</i> var. <i>yunnanensis</i> ", <i>Phytochemistry</i> , 201204 Publication	1 %
10	Tan, Qin-Gang, and Xiao-Dong Luo. "Meliaceous Limonoids: Chemistry and Biological Activities", <i>Chemical Reviews</i> , 2011. Publication	1 %
11	Tilak T. Raj, Maurice R. Eftink. "Synthesis of E and Z Diastereoisomers of Benzyl Methyl-2-[3-(N-Substituted)indolyl] Cyclopropane-1,1-dicarboxylates and Their Stereoselective Reductive Cleavage", <i>Synthetic Communications</i> , 1996 Publication	1 %

12	<a href="http://ir.kib.ac.cn:8080">ir.kib.ac.cn:8080</a> Internet Source	1 %
13	<a href="http://ijpt.iums.ac.ir">ijpt.iums.ac.ir</a> Internet Source	<1 %
14	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	<1 %
15	Qiu-Fen Hu, Bin Zhou, Xue-Mei Gao, Li-Ying Yang, Li-Dan Shu, Yanqiong Shen, Gan-Peng Li, Chun-Tao Che, Guang-Yu Yang. " Antiviral Chromones from the Stem of ", Journal of Natural Products, 2012 Publication	<1 %
16	<a href="http://studentsrepo.um.edu.my">studentsrepo.um.edu.my</a> Internet Source	<1 %
17	<a href="http://patents.justia.com">patents.justia.com</a> Internet Source	<1 %
18	Afifa Bergaoui, Saoussen Hammami, Hichem Ben Jannet, Zine Mighr. " Echiofruzine, a novel naringenin diglycoside from growing in Tunisia ", Natural Product Research, 2007 Publication	<1 %
19	C. -I. Chang. "Two new naphthoquinones from the stem of Diospyros maritima", Journal of Asian Natural Products Research, 3/2007 Publication	<1 %



20 Fiorentino, A.. "Carexanes from *Carex distachya* Desf.: revised stereochemistry and characterization of four novel polyhydroxylated prenylstilbenes", *Tetrahedron*, 20080811

<1 %

Publication

21 Wimal Herath, Daneel Ferreira, Ikhlas A. Khan. "Microbial metabolism. Part 7: Curcumin", *Natural Product Research*, 2007

<1 %

Publication

22 [patentscope.wipo.int](http://patentscope.wipo.int)

Internet Source

<1 %

23 Chen, Hua-Dong, Sheng-Ping Yang, Yan Wu, Lei Dong, and Jian-Min Yue. "Terpenoids from *Toona ciliata*", *Journal of Natural Products*, 2009.

<1 %

Publication

24 Chun-Shun Li. "Daphlongeranines A and B, Two Novel Alkaloids Possessing Unprecedented Skeletons from *Daphniphyllum longracemosum*", *Organic Letters*, 06/2007

<1 %

Publication

25 G. V. Gavrilova, A. A. Aleshkin, V. A. Nikanorov, D. P. Krut'ko, V. I. Rozenberg, O. A. Reutov. "Redox troponization as a novel method for the synthesis of stereoisomeric Eschenmoser's oximes and related non-

<1 %

benzenoid aromatic systems", Russian  
Chemical Bulletin, 1995

Publication

---

26

Zhengxi Hu, Ye Wu, Shuangshuang Xie,  
Weiguang Sun et al. " Phomopsterones A and  
B, Two Functionalized Ergostane-Type  
Steroids from the Endophytic Fungus sp.  
TJ507A ", Organic Letters, 2017

Publication

---

<1 %

27

[repository.kulib.kyoto-u.ac.jp](http://repository.kulib.kyoto-u.ac.jp)

Internet Source

---

<1 %

28

Hongwei Liu, Guanghui Wang, Michio  
Namikoshi, Hisayoshi Kobayashi, Xincheng  
Yao, Guoping Cai. "Sesquiterpene Quinones  
from a Marine Sponge Hippospongia sp. that  
Inhibit Maturation of Starfish Oocytes and  
Induce Cell Cycle Arrest with HepG2 Cells",  
Pharmaceutical Biology, 2008

Publication

---

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Ceramicines B–D, new antiplasmodial limonoids from *Chisocheton ceramicus*

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---