



COVID-19 Remote Access Support: Learn More about expanded access to ACS Publications research.



Editor-in-Chief:
Philip J. Proteau
About the Journal

Open Access
Submission & Review

Submit Manuscript

Editors & Editorial Board

Editor-in-Chief

Philip J. Proteau
Oregon State University
United States
E-mail: proteau-office@jnp.acs.org

Current Issue Editorial Masthead

View the Masthead in Current Issue

Associate Editors

Roberto G.S. Berlinc
Instituto de Química de São Carlos, Universidade de São Paulo
Brazil
E-mail: berlinc-office@jnp.acs.org

Joanna E. Burdette
University of Illinois at Chicago
United States
E-mail: burdette@jnp.acs.org

Cedric J. Pearce
Mycosynthetix, Inc.
United States
E-mail: pearce-office@jnp.acs.org

Editorial Advisory Board

Raymond Andersen
University of British Columbia
Canada

William Fenical
Scripps Institution of Oceanography
United States

Kerry L. McPhail
Oregon State University
United States

Yoshinori Asakawa
Tokushima Bunri University
Japan

Mary Garson
University of Queensland
Australia

Susan L. Mooberry
University of Texas Health Science Center
United States

Marcy Balunas
University of Connecticut
United States

William H. Gerwick
University of California
United States

Dale Nagle
University of Mississippi
United States

Vandertan da Silva Bozani
Sao Paulo State University
Brazil

Simon Gibbons
University College of London
United Kingdom

Guido Pauli
University of Illinois at Chicago
United States

Robert J. Capon
The University of Queensland
Australia

James B. Gloer
University of Iowa
United States

Mohammad A. Rashid
University of Dhakka
Bangladesh

Guy T. Carter
Carter-Brennan Consulting
United States

Kirk R. Gustafson
National Cancer Institute at Frederick
United States

Christine E. Salomon
University of Minnesota
United States

Nadja Cech
University of North Carolina—Greensboro
United States

Mark T. Hamann
Medical University of South Carolina
United States

Eric W. Schmidt
University of Utah
United States

Yongsheng Che
Chinese Academy of Medical Sciences and Peking Union Medical
College
China

Jianhua Ju
South China Sea Institute of Oceanology
China

Sheo B. Singh
Merck Research Laboratories, Rahway
United States

Robert Cichewicz
University of Oklahoma
United States

Biik San Clara Lau
Chinese University of Hong Kong
China

Amy E. Wright
Harbor Branch Oceanographic Institution
United States

Nunziatina de Tommasi
University of Salerno
Italy

Rachel Mata
National Autonomous University of Mexico
Mexico

Abiy Yenesew
University of Nairobi
Kenya

Honorary Editorial Advisory Board

Susan Band Horwitz
Albert Einstein College of Medicine
United States

Phillip Crews
University of California – Santa Cruz
United States

Jon C. Clardy
Harvard Medical School
United States

David G.J. Kingston
Virginia Polytechnic Institute & State University
United States

Identification of Antiadipogenic Constituents of the Rhizomes of *Anemarrhena asphodeloides*


Ui Joung Youn, Ye Seul Lee, Hana Jeong, Jun Lee, Joo-Won Nam, Yoo Jin Lee, Eun Sook Hwang, Je-Hyun Lee, Donghe Lee, Sam Sik Kang, and Eun-Hyoung Seo*

Journal of Natural Products 2009, 72, 10, 1895-1898 (Note)

Publication Date (Web): September 17, 2009

[Abstract](#) [Full text](#) [PDF](#)

ABSTRACT



Cassiarins C-E, Antiplasmodial Alkaloids from the Flowers of *Cassia siamea*


Shiori Oshimi, Jun Deguchi, Yasuke Hirasawa, Wijiwed Ekasari, Aty Widiyawanjanti, Tutik Sri Wanyuni, Noor Cholles Zaini, Osamu Shirota, and Hiroshi Morita*

Journal of Natural Products 2009, 72, 10, 1902-1903 (Note)

Publication Date (Web): September 16, 2009

[Abstract](#) [Full text](#) [PDF](#)

ABSTRACT



Ferene Triterpenoids from the Lichen *Pyxine berteriana*


Marta S. Meier*, Maria L. Rosso, Alejandra T. Fazio, Mônica T. Adler, and Maria D. Bertoni

Journal of Natural Products 2009, 72, 10, 1902-1903 (Note)

Publication Date (Web): September 16, 2009

[Abstract](#) [Full text](#) [PDF](#)

ABSTRACT



Xantheponone, an Antimicrobial Polyketide from a Soil Fungus Closely Related to *Phoma medicaginis*


Johannes C. Liermann, Heinz Kolshorn, Till Opatz*, Eckhard Thines, and Heidrun Anke*

Journal of Natural Products 2009, 72, 10, 1905-1907 (Note)

Publication Date (Web): October 1, 2009

[Abstract](#) [Full text](#) [PDF](#)

ABSTRACT



Revised Structure of the Alkaloid Drymaritin


Isolde Wetzel, Luis Allmendinger, and Franz Breche*

Journal of Natural Products 2009, 72, 10, 1908-1910 (Note)

Publication Date (Web): September 21, 2009

[Abstract](#) [Full text](#) [PDF](#)

ABSTRACT



Cassiarins C–E, Antiplasmodial Alkaloids from the Flowers of *Cassia siamea*


Shiori Oshimi,[†] Jun Deguchi,[†] Yusuke Hirasawa,[†] Wiwied Ekasari,[‡] Aty Widyawaruyanti,[‡] Tutik Sri Wahyuni,[‡] Noor Cholies Zaini,[‡] Osamu Shiota,[§] and Hiroshi Morita^{*,†}

Faculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41 Shinagawa, Tokyo 142-8501, Japan, Faculty of Pharmacy, Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia, and Faculty of Pharmaceutical Sciences at Kagawa Campus, Tokushima Bunri University, 1314-1 Shido, Sanuki City, Kagawa 769-2193, Japan


Received July 13, 2009

Three new alkaloids, cassiarins C–E (**1–3**), and a new chromone, 10,11-dihydroanhydrobarakol (**4**), which showed moderate antiplasmodial activity against *Plasmodium falciparum* 3D7, were isolated from flowers of *Cassia siamea*, and the structures of **1–4** were elucidated by 2D NMR analysis and chemical transformation. Cassiarin D (**2**) was a dimeric compound consisting of 5-acetylonyl-7-hydroxy-2-methylchromone and cassiarin C (**1**), and cassiarin E (**3**) was a dimer of cassiarins A and C (**1**).


Cassia siamea Lam. (Leguminosae) has been used widely in traditional medicine, particularly for treatment of periodic fever and malaria in Indonesia.¹ Earlier we isolated cassiarin A,² an alkaloid with an unprecedented tricyclic skeleton and potent antiplasmodial activity, and chrobisiamone A,³ a bischromone, from leaves of *C. siamea*. We then reported an efficient total synthesis of cassiarin A,⁴ which also showed in vivo antimalarial activity.⁵ Our continuing search for bioactive constituents of *C. siamea* has resulted in the isolation of three additional new alkaloids, cassiarins C–E (**1–3**), and a new chromone, 10,11-dihydroanhydrobarakol (**4**), from the flowers of *C. siamea*. Also isolated were cassiarin A, anhydrobarakol (**5**),⁶ and 5-acetylonyl-7-hydroxy-2-methylchromone (**6**).⁷ This paper describes the isolation and structure elucidation of **1–4** and their antiplasmodial activity against *Plasmodium falciparum* 3D7.




1



2



3



4

Cassiarin C (**1**), $[\alpha]_D^{20} -70$ (c 0.3, MeOH), had the molecular formula $C_{13}H_{13}NO_2$, which was determined by HRESITOFMS [m/z 216.1025, $(M + H)^+$, $\Delta +1.7$ mmu]. IR absorptions implied the presence of hydroxy and/or amino (3401 cm^{-1}) functionalities. ^1H and ^{13}C NMR data are presented in Table 1. The ^{13}C NMR spectrum revealed 13 signals due to six sp^2 quaternary carbons, three sp^2 methines, one sp^3 methine, one sp^3 methylene, and two methyl groups. Among them, four quaternary carbons (δ_C 149.9, 153.6, 156.9, and 159.1) and one sp^3 methine (δ_C 73.7) were ascribed to those bearing nitrogen or oxygen atoms. Two partial structures,

C-10 to C-12 and C-9 to C-3, were deduced from analysis of the $^1\text{H}-^1\text{H}$ COSY spectrum, which included a long-range coupling from H₃-12 to H-10 of **1**. The HMBC correlations of H-10 to C-4a (δ_C 110.6) and C-5 (δ_C 139.1) and of H₂-3 to C-4 (δ_C 153.6) and C-4a indicated connectivity of two partial structures through a nitrogen and C-4 atoms. Connection between the partial structure and a benzene ring was determined by a HMBC correlation between H-10 and C-6 (δ_C 100.0). The presence of an ether linkage between C-2 (δ_C 73.7) and C-8a (δ_C 156.9) to form a dihydropyran ring was also evident. Thus, cassiarin C (**1**) was concluded to be a 2,3-dihydrocassiarin A.

Cassiarin D (**2**), $[\alpha]_D^{20} -10$ (c 0.1, MeOH), had the molecular formula $C_{26}H_{23}NO_6$ (by HRESITOFMS). IR absorptions implied the presence of hydroxy and/or amino (3673 cm^{-1}) and carbonyl (1713 and 1632 cm^{-1}) functionalities. ^1H and ^{13}C NMR data are presented in Table 2. The ^{13}C NMR spectrum revealed 26 signals due to two carbonyl carbons, 11 sp^2 quaternary carbons, one sp^3 quaternary carbon, three sp^3 methylenes, six sp^2 methines, and three methyl groups. Among them, 10 quaternary carbons (δ_C 78.9, 152.4, 152.5, 154.9, 160.1, 162.5, 162.9, 163.1, 178.4, and 204.6) were ascribed to those bearing nitrogen or oxygen atoms. The gross structure of **2** consisted of two units, a chromone, and cassiarin C, as deduced from extensive analyses of the two-dimensional NMR data, including the $^1\text{H}-^1\text{H}$ COSY and HMBC spectra in pyridine- d_5 . The presence of an acetylonyl group at C-5 was supported by HMBC correlations of H₃-13 (δ_H 2.37) to C-11 (δ_C 49.8) and C-12 (δ_C 204.6) and H₂-11 (δ_H 4.37) to C-5 (δ_C 139.2), C-6 (δ_C 119.4), and C-10 (δ_C 115.1). HMBC correlations of H₂-14 (δ_H 2.83 and 2.96) to C-2 (δ_C 163.1), C-2' (δ_C 78.9), C-3 (δ_C 114.0), and C-3' (δ_C 42.3) and those of H₃-9' (δ_H 1.42) to C-14 (δ_C 42.9), C-2', and C-3' indicated connectivity of two units through C-14 and C-2'. Thus, cassiarin D (**2**) was concluded to be a unique dimeric compound consisting of 5-acetylonyl-7-hydroxy-2-methylchromone² and cassiarin C (**1**).

Cassiarin E (**3**), $[\alpha]_D^{20} -20$ (c 0.3, MeOH), had the molecular formula $C_{26}H_{22}N_2O_4$ (by HRESITOFMS). IR absorptions implied the presence of hydroxy and/or amino (3743 cm^{-1}) and carbonyl (1622 cm^{-1}) functionalities. The gross structure of **3** was elucidated by analyses of 2D NMR data including $^1\text{H}-^1\text{H}$ COSY, HOHAHA, HMQC, and HMBC spectra in pyridine- d_5 (Figure S3). Each pair of observed ^1H and ^{13}C NMR signals seemed to be due to each half moiety of a dimeric compound of cassiarin A.² Methylene signals (δ_H 2.79 and 2.90; δ_C 43.1) were observed in place of a methyl signal (C-12) in cassiarin A. Methylene signals (δ_H 3.38 and 3.52; δ_C 42.3) were observed as in the case of chrobisiamone A.³ The connection between the two units, C-9 to C-2', was provided by HMBC correlations of H₂-9 to C-2' (δ_C 79.1) and C-3'

* To whom correspondence should be addressed. Tel: (03)5498-5778. Fax: (03)5498-5778. E-mail: moritah@hoshi.ac.jp.

[†] Hoshi University.

[‡] Airlangga University.

[§] Tokushima Bunri University.

Table 1. ^1H NMR Data [δ_{H} (J , Hz)] and ^{13}C NMR Data [δ_{C}] for Cassiarin C (**1**) and 10,11-Dihydroanhydrobarakol (**4**) in CD_3OD at 300 K^a

position	1		4	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
2	4.49 (1H, m)	73.7		165.3
3a	3.06 (1H, dd, 16.8, 10.9)	37.8	6.04 (1H, d, 0.6)	110.0
3b	3.13 (1H, dd, 16.8, 3.2)			
4		153.6		160.2 ^b
4a		110.6		111.8
5		139.1		143.0
6	6.57 (1H, d, 2.0)	100.0	6.70 (1H, d, 2.5)	118.1
7		159.1		180.5
8	6.48 (1H, d, 2.0)	103.1	6.71 (1H, d, 2.5)	110.9
8a		156.9		161.6 ^b
9	1.56 (3H, d, 6.0)	19.9	2.34 (3H, d, 0.6)	18.4
10a	7.20 (1H, s)	115.6	3.05 (1H, dd, 12.4, 7.6)	43.9
10b			3.51 (1H, dd, 12.4, 4.4)	
11		149.9	4.00 (1H, m)	68.2
12	2.53 (3H, s)	22.1	1.22 (3H, d, 6.4)	22.2

^a δ in ppm. ^b Interchangeable.**Table 2.** ^1H NMR Data [δ_{H} (J , Hz)] and ^{13}C NMR Data [δ_{C}] for Cassiarins D and E (**2** and **3**) in Pyridine-*d*₅ at 300 K^a

position	2		3	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
2		163.1		158.0
3	6.24 (1H, s)	114.0	6.44 (1H, s)	110.0
4		178.4		150.7
4a				112.9
5		139.2		139.2
6	7.02 (1H, d, 1.5)	119.4	6.91 (1H, s)	101.6
7		162.9		162.2
8	7.04 (1H, d, 1.5)	102.7	6.91 (1H, s)	100.0
8a				155.9
9a		160.1	2.79 (1H, d, 14.2)	43.1
9b			2.90 (1H, d, 14.2)	
10		115.1	6.84 (1H, s)	113.3
11	4.37 (2H, s)	49.8		153.9
12		204.6	2.50 (3H, s)	24.6
13	2.37 (3H, s)	30.1		
14a	2.83 (1H, d, 14.2)	42.9		
14b	2.96 (1H, d, 14.2)			
2'		78.9		79.1
3'a	3.32 (1H, d, 16.6)	42.3	3.38 (1H, d, 16.6)	42.3
3'b	3.42 (1H, d, 16.6)		3.52 (1H, d, 16.6)	
4'		152.5		152.7
4'a		110.3		110.4
5'		139.0		139.1
6'	7.02 (1H, d, 1.0)	101.0	7.02 (1H, d, 2.0)	100.9
7'		162.5		162.6
8'	6.98 (1H, d, 1.0)	103.8	7.04 (1H, d, 2.0)	103.8
8'a		154.9		155.2
9'	1.42 (3H, s)	25.1	1.54 (3H, s)	25.2
10'	7.17 (1H, brs)	115.8	7.19 (1H, s)	115.7
11'		152.4		152.5
12'	2.61 (3H, s)	24.4	2.62 (3H, s)	24.5

^a δ in ppm.

(δ_{C} 42.3), and H₃-9' (δ_{H} 1.54) to C-9 and C-3'. Thus, the gross structure of cassiarin E (**3**) was assigned to be a unique dimeric compound consisting of cassiarin A and cassiarin C (**1**).

Compound **4**, 10,11-dihydroanhydrobarakol, [α_{D}^{20} +47 (*c* 1.0, MeOH), had the molecular formula C₁₃H₁₂O₃, which was determined by HRESITOFMS. The IR spectrum implied the presence of a carbonyl group (1652 cm⁻¹). ^1H and ^{13}C NMR data are presented in Table 1. The ^{13}C NMR spectrum revealed 13 carbon signals due to one carbonyl carbon, five sp² quaternary carbons, three sp² methines, one sp³ methine, one sp³ methylene, and two methyl groups. Among them, four quaternary carbons (δ_{C} 160.2, 161.6, 165.3, and 180.5) and one sp³ methine (δ_{C} 68.2) were

ascribed to those bearing an oxygen atom. Two partial structures, C-10 to C-12, and C-9 to C-2 and C-3, were deduced from ^1H - ^1H COSY analysis of **4** in CD₃OD. Connectivity of partial structures through an oxygen and C-4 atoms in the tricyclic skeleton was supported by HMBC correlations of H-3, H-6, H-8, and H-10 to C-4a (δ_{C} 111.8). Connection between partial structure and the dienone ring could be assigned by a HMBC correlation of H-10 to C-6 (δ_{C} 118.1). In addition, the presence of an ether linkage between C-2 (δ_{C} 165.3) and C-8a to form a pyran ring was also indicated.

A plausible biogenetic pathway for cassiarins C–E (**1**–**3**) and 10,11-dihydroanhydrobarakol (**4**) was proposed as shown in Scheme 1, Supporting Information. Cassiarin C (**1**) could be derived through an imine intermediate of 2,3-dihydro-5-acetyl-7-hydroxy-2-methylchromone followed by cyclization with the acetyl group as shown in Scheme 1, whereas 10,11-dihydroanhydrobarakol (**4**) might be produced by way of anhydrobarakol (**5**) from 5-acetyl-7-hydroxy-2-methylchromone (**6**). On the other hand, cassiarins D and E (**2** and **3**) could be derived by way of chrobisiamone A from 5-acetyl-7-hydroxy-2-methylchromone (**6**), as shown in Scheme 1. Treatment of cassiarin D (**2**) with ammonium acetate as the nitrogen source caused ring cyclization, giving cassiarin E (**3**) in 63% yield. This biomimetic transformation supports the biogenetic pathway proposed for cassiarins D and E.

Cassiarins C–E (**1**–**3**) and 10,11-dihydroanhydrobarakol (**4**) showed moderate in vitro antiplasmodial activity against *Plasmodium falciparum* 3D7 (IC₅₀ **1**: 24.2 μM ; **2**: 3.6 μM ; **3**: 7.3 μM ; **4**: 2.3 μM) and no cytotoxicity against HL-60 (human blood premyelocytic leukemia) cells (IC₅₀ >100 μM for **1**–**4**). Anhydrobarakol (**5**)⁶ and 5-acetyl-7-hydroxy-2-methylchromone (**6**)⁷ also showed moderate antiplasmodial activity (IC₅₀ **5**: 4.7 μM ; **6**: 8.6 μM) and no cytotoxicity against HL-60 cells (IC₅₀ >100 μM for **5** and **6**). Dimeric-type compounds such as cassiarins D (**2**) and E (**3**) were less active than cassiarin A (IC₅₀ 0.023 μM), although they contained the cassiarin A-type skeleton.² 10,11-Dihydroanhydrobarakol (**4**) showed activity similar to that of anhydrobarakol (**5**).

Experimental Section

General Experimental Procedures. Optical rotations were measured on a JASCO DIP-1000 automatic digital polarimeter. UV spectra were obtained on an Ultrospec 2100 pro spectrophotometer, and IR spectra were recorded on a JASCO FT/IR-4100 spectrophotometer. High-resolution ESIMS were obtained on a LTQ Orbitrap XL (Thermo Scientific). ^1H and 2D NMR spectra were recorded on Bruker AV 400 and Inova 500 spectrometers, and chemical shifts were referenced to the residual solvent peaks (δ_{H} 3.31 and δ_{C} 49.0 for methanol-*d*₄ and δ_{H} 7.71 and δ_{C} 135.5 for pyridine-*d*₅). Standard pulse sequences were employed for the 2D NMR experiments. ^1H - ^1H COSY, HOHAHA, and NOESY spectra were measured with spectral widths of both dimensions of 4800 Hz, and 32 scans with two dummy scans were accumulated into 1K data points for each of 256 t_1 increments. NOESY spectra in the phase-sensitive mode were measured with a mixing time of 800 ms. For HMQC spectra in the phase-sensitive mode and HMBC spectra, a total of 256 increments of 1K data points were collected. For HMBC spectra with Z-axis PFG, a 50 ms delay time was used for long-range C–H coupling. Zero-filling to 1K for F_1 and multiplication with squared cosine-bell windows shifted in both dimensions were performed prior to 2D Fourier transformation.

Material. Flowers of *C. siamea* were collected at Purwodadi Botanical Garden in 2007. The botanical identification was made by Ms. Sri Wuryanti, Purwodadi Botanical Garden, Indonesia. A voucher specimen (no. PU080306) has been deposited at Purwodadi Botanical Garden, Pasuruan, Indonesia.

Extraction and Isolation. Dried flowers of *C. siamea* (1.0 kg) were extracted with MeOH, and the extract was partitioned between CHCl₃ and H₂O. CHCl₃-soluble materials were subjected to a silica gel column (CHCl₃/MeOH, 1:0 \rightarrow 0:1) to afford cassiarin A (0.04%),² anhydrobarakol (0.006%),⁶ and 5-acetyl-7-hydroxy-2-methylchromone (0.007%).⁷ A fraction eluted by CHCl₃/MeOH (30:1) was further separated on an ODS column (MeOH/H₂O, 0:1 \rightarrow 1:0) and on a silica

gel column (CHCl₃/MeOH, 1:0 → 0:1) to afford cassiarin C (**1**, 0.00007%). A fraction eluted by CHCl₃/MeOH (25:1) was further purified on a silica gel column (CHCl₃ saturated with NH₃/MeOH, 1:0 → 0:1) to afford 10,11-dihydroanhydrobarakol (**4**, 0.00050%), and a fraction eluted by CHCl₃/MeOH (20:1) was dissolved in MeOH and then centrifuged to afford cassiarin D (**2**, 0.0013%). The CHCl₃/MeOH (9:1)-soluble fraction was separated on a silica gel column (CHCl₃ saturated with NH₃/MeOH, 1:0 → 0:1) to afford cassiarin E (**3**, 0.00078%).

Cassiarin C (1): brown solid; $[\alpha]_D^{20}$ -70 (*c* 0.3, MeOH); IR (KBr) ν_{\max} 3401, 2924, and 1618 cm⁻¹; UV (MeOH) λ_{\max} 218 (ϵ 16 000), 243 (21 000), 289 (sh, 4300), and 330 (4000) nm (ϵ 13 000); ¹H and ¹³C NMR data (Tables 1 and 2); ESIMS *m/z* 216 (M + H)⁺; HRESITOFMS *m/z* 216.1025 [(M + H)⁺] (calcd for C₁₃H₁₄NO₂, 216.1008).

Cassiarin D (2): brown solid; $[\alpha]_D^{20}$ -10 (*c* 0.1, MeOH); IR (KBr) ν_{\max} 3673, 2992, 1713, 1632, and 1588 cm⁻¹; UV (MeOH) λ_{\max} 243 (ϵ 24 000), 298 (6000), and 330 (3000) nm (ϵ 2400); ¹H and ¹³C NMR, see Tables 1 and 2; ESIMS *m/z* 446 (M + H)⁺; HRESITOFMS *m/z* 446.1604 [(M + H)⁺] (calcd for C₂₆H₂₄NO₆, 446.1611).

Cassiarin E (3): brown solid; $[\alpha]_D^{20}$ -20 (*c* 0.3, MeOH); IR (KBr) ν_{\max} 3743, 2986, and 1622 cm⁻¹; UV (MeOH) λ_{\max} 219 (ϵ 10 000), 243 (11 600), and 338 (2400) nm; ¹H and ¹³C NMR, see Tables 1 and 2; ESIMS *m/z* 427 (M + H)⁺; HRESITOFMS *m/z* 427.1648 [(M + H)⁺] (calcd for C₂₆H₂₃N₂O₄, 427.1658).

10,11-Dihydroanhydrobarakol (4): brown solid; $[\alpha]_D^{20}$ +47 (*c* 1.0, MeOH); IR (KBr) ν_{\max} 2924, 1652, and 1589 cm⁻¹; UV (MeOH) λ_{\max} 210 (ϵ 11 200), 242 (8600), 250 (8500), and 290 (6000) nm; ¹H and ¹³C NMR, see Tables 1 and 2; ESIMS *m/z* 217 (M + H)⁺; HRESITOFMS *m/z* 217.0859 [(M + H)⁺] (calcd for C₁₃H₁₃O₃, 217.0862).

Conversion of Cassiarin D (2) to Cassiarin E (3). Ammonium acetate (1.0 mg) was added to a solution of cassiarin D (**2**, 1.0 mg) in AcOH (0.2 mL), and the mixture was kept at 60 °C for 24 h. After evaporation, the residue was applied to a silica gel column (CHCl₃/MeOH, 9:1) to give a compound (0.6 mg) whose spectroscopic data and $[\alpha]_D$ value were identical to those of natural cassiarin E (**3**).

Antiplasmodial Activity. Human malaria parasites were cultured according to the method of Trager and Jensen.⁸ The antimalarial activity of the isolated compounds was determined by the procedure described by Budimulya et al.⁹ In brief, stock solutions of the samples were prepared in DMSO (final DMSO concentrations of <0.5%) and were diluted to the required concentration with complete medium (RPMI 1640 supplemented with 10% human plasma, 25 mM HEPES, and 25 mM NaHCO₃) until the final concentrations of samples in culture plate wells were 10, 1, 0.1, 0.01, and 0.001 μg/mL. The malarial parasite *P.*

falciparum 3D7 clone was propagated in a 24-well culture plate. Growth of the parasite was monitored by making a blood smear fixed with MeOH and stained with Geimsa stain. The antimalarial activity of each compound was expressed as an IC₅₀ value, defined as the concentration of the compound causing 50% inhibition of parasite growth relative to an untreated control.

The percentage of growth inhibition was expressed according to the following equation: Growth inhibition % = 100 - [(test parasitemia/control parasitemia)] × 100. Chloroquine: IC₅₀ 0.011 μM.

Acknowledgment. This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and by a grant from the Open Research Center Project. We also acknowledge the financial support provided by the Faculty of Pharmacy, Airlangga University, for antiplasmodial (antimalarial) activity.

Supporting Information Available: Scheme 1: Proposed Biogenetic Pathway of Cassiarins C–E (**1–3**) and 10,11-Dihydroanhydrobarakol (**4**). ¹H and ¹³C NMR spectra and 2D NMR correlations of **1–4**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- (a) Mbatchi, S. F.; Mbatchi, B.; Banzouzi, J. T.; Bansimba, T.; Nsonde Ntandou, G. F.; Ouamba, J. M.; Berry, A.; Benoit-Vical, F. *J. Ethnopharmacol.* **2006**, *104*, 168–174. (b) Sanon, S.; Ollivier, E.; Azas, N.; Mahiou, V.; Gasquet, M.; Ouattara, C. T.; Nebie, I.; Traore, A. S.; Esposito, F.; Balansard, G.; Timon-David, P.; Fumoux, F. *J. Ethnopharmacol.* **2003**, *86*, 143–147.
- Morita, H.; Oshimi, S.; Hirasawa, Y.; Koyama, K.; Honda, T.; Ekasari, W.; Indrayanto, G.; Zaini, N. C. *Org. Lett.* **2007**, *9*, 3691–3693.
- Oshimi, S.; Tomizawa, Y.; Hirasawa, Y.; Honda, T.; Widyawaruyanti, A.; Rudyanto, M.; Ekasari, W.; Indrayanto, G.; Zaini, C. N.; Morita, H. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 3761–3763.
- Rudyanto, M.; Tomizawa, Y.; Morita, H.; Honda, T. *Org. Lett.* **2008**, *10*, 1921–1922.
- Ekasari, W.; Widyawaruyanti, A.; Zaini, N. C.; Syafruddin, D.; Honda, T.; Morita, H. *Heterocycles* **2009**, *78*, 1831–1836.
- Teeyapant, R.; Srikun, O.; Wray, V.; Writte, L. *Fitoterapia* **1998**, *69*, 475–476.
- Arora, S.; Deymann, H.; Tiwri, R. D.; Winterfeldt, E. *Tetrahedron* **1971**, *27*, 981–984.
- Trager, W.; Jensen, J. B. *Science* **1976**, *193*, 673–675.
- Budimulya, A. S.; Syafruddin, T. P.; Wilariat, P.; Marzuki, S. *Mol. Biochem. Parasitol.* **1997**, *84*, 137–141.

NP9004213

Cassiarins C–E, Antiplasmodial Alkaloids from the Flowers of *Cassia siamea*

Shiori Oshimi,[†] Jun Deguchi,[†] Yusuke Hirasawa,[†] Wiwied Ekasari,[‡] Aty Widyawaruyanti,[‡] Tutik Sri Wahyuni,[‡] Noor Cholies Zaini,[‡] Osamu Shirota,[§] and Hiroshi Morita^{†,*}

Faculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41 Shinagawa, Tokyo 142-8501, Japan, Faculty of Pharmacy, Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia, and Faculty of Pharmaceutical Sciences at Kagawa Campus, Tokushima Bunri University, 1314-1 Shido, Sanuki City, Kagawa 769-2193, Japan

Supporting Information

Scheme S1. Proposed Biogenetic Pathway of Cassiarins C–E (**1–3**) and 10,11-Dihydroanhydrobarakol (**4**).

Figure S1. Selected 2D NMR Correlations for cassiarin C (**1**) in CD₃OD.

Figure S2. Selected 2D NMR Correlations for cassiarin D (**2**) in pyridine-*d*₅.

Figure S3. Selected 2D NMR Correlations for cassiarin E (**3**) in pyridine-*d*₅.

Figure S4. Selected 2D NMR Correlations for 10,11-dihydroanhydrobarakol (**4**) in CD₃OD.

Figure S5. ¹H NMR spectrum of cassiarin C (**1**) in CD₃OD.

Figure S6. ¹³C NMR spectrum of cassiarin C (**1**) in CD₃OD.

Figure S7. ¹H NMR spectrum of cassiarin D (**2**) in pyridine-*d*₅.


Figure S8. ¹³C NMR spectrum of cassiarin D (**2**) in pyridine-*d*₅.

Figure S9. ¹H NMR spectrum of cassiarin E (**3**) in pyridine-*d*₅.

Figure S10. ¹³C NMR spectrum of cassiarin E (**3**) in pyridine-*d*₅.

Figure S11. ¹H NMR spectrum of 10,11-dihydroanhydrobarakol (**4**) in CD₃OD.

Figure S12. ¹³C NMR spectrum of 10,11-dihydroanhydrobarakol (**4**) in CD₃OD.



Scheme S1. Proposed Biogenetic Pathway of Cassiarins C–E (1–3) and 10,11-Dihydroanhydrobarakol (4).




Figure S1. Selected 2D NMR Correlations for cassiarin C (1) in CD₃OD.




Figure S2. Selected 2D NMR Correlations for cassiarin D (**2**) in pyridine- d_5 .




Figure S3. Selected 2D NMR Correlations for cassiarin E (**3**) in pyridine- d_5 .




Figure S4. Selected 2D NMR Correlations for 10,11-dihydroanhydrobarakol (**4**) in CD_3OD .




Figure S5. ^1H NMR spectrum of Cassiarin C (1) in CD_3OD .




Figure S6. ^{13}C NMR spectrum of Cassiarin C (**1**) in CD_3OD .




Figure S7. ^1H NMR spectrum of cassiarin D (**2**) in pyridine- d_5 .




Figure S8. ^{13}C NMR spectrum of cassiarin D (**2**) in pyridine- d_5 .




Figure S9. ^1H NMR spectrum of cassiarin E (**3**) in pyridine- d_5 .




Figure S10. ^{13}C NMR spectrum of cassiarin E (**3**) in pyridine- d_5 .




Figure S11. ^1H NMR spectrum of 10,11-dihydroanhydrobarakol (**4**) in CD_3OD .




Figure S12. ^{13}C NMR spectrum of 10,11-dihydroanhydrobarakol (**4**) in CD_3OD .

Journal of Natural Products

Q1 Analytical Chemistry
best quartile

SJR 2021
0.87

powered by scimagojr.com

also developed by scimago **SCIMAGO INSTITUTIONS RANKINGS**

SJR Scimago Journal & Country Rank

Home Journal Rankings Country Rankings Viz Tools Help About Us

Enter Journal Title, ISSN or Publisher Name

Journal of Natural Products

COUNTRY United States Universities and research institutions in United States	SUBJECT AREA AND CATEGORY Biochemistry, Genetics and Molecular Biology Molecular Medicine Chemistry Analytical Chemistry Organic Chemistry Medicine Complementary and Alternative Medicine Pharmacology, Toxicology and Pharmaceutics Drug Discovery Pharmaceutical Science Pharmacology	PUBLISHER American Chemical Society American Chemical Society in Scimago Institutions Rankings	H-INDEX 147
PUBLICATION TYPE Journals	ISSN 01633864, 15206025	COVERAGE 1949, 1971, 1973, 1978-2021	INFORMATION Homepage How to publish in this journal eic@jnp.acs.org

Scopus Preview

Author Search Sources Create account Sign in

Source details

Feedback > Compare sources >

Journal of Natural Products
Formerly known as: Lloydia: A Quarterly Journal of Biological Science
Scopus coverage years: 1949, 1971, 1973, from 1978 to Present
Publisher: American Chemical Society
ISSN: 0163-3864 E-ISSN: 1520-6025

Subject area: Medicine: Complementary and Alternative Medicine Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science
Chemistry: Analytical Chemistry Chemistry: Organic Chemistry Pharmacology, Toxicology and Pharmaceutics: Drug Discovery
Pharmacology, Toxicology and Pharmaceutics: Pharmacology Biochemistry, Genetics and Molecular Biology: Molecular Medicine

View less ^

Source type: Journal

View all documents > Set document alert Save to source list Source Homepage

CiteScore 2021 7.1
SJR 2021 0.866
SNIP 2021 1.593

CiteScore CiteScore rank & trend Scopus content coverage

Improved CiteScore methodology
CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$7.1 = \frac{10,504 \text{ Citations } 2018 - 2021}{1,485 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$6.4 = \frac{8,080 \text{ Citations to date}}{1,266 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Medicine Complementary and Alternative Medicine	#7/93	93rd
Pharmacology, Toxicology and Pharmaceutics Pharmaceutical Science	#28/171	83rd
Chemistry Analytical Chemistry	#22/130	83rd
Chemistry Organic Chemistry	#37/192	80th
Pharmacology, Toxicology and Pharmaceutics Drug Discovery	#35/154	77th
Pharmacology, Toxicology and Pharmaceutics Pharmacology	#73/303	75th
Biochemistry, Genetics and Molecular Biology Molecular Medicine	#56/167	66th