


*Organic*  
**LETTERS**

AUGUST 19, 2011

Volume 13, Number 16 [pubs.acs.org/OrgLett](http://pubs.acs.org/OrgLett)



ACS Publications  
MOST TRUSTED. MOST CITED. MOST READ.

[www.acs.org](http://www.acs.org)

## Editors & Editorial Board

### Editor-in-Chief

**Marisa C. Kozlowski**  
University of Pennsylvania  
United States  
E-mail: [eic@orglett.acs.org](mailto:eic@orglett.acs.org)

### Current Issue Editorial Masthead

[View the Masthead in Current Issue](#)

### Associate Editors

**Margaret A. Brimble**  
The University of Auckland  
New Zealand  
E-mail: [brimble-office@orglett.acs.org](mailto:brimble-office@orglett.acs.org)

**Pauline Chiu**  
The University of Hong Kong  
China  
E-mail: [chiu-office@orglett.acs.org](mailto:chiu-office@orglett.acs.org)

**Kuilong Ding**  
Shanghai Institute of Organic Chemistry  
China  
E-mail: [ding-office@orglett.acs.org](mailto:ding-office@orglett.acs.org)

**Margaret M. Faul**  
Amgen, Inc.  
United States  
E-mail: [faul-office@orglett.acs.org](mailto:faul-office@orglett.acs.org)

**Tanja Gaich**  
Universität Konstanz  
Germany  
E-mail: [gaich-office@orglett.acs.org](mailto:gaich-office@orglett.acs.org)

**Nuno Maulide**  
University of Vienna  
Austria  
E-mail: [maulide-office@orglett.acs.org](mailto:maulide-office@orglett.acs.org)

**Bradley S. Moore**  
University of California, San Diego  
United States  
E-mail: [moore-office@orglett.acs.org](mailto:moore-office@orglett.acs.org)

**Kavirayani R. Prasad**  
Indian Institute of Science  
India  
E-mail: [prasad-office@orglett.acs.org](mailto:prasad-office@orglett.acs.org)

**Tobias Ritter**  
Max-Planck-Institut für Kohlenforschung  
Germany  
E-mail: [ritter-office@orglett.acs.org](mailto:ritter-office@orglett.acs.org)

**Mariola Tortosa**  
Autonomous University of Madrid  
Spain  
E-mail: [tortosa-office@orglett.acs.org](mailto:tortosa-office@orglett.acs.org)

**Zhenfeng Xi**  
Peking University  
China  
E-mail: [xi-office@orglett.acs.org](mailto:xi-office@orglett.acs.org)

**Shigehiro Yamaguchi**  
University of Nagoya  
Japan  
E-mail: [yamaguchi-office@orglett.acs.org](mailto:yamaguchi-office@orglett.acs.org)

**Jin-Quan Yu**  
Scripps Research  
United States  
E-mail: [yu-office@orglett.acs.org](mailto:yu-office@orglett.acs.org)

### Editorial Advisory Board

**Gregory L. Beutner**  
Bristol-Myers Squibb Company  
United States

**David Nicewicz**  
University of North Carolina at Chapel Hill  
United States

**Donna Blackmond**

Scripps Research  
United States

**Cynthia J. Burrows**

University of Utah  
United States

**Erick Carreira**

ETH Hönggerberg, HCI  
Switzerland

**Sherry R. Chemler**

The State University of New York at Buffalo  
United States

**Rick L. Danheiser**

Massachusetts Institute of Technology  
United States

**Scott E. Denmark**

University of Illinois at Urbana-Champaign  
United States

**Vy M. Dong**

University of California at Irvine  
United States

**Neil K. Garg**

UCLA  
United States

**Véronique Gouverneur**

University of Oxford  
United Kingdom

**Kendall N. Houk**

University of California, Los Angeles  
United States

**Koichi Komatsu**

Kyoto University  
Japan

**Ang Li**

Shanghai Institute of Organic Chemistry  
China

**Géraldine Masson**

Université de Paris-Saclay  
France

**Scott J. Miller**

Yale University  
United States

**Cristina Nevado**

University of Zurich  
Switzerland

**Larry E. Overman**

University of California, Irvine  
United States

**Nicola L. B. Pohl**

Indiana University  
United States

**Joëlle Prunet**

University of Glasgow  
United Kingdom

**William R. Roush**

Scripps Research  
United States

**Corinna Schindler**

University of Michigan  
United States

**Franziska Schoenebeck**

RWTH Aachen University  
Germany

**Jennifer M. Schomaker**

University of Wisconsin-Madison  
United States

**Vinod K. Singh**

Indian Institute of Technology  
India

**Amos B. Smith, III**

University of Pennsylvania  
United States

**J. Fraser Stoddart**

Northwestern University  
United States

**Raghavan B. Sunoj**

Indian Institute of Technology, Bombay  
India

**Barry M. Trost**

Stanford University  
United States

**Jianbo Wang**

Peking University  
China

**Helma Wennemers**

ETH Zürich  
Switzerland



August 19, 2011  
Volume 13, Issue 16  
Pages 4133-4478

Download Cover

OL Organic Letters

In this issue:

> Letters  
> Additions and Corrections

< PREVIOUS ISSUE  
> NEXT ISSUE  
VIEW ALL ISSUES  
ASAPs

Get e-Alerts

Submit Manuscript

Sort By: Page

## LETTERS

### Ruthenium-Catalyzed Oxidative C–H Bond Alkenylations in Water: Expedient Synthesis of Annulated Lactones

Lutz Ackermann and Jola Pospech

Organic Letters 2011, 13, 16, 4153-4155 (Letter)  
Publication Date (Web): July 12, 2011

Abstract Full text PDF

ABSTRACT




### Protection-Free One-Pot Synthesis of 2-Deoxynucleoside 5'-Triphosphates and DNA Polymerization

Julianne Caton-Williams, Matthew Smith, Nicolas Carrasco, and Zhen Huang

Organic Letters 2011, 13, 16, 4156-4159 (Letter)  
Publication Date (Web): July 26, 2011

Abstract Full text PDF

ABSTRACT




### One- and Two-Photon Live Cell Imaging Using a Mutant SNAP-Tag Protein and Its FRET Substrate Pairs

Chong-Jing Zhang, Lin Li, Grace Y. J. Chen, Qing-Hua Xu, and Shao Q. Yao\*

Organic Letters 2011, 13, 16, 4160-4163 (Letter)  
Publication Date (Web): July 25, 2011

Abstract Full text PDF

ABSTRACT




### TPAP-Catalyzed Direct Oxidation of Primary Alcohols to Carboxylic Acids through Stabilized Aldehyde Hydrates

Andrea-Katharina C. Schmidt and Christian B. W. Stark

Organic Letters 2011, 13, 16, 4164-4167 (Letter)  
Publication Date (Web): July 27, 2011

Abstract Full text PDF

ABSTRACT




### Three-Component Synthesis of Neoglycopeptides Using a Cu(II)-Triggered Aminolysis of Peptide Hydrazide Resin and an Azide-Alkyne Cycloaddition Sequence

Jean-Philippe Ebran, Nabil Dendane, and Oleg Melnyk\*

Organic Letters 2011, 13, 16, 4336-4339 (Letter)  
Publication Date (Web): July 18, 2011

Abstract Full text PDF

ABSTRACT




### Highly Efficient Synthesis of Phenols by Copper-Catalyzed Hydroxylation of Aryl Iodides, Bromides, and Chlorides

Kai Yang, Zheng Li, Zhaoyang Wang, Zhiyi Yao, and Sheng Jiang\*

Organic Letters 2011, 13, 16, 4340-4343 (Letter)  
Publication Date (Web): July 25, 2011

Abstract Full text PDF

ABSTRACT




### Total Synthesis of A Novel Tetracyclic Alkaloid, Cassiarin F from the Flowers of *Cassia siamea*

Jun Deguchi, Tomoe Hirahara, Shiori Oshimi, Yusuke Hirasawa, Wivied Ekasari, Osamu Shiota, Toshio Honda, and Hiroshi Morita

Organic Letters 2011, 13, 16, 4344-4347 (Letter)  
Publication Date (Web): July 14, 2011

Abstract Full text PDF

ABSTRACT




### Highly Effective Asymmetric Hydrogenation of Cyclic *N*-Alkyl Imines with Chiral Cationic Ru-MsDPEN Catalysts

Fei Chen, Ziyuan Ding, Jie Qin, Tianli Wang, Yanmei He, and Qing-Hua Fan\*

Organic Letters 2011, 13, 16, 4348-4351 (Letter)  
Publication Date (Web): July 19, 2011

Abstract Full text PDF

ABSTRACT




### Synthesis of the Pyridine Core of Cyclothiazomycin

Yan Zou, Qingyang Liu, and Alexander Deters\*

Organic Letters 2011, 13, 16, 4352-4355 (Letter)  
Publication Date (Web): July 15, 2011

Abstract Full text PDF

ABSTRACT




### Photoresponsive Formation of Pseudo[2]rotaxane with Cyclodextrin Derivatives

Zhibin Wang, Yoshinori Takashima, Hiroyasu Yamaguchi, and Akira Harada

Organic Letters 2011, 13, 16, 4356-4359 (Letter)  
Publication Date (Web): July 20, 2011

Abstract Full text PDF

ABSTRACT



# Total Synthesis of A Novel Tetracyclic Alkaloid, Cassiarin F from the Flowers of *Cassia siamea*


Jun Deguchi,<sup>†</sup> Tomoe Hirahara,<sup>†</sup> Shiori Oshimi,<sup>†</sup> Yusuke Hirasawa,<sup>†</sup> Wiwied Ekasari,<sup>‡</sup> Osamu Shiota,<sup>§</sup> Toshio Honda,<sup>†</sup> and Hiroshi Morita<sup>\*†</sup>

Faculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41 Shinagawa-ku, Tokyo 142-8501, Japan, Faculty of Pharmacy, Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia, and Faculty of Pharmaceutical Sciences at Kagawa Campus, Tokushima Bunri University, 1314-1 Shido, Sanuki City, Kagawa 769-2193, Japan

moritah@hoshi.ac.jp

Received June 22, 2011

## ABSTRACT



A novel alkaloid, cassiarin F (1), which showed potent antiplasmodial activity against *Plasmodium falciparum* in vitro, was isolated from the flowers of *Cassia siamea*, and its structure was elucidated on the basis of 2D NMR analyses. A total synthesis of 1 was also achieved by employing the Suzuki coupling constructing biaryl unit, nucleophilic aromatic substitution, and Houben–Hoesch type ring construction as key steps.

*Cassia siamea* Lam. (Leguminosae) has been used widely in traditional medicine, particularly for treatment of periodic fever and malaria in Indonesia.<sup>1</sup> Recently we isolated cassiarin A (2),<sup>2</sup> an alkaloid with an unprecedented tricyclic skeleton exhibiting potent antiplasmodial activity, and chrobisiamone A,<sup>3</sup> a bischromone, from the leaves of *C. siamea*.

Owing to the attractive biological activity and the unique structural feature of cassiarin A, which showed *in vivo* antimalarial<sup>4</sup> and vasorelaxant activities,<sup>5</sup> we investigated its total synthesis and succeeded in establishing an efficient synthetic strategy.<sup>6</sup> In 2009 we reported the isolation of new types of dimeric alkaloids, cassiarins D and E from the flowers of *C. siamea*.<sup>7</sup>

<sup>†</sup> Hoshi University.

<sup>‡</sup> Airlangga University.

<sup>§</sup> Tokushima Bunri University.


(1) (a) Mbatchi, S. F.; Mbatchi, B.; Banzouzi, J. T.; Bansimba, T.; Nsonde Ntandou, G. F.; Ouamba, J. M.; Berry, A.; Benoit-Vical, F. *J. Ethnopharmacol.* **2006**, *104*, 168–174. (b) Sanon, S.; Ollivier, E.; Azas, N.; Mahiou, V.; Gasquet, M.; Ouattara, C. T.; Nebie, I.; Traore, A. S.; Esposito, F.; Balansard, G.; Timon-David, P.; Fumoux, F. *J. Ethnopharmacol.* **2003**, *86*, 143–147.

(2) Morita, H.; Oshimi, S.; Hirasawa, Y.; Koyama, K.; Honda, T.; Ekasari, W.; Indrayanto, G.; Zaini, N. C. *Org. Lett.* **2007**, *9*, 3691–3693.

(3) Oshimi, S.; Tomizawa, Y.; Hirasawa, Y.; Honda, T.; Widyawaruyanti, A.; Rudyanto, M.; Ekasari, W.; Indrayanto, G.; Zaini, C. N.; Morita, H. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 3761–3763.

(4) Ekasari, W.; Widyawaruyanti, A.; Zaini, N. C.; Syafruddin, D.; Honda, T.; Morita, H. *Heterocycles* **2009**, *78*, 1831–1836.

(5) Morita, H.; Tomizawa, Y.; Deguchi, J.; Ishikawa, T.; Arai, H.; Zaima, K.; Hosoya, T.; Hirasawa, Y.; Matsumoto, T.; Kamata, K.; Ekasari, W.; Widyawaruyanti, A.; Wahyuni, T. S.; Zaini, N. C.; Honda, T. *Bioorg. Med. Chem.* **2009**, *17*, 8234–8240.



Further investigation on extracts from the flowers of *C. siamea* resulted in the isolation of a novel tetracyclic alkaloid, cassiarin F (1). In this paper, we would like to

(6) Rudyanto, M.; Tomizawa, Y.; Morita, H.; Honda, T. *Org. Lett.* **2008**, *10*, 1921–1922.

report the structure elucidation based on spectroscopic analyses and a total synthesis of **1**.

Cassarin F (**1**)<sup>8</sup> was obtained as a yellow amorphous solid, and its molecular formula was established to be C<sub>26</sub>H<sub>21</sub>NO<sub>5</sub> by HRESITOFMS analysis. In addition to <sup>1</sup>H and <sup>13</sup>C NMR data (Table 1), 2D NMR correlations indicated that **1** had a cassiarin A type skeleton with a biaryl moiety, which is the first example of *Cassia* alkaloid (Figure 1). The biaryl moiety of **1** was considered to be racemic from its small optical rotation value and CD spectrum, in which no Cotton effect was observed. Cassiarin F (**1**) showed potent antiplasmodial activity against the chloroquine-sensitive *P. falciparum* strain 3D7 (IC<sub>50</sub> 3.3 μM), while showing no cytotoxicity against human blood premyelocytic leukemia (HL-60, > 50 μM).


**Table 1.** <sup>1</sup>H [ $\delta_{\text{H}}$  (*J*, Hz)] and <sup>13</sup>C NMR Data [ $\delta_{\text{C}}$ ] of Cassiarin F (**1**) in Pyridine-*d*<sub>5</sub> at 300 K

Position		
2		155.7
3		119.1
4		148.3
5		139.9
6	6.91 (1H, d, 2.2)	101.2
7		161.8
8	6.98 (1H, d, 2.2)	100.1
9		154.1
10		111.6
11	6.89 (1H, s)	114.2
12		153.1
13	2.26 (3H, s)	24.6
14	7.01 (1H, d, 1.0)	117.0
2'		141.2
3'	7.09 (1H, d, 1.0)	130.5
4'		138.0
5'		134.0
6'	6.99 (1H, d, 2.4)	108.6
7'		158.7
8'	7.08 (1H, d, 2.4)	102.6
9'		157.2
10'		120.0
11'a	3.60 (1H, d, 13.3)	49.8
11'b	3.65 (1H, d, 13.3)	
12'		206.0
13'	1.97 (3H, s)	29.2
14'	2.20 (3H, s)	21.0

In order to confirm its unique tetracyclic structure, a total synthesis of cassiarin F (**1**) was undertaken. Our retrosynthetic analysis of **1** is outlined in Figure 2. In this approach, the two heterocyclic rings would be constructed at a late stage in the synthesis, because of their susceptibility to acidic and basic reaction conditions.


(7) Oshimi, S.; Deguchi, J.; Hirasawa, Y.; Ekasari, W.; Widyawaruyanti, A.; Wahyuni, T. S.; Zaini, N. C.; Shirota, O.; Morita, H. *J. Nat. Prod.* **2009**, *72*, 1899–1901.

(8) Cassiarin F (**1**): yellow amorphous solid; IR (KBr)  $\nu_{\text{max}}$  3430, 3270, 1710, and 1600 cm<sup>-1</sup>; UV (MeOH)  $\lambda_{\text{max}}$  202 ( $\epsilon$  13000), 245 ( $\epsilon$  3200) and 290 ( $\epsilon$  1400) nm; <sup>1</sup>H and <sup>13</sup>C NMR (Table 1); ESIMS (pos.) *m/z* 428 (M + H)<sup>+</sup>; HRESITOFMS *m/z* 428.1473 (M + H)<sup>+</sup>, calcd for C<sub>26</sub>H<sub>22</sub>NO<sub>5</sub> 428.1498.



**Figure 1.** Selected 2D NMR correlations for cassiarin F (**1**) in pyridine-*d*<sub>5</sub>.

Making the two disconnections shown to **1** led to **3** as the key intermediate for the Houben–Hoesch type ring closure. Two further disconnections generated the three building blocks (phenol **4**, benzonitrile **5**, and boronic acid **6**) needed for the assembly of **3** (Figure 2). It was envisaged that a Suzuki coupling of **5** with **6** would lead to formation of the desired biphenyl unit.<sup>9</sup> Then selective nucleophilic aromatic substitution of the resulting halogenated biaryl derivative with **4** would give the key intermediate **3**.



**Figure 2.** Retrosynthetic analysis for cassiarin F (**1**).

Our synthesis began with preparation of the boronic acid **6**, which was readily accessible from known propenone **7**.<sup>10</sup>


Ketalization of **7** with ethylene glycol and subsequent bromination with *N*-bromosuccinimide (NBS) gave bromide **8** in good yield. The boronic ester was obtained by halogen–metal interchange with *n*-butyllithium and trapping the generated compound with triisopropyl borate. The desired boronic acid **6** was obtained by hydrolysis of the ester with hydrochloric acid. Suzuki coupling of **6** with **5**<sup>11</sup> was carried out in the usual manner to provide the biaryl **9** in 84% yield (Scheme 1).

(9) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483.

(10) Bringmann, G.; Gulder, T.; Reichert, M.; Meyer, F. *Org. Lett.* **2006**, *8*, 1037–1040.


(11) Kristensen, J. L. *Tetrahedron Lett.* **2008**, *49*, 2351–2354.

### Scheme 1. Synthesis of Key Biaryl Intermediate 9



The phenol **4** was readily accessed in a five-step process from the known carboxylic acid **10**.<sup>12</sup> Treatment of **10** with oxalyl chloride provided the corresponding acyl chloride, which was directly reacted with *N,O*-dimethylhydroxylamine in the presence of triethylamine to give the Weinreb amide. The resulting Weinreb amide was treated with MeMgBr to generate the desired methyl ketone **11** in 80% yield from **10**. Debenzylation of **11** via Pd-catalyzed hydrogenation, followed by ketalization of carbonyl group with ethylene glycol, gave the desired phenol **4**. Selective nucleophilic aromatic substitution of **9** with **4** in the presence of cesium carbonate in DMSO gave **3** in excellent yield (Scheme 2).

### Scheme 2. Synthesis of Key Intermediate 3 for Houben–Hoesch Type Ring Closure




Having the desired cyano compound in hand, our attention was focused on construction of the two heterocyclic rings as follows. It has been recognized that both a Brønsted acid and Lewis acid generate ketimine species from the corresponding nitriles in a Houben–Hoesch type cyclization.<sup>13</sup> Therefore, treatment of **3** with acid could lead to the formation of two heterocyclic rings, simultaneously.

(12) Pearson, A. J.; Belmont, P. O. *Tetrahedron Lett.* **2000**, *41*, 1671–1675.

Our first trial, starting from **3** in concentrated sulfuric acid, expectedly only led to extensive decomposition. Fortunately, treatment of **3** with sulfuric acid in acetic acid produced the tetracyclic core **12**. Trials with different combinations of acids or solvents are summarized (Scheme 3). The best result was obtained with the reaction conditions of entry 5. Thus a 30% yield of the tetracyclic core **12** with a side product **13**<sup>14</sup> was obtained using sulfuric acid and acetic acid in a ratio of 1:1.

### Scheme 3. Construction of the Tetracyclic Core of Cassiarin F (1) via a Houben–Hoesch Type Ring Closure<sup>a</sup>



entry	acid or Lewis acid	solvent	ratio <sup>a</sup>	time (h)	yield (%) <sup>b</sup>	
					12	13
1	H <sub>2</sub> SO <sub>4</sub>	AcOH	1:4	12	12	0
2	H <sub>2</sub> SO <sub>4</sub>	CH <sub>2</sub> Cl <sub>2</sub>	1:1	1	12	0
3	H <sub>2</sub> SO <sub>4</sub>	AcOH	2:1	1	23	17
4	TfOH	AcOH	2:1	1	17	19
5	H <sub>2</sub> SO <sub>4</sub>	AcOH	1:1	1	30	10
6	H <sub>2</sub> SO <sub>4</sub>	AcOH	9:10	1	26	7
7	BF <sub>3</sub> ·OEt <sub>2</sub> (1 equiv)	CH <sub>2</sub> Cl <sub>2</sub>		3	0	0
8	TiCl <sub>4</sub> (1 equiv)	CH <sub>2</sub> Cl <sub>2</sub>		3	0	0

<sup>a</sup>Concentration; 1 mL for 10 mg of **3**. <sup>b</sup>Isolated yields.

The tris-*O*-demethylation of aryl methyl ether **12** was achieved by treatment of iodocyclohexane in DMF.<sup>15</sup>

Spectroscopic data (<sup>1</sup>H and <sup>13</sup>C NMR, MS, and IR) and TLC behavior of the synthesized compound were identical to those of the authentic sample of isolated compound.

In summary, we have isolated a novel alkaloid, cassiarin F (**1**), a hybrid compound consisting of cassiarin A and a biphenyl unit with an acetyl moiety. The occurrence of a tetracyclic polysubstituted skeleton with a biaryl unit is very rare. Cassiarin F (**1**) might be biosynthetically produced by a transformation of a

(13) (a) Cameron, D. W.; Deutscher, K. R.; Feutrill, G. I.; Hunt, D. E. *Aust. J. Chem.* **1982**, *35*, 1451–1468. (b) Auricchio, S.; Antonella, B.; Pastormerlo, E.; Ricca, A.; Truscillo, A. M. *Tetrahedron* **1994**, *50*, 7589–7596. (c) Sato, Y.; Yato, M.; Ohwada, T.; Saito, S.; Shudo, K. *J. Am. Chem. Soc.* **1995**, *117*, 3037–3043. (d) Yato, M.; Ohwada, T.; Shudo, K. *J. Am. Chem. Soc.* **1991**, *113*, 691–692. (e) Amer, M. I.; Booth, B. L.; Noori, G. F. M.; Proenca, M. F. J. R. P. *J. Chem. Soc., Perkin Trans. 1* **1983**, 1075–1081.

(14) **13** should be produced by the Friedel–Crafts type reaction with acetic acid. **13** was obtained in the way we predicted by the treatment of **12** with sulfuric acid in acetic acid. The position of methyl ketone of **13** was determined by NOESY correlations.

(15) (a) Zuo, L.; Yao, S.; Wang, W.; Duan, W. *Tetrahedron Lett.* **2008**, *49*, 4054–4056. (b) Since **12** has a labile functionality, methyl ketone, acidic conditions utilizing Lewis acids such as BBr<sub>3</sub> cannot be applied for this purpose. The use of Lewis acids led to significant decomposition of the material.



bischromone, chrobisiamone A. We have also achieved the total synthesis of cassiarin F (**1**) (8 steps, 10% overall yield) by combination of the Suzuki coupling reaction, nucleophilic aromatic substitution, and a Houben–Hoesch type ring construction as key reactions. As such we have confirmed its structure; the strategy with a Houben–Hoesch type ring construction seems to be an innovative approach to the synthesis of the tetracyclic core of **1**.

**Acknowledgment.** This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of

Education, Culture, Sports, Science, and Technology of Japan, and by grants from the Research Foundation for Pharmaceutical Sciences and The Open Research Center Project.

**Supporting Information Available.** Experimental procedures for isolation of cassiarin F; preparation of compounds **1**, **3**, **4**, **6**, **8**, **9**, **11**, **12**, and **S1**; copies of  $^1\text{H}$ ,  $^{13}\text{C}$  NMR, and 2D spectral data for natural cassiarin F; and  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data for compounds **1**, **3**, **4**, **8**, **9**, **11**, **12**, **13**, and **S1**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

# Organic Letters

## COUNTRY

United States

Universities and research institutions in United States

## SUBJECT AREA AND CATEGORY

Biochemistry, Genetics and Molecular Biology  
└ Biochemistry

Chemistry  
└ Organic Chemistry  
└ Physical and Theoretical Chemistry

## PUBLISHER

American Chemical Society

American Chemical Society in Scimago Institutions Rankings

## H-INDEX

**246**

## PUBLICATION TYPE

Journals

## ISSN

15237052, 15237060


## COVERAGE

1999-2021

## INFORMATION

[Homepage](#)  
[How to publish in this journal](#)  
[eic@orglett.acs.org](mailto:eic@orglett.acs.org)


## Quartiles



## SJR

Year	SJR
2008	3.319
2009	3.061
2010	3.014
2011	3.267
2012	3.338
2013	3.157
2014	2.964
2015	3.005
2016	2.985
2017	2.952

## Total Documents





## Source details

Feedback > Compare sources >

### Organic Letters

Scopus coverage years: **from 1999 to Present**

Publisher: American Chemical Society

ISSN: 1523-7060 E-ISSN: 1523-7052

Subject area: Chemistry: Organic Chemistry Chemistry: Physical and Theoretical Chemistry

Biochemistry, Genetics and Molecular Biology: Biochemistry

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

CiteScore 2021 ⓘ  
**10.4**

SJR 2021 ⓘ  
**1.669**

SNIP 2021 ⓘ  
**1.135**

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

**i** Improved CiteScore methodology ×

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▾

$$10.4 = \frac{77,699 \text{ Citations 2018 - 2021}}{7,474 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$8.7 = \frac{53,808 \text{ Citations to date}}{6,201 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly

### CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Chemistry		
└ Organic Chemistry	#14/192	92nd
Chemistry		
└ Physical and Theoretical Chemistry	#13/174	92nd
Biochemistry, Genetics and Molecular Biology		
└ Biochemistry	#48/425	88th