

ISOLATION AND PRESENCE OF ANTIMALARIAL ACTIVITIES OF MARINE SPONGE *Xestospongia* *sp*

by Wiwied Ekasari

Submission date: 17-May-2021 11:07AM (UTC+0800)

Submission ID: 1587569791

File name: Indones J Chem_naskah.pdf (659.48K)

Word count: 3211

Character count: 16897

ISOLATION AND PRESENCE OF ANTIMALARIAL ACTIVITIES OF MARINE SPONGE *Xestospongia* sp.

Murtihapsari^{1*}, Apriani Sulu Parubak¹, Bertha Mangallo¹,
Wiwied Ekasari², Puji Budi Asih³, and Ayu Indah Lestari¹

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Papua State University,
Jl. Gunung Salju Kampus Amban Manokwari, West Papua 98314, Indonesia

²Faculty of Pharmacy, Airlangga University, Kampus B Unair,
Jl. Dharmawangsa Dalam Surabaya East Java 60286, Indonesia

³Eijkman Institute for Molecular Biology, Jl. Diponegoro 69, Jakarta Indonesia

Received April 27, 2013; Accepted August 28, 2013

ABSTRACT

Plasmodium falciparum, the agent of malignant malaria, is one of mankind's most severe scourges, mainly in the tropic world. Efforts to develop preventive vaccines or remedial drugs are handicapped by the parasite's rapid evolution of drug resistance. Here, we presented an advance work on examination of antimalarial component from marine life of *Xestospongia* sp., the study is based on hexane extraction method. The premier result, we obtained five fractions. Among these five fractions, the fourth has the most potent inhibitory against the growth of *P. falciparum* 3D7 with an IC₅₀: 7.13 µg/mL. A compiled spectrum analysis, FTIR, ¹H-NMR and GC-MS, revealed that the fourth fraction consisted abundantly of two secondary metabolites such as flavonoids and triterpenoids. Finally, our results suggest a plausible structure rooted to the base of ibuprofen.

Keywords: sponge; *xestospongia*; papua; antimalarial

ABSTRAK

Plasmodium falciparum adalah parasit penyebab utama penyakit malaria, yang merupakan salah satu penyakit terparah di negara-negara tropis. Sayangnya, upaya pengembangan vaksin preventif atau obat remedial lainnya sedang menghadapi kendala besar karena cepatnya evolusi resistensi obat malaria. Berikut, kami ketengahkan hasil riset kali ini tentang uji aktivitas antimalaria dari spons laut *Xestospongia* sp. yang diekstraksi dengan menggunakan pelarut n-heksana. Hasil pengamatan menghasilkan lima fraksi. Hasil identifikasi menggunakan FTIR, ¹H-NMR dan GC-MS, menunjukkan bahwa fraksi keempat menghasilkan daya hambat tertinggi terhadap pertumbuhan *P. falciparum* 3D7 (IC₅₀: 7,13 µg/mL). Dapat kami laporkan pula bahwa fraksi keempat terdeteksi mengandung senyawa metabolit sekunder, flavonoid dan triterpenoid. Terakhir, penelitian ini menghasilkan suatu struktur dugaan dengan basal ibuprofen.

Kata Kunci: spons; *xestospongia*; papua; antimalaria

INTRODUCTION

Malaria is the most world severe and infectious disease infected 2.4 billion people, across over 100 countries [1-3]. Global cases are reported to about 515 million a year [4-5]. This disease also is responsible of the human deaths about 1.5-3.0 million people per year [6-5], most of the death cases occurred in Africa and the other tropical world [7-8].

A compiled recent investigation have stipulated that Indonesia is home of 20 vectors of malaria associating with four species known of *Plasmodium* [9]. Cases of malaria in Indonesia are estimated to about 89 millions cases by 2011, one people died per a hundred of

thousand cases [5]. This infectious disease has spread across the archipelago including Sumatera, Borneo, Java, Sulawesi, Maluku and Indonesian New Guinea Island. In the eastern most province of archipelago, Papua and West Papua are well known to have an important case of malaria [10]. Current report revealed that malaria is being infected seventy people per a hundred populations; most of them are women and children [5].

In West Papua particularly, searching for a credible drug for antimalarial treatment is scarce. In addition, the most commonly drugs such as chloroquine, primaquine and sulfadoxine-pyrimethamine already reported to be resistance to

* Corresponding author. Tel/Fax : +62-8124863848
Email address: murtihapsari@pnsmail.go.id

P. falciparum and *P. vivax* [9,11-12], elevating the complexity of the malaria treatments [13-14]. Related to this meticulous fact, the chloroquine is now listed to be unrecommended drug by WHO [15-16] since its important toxicities to the woman pregnancies [17-18].

It's generally admitted that the most antimalarial drugs used were extracted from plant herbs and some of their derive medicaments are reportedly resistance to malaria [19-21]. In respond to the drugs already resistance, many researchers have focus to explore a wide range of the habitats including all spectrum of fresh, brackish and salt water [22]. In recent decades, marine sponge (e.g *Xestospongia*, *Haliclona*) are largely used as the potential source of the new drugs, a dozen patenting already obtained from this unique biota, many others researchers are continuing to discover of new compound from other taxa [23-24]. In this context, Indonesia harbors more than 800 species of marine sponge [25-26], in other words this country represents more than 70% of the total marine sponge species in the globe [27-28].

As a part of the global encourage in finding solution for malaria, we are characterizing the potential components of the marine sponge for antimalarial agents. In the present work, we are initially commenced an examination of antimalarial activities based on Papuan sponge of *Xestospongia* sp., a taxon of marine life that has widely distribution in the country.

EXPERIMENTAL SECTION

Materials

The study was based on specimen of marine sponge *Xestospongia* sp., collected in Yapen Island, Papua, Indonesia (1°53'56"S-136°13'14"E). The following chemicals were used including ethyl acetate (p.a), hexane (p.a), gel silica, TLC plat, Liebermann-Burchard, Dragendorff, Bouchard and Mayer, potassium hydroxide, *P. falciparum* strain 3D7 developed by the Eijkman Institute for Molecular Biology and Airlangga University, Roswell Park Memorial Institute (RPMI) 1640 containing the L-glutamine, N-(2-Hydroxyethyl) piperazine-ethanesulfonate acid (HEPES), NaHCO₃ 5%, antibiotic gentamicin sulfate injection, NaCl 0.9% and 3.5%, sorbitol 5%, serum, RBC "O", the anticoagulant Citrate Phosphate Dextrose (CPD), aqua-bides, giemsa, buffer phosphate pH 7.2, alcohol 70% and Bunsen burner.

Instrumentation

The spectrums were performed by using the following instruments: the Fourier Transform Infrared Spectroscopy (FTIR)-Shimadzu Prestige-21; ¹H-NMR. The ¹³C-NMR spectrums were obtained on JEOL JNM

ECA at 500 MHz; MS spectrums were recorded on GC-MS (Agilent MS_5971).

Procedure

The obtained specimens of marine sponge were collected by scuba in about 10 m depth. Six kilograms of sponge were extracted following maceration method by using about twelve liter of hexane. Removing the solvents from samples have been performed through the evaporation method that were used rotary evaporator, afterwards the extracts have been stored at the freezer.

The fractionation method following a previous work [29] with some modification, in the current work, fractionation was carried out in vials by using a combination hexane and ethyl acetate (3:1). The purification was performed by column chromatography. Obtained eluates were separated and identified with Thin Layer Chromatography (TLC) at the wave lengths between 254 and 366 nm.

The culture of *P. falciparum* 3D7 strain (a chloroquine-sensitive clone) was used in the experiment, were maintained by *in vitro* techniques adapted by Trager and Jensen (1976) in human red blood cells (blood type O+) with 5% hematocrit in RPMI 1640 (GIBCO BRL, USA) supplemented with 22.3 mM HEPES (Sigma), Hypoxanthine and Sodium bicarbonate) plus 10% human O+ plasma.

Test were performed in duplicate of 24-well culture plates with an initial and experimental parasitemia of 1% (1 mL suspension each well). Sample was prepared at various concentrations for analysis. Test culture was then incubated in a multigas incubator at 37 °C for 48 h. After incubation, thin blood films were made on glass slide, air dried are fixed in methanol and stained with Giemsa. Fifty percent inhibitory concentration (IC₅₀) was estimated with the approach of Probit Analysis.

The secondary metabolites such as the flavonoids and steroids/triterpenoids, were identified using the phytochemistry method, those method is derived and adapted from classical phytochemistry protocol [30].

Structure of molecule was reconstructed using the ChemDraw Ultra 10.0 [31]. The structure was identified using the Dictionary of Natural Products [32-33] and then performed by ChemDrawUltra 10.0 [31].

RESULT AND DISCUSSION

A total of 15.8 g of extraction was obtained, using hexane solvent, has produced a rendement of 0.003%. Comparing to the same solvent were used on other works, the authors obtained a total rendement about 0.96%. Lowly rendement yielded in the present study is




Fig 1. Percentage inhibition of the three fractions against *Plasmodium falciparum*, fraction II: IC₅₀ 22.45 µg/mL; IV: IC₅₀ 7.13 µg/mL; and V: IC₅₀ 16.69 µg/mL

probably due to semi polar properties of the hexane solvent [34-35].

Our results revealed that *Xestospongia* sp., contained more triterpenoid than steroid, and no alkaloid was detected. Extraction and isolation for herbal products may fully be affected by the solvents were used [36], a non polar solvent (e.g. hexane) may lead to locate the secondary metabolites such as steroid, peptide, quinone and polypropionate [37]. Moreover, a semi polar solvent may contain saponine, alkaloid salt, amino acid [37] and polyhydroxysteroids [38].

In the present study, we obtained an IC₅₀ for the extraction step of the *Xestospongia* is 8.21×10^{-13} µg/mL, these concentrate inhibition are lesser than the strain for control, W2: 3.11 µg/mL, and a strain on D6: 0.013 µg/mL. It's generally admitted that low value of IC₅₀ means that the concentrate has highly potential to inhibit *P. falciparum* [39].

In the fractionation step, we obtained five fractions: I:0.25 g, II:1.46 g, III:1.19 g, IV:1.40 g and fraction V: 0.73 g. The inhibition assay against the parasite of *P. falciparum*, we reported that the fraction IV has highly inhibit *P. falciparum* with a concentration 100 µg/mL (90.6%), while fraction I seems to have less inhibited

(44.62%). Others fractions were unfairly impeded *Plasmodium*.

Analysis of quantitative measurement (IC₅₀), of these three fractions (i.e II, IV, and V) fairly displayed high inhibition against *Plasmodium*. The half maximal inhibitory concentration (IC₅₀) is commonly used as a measure of antagonist compound in inhibiting biological function [40].

The confrontation of percentage inhibition versus log concentration value of each fraction (Fig. 1) demonstrates a linear curve with the range value between 0.56 and 1.34, manifested a correlation level to about > 90. Regarding the data obtained in inhibition assay, we clearly reported that fraction IV exhibits more slightly with IC₅₀ (7.13 µg/mL). The smallest value of IC₅₀ indicates the effectiveness to inhibit a given parasite of *Plasmodium* [40].

A slight value IC₅₀ designates also that a fraction has highly potential to be antimalarial agent. The obtained value of IC₅₀ in the present work was lower than the common standard used worldwide, stipulating the value < 25 µg/mL [41].

An advanced examination through phytochemistry assay using hexane solvent revealed that fraction IV




Fig 2. FTIR spectra of the fraction IV using hexane solvent, displayed evidence of the various type of vibrations




Fig 3. The fragmentation pattern of the GC-MS m/z have obtained from isolation record

manifests more profusion of triterpenoid than flavonoid, while steroid was detected negatively.

The Fourier Transform Infrared Spectroscopy (FTIR) of fraction IV (Fig. 2) revealed four typical substituents as follow, first, wavelength ranging between 2854 and 2924 cm^{-1} , related to the type of vibration as carboxylic acids: O-H, frequencies 3400-2400 cm^{-1} (see Pavia et al. 2001 for designation). Second, wavelength 1705 cm^{-1} , related to carboxylic acids: C=O, frequencies 1725-1700 cm^{-1} . Third, wavelength 3109 cm^{-1} , related to aliphatic double bond or aromatic stretch: C-H, frequencies 3100-3000 cm^{-1} . Fourth, wavelength ranging between 1627-1458 cm^{-1} , related to aromatic ring: C=C, frequencies 1600-1450 cm^{-1} .

The $^1\text{H-NMR}$ spectra of fraction IV (not presented on the figure), showed three substituents metals at the distinct region are δ 0.8 ppm; one pair of geminal proton

δ 1.6 & 2.3 ppm; and one substituent of methylene at δ 4.1 ppm.

We also recorded three methylenes substituents (CH_2) performed by the $^{13}\text{C-NMR}$ spectra, these three recognized alkanes were found at the difference peak regions that are δ 11.6; 14.31 and 14.37 ppm. We documented also the presence of a substituent alcohol at δ 60.6 ppm, a substituent carbonyl δ 171.4 ppm, and the peaks of methylene at δ 20-34 ppm [42].

The analysis of Gas Chromatography-Mass Spectrometry (GC-MS) from fraction IV, represents 25 fragmentation patterns m/z as 41, 55, 65, 71, 77, 83, 91, 101, 107, 113, 119, 129, 135, 145, 152, 161, 171, 177, 185, 191, 199, 206, 212, 219, and 232 (Fig. 3). A comprehensive analysis based on trio spectras dissections, revealed that these identified substituents are the 2-(4-isobutylphenyl) propanoic acid (Fig. 4).




Fig 4. An identified plausible structure, isolated from Papuan marine sponge *Xestospongia* sp., 2-(4-isobutylphenyl) propanoic acid

The plausible structure forming a typical fragment (Fig. 4) was a part from terpenoid, identified with the molecular weight of 206. The base structure of those fragments is ibuprofen with a similarity index reaching to about 99% (time retention 9.22). Those molecular mass can be traced in the fragmentation spectrum of GC-MS [31-33,43].

CONCLUSION

The Papuan marine sponge of *Xestospongia* sp., originate from Yapen Island, diagnosed to have an interesting chemicals components in response to the acute human disease such as malarial. The Yapen Island's sponge also contained of plenty of secondary metabolites including flavonoids and triterpenoids. By using the hexane solvent, we obtained unexpected results, stipulating that fraction IV can inhibit *Plasmodium* with a lower value of IC_{50} . Another important result, we have identified a plausible structure, it proved that the molecule is partly derived from terpenoid properties. Further investigation of the various taxon of the sponge in the area, with a large range collection could be a promising effort to discover much more of the newly compounds for antimalarial.

ACKNOWLEDGEMENT

This study was funded by the grant of Hibah Bersaing UNIPA (1082/42/KU/2011). We are extremely grateful to H. Werimon, T. Pihahai, E. Rumayomi and P. Surbakti for their supports during the field collection, their efforts are greatly appreciated. The authors also thank all staff of Pharmacognosy and Phytochemistry Laboratory UNAIR Surabaya for *in vitro* malarial examination, and the colleagues of Eijkman Institute for Molecular Biology Jakarta, for their carefully and excellent results of extractions. Without their helps we could not have carried out this study.

REFERENCES

- Hay, S.I., Guerra, C.A., Tatem, A.J., Atkinson, P.M., and Snow, R.W., 2005, *Nat. Rev. Microbiol.*, 3, 1, 81–90.
- Kager, P.A., 2002, *Trop. Med. Int. Health*, 7, 12, 1042–1046.
- Bagavan, A., Rahuman, A.A., Kaushik, N.K., and Sahal, D., 2011, *Parasitol. Res.*, 108, 15–22.
- White, N.J., 1992, *Br. J. Clin. Pharmacol.*, 34, 1, 1–10.
- WHO, 2011, <http://www.who.int/malaria.20WHO.htm>, accessed 10 October 2011.
- Hay, S.I., Rogers, D.J., Toomer, J.F., and Snow, R.W., 2000, *Trans. R. Soc. Trop. Med. Hyg.*, 94, 2, 113–127.
- Snow, R.W., Guerra, C.A., Noor, A.M., Myint, H.Y., and Hay, S.I., 2005, *Nature*, 434, 214–217.
- Nayyar, G.M., Breman, J.G., Newton P.N., and Herrington, J., 2012, *Lancet Infect. Dis.*, 12, 6, 488–496.
- Elyazar, I.R.F., Gething, P.W., Patil A.P., Rogayah, H., Kusriastuti, R., Wismarini, D.M., Tarmizi, S.N., Baird, J.K., and Hay, S.I., 2011, *PLoS ONE*, 6, 6, e21315.
- Nagesha, H.S., Din-Syafuruddin, Casey, G.J., Susanti, A.I., Fryauff, D.J., Reeder, J.C., and Cowman, A.F., 2001, *Trans. R. Soc. Trop. Med. Hyg.*, 95, 1, 43–49.
- Suwanarusk, R., Russell, B., Chavchich, M., Chalfein, F., Kenangalem, E., Kosaisavee, V., Prasetyorini, B., Piera, K.A., Barends, M., Brockman, A., Lek-Uthai, U., Anstey, N.M., Tjitra, E., Nosten, F., Cheng, Q., and Price, R.N., 2007, *PLoS ONE*, 2, 10, e1089.
- Baird, J.K., 2004, *Antimicrob. Agents Chemother.*, 48, 11, 4075–4083.
- Klein, E.Y., Smith, D.L., Ramanan, L.R., and Simon, L., 2012, *Proc. R. Soc. B*, 279, 1743, 3834–3842.
- Goldberg, D.E., Siliciano, R.F., and Jacobs, W.R.Jr., 2012, *Cell*, 148, 6, 1271–1283.
- Plowe, C.V., 2005, *Curr. Top. Microbiol. Immunol.*, 295, 55–79.
- Martin, R.E., Marchetti, R.V., Cowan, A.I., Howitt, S.M., Bröer, S., and Kirk, K., 2009, *Science*, 325, 5948, 1680–1682.
- McGready, R., Cho, T., Keo, N.K., Thwai, K.L., Villegas, L., Looareesuwan, S., White, N.J., and Nosten, F., 2001, *Clin. Infect. Dis.*, 33, 12, 2009–2016.
- Tang, C., Godfrey, T., Stawell, R., and Nikpour, M., 2012, *Intern. Med. J.*, 42, 9, 968–978.
- Hyde, J.E., 2005, *Trends Parasitol.*, 21, 11, 494–498.
- Schlitzer, M., 2007, *ChemMedChem*, 2, 944–986.
- Ross and Flaningan, 2006, <http://www.healthatoz.com>, accessed 1 January 2010.

22. Summers, R.L., Nash, M.N., and Martin, R.E., 2012, *Cell. Mol. Life Sci.*, 69, 12, 1967–1995.
23. Inbaneson, S.J., and Ravikumar, S., 2012, *Parasitol. Res.*, 110, 6, 2255–2262.
24. Davis, R.A., Buchanan, M.S., Duffy, S., Avery, V.M., Charman, S.A., Charman, W.N., White, K.L., Shackelford, D.M., Edstein, M.D., Andrews, K.T., Camp, D., and Quinn, R.J., 2012, *J. Med. Chem.*, 55, 12, 5851–5858.
25. McKenna, S.A., Allen, G.R., Suryadi, S., and Rapid Assessment Program 2002, *A marine rapid assessment of the Raja Ampat Islands, Papua Province, Indonesia*, Washington DC: Conservation International, Center for Applied Biodiversity Science, Dept. of Conservation Biology.
26. McCarthy, P.J., and Pomponi, S.A., 2004, *Mar. Biomed. Res.*, 1–2.
27. Allen, G.R., 2008, *Aquat. Conserv.*, 18, 5, 541–556.
28. Treml, E.A., and Halpin, P.N., 2012, *Conserv. Lett.*, 5, 6, 441–449.
29. Swantara, I.M.D., Supriyono, A., and Trinoviani, M., 2007, *J. Kimia*, 1, 1, 67–79.
30. Harborne, J.B., 1987, *Metode Fitokimia*, 2nd ed., ITB Bandung, pp.354.
31. Mills, N., 2006, *J. Am. Chem. Soc.*, 128, 41, 13649–13650.
32. Buckingham, J., 1993, *Dictionary of Natural Products*, Vol. 1, Chapman & Hall/CRC, London, p. 8584.
33. Jin, Z., 2008, *J. Med. Chem.*, 51, 5, 1501.
34. Morris, S., Bersuder, P., Allchin, C.R., Zegers, B., Boon, J.P., Leonards, P.E., and de Boer, J., 2006, *TrAC, Trends Anal. Chem.*, 25, 4, 343–349.
35. Bintang M., 2010, *Biokimia Teknik Penelitian*, Erlangga, Jakarta, Ed. xvi, p.256.
36. Ebada, S., Edrada, R.A., Lin, W., and Proksch, P., 2008, *Nat. Protoc.*, 3, 12, 1820–1831.
37. Rivera, A.P., and Uy, M.M., 2012, *J. Chem.*, 9, 1, 354–358.
38. Chaudary, A., Singla, S.K., and Tandon, C., 2010, *Indian J. Pharmacol. Sci.*, 72, 3, 340–345.
39. Wet, D.H., 2005, An Ethnobotanical and Chemataxonomix Study of South African Menispermaceae, *Thesis*, University of Johannesburg.
40. Baniecki, M.L., Wirth, D.F., and Clardy, J., 2007, *Antimicrob. Agents Chemother.*, 51, 2, 716–723.
41. Ringwald, P., Bickii, J., and Basco, L.K., 1999, *Am. J. Trop. Med. Hyg.*, 61, 2, 187–192.
42. Pavia, D.L., Lampman, G.M., Kriz G.S., 2001, *Introduction to Spectroscopy*, 3rd ed., Brooks/Cole Thomson Learning, p. 390.
43. Blunt, J.W., and Munro, M.H.G., 2007, *Dictionary of Marine Natural Products with CD-ROM*, Boca Raton: Chapman and Hall/CRC.

ISOLATION AND PRESENCE OF ANTIMALARIAL ACTIVITIES OF MARINE SPONGE *Xestospongia* sp

ORIGINALITY REPORT

7%

SIMILARITY INDEX

5%

INTERNET SOURCES

5%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	Nik Najib Nik A Rahman, Takahisa Furuta, Somei kojima, Kikuchi Takane, Mustafa Ali Mohd. "Antimalarial activity of extracts of Malaysian medicinal plants", Journal of Ethnopharmacology, 1999 Publication	2%
2	escholarship.org Internet Source	1%
3	www.hindawi.com Internet Source	1%
4	academicjournals.org Internet Source	<1%
5	ira.le.ac.uk Internet Source	<1%
6	www.paper.edu.cn Internet Source	<1%
7	Malariajournal.biomedcentral.com Internet Source	<1%

8

espace.library.uq.edu.au

Internet Source

<1 %

9

www.isca.in

Internet Source

<1 %

10

www.pnas.org

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

ISOLATION AND PRESENCE OF ANTIMALARIAL ACTIVITIES OF MARINE SPONGE *Xestospongia* sp

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
