

CALLUS INDUCTION OF Sonchus arvensis L. AND ITS ANTIPLASMODIAL ACTIVITY

by Wiwied Ekasari

Submission date: 11-May-2021 02:44PM (UTC+0800)

Submission ID: 1583407754

File name: C-29 - Afr J Infect Dis_naskah.pdf (197.75K)

Word count: 4810

Character count: 25108

CALLUS INDUCTION OF *Sonchus arvensis* L. AND ITS ANTIPLASMODIAL ACTIVITY

Dwi Kusuma Wahyuni^{1*}, Hery Purnobasuki¹, Eko Prasetyo Kuncoro¹, and Wiwied Ekasari²

¹Department of Biology, Faculty of Science and Technology, Airlangga University Surabaya, East Java, 60115, Indonesia. ²Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Airlangga University Surabaya, East Java, 60115, Indonesia.

*Corresponding Author's E-mail: kusumaanwar@yahoo.com

Article History

Received: 8th, May, 2019.

Revised Received: 3rd, Aug, 2019

Accepted: 8th, Aug, 2019

Published Online: 8th, Jan, 2020

Abstract

Background: Malaria is a global health problem that requires urgent need for new drugs. Tempuyung (*Sonchus arvensis* L.) possesses many potential medicinal compounds. As the plant is originally found wild, it is important to reproduce its secondary metabolites by tissue culture. The objectives of this study were to look for effective methods to induce callus from leaf explants of *Sonchus arvensis* L. and to test its *in vitro* antiplasmodial activity.

Materials and Methods: The leaves and petioles of the plant were cultured on Murashige and Skoog (MS) solid medium supplemented with indole acetic-3-acid (IAA), indole-3-butyric acid (IBA), naphthalene acetic acid (NAA), 2,4-dichlorophenoxyacetic acid (2,4-D) and benzyl amino purine (BAP), in light and dark incubations. The best results obtained from callus induction were then treated by with several concentrations of sucrose (1- 5%). The best results from callus induction were then extracted with methanol for antiplasmodial test by Trager and Jensen's method. It was also tested against 3D7 strain of *Plasmodium falciparum*.

Results: The combination of 1mg/L 2,4-D and 0.5 mg/L BAP in dark incubation was the best treatment for callus induction of tempuyung. It produced the best quality of callus and the shortest period for callusing. Sucrose treatment had various effects on leaves callusing, but had no effect on petioles callusing, whereby 4% sucrose was the best treatment for leaves callusing in dark incubation. The methanol extract of the best callus had anti-plasmodial activity with $IC_{50}=0.343 \mu\text{g/mL}$.

Conclusion: Methanol extract of tempuyung callus shows potential as an antimalarial drug but more studies would be required.

Keyword: Antiplasmodial activity, Callus, Growth regulator, Methanol extract, *Plasmodium falciparum*, *Sonchus arvensis*, tempuyung.

List of Abbreviations: MS: Murashige and Skoog, IAA: Indole Acetic-3-Acid, IBA: Indole-3-Butyric Acid, NAA: Naphthalene Acetic Acid, BAP: Benzyl Amino Purine, 2,4-D: 2,4- Dichlorophenoxyacetic Acid, API: Annual Parent Incidence, WHO: World Health Organization, ACT: Artemisinin-Based Combination Therapy, MDG: Millennium Developmental Goal.

Introduction

Plasmodium is a single-cell protozoan parasite that causes malarial disease and transmitted to human through female *Anopheles mosquito* vector. Malaria is well known as a major disease in the world, especially in tropical areas, and endemic in Asia, Africa and Latin American countries. The spread of malaria is stable in populations with a high frequency of malaria inoculation (WHO, 2015). Malaria is a contagious disease that can cause death, anaemia and reduce work productivity (Permenkes, 2013). During the period 2005-2013, malarial incidence throughout Indonesia tended to decline. Eighty per cent (80%) of malarial cases occurred in East Indonesia region where the Annual Parent Incidence (API) value was still highest. Residents living in the Eastern region of Indonesia had a high potential of contracting (Kesehatan, 2016). Millennium Developmental Goal (MDG) aims to stop the spread of malaria and decrease its incidence through indicators that decrease the morbidity and mortality caused by the disease (WHO-World Malaria Report, 2015). Malaria eradication efforts are very important, including malaria prevention, diagnosis and treatment. The first malaria treatment approach used chloroquine. After many reports of drug resistance, malaria treatment was changed to artemisinin-based combination therapy (ACT). Malaria is a complex disease because it depends on 1) parasite aspect, *Plasmodium*; 2) vector aspect, *Anopheles*; 3) environmental aspect; 4) population behaviour; and 5) climate change that helps malarial parasites to be resistant to antimalarial drugs, and

Anopheles line to be resistant to insecticides (Ekasari, 2010). Resistance problems to first line antimalarial drugs caused various obstacles to solving malarial disease treatments. Recently, there are so many challenges and also an opportunity to overcome the spread of the disease. To reduce mortality due to malaria, it is imperative to develop new antimalarial drugs. This condition drove many researchers to find new potential drugs from both synthetic and natural sources.

Eighty per cent (80%) of the world's population still use natural products as antimalarial remedies, and 75% of malaria patients chose traditional medicines to treat the disease (Bagavan *et al.*, 2011). Previously, malarial medicine was derived from *Cinchona succirubra* L., while new antimalarial drug generation, artemisinin, was obtained from *Artemisia annua* L. (Wright, 2005). Tempuyung (*Sonchus arvensis* L.) is an Indonesian traditional medicinal plant that contains various active compounds, such as flavonoids, saponins and polyphenols, which are widely used as antioxidants, hepato-protective, diuretic and potential antimalarial agents (Deylan, 2016; Sukandar and Safitri 2016). Tempuyung can be found easily, but the contents of its active compounds can vary, based on the plantation area. Furthermore, using plant material directly from nature can cause genetic loss (Thomas and Yoichiro, 2010). Tissue culture can be the best option to standardize the products, which may suggest that the amount of its secondary metabolites production is identical per mass unit. *In vitro* culture was one of the technologies used to improve the quality of plants, increase biomass and plant productivity (Sitorus *et al.*, 2011; Ariati *et al.*, 2012).

Plant tissue culture can produce secondary metabolites with a high economic value in a relatively short time, continuously with more consistent and controlled quality, and a higher level of content compared to direct harvest (Sitorus, 2011; Ariati *et al.*, 2012). The factors that influence its success are basic medium (Hashemabadi and Kaviani, 2010), the combination of growth regulators (Shirin *et al.*, 2007; Hoesen *et al.*, 2008; Jahan *et al.*, 2009), plant external conditions, and explant genotypes (Ibrahim *et al.*, 2010; Reddy *et al.*, 2011). Conservation is also the main consideration in using *in vitro* techniques, which is the way we can preserve natural genetic resources. For commercial purposes, secondary metabolites can be produced by using a bioreactor method (Radji, 2005). Based on this background, it is important to find an effective protocol for callus induction from *Sonchus arvensis* L. leaf, and examine its antimalarial activity by *in vitro* assay.

3 Materials and Methods

Plant material

Leaves and petioles were obtained from *Sonchus arvensis* L. located at the park of Faculty of Science and Technology, Airlangga University, and authenticated by Purwodadi Botanical Garden, Indonesian Science Institute. The 2nd and 3rd leaves of the plants were collected to be used as explants.

Explants surface sterilization

Fresh leaves and leaf stalks were washed carefully with running tap-water for 5 minutes. Then, emphasized using a fungicide (500 mg fungicide/500 ml distilled water), shaken continuously for 10 minutes and immediately rinsed twice using distilled water. Finally, the leaves were sterilized for 10 minutes in 10% Chlorox (NaOCl). Explants were washed with sterile distilled water three times.

Media preparation for growth regulator on callus induction and sucrose level on callus biomass

Medium MS (Murashige and Skoog, 1962) was used in the study. For callus induction, MS medium was enriched with 3% sucrose and 0.8 g agar. Furthermore, growth regulators of 1 mg/L Indolyl-3-acetic acid (IAA), Indolyl-3-butyric acid (IBA), naphthalene acetic acid (NAA) and 2,4-dichlorophenoxy-acetic acid (2,4-D) alone and their combinations yellowish-green with 0.5 mg/L benzyl amino purine (BAP). Media acidity was adjusted to pH 5 – 6 by pH paper. Then autoclaved at 121° C and 1 atm pressure for 15 minutes.

The best growth regulator treatment was used for the next callusing and supplemented with various sucrose levels. They were 1%, 2%, 3%, 4%, and 5% of sucrose. All experimental treatments were repeated 5 times. Callus formation time data and callus-forming percentages were recorded every week until the 4th week of culture.

Inoculation of explants material

Inoculation was conducted in the Laminar Air Flow Cabinet. Sterile leaves and petioles were initially cut into many small pieces. Explant size was 1 cm². The surface-sterilized explants were then planted on the media with an abaxial position exposed to the media. The culture bottle was closed using aluminium foil. Next, the culture was incubated in a dark and light place. All callus cultures were harvested at 4 weeks old.

Extractions and In Vitro Antiplasmodial Assays

The best callus was extracted with methanol at room temperature. The methanol extracts were evaporated and used in the *in vitro* anti-plasmodial assays at concentration of 100, 10, 1, 0.1, 0.01 µg/L. The 3D7 strain of *Plasmodium falciparum* was used in the *in vitro* anti-plasmodial assays by Trager and Jensen's method (Trager and Jensen, 1967), that was adopted by Ekasari *et al.* (2009).

Results

Callus Induction

The combinations of plant growth regulators affected tempuyung culture. Various morphologies of the cultures were shown in different combinations. Almost all combinations induced callus and the callus grew into plantlet at the end of the culture. Combination of 2,4D 1 mg/L and BAP 0,5 mg/L was the best combination for tempuyung callusing, and it resulted in the shortest callusing time of 2 weeks in dark incubation (Table 1, Figure 1). At the first growth, callus was coloured yellow with friable texture, but at the end of growth, callus was coloured brown.

Table 1: Time and percentage degree of callusing at 4th week of culture

Growth regulator hormone treatment	Time of callusing (week)		Percentage of callusing (%)	Degree of callusing	Morphology of callus
	Dark incubation	Light incubation			
MS ₀	0	0	0	-	No callus
MS ₁	2	3	100	++	Compact callus, green, the end of the week callus became plantlet
MS ₂	2	2	100	++	Compact callus, green
MS ₃	2	3	100	++	Compact callus, green, at the end of the week callus became plantlet
MS ₄	2	3	100	++	Compact callus, green, at the end of the week callus become plantlet
MS ₅	2	2	100	++	Compact callus, green, at the end of the week callus became plantlet
MS ₆	2	3	100	+	Compact callus, green
MS ₇	2	3	100	++	Compact callus, at the end of the week callus became plantlet
MS ₈	2	3	100	+	Compact callus, green
MS ₉	2	3	100	++++	Friable callus, yellow

Note: MS₁= BAP 0,5 mg/L; MS₂ = IAA 1 mg/L; MS₃=IAA 1mg/L +BAP 0,5 mg/L; MS₄ = IBA 1 mg/L; MS₅ = IBA 1 mg/L + BAP 0,5 mg/L; MS₆ = NAA 1 mg/L; MS₇=NAA 1 mg/L + BAP 0,5 mg/L; MS₈ = 2,4D 1 mg/L; MS₉ = 2,4D 1 mg/L + BAP 0,5 mg/L; MS₀ = without growth regulator hormon




Figure 1: Callus growth of *Sonchus arvensis* L. on MS medium supplemented with 2,4D 1 mg/L + BAP 0,5 mg/L treatment. (A) the first week of culture; (B) the 4th week of culture; (C) the 4th week of culture on MS medium supplemented with BAP 0,5 mg/L

Effect of sucrose level on callus biomass

Sucrose treatment affected callus biomass from leaves explant significantly but no significant difference was obtained on petiole explant according to Duncan Test ($\alpha = 0.05$). The fresh and dry weights of leaves callus varied according to the sucrose concentration added (Table 2 and Table 3). Table 2 shows that 4% sucrose produced the highest of callus dry weight (0.07 ± 0.01 g) in dark incubation, followed by 3% sucrose (0.06 ± 0.02 g), 5% sucrose (0.05 ± 0.00 g), 2% sucrose (0.04 ± 0.01 g) and 1% sucrose (0.03 ± 0.01 g). The average dry weight of leaves in light incubation was 0.15 ± 0.09 g, 0.05 ± 0.02 g, 0.03 ± 0.00 g, 0.02 ± 0.00 g, and 0.02 ± 0.00 g for 5%, 4%, 2%, 3%, and 1% sucrose, respectively.

In vitro antiplasmodial activity of *Sonchus arvensis* L. callus against 3D7 strain of *Plasmodium falciparum*

Results of the *in vitro* anti-plasmodial assays of *Sonchus arvensis* L. callus methanol extract is presented in Table 4. Data of the *in vitro* anti-plasmodial assays were analyzed using Probit analysis to obtain the IC₅₀ value, that is IC₅₀= 0.343 µg/ml.

Discussion

Callus induction is indicated by using the parameters of callus formation time, and the percentage of explant forming callus. The parameter of callus formation time was intended to find out whether the combination growth regulator is faster in callus induction and proliferation. Based on this observation, the callus formation started on 2nd week in dark condition, and 2nd-3rd weeks in the light condition. Callus formation time in the dark is faster than in the light, because callus induction is triggered by auxin that is inactive by light (Taiz and Zieger, 2013).

Table 2: Fresh and dry weights of *Sonchus arvensis* L. by 2,4D 1 mg/L + BAP 0,5 mg/L and sucrose treatment in dark incubation

Treatment	Leaf		Petiole	
	Fresh Weight (g)	Dry weight (g)	Fresh weight(g)	Dry weight(g)
N1	0.47 ^a ±0,05	0.03 ^a ±0,01	0.11 ^a ±0,01	0.01 ^a ±0,00
N2	0.70 ^{bc} ±0,04	0.04 ^b ±0,01	0.33 ^a ±0,23	0.02 ^a ±0,01
N3	0.86 ^c ±0,27	0.06 ^b ±0,02	0.23 ^a ±0,24	0.02 ^a ±0,02
N4	1.07 ^d ±0,16	0.07 ^c ±0,01	0.16 ^a ±0,02	0.02 ^a ±0,00
N5	0.61 ^{ab} ±0,03	0.05 ^b ±0,00	0.13 ^a ±0,03	0.02 ^a ±0,01

Note: N₁ = 1% sucrose; N₂ = 2% sucrose; N₃ = 3% sucrose; N₄ = 4% sucrose; N₅ = 5% sucrose. Number followed by the same superscript letter show no significant differences according to Duncan Test ($\alpha = 0.05$)

Table 3: Fresh and dry weights of *Sonchus arvensis* L. by 2,4D 1 mg/L + BAP 0,5 mg/L and sucrose treatment in light incubation

Treatment	Leaf		Petiole	
	Fresh weight (g)	Dry weight (g)	Fresh weight (g)	Dry weight (g)
N1	0.31 ^{ab} ±0,07	0.02 ^a ±0,00	0.72 ^c ±0,21	0.03 ^a ±0,01
N2	0.54 ^{bc} ±0,13	0.03 ^a ±0,00	0.49 ^{bc} ±0,33	0.02 ^a ±0,02
N3	0.21 ^a ±0,00	0.02 ^a ±0,00	0.56 ^{bc} ±0,29	0.03 ^a ±0,01
N4	0.70 ^c ±0,26	0.05 ^a ±0,02	0.35 ^{ab} ±0,08	0.03 ^a ±0,01
N5	1.30 ^d ±0,25	0.15 ^b ±0,09	0.18 ^a ±0,05	0.02 ^a ±0,00

Note: N₁ = 1% sucrose; N₂ = 2% sucrose; N₃ = 3% sucrose; N₄ = 4% sucrose; N₅ = 5% sucrose. Number followed by the same superscript letter show no significant differences according to Duncan Test ($\alpha = 0.05$).

Table 4: Percentage growth and inhibition of *Sonchus arvensis* L. callus extract toward to 3D7 strain of *Plasmodium falciparum*.

Concentration (µg/mL)	R	% Parasitemia		Growth percentage	inhibition percentage	Average inhibition percentage
		0 hour	48 hours			
control (-)	1	1,22	9,08	7,86	-	-
	2	1,22	9,19	7,97	-	
100	1	1,22	1,45	0,23	97,47	96,59
	2	1,22	1,56	0,34	95,71	
10	1	1,22	2,34	1,12	85,86	83,33
	2	1,22	2,74	1,52	80,81	
1	1	1,22	3,99	2,77	65,03	66,48
	2	1,22	3,76	2,54	67,93	
0.1	1	1,22	6,50	5,28	33,33	33,90
	2	1,22	6,41	5,19	34,47	
0,01	1	1,22	8,12	6,90	12,88	13,13
	2	1,22	8,08	6,86	13,38	

Formation of callus is marked by the presence of cell masses, yellowish-green or light-green, on the wound. In addition, this clot will form a cell mass called "callus". Callus formation is the result of cell division in response to wounds controlled by endogenous and exogenous explants (George and Sherrington, 1992).

Combination of plant growth regulator 2,4-D and BAP was the best choice for callusing. Sen *et al.* (2014) reported that *Achyranthes aspera* L leaf explants supplemented by various concentrations of 2,4-D and BAP showed the best result in callus formation. The results of this study are also consistent with the previous work reported by Rashmi and Trivedi, 2014, which used 2,4-D and BAP at various combinations, ranging from 0.5 to 10 mg/L on *Nerium odorum* leaf explants. Arianto *et al.* (2013) reported that 2,4-D is a growth regulator substance which is commonly utilized in callus induction because its strong action stimulates differentiation of cell, suppresses organogenesis, and maintains growth of callus. This is in agreement with the statement that 2,4-D shows stronger and more optimal activities than other auxins for callus induction (Manuhara, 2014).

Callus morphology is indicated by the colour and texture of the callus. At the beginning of the appearance, the callus is clear yellowish-green and later turns brown in the fourth week of treatment with 1 mg/L of 2,4-D and 0.5 mg/L of BAP, while in the other treatments, callus develops into buds. Changes in callus were budded in treatments other than 2,4-D, because 2,4-D is able to maintain callus growth until the end of culture. The colour change of callus indicates cell activity during cell division (Rasud, 2012). George and Sherintong (1992) also reported that the colour change of callus becomes brown because callus produces phenolic compounds which can be toxic to plants and stop the growth (Hayati *et al.*, 2012). Further confirmation of discoloration of the callus from yellowish-white to brown shows the low cleavage activity of callus cells, so that their growth power drops (Widayanto, 2004).

Moreover, about the texture of callus, callus texture shows the quality of callus which is in accordance with the purpose of this study. Clear white callus shows cell activity that is actively dividing and is embryogenic. Furthermore, the callus will change colour to greenish and grow buds or turn brown and yellow because of its secondary metabolite contents (Manuhara, 2014). Callus texture can vary from compact texture to friable, depending on explant variant, basic medium, growth regulator, and the biotic and abiotic environments of the culture (Sitorus *et al.*, 2011)

Effect of sucrose levels on callus growth varied according to the concentration added (Table 2 and Table 3). The high dry weight of callus was 0.6 g, which came from addition of 4% sucrose given to leaf explant in dark incubation. The lowest dry weight was found to be 0.01g from addition of 1% and 5% sucrose given to petiole explant. Both were incubated without light. In earlier studies, 4% sucrose was found to be the best concentration for callusing of *Tagetes spp.* (Thaneshwari and Aswath, 2018), and on 3% sucrose treatment, fresh weight of *Ficus religiosa* L. callus growth started to increase (Siwach *et al.*, 2011). Conditions of lack of light increase biomass. Light affects the performance of plant hormones. Auxin hormones in the form of auxin growth regulators can work optimally to affect cell division without being blocked by light (George and Sherintong, 1992).

The resulting IC₅₀ value of methanol extracts of callus of *Sonchus arvensis* L. was found at 0.343 µg/ml for 1mg/L 2,4-D and 0.5 mg/L BAP incubated without light. The extract which has IC₅₀ value at the range of 1-10 (µg/ml) was categorized as a material with good anti-plasmodial activity (Weenen, 1990). In China, a plant extract with IC₅₀ value at 0.008-15.38 (µg/ml) is applied as anti-plasmodial material (Aryanti *et al.*, 2006). Kayano *et al.* (2011) stated that an extract is said to be active as anti-malarial agent if it possesses IC₅₀<5 µg/ml, moderate if the IC₅₀ value is 10>IC₅₀>5 µg/ml, and not active if the IC₅₀ is IC₅₀>10 µg/ml.

The minimum inhibitory concentration (IC₅₀) value of methanol extract of *Sonchus arvensis* L. callus was found to be lower than the IC₅₀ value of *Tamarindus indica* fruit methanol extract (IC₅₀=4.786µg/ml); aerial part of *Pavetta corymbosa* methanol extract (IC₅₀=2.024 µg/ml) (Koudouvo *et al.*, 2011); *Chaetomorpha antennina* (IC₅₀=26.37µg/ml) (Ravikumar *et al.*, 2011); *Phyllanthus* at leaf methanol extract (IC₅₀=4.76µg/ml) (Bagavan *et al.*, 2011) against *Plasmodium falciparum*; purified compound of *Cassia fistula* against the chloroquine-sensitive strain of *Plasmodium falciparum* (phytol (IC₅₀=18.9 ± 0.60 µM), lutein (IC₅₀=12.5 ± 0.35 µM), and di-The methanol-glycerol (DLGG) (IC₅₀=5.8 ± 0.27 µM) (Gracea *et al.*, 2013); and methanolic leaf extract of *Christia vespertilionid* (IC₅₀=32.0 µg/mL) against *Plasmodium falciparum* NF-54 (Upadhyaya *et al.*, 2013).

The best combination of growth regulators will be used to develop methods for the production of secondary metabolites from *Sonchus arvensis* L. callus by tissue culture as an anti-malarial agent. Each type of plant will respond differently to the hormones given. The compounds produced are also different. As reported by Zhao *et al.* (2012), callus induction in *Sonchus oleraceus* L. used 2 mg/L of NAA. The utilization of callus *Sonchus arvensis* L. as an antimalarial ingredient still needs to be supported by further research. Screening of metabolites contained in the culture and strengthening the method of production of metabolites in tissue culture in order to replace the original plants that have not been cultivated also require further studies.

Conclusion

From the above discussion, it could be concluded that combination of 1mg/L 2,4-D and 0.5 mg/L BAP produced the highest callus in the shortest time, and that *in vitro* anti-plasmodial activity of methanol extract of *Sonchus arvensis* L. callus has potential to be applied as an anti-plasmodial material.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the study presented in this paper.

Acknowledgments

Financial support for this study is from Kementerian Ristek dan Dikti Republik Indonesia, for which the authors are grateful. We also thank all our colleagues for their excellent assistance.

References

1. Ariati, SN., Waeniati, Muslimin, Suwastika, IN. (2012). Induksi kalus tanaman kakao (*Theobroma cacao* L.) pada media MS dengan penambahan 2,4-D, BAP dan air kelapa. *Natural Science: Journal of Science and Technology*, 1(1): 74-84.
2. Arianto, D., Basri, Z., Bustamil, MU. (2013). Induksi kalus dua klon kakao (*Theobroma cacao* L.) unggul Sulawesi pada berbagai konsentrasi 2,4 dichlorophenoxy acetic acid secara in vitro. *Agrotekbis*, 1(3): 211-220.
3. Aryanti, Tri, ME., Kartika, IP., Rita, MD. (2006). Uji daya antimalarial *Artemisia spp.* terhadap *Plasmodium falciparum*. *Majalah Farmasi Indonesia*, 17 (2): 81-84.
4. Bagavan, A., Rahuman, AA., Kamaraj, C., Kaushik, NK., Mohanakrishnan, D., Sahal, D. (2011). Antiplasmodial activity of botanical extracts against *Plasmodium falciparum*. *Parasitologi Res*, 108: 1099-1109.
5. Delyan, E. (2016). Analisis of composition of volatile compounds of field sow thistle (*Sonchus arvensis* L.) leaves using the method of gas chromatography with mass-detection. *The pharma innovation*. 5(10): 118-121.
6. Ekasari, E. (2010). Identifikasi target biokimiawi senyawa antimalaria hasil isolasi daun *Cassia siamea* Lamk. Ringkasan Disertasi. Program Pasca Sarjana. Universitas Airlangga. Surabaya.
7. Ekasari, W., Widyawaruyanti, A., Zaini, NC., Syafruddin, A., Honda, T., Morita, H. (2009). Antimalarial activity of cassiarin A from the leaves of *Cassia siamea*. *Heterocycles*, 78(7): 1831-1836.
8. Hashemabadi, D., Kaviani, B. (2010). In vitro proliferation of an important medicinal plant Aloe; a method for rapid production. *Australian Journal of Crop Science*, 4(4): 216-222.
9. Hayati, SK., Nurchayati, Y., Setiari, N. (2010). Induksi kalus dari hipokotil Alfafa (*Medicago sativa*) secara in vitro dengan penambahan benzyl amino purine (BAP) dan a-naphtalene acetic acid (NAA). *Bioma*, 12(1): 6-12.
10. Hoesen, DSH., Witjaksono, Sukanto, LA. (2008). Induksi kalus dan organogenesis kultur in vitro *Dendrobium lineale* Rolfe. *Berita Biologi*, 9(3): 333-341
11. George, EF., Sherrington, P. D. (1992). *Plant propagation by tissue culture: Handbook and directory of commercial laboratories*. London: Cambridge University Press.
12. Gracea, MH., Lateganb, C., Graziose, R., Smithb, PJ., Raskinc, I., Lila, MA., (2012). Antiplasmodial activity of the ethnobotanical plant *Cassia fistula*. *Natural Product Communications*, 7: 1-4.
13. Ibrahim, MSD., Rostiana, O., Khumaidah, N. (2010). Pengaruh umur eksplan terhadap keberhasilan pembentukan kalus embriogenik pada kultur meristem jahe (*Zingiber officinale* Rosc.). *Jurnal Litri*, 16(1), 37-42.
14. Jahan, MT., Islam, MR., Khan, R., Mamun, ANK., Ahmed, G., Hakim, L. (2009). In vitro clonal propagation of *Anthurium andraeanum* L.) using callus culture. *Plant Tissue Culture and Biotechnology*, 19(1): 61-69.
15. Kayano, CAV., Lopes, SCP., Bueno, FG., Cabral, EC., Souza-Neiras, W.C., Yamauchi, L.M., Foglio, MA., Eberlin, MN., Mello, CJP., Costa, FTM. (2011). In vitro and in vivo assesment of the antimalarial activity of *Caesalpinia pluviosa*. *Malaria Journal*. 1-12.
16. Kemenkes RI. (2016). Kementerian Kesehatan RI, 2016. Info DATIN Pusat Data dan Informasi Kesehatan Kementerian Kesehatan RI: Malaria. Jakarta: Kementerian Kesehatan Republik Indonesia.
17. Koudouvo, K., Karou, SD., Ilboudo, DP., Kokou, K., Essien, K., Aklikokou, K., de Souza, C., Simpore, J., Gbeassor, M. (2011). In vitro antiplasmodial activity of crude extracts from Togolose medicinal plants. *Asian Pasific Journal of Tropical Medicine*, 129-132.
18. Manuhara, YSW. (2014). *Kapita selekta kultur jaringan tumbuhan*. Surabaya. Airlangga University Press.
19. Murashige, T., Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiologia Plantarum*, 15(3): 437-497.
20. Permenkes. (2013). Peraturan Menteri Kesehatan Republik Indonesia Nomor 5 Tahun 2013 tentang Pedoman tata laksana malaria. Jakarta.
21. Radji, M. (2005). Peranan bioteknologi dan mikroba endofit dalam pengembangan obat herbal. *Majalah Ilmu Kefarmasian*, 2(3): 113 – 126.
22. Rashmi, R., Trivedi, MP. (2014). Effect of various growth hormone concentration and combination on callus induction, nature of callus and callogenic response of *Nerium odorum*. *Applied Biochemistry and Biotechnology*, 172(5): 2562-2570.
23. Ravikumar, S., Ramanathan, G., Gnanadesigan, M., Ramu, A., Vijaykumar, V. (2011). In vitro antiplasmodial activity of methanolic extracts from seaweeds of South West Coast of India. *Asian Pasific Journal of Tropical Medicine*, 862-865.
24. Reddy, JM., Bopaiah, AK., Abhilash, M. (2011). In vitro micropropagation of *Anthurium digitatum* using leaf as explant. *Asian Journal of Pharmaceutical and Health Sciences*, 2(1), 70-74.
25. Rasud, Y. (2012). Induksi kalus dan inisiasi tunas cengkeh (*Syzygium aromaticum* L.) secara in vitro. Palu: Program Pasa Sarjana Universitas Tadulako.
26. Sen, MK., Nasrin, S., Rahman, S., Jamal, AH. (2014). In vitro callus induction and plantlet regeneration of *Achyranthes aspera* L., a high value medicinal plant. *Asian Pacific Journal of Tropical Biomedicine*, 4(1): 40-46.
27. Shirin, F., Hossain, M., Kabir, MF., Roy, M., Sarker, SR. (2007). Callus induction and plant regeneration from internodal and leaf explant of four potato (*Solanum tuberosum* L.) cultivars. *World Journal of Agricultural Sciences*, 3(1):1-6.
28. Siwach, P., Gill, AR., Kumari, K. (2011). Effect of season, explants, growth regulators and sugar level on induction and long term maintenance of callus cultures of *Ficus religiosa* L.. *African Journal of Biotechnology*, 10(24): 4879-4886.

29. Sitorus, EN., Hastuti, ED., Setiari, N. (2011). Induksi kalus binahong (*Basella rubra* L.) secara *in vitro* pada media Murashige& Skoog dengan konsentrasi sukrosa yang berbeda. *Bioma*,13(1): 1-7.
30. Sukandar, EY., Safitri, D. (2016). Evaluation of teratogenic effect of tempuyung (*Sonchus arvensis*) extract on Wistar Rats. *International journal of farmakognosy and phytochemical research*. 8(5):761-766.
31. Taiz L. Zieger E. (2013) *Plant physiology*, 3rd Edition. Sinauer Associates, Inc. p.317.
32. Thaneshwari, Aswath C. (2018) Effect of plant growth regulators and sucrose concentration on callus induction and shoot differentiation from ovary culture of marigold (*Tagetes* spp). *International Journal of Chemical Studies*, 6(1): 618-623.
33. Thomas, TD., Yoichiro, H. (2010). *In vitro* propagation for the conservation of a rare medicinal plant *Justicia gendarussa* Burm.f. by nodal explants and shoot regeneration from callus. *Acta Physiologiae Plantarum*, 32(5): 943-950
34. Trager, W., Jensen, JB. (1976). Human malarial parasites in continous culture. *Science*, 193: 673-676.
35. Upadhyaya, HC., Sisodiab, BS., Cheemab, HS., Agrawalb, J., Palb, A., Darokarb, MP., Srivastavaa, SK. (2013). Novel antiplasmodial agents from *Christia vespertilionis*. *Natural Product Communications*, 8 (11): 1591 – 1594.
36. Weenen, H. (1990). Antimalarial activity of Tanzanian medicinal plants. *J. Planta Medica*, 56: 386-370.
37. Widayanto, W. (2004). Pengaruh 2,4-D dan kinetin terhadap pertumbuhan dan perkembangan eksplan serta kandungan metabolit sekunder kalus jati Belanda (*Guazuma ulmifolia* Lamk.) secara *in vitro*. Surakarta: Fakultas Pertanian Universitas Sebelas Maret Surakarta.
38. WHO. (2015). World Health Organization: Guidelines for the treatment of malaria 3rd Ed. Italy: WHO Press.
39. WHO. (2015). World Health Organization: World Malaria Report 2015. Geneva. WHO Press.
40. Wright, CW. (2005). Traditional antimalarials of the active constituents of some traditional in: traditional medicinal plants and malaria. Edited Willcox. CRC Press. New York.
41. Zhao, M., Zhao, Y., Xing, F., Bai, L. (2012). Components from callus cultures of *Sonchus oleraceus* L. and their cytotoxicity *in vitro*. In: Zhu, E., Sambath, S. (eds) *Information Technology and Agricultural Engineering. Advances in Intelligent and Soft Computing*, vol 134. Springer, Berlin, Heidelberg, 107-112.

CALLUS INDUCTION OF *Sonchus arvensis* L. AND ITS ANTIPLASMODIAL ACTIVITY

ORIGINALITY REPORT

15%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nurul Hidayah Nasution, Irda Wahidah Nasution. " The Effect Of Plant Growth Regulators On Callus Induction Of Mangosteen (L.) ", IOP Conference Series: Earth and Environmental Science, 2019
Publication 2%
- 2 worldwidescience.org
Internet Source 1%
- 3 www.thefreelibrary.com
Internet Source 1%
- 4 Chinnaperumal Kamaraj, Abdul Abdul Rahuman, Selvaraj Mohana Roopan, Asokan Bagavan et al. "Bioassay-guided isolation and characterization of active antiplasmodial compounds from *Murraya koenigii* extracts against *Plasmodium falciparum* and *Plasmodium berghei*", Parasitology Research, 2014
Publication 1%

5

Internet Source

1 %

6

Mary H. Grace, Carmen Lategan, Rocky Graziose, Peter J. Smith, Ilya Raskin, Mary Ann Lila. " Antiplasmodial Activity of the Ethnobotanical Plant ", Natural Product Communications, 2012

Publication

1 %

7

uad.portalgaruda.org

Internet Source

<1 %

8

Christopher P. Keller. "Leaf expansion in Phaseolus: transient auxin-induced growth increase", Physiologia Plantarum, 8/2007

Publication

<1 %

9

link.springer.com

Internet Source

<1 %

10

Seied Mehdi Miri. " Micropropagation, Callus Induction and Regeneration of Ginger (Rosc.) ", Open Agriculture, 2020

Publication

<1 %

11

Hemant Lata, Suman Chandra, Ikhlas Khan, Mahmoud ElSohly. " High Frequency Plant Regeneration from Leaf Derived Callus of High-Tetrahydrocannabinol Yielding L. ", Planta Medica, 2010

Publication

<1 %

12 Brahim Bouzdoudi, R Saïdi, Z Ansari, M Bouras, A Badoc, A Lamarti. "Callus Induction from Carob (*Ceratonia siliqua* L.) Seedlings and Leaves of Mature Tree", Annual Research & Review in Biology, 2017
Publication <1 %

13 notulaebiologicae.ro
Internet Source <1 %

14 Sajad Jafari, Mohammadreza Hassandokht, Madi Taheri. "Optimizing proliferation and assessment of valak morpho-phenological traits; an endangered nutritious *Allium* endemic to Iran", Research Square, 2021
Publication <1 %

15 journals.ashs.org
Internet Source <1 %

16 ri.ufs.br
Internet Source <1 %

17 www.cropj.com
Internet Source <1 %

18 www.tandfonline.com
Internet Source <1 %

19 Bharat Singh, Ram Avtar Sharma. "Secondary Metabolites of Medicinal Plants", Wiley, 2020
Publication <1 %

krishikosh.egranth.ac.in

20

Internet Source

<1 %

21

www.idosi.org

Internet Source

<1 %

22

www.ishs.org

Internet Source

<1 %

23

academicjournals.org

Internet Source

<1 %

24

"Medicinal and Aromatic Plants V", Springer Science and Business Media LLC, 1993

Publication

<1 %

25

Lissette Valverde-Cerdas, Magaly Dufour, Victor M. Villalobos. "In vitro propagation of Pithecellobium saman (Raintree)", In Vitro Cellular & Developmental Biology - Plant, 1997

Publication

<1 %

26

Yadi Wen, Tianming Yao, Yijuan Xu, Harold Corke, Zhongquan Sui. "Pasting, thermal and rheological properties of octenylsuccinylate modified starches from diverse small granule starches differing in amylose content", Journal of Cereal Science, 2020

Publication

<1 %

27

www.acsad.org

Internet Source

<1 %

28

www.biotech-asia.org

<1 %

29

www.innspub.net

Internet Source

<1 %

30

www.stcp.ac.in

Internet Source

<1 %

31

Tasiu Isah. "Influence of plant growth regulators on morphogenic response, biomass and camptothecin production in the callus cultures of *Chonemorpha fragrans* (Moon) Alston", Research Square, 2019

Publication

<1 %

32

Andreza R. Garcia, Ana Claudia F. Amaral, Mariana M. B. Azevedo, Suzana Corte-Real et al. " Cytotoxicity and anti- activity of leaf extracts ", *Pharmaceutical Biology*, 2017

Publication

<1 %

33

Behzad Kaviani, Behnaz Bahari. "In vitro proliferation and ex vitro rooting of microshoots of *lisianthus* using auxin and cytokinin on the solid, liquid and double-phase culture systems", *Acta Scientiarum Polonorum Hortorum Cultus*, 2019

Publication

<1 %

34

Shahira M. Ezzat, Mohamed A. Salem, Ahmed Zayed. "Plants against malarial and typhoid fever", Elsevier BV, 2021

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

CALLUS INDUCTION OF *Sonchus arvensis* L. AND ITS ANTIPLASMODIAL ACTIVITY

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
