

ISSN 2548-5660

BERKALA KEDOKTERAN

MEDICAL AND HEALTH JOURNAL

September 2021, Volume 17, Number 2

Berkala
Kedokteran

Volume 17

Number 2

Halaman

Banjarmasin,
September, 2021

ISSN
2548-5660

BERKALA

KEDOKTERAN

JURNAL KEDOKTERAN & KESEHATAN

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [ARTICLE IN PRESS](#)

[Home](#) > [About the Journal](#) > **Editorial Team**

EDITORIAL TEAM

EDITOR IN CHIEF

Mr. **Eko Suhartono**, (Scopus ID: 56951281500): Lambung Mangkurat University, Banjarmasin, Indonesia, Indonesia

ASSOCIATE EDITOR

Mrs **Ida Yuliana**, Lambung Mangkurat University, Banjarmasin, Indonesia, Indonesia

Mrs **Farida Heriyani**, Lambung Mangkurat University, Banjarmasin, Indonesia, Indonesia

LAYOUT EDITOR

Ms. **Hasni Syahida**, Lambung Mangkurat University, Indonesia

Indexed by :



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

00132383 [View My Stats](#)

Publication Ethic

ISSN Print P-ISSN **1412-0550**

ISSN ONLINE E-ISSN **2548-5660**



USER

Username

Password

Remember me

NOTIFICATIONS

[View](#)

[Subscribe](#)

LANGUAGE

Select Language

English

JOURNAL CONTENT

Search

Search Scope

All

Browse

[By Issue](#)

[By Author](#)

[By Title](#)

[Other Journals](#)

INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)



[Home](#) > [Archives](#) > **Vol 17, No 2 (2021)**

VOL 17, NO 2 (2021)

DOI: <http://dx.doi.org/10.20527/jbk.v17i2>

TABLE OF CONTENTS

ORIGINAL RESEARCH ARTICLES

Ethanol Extract of Areca Nuts was Able to Improve the Histopathological Features of P. Berghei-Induced Liver Damage in MicePDF
79-86

Abstract view : 53 times

DOI: [10.20527/jbk.v17i2.11664](https://doi.org/10.20527/jbk.v17i2.11664)*Alissa Anna Safira, Ika Fidianingsih, Dwi Nur Ahsani***Hand Soap Activity Against the Number of Bacterial Colonies from the Housewife's Hand Swab Samples in a Temporary Landfill in Kelurahan Gadang Banjarmasin**PDF
87-94

Abstract view : 61 times

DOI: [10.20527/jbk.v17i2.11665](https://doi.org/10.20527/jbk.v17i2.11665)*Farida Heriyani, Lia Yulia Budiarti, Widya Nursantari, Ayu Apriliani***Stabilization of the Interaction of the Compounds of *Mallotus paniculatus* Müll.Arg. on Gluthathions S-Transferase (GSTs)**PDF
95-102

Abstract view : 52 times

DOI: [10.20527/jbk.v17i2.11668](https://doi.org/10.20527/jbk.v17i2.11668)*Samsul Hadi***The Difference in Antibacterial Potency of the Infusion of Akar Kuning (*Fibraurea tinctoria* Lour.) against *Shigella* dysenteria and *Salmonella typhi* In Vitro**PDF
103-112

Abstract view : 30 times

DOI: [10.20527/jbk.v17i2.11669](https://doi.org/10.20527/jbk.v17i2.11669)*Mohammad Bakhriansyah, Divae Sandrainy, Agung Biworo***The Knowledge of Lombok Praya-Indonesia Regional Community on the Safety Use of Medicinal Plants**PDF
113-124

Abstract view : 67 times

DOI: [10.20527/jbk.v17i2.11671](https://doi.org/10.20527/jbk.v17i2.11671)*Wiwied Ekasari, Suko Hardjono, Sugijanto Sugijanto, Juni Ekowati, I Nyoman Wijaya, Tri Widlandani***Play Therapy as a Method of Trauma Healing in PTSD Children Victims of Flood Disaster in West Martapura, South Kalimantan**PDF
125-132

Abstract view : 75 times

DOI: [10.20527/jbk.v17i2.11673](https://doi.org/10.20527/jbk.v17i2.11673)*Endang Pertiwiwati, Irfan Maulana, Fitriani Az Zahra, Ida Yuliana***Correlation between Hemoglobin Reticulocytes and Ferritin in Chronic Kidney Disease Patients Undergoing Hemodialysis at PKU Bantul**PDF
133-142

Abstract view : 65 times

DOI: [10.20527/jbk.v17i2.11674](https://doi.org/10.20527/jbk.v17i2.11674)*Muhammad Salman Shalahuddin, Linda Rosita***Correlation of Vitamin D with Ferritin in Pregnant Mothers Chronic Energy Deficiency of the Second Trimester**PDF
143-150

Abstract view : 115 times

DOI: [10.20527/jbk.v17i2.11675](https://doi.org/10.20527/jbk.v17i2.11675)*Hendera Henderi, Salmon Charles P T Siahaan, Irwin Priyatna Kusumah, Heru Cahjono, Ferdinand A Tannus, Nimas Pristiawanto D S, Helen Octavia***Comparison of the Colonies Number of the Germs Formed Post Operation between the Utilization of Single, Double, and Orthopedic Gloves in Closed Fracture Operations More Than 2 Hours**PDF
151-158

Abstract view : 34 times

DOI: [10.20527/jbk.v17i2.11676](https://doi.org/10.20527/jbk.v17i2.11676)**Publication Ethic**ISSN Print P-ISSN **1412-0550**ISSN ONLINE E-ISSN **2548-5660****USER**Username
Password
 Remember me
NOTIFICATIONS[View](#)
[Subscribe](#)**LANGUAGE**

Select Language

English
JOURNAL CONTENTSearch

Search Scope

All

Browse

[By Issue](#)
[By Author](#)
[By Title](#)
[Other Journals](#)**INFORMATION**[For Readers](#)
[For Authors](#)
[For Librarians](#)

THE KNOWLEDGE OF LOMBOK PRAYA-INDONESIA REGIONAL COMMUNITY ON THE SAFETY USE OF MEDICINAL PLANTS

Wiwied Ekasari¹, Suko Hardjono¹, Sugijanto¹, Juni Ekowati¹,
I Nyoman Wijaya¹, Tri Widiandani¹

¹Faculty of Pharmacy, University of Airlangga, Surabaya

Correspondence Email: wiwied-e@ff.unair.ac.id

Abstract: *The use of medicinal plants to maintain or overcome health problems has long been carried out by worldwide community. Along with the development of knowledge, data about the existence of undesirable effects or side effects associated with the use of medicinal plants that are not appropriate are obtained. This research is conducted to determine public knowledge about the safety several medicinal plants use. The study was conducted with a questionnaire technique in Lombok Praya, Indonesia. The results show that knowledge about the safety of the use of medicinal plants in breastfeeding mothers obtains the correct answer by 48%, knowledge about the parts of medicinal plants that are dangerous for consumption by 39% and knowledge about medicinal plants that can increase health risks by 21%. Whereas, the knowledges for the use of medicinal plants in pregnant women and the use of medicinal plants that are not right obtains very good results with correct answers of 79% and 81%. Based on further interviews, these good results are obtained because they are supported more by traditional use of the plant that has already existed in the area. Public understanding of the safety of some medicinal plants in Lombok Praya is still lacking. It is necessary to add knowledge about the use of medicinal plants that requires special attention, especially on plants that are often used by people in the area. Local health agencies need to disseminate information on the safety of medicinal plants that is wider and more sustainable.*

Keywords: *Community, medicinal plant, safety, public understanding.*

INTRODUCTION

The use of medicinal plants to maintain or overcome health problems has long been carried out by Indonesian people. The legacy of the ancestor known as *Jamu*, continues to be preserved and used today. As science advances, research on medicinal plants has increased quite rapidly. In addition to information on pharmacological property that is scientifically proven, another information that develops is the discovery that there are compounds contained in plants that have a detrimental effect on humans under certain conditions. Until now, the world community in general that functions medicinal plants believe that using medicinal plants will always be safer than conventional medicine. However, with the development of knowledge, data obtained show that there are undesirable effects or side effects associated with the use of medicinal plants that are not appropriate.¹ Some undesirable effects, for example, are harmful to pregnant women and breastfeeding mothers, are not for internal use, can increase health risks.²

For this reason, this research will examine public understanding of the use of medicinal plants that need special attention because of the content of the compounds they have which can cause harmful effects to humans.

Until now, knowledge about medicinal plants in the community is more focused on aspects of the benefits or efficacy. Information about the adverse effects that must be considered by the public is not conveyed even many people do not know. For this reason, survey is needed to determine people understanding of this matter, so that adverse effects that are harmful to health can be prevented and can maximize the desired treatment properties. In addition, the results of this study are hoped to be used by authorized health agencies to design policies relating to medicinal plants.

This research will be conducted in the Praya Lombok area. Praya is a sub-district in Central Lombok district, West Nusa Tenggara, Indonesia. The capital of Central Lombok Regency is located between 115°46 – 119°05 East Longitude and 08°10 – 09°05 South Latitude. Lombok district area is known to have Sasak tribe as a native tribe from West Nusa Tenggara, in which its traditions and customs are still preserved, including its medical treatment. Thus, until now the use of traditional medicine is still very entrenched in the area of Lombok which of course is largely colored by the customs of the Sasak tribe.³ This research will focus on medicinal plants that are often used by people in certain conditions. These conditions include the use of medicinal plants in pregnant women and breastfeeding mothers, parts of plants used, how to use, and what conditions are not allowed.

RESEACRH METHOD

The research design used was a non-experimental research design and was descriptive in nature. In this study, a questionnaire was used as a medium to obtain information in the form of data. This research was limited to only being carried out in Lombok Praya, West Nusa Tenggara-Indonesia in July 2019. Demographic physiognomies of the informants were determined and documented through face-to face conversations. Research using that questionnaire was conducted on women in Lombok Praya area, Indonesia, with as many as 62 people with ages of ≥ 30 years old.

RESULTS AND DISCUSSION

All respondents were given a questionnaire containing questions about plants with their respective criteria and choice of answers. Plants given in each criterion were plants that were commonly used in traditional ingredients of Lombok community.

Table 1. Distribution of respondents based on community age

Age group (year)	Total	Percentage (%)
≤ 30	0	0.0
31-40	15	24.2
41-50	36	58.1
51-60	10	16.1
≥ 61	1	1.6
Total	62	100

Table 2. Distribution of respondents based on community education in Lombok Praya in 2019

Highest education	Total	Percentage (%)
High school/equivalent	24	38.7
Diploma I/II/III	4	6.4
Bachelor degree	28	45.2
Master degree	1	1.6
Others	5	8.1
Total	62	100

Based on Table 1, it can be seen that the distribution of the most respondents is in respondents aged 41-50 years (58.1%). This illustrates that the majority of traditional drug consumers in Lombok Praya area are adults. It is known that the older a person is, he or she will be wiser and has more experiences or things figured out and done.⁴ However, from the level of education can be seen in Table 2, the majority of respondents have a good level of education, which is bachelor degree as much as 45.2%. This illustrates that from the

level of education, the respondents adequately know and understand the use of traditional medicines.




Figure 1. Percentage distribution of respondents based on age




Figure 2. Percentage distribution of respondents based on level of education

Table 3. Respondents' Answers on the Questionnaire Given

No.	Plant criteria	Respondents' Answers	
		True (%)	False (%)
1.	Dangerous for pregnant women (fetus/baby)	79	21
2.	Dangerous for breastfeeding mothers	48	52
3.	Plant parts that are dangerous for consumption	39	61
4.	Improper use	81	19
5.	Increase health risks	21	79

The respondents are given a choice of four kinds of medicinal plants that they are familiar with and commonly consume namely papaya fruit (*Carica papaya*), onion

bulb (*Allium cepa*), green tea (*Camellia sinensis*), and cucumber (*Cucumis sativus*). Then, respondents are required to choose medicinal plant that is dangerous to be

consumed for pregnant women because it is dangerous for the fetus or baby. As many as 79% of respondents give the correct answer that is green tea.

Consuming green tea is very beneficial for health, including to help overcome cancer, rheumatoid arthritis, high cholesterol level, cardiovascular disease, infection, and impaired immune function.⁵ Green tea can also reduce body weight, as an anti-inflammatory for arthritis, reduce stress, has anticariogenic and preventive effects of ultraviolet-induced (UV-induced) skin carcinogenesis, and as a topical skin protector against UV radiation.⁶

Green tea is a plant that is considered dangerous if consumed too much by pregnant women. This is because it is related to the availability of folic acid for the fetus. Folic acid supplementation will significantly reduce the small risk of pregnancy and at birth.⁷ Folic acid plays a key role in epigenetic regulation of fetal development programming.⁸ The initial supplementation of folic acid is consistent with the role of folic acid recently in cognitive function and autism spectrum disorders (ASD) prevention.⁹ Folic acid intake is 0–800 µg/day. The results show that the average daily intake of folic acid in excess of 600 µg compared to less than 600 µg during the first month of pregnancy is associated with a reduced risk of ASD.¹⁰ Thus, folic acid deficiency in the fetus or baby can be a risk factor for birth defects and neural tube defects (anencephaly and spina bifida).⁸ Several studies have shown that folic acid deficiency and/or imbalance in folic acid and vitamin B12 intake together with changes in folic acid or vitamin B12 in the blood during and at the end of pregnancy are associated with low birth weight, spontaneous abortion, placental abruption, and congenital abnormalities.^{11,12}

Research shows that catechin, an antioxidant found in tea, can reduce the bioavailability of folic acid. Because the

intake of periconception folic acid has been proven to reduce the risk of spina bifida, tea consumption can put pregnant women at risk because of the possibility of having antifolate properties. Among women with a total intake of folic acid of more than 400 µg, consumption of three or more cups of tea per day is associated with an increased risk of spina bifida.¹³ Another result of study on healthy volunteers aims to investigate the possibility of pharmacokinetic interactions between tea and folic acid shows that average reductions of 0.4 mg of folic acid, green, and black tea of C_{max} of folate serum are 39.2% and 38.6%, and the average $AUC_{0 \rightarrow \infty}$ are respectively 26.6% and 17.9%. At a dose of 5mg of folic acid, average C_{max} of folate serum is reduced by 27.4% and the average $AUC_{0 \rightarrow \infty}$ is significantly reduced by 39.9% by the joint application of green tea. This result shows in vivo interaction between tea and folic acid with low concentration of green and black tea which results in decreased bioavailability of folic acid.¹⁴ Therefore, consuming tea, especially green tea, can cause a decrease in the availability of folic acid for babies.

The survey results show that the respondents in Praya Lombok know that green tea is not good for pregnant women. However, further interview reveals that this good result is further supported by respondents' knowledge that green tea is widely used for weight loss,⁶ so according to them it is definitely not good for pregnant women.

For plants that are harmful for breastfeeding mothers, respondents are given four kinds of plant choices, namely ginger rhizome (*Curcuma xanthorrhiza*), betel leaf (*Piper betel*), carrot (*Daucus carota*) and aloe vera leaves (*Aloe vera*). The survey results state that only 48% of respondents give the correct answer, that is aloe vera.

It is known that fresh *Aloe vera* leaf consists of about 98.5% water, and the rest

are various compounds, including nutrients.¹⁵ It also contains 12 different types of anthraquinone, namely aloin, isobarbaloin, anthracene, emodin, ester of cinnamonic acid, chrysophanic acid, barbaloin, anthranol, aloetic acid, aloe emodin, ethereal oil, and resistanol¹⁶ and also various acids and compounds.¹⁷

Aloe vera has many health benefits. People are very familiar with functioning this plant to maintain health. The benefits of *Aloe vera* include wound healing, protective effect against radiation damage to the skin, anti-inflammatory, effects on the immune system, laxatives, antiviral, antitumor, moisturizer, anti-aging, and antiseptics. Clinical use based on scientific evidence is for seborrheic dermatitis, psoriasis vulgaris, genital herpes, skin burns, diabetes (type 2), HIV infection, cancer prevention, and ulcerative colitis.^{18,19} Several implementations based on traditional use are for alopecia, bacterial and fungal skin infections, chronic foot injuries, parasitic infections, systemic lupus erythematosus, arthritis and tic douloureux.¹⁸ About 23 polypeptides exist in aloe juice help controlling the broad spectrum of diseases and disorders of the immune system. Polypeptides plus anti-tumor agents, aloe emodin, and aloe lectins are now also used in the treatment of cancer.²⁰

However, *Aloe vera* is included in plants that are dangerous for breastfeeding mothers. This is because the anthraquinone glycoside constituent compound from aloe vera leaf juice will be secreted into breast milk.²¹ It is known from the results of the experiment Aloe-emodin-9-anthrone (AE-anthrone) that the decomposition product of barbaloin produced in the rat large intestine causes not only an increase in intestinal water level but also stimulating mucus secretion, which may play an important role in the occurrence of diarrhea.²² Therefore, consumption of aloe vera (aloe and aloin) should be avoided during breastfeeding. A case in which two

days old baby has reported to experience severe diarrhea and dehydration and lose 10-16% of body weight after consuming breast milk. It is known that the baby's mother has consumed 2 Liter of Aloe Vera King® (OKF Corporation, Seoul, Korea) during pregnancy and breastfeeding.²³

The survey results show that 52% of Lombok Praya respondents do not yet know that *Aloe vera* cannot be consumed by breastfeeding mothers. This shows that there needs to be an increase in community knowledge regarding to the consumption of *Aloe vera* in nursing mothers.

In this survey, bitter melon fruit (*Momordica charantia*) is selected as the example of the presence of dangerous parts of plants for consumption, because bitter melon has long been used traditionally to treat diabetes. The respondents are given four kinds of choices of bitter melon, namely the fruit, seeds, roots, and flowers. The survey results mention only 39% of respondents give the correct answer, namely the seed.

Bitter melon fruit has many health benefits, such as hypoglycemic, antiviral, and antineoplastic. Moreover, it can also be functioned for psoriasis, infertility, gastrointestinal cramp, infection, abortifacient, wound healing, antimicrobial, anticancer, and immunomodulator.^{24,25} Bitter melon fruit is also consumed as food and is found as an ingredient in several South Asian curry dishes.²⁴ Fruit extracts have also been shown to have activity in fighting *Helicobacter pylori*, which can cause stomach ulcers.²⁶ However, its seeds are different.

It is known that bitter melon seeds contain twenty-five compounds identified in *M. charantia* seed oil amounting to 90.9% of total oil. The main compound is sesquiterpenes (71.7%), phenylpropanoids (11.0%), and monoterpenes (7.6%), trans nerolidol becomes the main constituent (61.6%).²⁷ It also contains β -sitosterol- β -D-

glucoside and other compounds.²⁸ The results show that two basic glycoproteins isolated from *M. charantia* seed, which are α - and β -momorcharin, are proven effective in inducing early and medium term (post-implantation) of abortion in mice.²⁹⁻³¹ An in vivo study towards Wistar male rats shows that 400-800 mg/kg/day of *M. charantia* (bitter melon) seed ethanol extract causes infertility, which is associated with direct toxic effects on the seminiferous tubules and epididymis, as well as the reduction in seminal and plasma testosterone levels that may significantly impact sperm motility, spermatozoa life, normal morphology of spermatozoa, and the activity of the acrosome membrane.²⁸

In a clinical case, it has been reported that aryl red around bitter melon seeds has produced toxicity; vomiting, diarrhea, and death in a child have been reported.³² There are other reports of children aged three and four who need immediate medical attention after swallowing *M. charantia* leaf water extracts and vines. In both cases, the Sorrosi tea (cerasee) has been given by their mother early in the morning before another food is consumed. Between 1-2 hours after consumption, children experience seizures followed by coma. Blood glucose is in the region of 1 mM (normal range 3.8-5.5 mM). Both patients recover after getting treatment.³³

The survey results show that as many as 61% of respondents in Lombok Praya do not yet know that bitter melon seeds are dangerous for consumption. This shows that there needs to be an increase in public knowledge about the use of hazardous plant parts for consumption even though the other parts are indeed very beneficial for health, in this case especially the use of bitter melon fruit without involving the seeds.

To find out respondents' knowledge about inappropriate use of traditional medicines that can cause harmful effects,

respondents are given four kinds of plant choices, namely celery (*Apium graveolens*), amethyst (*Datura stramonium*), ginger (*Zingiber officinale*), and *sambiloto* (*Andrographis paniculata*). Respondents are told to choose medicinal plant that is consumed not as a drinkable medicine, but for external use. The survey results state that 81% of respondents give the correct answer, namely amethyst.

It is known in amethyst plants that the main active ingredients in this plant are atropine alkaloid and scopolamine.³⁴ As a matter of fact, sixty-four tropane alkaloids have been detected from *D. stramonium*³⁵ besides other compounds.³⁶ Although it is reported that the use of this plant in small amounts has health benefits,^{37,38} but it is more recommended for external use because traditionally it is also more widely used for external use and has been proven scientifically.^{35,39-42}

The use of amethyst plant that is used traditionally is more recommended for use on external use rather than internal use. This is supported by various research results including Diker⁴³ which state that consuming any part of this plant can cause severe anticholinergic reactions which can cause poisoning. *D. stramonium* contains various alkaloids which can cause anticholinergic poisoning if consumed in large concentration⁴⁴ This statement is supported by Kurzbaum⁴⁵ which state that all parts of the *D. stramonium* plant are toxic because of the high level of tropane alkaloids.

In a clinical case, there is even a report of a 6-year-old boy who is diagnosed with anticholinergic poisoning.⁴⁶ Another unusual case of acute *D. stramonium* poisoning in a 3-year-old girl who experiences toxic delirium an hour after swallowing *D. stramonium* plant leaves. This plant poisoning is characterized mainly by toxic delirium which occurs quickly after consumption.⁴⁷

Survey results show that 81% of respondents in Lombok Praya have already known that *D. stramonium* plant should be used for external use and not for consumption. After further observation, this turns out to be because this plant has long been traditionally used in Lombok for external use,³ so people are accustomed to using this plant for external use.

For medicinal plants that can increase health risks by increasing the occurrence of bleeding, respondents are given four kinds of plant choices namely garlic (*Allium sativum*), noni (*Morinda citrifolia*), turmeric (*Curcuma domestica*), and aloe vera (*Aloe vera*). The survey results show that only 21% of the respondents give the correct answer, which is garlic.

It is known that garlic has many benefits for human health. Research that has been conducted, among others, states that garlic has a pharmacological effect on diabetic nephropathy, antibacterial, spermatogenesis, neurological abnormalities, coagulation, and blood pressure. Active against cardiovascular disease, chemo preventive cancer (colorectal cancer, prostate cancer, breast cancer, stomach cancer, oral cancer, liver cancer, leukemia, and anticancer toxicity), hepatotoxicity, cardioprotective (antihypertensive, anti-atherosclerosis, antithrombotic, antithrombotic, and lipid-lowering), hepatotoxicity, cardioprotective (antihypertensive, anti-atherosclerosis, antithrombotic, and lipid-lowering), antimicrobial, antiprotozoal, antifungal, and antiviral.⁴⁸⁻⁵⁰

The content of compounds presents in garlic bulbs include essential oils which are dominated by compounds containing sulfur, especially allyl polysulfides, diallyl disulfide, and diallyl trisulfide.⁵¹ Garlic contains about 65% water, 28% carbohydrates (fructans), 2.3% organosulfur compounds, 2% protein (allinase), 1.2% free amino acids (arginine), and 1.5% fiber, and also contains active

compounds like allicin, diallyl disulphide, diallyl sulphide, diallyl trisulfide, vinyl dithiols, ajoenes, and S-allyl-L-cysteine.^{48,52}

The results show that garlic extract is able to increase fibrinolysis activity compared to controls, with an increase in the time needed for blood clotting.⁵³ Garlic (*Allium sativum* L.) can also increase the pharmacological effects of anticoagulants such as warfarin or fluindione.⁵⁴ The antiplatelet effect of garlic is manifested as a reduction in the adhesion of platelets against collagen, and fibrinogen as a reduction in platelet aggregation in response to epinephrine.⁵⁵ Based on the results of the study above, the consumption of garlic in certain cases can increase the risk of bleeding.

In a clinical case, it is reported that a 55-year-old man who undergoes by-off bypass surgery and experiences diffusion and bruising after surgery. The hematological profile is normal before surgery and the patient has not used antiplatelet for a week before surgery. Based on a known investigation, the patient has been consuming dietary supplements containing fish oil and garlic, where both are known to affect the function of platelets. This requires the surgeon and anesthesiologist to screen all patients before surgery for possible intake of any food supplements taken by the patients.⁵⁶

The survey results show that 79% of respondents in Lombok Praya are not aware of the effect of increasing the risk of bleeding due to consuming garlic. For this reason, there is a need for socialization to reduce the consumption of garlic, especially as traditional medicine for patients who will undergo surgical operations.

CONCLUSION

Public understanding of the safety of some medicinal plants in Lombok Praya is still lacking. It is necessary to increase

knowledge about the safety of using medicinal plants that are often used by the community. Local health agencies need to disseminate information on the safety of medicinal plants that is wider and more sustainable. Thus, it is hoped that the public will benefit from the use of medicinal plants to the maximum and undesirable effects or side effects associated with the use of medicinal plants that are not properly eliminated.

ACKNOWLEDGEMENTS

We would like to thank “DANA RKAT 2019” of Faculty of Pharmacy, Universitas Airlangga scheme for financial support.

REFERENCES

1. Haq I. Safety of medicinal plants. *Pak J Med Res* 2004; 43(4): 203-210.
2. Nasri H, Shirzad H. Toxicity and safety of medicinal plants. *J HerbMed Pharmacol* 2013; 2(2): 21-22.
3. Yamin M, Burhanudin, Jamaluddin, Nasruddin. Treatment and traditional medicine of the Sasak tribe in Lombok. *J Biol Trop* 2018; 18(1): 1-12.
4. Notoatmodjo S. Introduction to health education and health behavioral sciences. Yogyakarta: Andi Offset, 1993.
5. Sinija VR, Mishira HN. Green tea: Health benefits. *J Nutr Environ Med* 2008; 17(4): 232-242.
6. Cooper R, Morre DJ, Morre DM. Medicinal benefits of green tea: part I. Review of noncancer health benefits. *J Altern Complement Med* 2005; 11(3): 521-528.
7. Hodgetts VA, Morris RK, Francis A, Gardosi J, Ismail KM. Effectiveness of folic acid supplementation in pregnancy on reducing the risk of small-for-gestational age neonates: a population study, systematic review and meta-analysis. *Br J Obstet Gynaecol* 2015; 122(4): 478-490.
8. Barua S, Kuizon S, Junaid MA. Folic acid supplementation in pregnancy and implications in health and disease. *J Biomed Sci* 2014; 21(1): 77-85.
9. Meador KJ, Baker GA, Browning N, Cohen MJ, Bromley RL, Clayton-Smith J, et al. Fetal antiepileptic drug exposure and cognitive outcomes at age 6 years (NEAD study): a prospective observational study. *Lancet Neurol* 2013; 12(3): 244-252.
10. Schmidt RJ, Tancredi DJ, Ozonoff S, Hansen RL, Hartiala J, Allayee H, et al. Maternal periconceptional folic acid intake and risk of autism spectrum disorders and developmental delay in the CHARGE (Childhood Autism Risks from Genetics and Environment) case-control study. *Am J Clin Nutr* 2012; 96(1): 80-89.
11. Antony AC. In utero physiology: the role of folic acid in nutrient delivery and fetal development. *Am J Clin Nutr* 2007; 85(2): 598S-603S.
12. Molloy AM, Kirke PN, Brody LC, Scott JM, Mills JL. Effects of folate and vitamin B12 deficiency during pregnancy on fetal, infant, and child development. *Food Nutr Bull* 2008; 29(2 Suppl): S101-115.
13. Yazdy MM, Tinker SC, Mitchell AA, Demmer LA, Werler MM. Maternal tea consumption during early pregnancy and the risk of spina bifida. *Birth Defects Res A Clin Mol Teratol* 2012; 94(10): 756-761.
14. Alemdaroglu NC, Dietz U, Wolfram S, Spahn-Langguth H, Langguth P. Influence of green and black tea on folic acid pharmacokinetics in healthy volunteers: potential risk of diminished folic acid bioavailability. *Biopharm Drug Dispos* 2008; 29(6): 335-348.

15. Guo X, Mei N. *Aloe vera*: a review of toxicity and adverse clinical effects. *J Environ Sci Health C* 2016; 34(2): 77-96.
16. Mahor G, Ali SA. Recent update on the medicinal properties and use of *Aloe vera* in the treatment of various ailments. *Biosci Biotechnol Res Commun* 2016; 9(2): 273-288.
17. Maan AA, Nazir A, Khan MKI, Ahmad T, Zia R, Murid M, et al. The therapeutic properties and applications of *Aloe vera*: a review. *J Herb Med* 2018; 12: 1-10.
18. Surjushe A, Vasani R, Saple DG. *Aloe vera*: a short review. *Indian J Dermatol* 2008; 53(4): 163-166.
19. Zagórska-Dziok M, Furman-Toczek D, Dudra-Jastrzębska M, Zygo K, Stanisławek A, Kapka-Skrzypczak L. Evaluation of clinical effectiveness of *Aloe vera* - a review. *JPCCR* 2017; 11(1): 86-93.
20. Rajeswari R, Umadevi M, Rahale CS, Pushpa R, Selvavenkadesh S, Kumar KPS, et al. *Aloe vera*: the miracle plants its medicinal and traditional uses in India. *J Pharmacogn Phytochem* 2012; 1(4): 118-124.
21. Woolf AD, Gardiner PM, Dvorkin-Camiel L, Maypole J. Constipation. In: Loo M editor. *Integrative medicine for children*. Philadelphia: Saunders Elsevier, 2009; p. 269-280.
22. Ishii Y, Tanizawa H, Takino Y. Studies of aloe. V. Mechanism of cathartic effect. *Biol Pharm Bull* 1994; 17(5): 651-653.
23. Finn D, Sett A, Ryan CA. *Aloe vera* induced toxic colitis in a breast-feeding baby: a case report. *Case Reports in Perinatal Medicine* 2013; 2(1-2): 69-70.
24. Basch E, Gabardi S, Ulbricht C. Bitter Melon (*Momordica charantia*): a review of efficacy and safety. *Am J Health Syst Pharm* 2003; 60(4): 356-359.
25. Tan SP, Kha TC, Parks SE, Roach PD. Bitter Melon (*Momordica charantia* L.) bioactive composition and health benefits: a review. *Food Rev Int* 2015; 32(2): 181-202.
26. Jia S, Shen M, Zhang F, Xie J. Recent advances in *Momordica charantia*: functional components and biological activities. *Int J Mol Sci* 2017; 18(12): 2555.
27. Braca A, Siciliano T, D'Arrigo M, Germano MP. Chemical composition and antimicrobial activity of *Momordica charantia* seed essential oil. *Fitoterapia* 2008; 79(2): 123-125.
28. Tumkiratiwong P, Ploypattarapinyo R, Pongchairerk U, Thong-asa W. Reproductive toxicity of *Momordica charantia* ethanol seed extracts in male rats. *Iran J Reprod Med* 2014; 12(10): 695-704.
29. Law LK, Tam PPL, Yeung HW. Effects of α -trichosanthin and α -momorcharin on the development of peri-implantation mouse embryos. *J Reprod Fertil* 1983; 69(2): 597-604.
30. Chan WY, Tam PPL, Yeung HW. The termination of early pregnancy in the mouse by β -momorcharin. *Contraception* 1984; 29(1): 91-100.
31. Yeung HW, Li WW, Law LK, Chan WY. Purification and partial characterization of momorcharins, abortifacient proteins from the Chinese drug, Kuguazi (*Momordica charantia* seeds). In: Chang HM, Yeung HW, Tso WW, Koo A, editors. *Advances in Chinese medicinal materials research*. Hackensack: World Scientific, 1985; p. 311-318.
32. Shane-McWhorter L. Biological complementary therapies: a focus on botanical products in diabetes. *Diabetes Spectr* 2001; 14(4): 199-208.

33. Raman A, Lau C. Anti-diabetes properties and phytochemistry of *Momordica charantia* L. (Cucurbitaceae). *Phytomedicine* 1996; 2(4): 349-362.
34. Ivancheva S, Nikolova M, Tsvetkova R. Pharmacological activities and biologically active compounds of Bulgarian medicinal plants. In: Imperato F editor. *Phytochemistry: advances in research*. Trivandrum: Research Signpost, 2006; p. 87-103.
35. Soni P, Siddiqui AA, Dwivedi J, Soni V. Pharmacological properties of *Datura stramonium* L. as a potential medicinal tree: an overview. *Asian Pac J Trop Biomed* 2012; 2(12), 1002-1008.
36. Berkov S, Zayed R, Doncheva T. Alkaloid patterns in some varieties of *Datura stramonium*. *Fitoterapia* 2006; 77(3): 179-182.
37. Maheshwari NO, Khan A, Chopade BA. Rediscovering the medicinal properties of *Datura sp.*: a review. *J Med Plants Res* 2013; 7(39): 2885-2897.
38. Sayyed A, Shah M. Phytochemistry, pharmacological and traditional uses of *Datura stramonium* L. review. *J Pharmacogn Phytochem* 2014; 2(5): 123-125.
39. Guarrera PM. Traditional antihelmintic, antiparasitic and repellent uses of plants in Central Italy. *J Ethnopharmacol* 1999; 68(1-3): 183-192.
40. Vijendra N, Kumar KP. Traditional knowledge on ethno-medicinal uses prevailing in the tribal pockets of Chhindwara and Betul districts, Madhya Pradesh, India. *Afr J Pharm Pharmacol* 2010; 4(9): 662-670.
41. Biswas KR, Khan T, Monalisa MN, Swarna A, Ishika T, Rahman M, et al. Medicinal plants used by folk medicinal practitioners of four adjoining villages of Narail and Jessore districts, Bangladesh. *Am Eurasian J Sustain Agric* 2011; 5(1): 23-33.
42. Jonasson M, Afshari R. Chronicle of *Datura* toxicity in 18th and 19th century. *Asia Pac J Med Toxicol* 2016; 5(4): 101-106.
43. Diker D, Markovitz D, Rothman M, Sendovski U. Coma as a presenting sign of *Datura stramonium* seed tea poisoning. *Eur J Intern Med* 2007; 18(4): 336-338.
44. Ertekin V, Selimoglu MA, Altinkaynak S. A combination of unusual presentations of *Datura stramonium* intoxication in a child: rhabdomyolysis and fulminant hepatitis. *J Emerg Med* 2005; 28(2): 227-228.
45. Kurzbaum A, Simsolo C, Kvasha L, Blum A. Toxic delirium due to *Datura stramonium*. *Isr Med Assoc J* 2001; 3(7): 538-539.
46. Korkmaz MF, Bostanci M, Onur H, Cagan E. *Datura stramonium* poisoning: a case report and review of the literature. *Eur Respir J* 2019; 5(1): 186-188.
47. Bouziri A, Hamdi A, Borgi A, Hadj SB, Fitouri Z, Menif K, et al. *Datura stramonium* L. poisoning in a geophagous child: a case report. *Int J Emerg Med* 2011; 4(1): 31.
48. Bhandari PR. Garlic (*Allium sativum* L.): a review of potential therapeutic applications. *IJGP* 2012; 6(2): 118-129.
49. Bayan L, Koulivand PH, Gorji A. Garlic: a review of potential therapeutic effects. *Avicenna J Phytomed* 2014; 4(1): 1-14.
50. Hashemi SA, Ghorbanoghli S, Manouchehri AA, Hatkehlouei MB. Pharmacological effect of *Allium sativum* on coagulation, blood pressure, diabetic nephropathy, neurological disorders, spermatogenesis, antibacterial effects. *AIMS Agric Food* 2019; 4(2): 386-398.

51. Satyal P, Craft JD, Dosoky NS, Setzer WN. The chemical compositions of the volatile oils of garlic (*Allium sativum*) and wild garlic (*Allium vineale*). *Foods* 2017; 6(8): 63.
52. Santhosha SG, Jamuna P, Prabhavathi SN. Bioactive components of garlic and their physiological role in health maintenance: a review. *Food Biosci* 2013; 3: 59-74.
53. Rama NRA, Srividya L, Swamy TP, Prasad VB. Effect of *Allium sativum* (Garlic) extract on blood coagulation and fibrinolysis. *Adv Pharmacol Clin Trials* 2017; 2(1): 000120.
54. Borrelli F, Capasso R, Izzo AA. Garlic (*Allium sativum* L.): adverse effects and drug interactions in humans. *Mol Nutr Food Res* 2007; 51(11): 1386-1397.
55. Steiner M, Li W. Aged garlic extract, a modulator of cardiovascular risk factors: a dose-finding study on the effects of AGE on platelet functions. *J Nutr* 2001; 131(3s): 980S-984S.
56. Bedi HS, Tewarson V, Negi K. Bleeding risk of dietary supplements: A hidden nightmare for cardiac surgeons. *Indian Heart J* 2016; 68(Suppl 2): 249-250.



**KEMENTERIAN RISET, TEKNOLOGI, DAN PENDIDIKAN TINGGI
DIREKTORAT JENDERAL PENGUATAN RISET DAN PENGEMBANGAN**

Jl. M.H Thamrin No. 8 Jakarta Pusat 10340 Gedung BPPT II Lt 19-20

Telepon (021) 316-9804/9805, Faksimil (021) 3101728, 3102368

www.ristekdikti.go.id

SALINAN

KEPUTUSAN DIREKTUR JENDERAL PENGUATAN RISET DAN PENGEMBANGAN
KEMENTERIAN RISET, TEKNOLOGI, DAN PENDIDIKAN TINGGI
REPUBLIK INDONESIA

NOMOR 14/E/KPT/2019

TENTANG

PERINGKAT AKREDITASI JURNAL ILMIAH PERIODE III
TAHUN 2019

DIREKTUR JENDERAL PENGUATAN RISET DAN PENGEMBANGAN
KEMENTERIAN RISET, TEKNOLOGI, DAN PENDIDIKAN TINGGI,

Menimbang : a. bahwa berdasarkan hasil akreditasi jurnal ilmiah yang ditetapkan oleh Tim Akreditasi Jurnal Ilmiah Kementerian Riset, Teknologi, dan Pendidikan Tinggi pada tanggal 7 Mei 2019 dan dalam rangka melaksanakan ketentuan Pasal 6 ayat (5) Peraturan Menteri Riset, Teknologi dan Pendidikan Tinggi Nomor 9 Tahun 2018 tentang Akreditasi Jurnal Ilmiah, perlu menetapkan Peringkat Akreditasi Jurnal Ilmiah Periode III Tahun 2019;

b. bahwa berdasarkan pertimbangan sebagaimana dimaksud pada huruf a, perlu menetapkan Keputusan Direktur Jenderal Penguatan Riset dan Pengembangan Kementerian Riset, Teknologi, dan Pendidikan Tinggi tentang Peringkat Akreditasi Jurnal Ilmiah Periode III Tahun 2019;

Mengingat : 1. Undang-Undang Nomor 12 Tahun 2012 tentang Pendidikan Tinggi (Lembaran Negara Republik Indonesia Tahun 2012 Nomor 158, tambahan Lembaran Negara Republik Indonesia Nomor 5336);

2. Peraturan Pemerintah Nomor 4 Tahun 2014 tentang Penyelenggaraan Pendidikan dan Pengelolaan Perguruan Tinggi (Lembaran Negara Republik Indonesia Tahun 2014, Nomor 16, tambahan Lembaran Negara Republik Indonesia Nomor 5500);

3. Peraturan Presiden Nomor 13 Tahun 2015 tentang Kementerian Riset, Teknologi, dan Pendidikan Tinggi (Lembaran Negara Republik Indonesia Tahun 2015 Nomor 14);

4. Keputusan Presiden Nomor 121/P Tahun 2014 tentang Pembentukan Kementerian dan Pengangkatan Menteri Kabinet Kerja Periode Tahun 2014-2019;

5. Keputusan Presiden Nomor 99/M Tahun 2015 tentang Pemberhentian dan Pengangkatan Dari dan Dalam Jabatan Pimpinan Tinggi Madya di Lingkungan Kementerian Riset, Teknologi, dan Pendidikan Tinggi;

6. Peraturan Menteri Keuangan Republik Indonesia Nomor 32/PMK.02/2018 tentang Standar Biaya Masukan Tahun Anggaran 2019;

7. Peraturan Menteri Riset, Teknologi dan Pendidikan Tinggi Nomor 15 Tahun 2015 tentang Organisasi dan Tata Kerja Kementerian Riset, Teknologi dan Pendidikan Tinggi sebagaimana telah diubah dengan Peraturan Menteri Riset, Teknologi dan Pendidikan Tinggi Nomor 23 Tahun 2019 tentang Perubahan Atas Peraturan Menteri Riset, Teknologi dan Pendidikan Tinggi Nomor 15 Tahun 2015 (Berita Negara Republik Indonesia Tahun 2019 Nomor 238);
8. Peraturan Menteri Riset, Teknologi dan Pendidikan Tinggi Nomor 9 Tahun 2018 tentang Akreditasi Jurnal Ilmiah (Berita Negara Republik Indonesia Tahun 2018 Nomor 428);

MEMUTUSKAN:

- Menetapkan : KEPUTUSAN DIREKTUR JENDERAL PENGUATAN RISET DAN PENGEMBANGAN KEMENTERIAN RISET, TEKNOLOGI, DAN PENDIDIKAN TINGGI TENTANG PERINGKAT AKREDITASI JURNAL ILMIAH PERIODE III TAHUN 2019.
- KESATU : Menetapkan Peringkat Akreditasi Jurnal Ilmiah Periode III Tahun 2019 sebagaimana tercantum dalam Lampiran yang merupakan bagian yang tidak terpisahkan dari Keputusan Direktur Jenderal ini.
- KEDUA : Akreditasi Jurnal Ilmiah sebagaimana dimaksud dalam Diktum KESATU berlaku selama 5 (lima) tahun mulai dari nomor yang ditetapkan dalam lampiran keputusan ini.
- KETIGA : Akreditasi Jurnal Ilmiah sebagaimana dimaksud dalam Diktum KESATU dapat mengajukan kembali kenaikan peringkat setelah menerbitkan minimal 1 (satu) nomor penerbitan.
- KEEMPAT : Setiap jurnal ilmiah wajib mencantumkan masa berlaku akreditasi dengan menuliskan tanggal penetapan dan tanggal akhir masa berlaku akreditasi.
- KELIMA : Apabila dikemudian hari ditemukan ketidaksesuaian dengan Pedoman Akreditasi Jurnal Ilmiah, maka status akreditasi jurnal ilmiah yang bersangkutan dapat dicabut atau diturunkan.
- KEENAM : Keputusan Direktur Jenderal ini mulai berlaku pada tanggal ditetapkan.

Ditetapkan di Jakarta
pada tanggal 10 Mei 2019
DIREKTUR JENDERAL
PENGUATAN RISET DAN PENGEMBANGAN,

TTD.

MUHAMMAD DIMYATI
NIP 195912171984041001

Salinan sesuai dengan aslinya,
Direktorat Jenderal Penguatan Riset dan Pengembangan
Kementerian Riset, Teknologi, dan Pendidikan Tinggi
Kepala Bagian Hukum, Kerjasama, dan Layanan Informasi,

TTD.

Syarip Hidayat
NIP 197306101997031004

3.	Agrovigor: Jurnal Agroekoteknologi	24770353	Program Studi Agroekoteknologi, Universitas Trunojoyo Madura	Usulan baru mulai volume 10, nomor 1, tahun 2017
4.	Akuntabilitas: Jurnal Ilmu Akuntansi	24611190	Fakultas Ekonomi dan Bisnis Universitas Islam Negeri Syarif Hidayatullah Jakarta	Reakreditasi naik peringkat dari peringkat 5 ke 4 Mulai Volume 11, Nomor 2 Tahun 2018
5.	Al-Urban: Jurnal Ekonomi Syariah Filantropi Islam	25812874	Fakultas Ekonomi dan Bisnis Universitas Muhammadiyah Prof. DR. HAMKA	Usulan baru mulai volume 1, nomor 1, tahun 2017
6.	Berkala Kedokteran : Jurnal Kedokteran dan Kesehatan	25485660	Fakultas Kedokteran Universitas Lambung Mangkurat	Reakreditasi tetap di peringkat 4 mulai volume 15, nomor 1, tahun 2019
7.	Biodik : Jurnal Ilmiah Pendidikan Biologi	25800922	Program Studi Pendidikan Biologi PMIPA FKIP Universitas Jambi	Reakreditasi tetap di peringkat 4 mulai volume 4, nomor 2, tahun 2018
8.	<i>Bioscience</i>	2579308X	Universitas Negeri Padang	Usulan baru mulai volume 1, nomor 1, tahun 2017
9.	Biosfer: Jurnal Tadris Biologi	25804960	Prodi Pendidikan Biologi Fakultas Tarbiyah Dan Keguruan Universitas Islam Negeri Raden Intan Lampung	Usulan baru mulai volume 9, nomor 1, tahun 2018
10.	Citizenship : Jurnal Pancasila dan Kewarganegaraan	25795740	Universitas PGRI Madiun	Reakreditasi tetap di peringkat 4 mulai volume 6, nomor 2, tahun 2018
11.	<i>Eduvelop: Journal of English Education and Development</i>	25977148	Universitas Sulawesi Barat	Usulan baru mulai volume 2, nomor 2, tahun 2019
12.	<i>EKONOMIS: Journal of Economics and Business</i>	25978829	Lembaga Penelitian dan Pengabdian pada Masyarakat Universitas Batanghari	Usulan baru mulai volume 2, nomor 1, tahun 2018
13.	<i>ENGLISH FRANCA : Academic Journal of English Language and Education</i>	25803689	Sekolah Tinggi Agama Islam Negeri Curup	Usulan baru mulai volume 2, nomor 1, tahun 2018
14.	Gema Teknologi	2656582X	Sekolah Vokasi, Universitas Diponegoro	Usulan baru mulai volume 19, nomor 2, tahun 2017
15.	GERAM (GERAKAN AKTIF MENULIS)	2580376X	Program Studi Pendidikan Bahasa dan Sastra Indonesia Fakultas Keguruan dan Ilmu Pendidikan Universitas Islam Riau	Usulan baru mulai volume 5, nomor 1, tahun 2017