

ANTICANCER ACTIVITY OF ISOLATED COMPOUNDS FROM *Syzygium aqueum* STEM BARK

by Alfinda Novi Kristanti

Submission date: 12-May-2022 06:46PM (UTC+0800)

Submission ID: 1834545825

File name: RJC-Ei_Ei_complete_-_Copy.pdf (17.01M)

Word count: 3575

Character count: 16579

ANTICANCER ACTIVITY OF ISOLATED COMPOUNDS FROM *Syzygium aqueum* STEM BARK

E.E. Aung^{1,2}, A. N. Kristanti^{3,4,✉}, N. S. Aminah^{3,4}, Y. Takaya⁵,
R. Ramadhan³ and H. T. Aung⁶

¹Ph.D. Student of Mathematics and Natural Sciences, Faculty of Science and Technology,
Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo-60115,
Surabaya, Indonesia

²Department of Chemistry, Yadanarbon University, Amarapura-05063, Mandalay, Myanmar

³Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga,
Komplek Kampus C UNAIR, Jl. Mulyorejo-60115, Surabaya, Indonesia

⁴Biotechnology of Tropical Medicinal Plants Research Group, Universitas Airlangga,
Surabaya, Indonesia

⁵Faculty of Pharmacy, Meijo University, Tempaku-468-8502, Nagoya, Japan.

⁶Department of Chemistry, University of Mandalay, Mandalay, Myanmar.

✉Corresponding Author: alfinda-n-k@fst.unair.ac.id

ABSTRACT

From stem bark of *Syzygium aqueum*, oleanolic acid, and β -sitosterol were isolated successfully. Their structure was identified using spectroscopic methods such as UV-Visible, Fourier Transform Infrared and Nuclear Magnetic Resonance. The isolated compounds were then tested for their anticancer activity against cervical, and breast cancer cell lines using the MTT method. The presence of these two compounds isolated from the stem bark of *S. aqueum* is the first to be reported.

Keywords: *Syzygium aqueum*, Stem Bark, Oleanolic Acid, β -sitosterol, T47D and HeLa Cancer Cell

© RASĀYAN. All rights reserved

INTRODUCTION

In the world, the second biggest killer disease for humans is cancer. According to the World Health Organization, fourteen million people have cancer, and eight million die worldwide.¹ Some cancer treatments are surgery, radiation therapy, chemotherapy, and radiotherapy, which depend on the specific type, location, and stage.² Chemotherapy and radiation therapy are primary clinical treatments at an early stage. Various anticancer medicines are also used in the present. However, these treatment methods and anticancer drugs have severe side effects such as destroying cancer cells and healthy cells.^{1,3}

In therapeutic, phytochemical constituents derived from nature or living organisms play a significant role. Numerous substances have converted into drug candidates as paclitaxel, theophylline, doxorubicin, morphine, penicillin G, digoxin and vitamin A, etc.⁴ Moreover, herbal medicines have been reported as an essential cancer treatment because this treatment is found to have high-quality effects and common side effects. Medicinal plants are recognized as one of the common alternatives for cancer treatment in many countries worldwide.⁵ Medicinal plants contain various kinds of critical phytochemical constituents such as alkaloids, terpenoids, flavonoids, pigments, tannins, etc., which have antioxidant, anti-inflammatory, anticancer, and other bioactivity properties.^{6,7}

The genus *Syzygium* has been reported to be a rich source of phytochemical constituents. This genus belongs to the Myrtaceae family, a woody flowering family consisting of 144 genera and 5500 species. Approximately 1,200 species of this family are species in the genus *Syzygium*.^{8,9} They are widely found in South Asia, Southeast Asia, Australia, New Caledonia, tropical Africa, sub-tropical, and tropical Asia.¹⁰ *S. samarangense*, *S. malaccensis*, *S. aqueum*, *S. aromaticum*, *S. jambos* and *S. polyanthum* are reported as public large. *Syzygium* species are well-known species in this genus, and they are cultivated because of the edible fruit or used as traditional medicine.^{10,11}

S. aqueum, which is native to Indonesia and Malaysia, is better known as watery rose apple or water apple.¹²⁻¹⁴ This species is also used as traditional medicine, especially bark, leaves, and roots.^{13,15} Most researchers have already presented the phenolic content, antioxidant, and antidiabetic activities of leaves and fruit extract of *S. aqueum* using various methods.¹⁶⁻¹⁸ Tannin and flavonoids from the leaves have already been reported in some international research papers.^{19,20} The bioactive components of the stem bark have never been noted. From this study, one terpenoid and one steroid were isolated and anticancer activity was evaluated using an MTT assay.

EXPERIMENTAL

Material and Methods

Plant Materials

The stem bark of *S. aqueum* was obtained from Wage, Taman, Sidoarjo, East Java, Indonesia. The plant material was determined by the Department of Biology, Faculty of Science and Technology, Universitas Airlangga. It was crushed into small pieces like a powder. The powdered sample was then extracted and separated.

Chemicals and Instrumentation

Silica gel 60 (700-200 mesh ASTM) was applied for column chromatography. Analytical TLC was conducted on a pre-coated silica gel 60 F254 (25 Aluminium sheets 20×20 cm, Merck). The U.V. spectrum was recorded on a UV-visible spectrometer (Shimadzu). FT-IR (Fourier Transform Infrared spectroscopy) spectra were measured on Tracer-100 spectrophotometer (Shimadzu). I.R. spectrum (KBr) was showed in cm⁻¹. Nuclear Magnetic Resonance (¹H-NMR and ¹³C-NMR) were recorded on BRUKER 600 Hz in CDCl₃. δ (ppm) was used to measure the chemical shift, and Hertz (Hz) was used to measure the coupling constant (*J*). Anticancer activity assay was measured on ELISA (Enzyme-Linked Immunosorbent Assay) reader

General Procedure

Extraction and Separation

The powdered stem bark of *S. aqueum* (1kg) was extracted with methanol (40L) at room temperature for 3×24 hr. The methanol extract was evaporated with Buchi rotary vacuum evaporator. The extract (450g) was partitioned with n-hexane (6L) three times. N-hexane extract (50g) was put into silica gel column chromatography and then eluted with n-hexane-ethyl acetate mixture, in which polarity was increased gradient. The similar fractions were combined to obtain 19 main fractions (SA-1 to SA-19). Fraction (SA-11) was re-chromatographed with eluent n-hexane: ethyl acetate (7:3), gave 13 fractions (SA-11-1 to SA-11-13). Fraction (SA-11-3) yielded a pure compound, namely compound 1 (22mg). Fraction SA-9 was separated more using column chromatography with a similar eluent (8:2), yielding 6 fractions (SA-9-1 to SA-9-6). Fraction SA-9-5 gave compound 2 (25mg).

Bioassay

Cell Culture

Cervical (HeLa) and breast (T47D) cell lines were collected from ATCC (American Type Culture Collection). Cells were cultured at 37 °C and 5% CO₂ for 24 hours and 100% humidified in medium supplemented with 1% L-glutamine, 10% FBS, and 1% penicillin/streptomycin. The process of cell culture (HeLa and T47D) for the anticancer test was carried out in Cancer Chemoprevention Research Center, Faculty of Pharmacy, Gajah Mada University, Indonesia.

MTT Assay

Anticancer activity test of isolated pure compounds was executed with MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay method. Cancer cells were cultured in a 96well plate at 213 x 10⁴ cell/well density. The compound was prepared as serial dilutions (1.5625-100 μ g/mL) dissolving in 0.1 M of DMSO, and it's 100 μ L were treated with cells at 37 °C in a CO₂ incubator for 24 hours. A 100 μ L of MTT reagent (50 mg in 10 mL phosphate buffered saline) was filled in each well, followed by incubation in CO₂ incubator for 2-4 h until the forming of the purple formazan crystals. The formed formazan was proportional to the total number of viable cells that read its absorbance value at the wavelength 560 nm using ELISA (Enzyme-Linked Immunosorbent Assay) reader. The following formula calculated the % cell viability and IC₅₀ value were calculated using SPSS 17, IBM Analytics.^{21,22} Doxorubicin is used as a standard drug.

$$\% \text{ Cell viability} = \frac{\text{Absorbance of treatment} - \text{Absorbance of media control}}{\text{Absorbance of negative control} - \text{Absorbance of media control}} \times 100\%$$

RESULTS AND DISCUSSION

Compound 1

White powder. The structure was determined according to FTIR spectrum, $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ (including 2D: HMBC, HSQC and COSY). FTIR spectra showed some bands at 3421 cm^{-1} (-OH group), $2931\text{-}2870 \text{ cm}^{-1}$ ($\text{sp}^3 \text{ C-H}$ stretching), 1689 cm^{-1} (C=O). The $^1\text{H-NMR}$ spectra indicated some signals at δH (ppm): 0.73 (m, 1H), 0.76 (s, 3H), 0.78 (s, 3H), 0.91 (s, 3H), 0.93 (s, 6H), 0.99 (s, 3H), 1.09 (m, 2H), 1.14 (s, 3H), 1.17 (m, 1H), 1.22 (m, 1H), 1.30 (m, 1H), 1.35 (m, 2H), 1.43 (m, 1H), 1.56 (m, 2H), 1.59 (m, 1H), 1.61 (m, 2H), 1.63 (m, 4H), 1.74 (m, 1H), 1.79 (m, 1H), 1.91 (m, 1H), 1.99 (m, 1H), 2.83 (dd, $J=4.2$ and 13.8 Hz , 1H), 3.22 (dd, $J=4.2$ and 11.4 Hz , 1H), 5.29 (t, $J=3.5 \text{ Hz}$, 1H). $^{13}\text{C-NMR}$ spectra revealed some signals at δC (ppm): 15.5, 15.3, 17.1, 18.3, 22.9, 23.4, 23.6, 25.9, 27.2, 27.7, 28.1, 30.6, 32.4, 32.6, 33.0, 33.8, 37.0, 38.4, 38.7, 39.2, 41.0, 42.6, 45.8, 46.5, 55.2, 79.0, 122.6, 143.5, 182.8. Moreover, $^1\text{H-}^1\text{H}$ COSY correlations and $^1\text{H-}^{13}\text{C}$ HMBC correlations were used for confirmation of compound 1 structure (Fig-2). Compound 1 was recognized as oleanolic acid ($\text{C}_{30}\text{H}_{48}\text{O}_3$) according to the above data and comparison with literature data.²³ All NMR spectra data (600 MHz, CDCl_3) were presented in Table-1.

Compound 2

White powder. UV spectrum, FTIR spectrum, $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectrum (including 2D: HMBC, HSQC and COSY) were used for identification of its structure. UV spectrum (in MeOH) showed the λ_{max} 260 nm. In FTIR spectrum, some bands were appeared at 3421 cm^{-1} (-OH), $2958\text{-}2868 \text{ cm}^{-1}$ ($\text{sp}^3 \text{ C-H}$ stretching). The signals on $^1\text{H-NMR}$ spectrum were appeared at chemical shift δH (ppm): 0.69 (s, 3H), 0.82 (d, $J=6.8 \text{ Hz}$, 3H), 0.83 (d, $J=6.8 \text{ Hz}$, 3H), 0.85 (m, 3H), 0.93 (m, 5H), 0.97 (m, 1H), 1.01 (s, 3H), 1.03 (m, 1H), 1.09 (m, 3H), 1.17 (m, 3H), 1.26 (m, 2H), 1.28 (m, 1H), 1.33 (m, 1H), 1.36 (m, 1H), 1.50 (m, 3H), 1.58 (m, 1H), 1.67 (m, 1H), 1.85 (m, 3H), 1.98 (m, 2H), 2.01 (m, 2H), 2.24 (m, 1H), 2.30 (m, 1H), 3.53 (m, 1H), 5.36 (m, 1H). The $^{13}\text{C-NMR}$ spectrum suggested as twenty-nine carbon at δC (ppm): 11.8, 11.9, 18.8, 19.0, 19.4, 19.8, 21.1, 23.1, 24.3, 26.1, 28.3, 29.2, 31.6, 31.9, 33.9, 36.2, 36.5, 37.3, 39.8, 42.3, 45.8, 50.2, 56.1, 56.8, 71.8, 121.7, 140.8. Furthermore, the structure of compound 2 was proved with the correlations of $^1\text{H-}^1\text{H}$ COSY and $^1\text{H-}^{13}\text{C}$ HMBC correlations (Fig-2). Compound 2 was recognized as β -sitosterol ($\text{C}_{29}\text{H}_{50}\text{O}$) according to the above data which were compared with literature.²⁴ All NMR spectra data (600 MHz, CDCl_3) were showed in Table-2.




Fig.-1: Molecular Structure of Compound 1 and Compound 2




Fig.-2: HMBC () and COSY () Correlations of Compound 1 and Compound 2

Table-1: NMR Spectral Data of Compound-1 (Oleanolic Acid)

Position	δ_H (ppm) {m, J (Hz)}	δ_C (ppm)	HMBC
1	1.63 2H (<i>m</i>)	38.4	C-25
2	1.61 2H (<i>m</i>)	27.2	-
3	3.22 1H (<i>dd</i> , $J=11.4$ & 4.2 Hz)	79.0	-
4	-	38.7	-
5	0.73 1H (<i>m</i>)	55.2	C-4
6	1.56 1H (<i>m</i>)	18.3	-
	1.35 1H (<i>m</i>)		
7	1.79 1H (<i>m</i>)	32.4	-
	1.59 1H (<i>m</i>)		
8	-	39.2	-
9	1.56 1H (<i>m</i>)	46.5	C-11
10	-	37.0	-
11	1.91 1H (<i>m</i>)	23.4	-
	1.09 1H (<i>m</i>)		
12	5.29 1H (<i>t</i> , $J=3.5$ Hz)	122.6	C-11; C-14
13	-	143.5	-
14	-	42.6	-
15	1.09 1H (<i>m</i>)	27.7	-
	1.74 1H (<i>m</i>)		
16	1.99 1H (<i>m</i>)	22.9	C-17; C-28
	1.63 1H (<i>m</i>)		
17	-	46.5	-
18	2.83 1H (<i>dd</i> , $J=4.2$ & 13.8 Hz)	41.0	-
19	1.63 1H (<i>m</i>)	45.8	C-20; C-18
	1.17 1H (<i>m</i>)		
20	-	30.6	-
21	1.35 1H (<i>m</i>)	33.8	-
	1.22 1H (<i>m</i>)		
22	1.43 1H (<i>m</i>)	32.6	-
	1.30 1H (<i>m</i>)		
23	0.99 3H (<i>s</i>)	28.1	C-3; C-4; C-5; C-24
24	0.78 3H (<i>s</i>)	15.5	C-3; C-4; C-5; C-23
25	0.93 3H (<i>s</i>)	15.3	C-5; C-9
26	0.76 3H (<i>s</i>)	17.1	C-7; C-8; C-9; C-14

Position	δ_H (ppm) {m, <i>J</i> (Hz)}	δ_C (ppm)	HMBC
27	1.14 3H (<i>s</i>)	25.9	C-8; C-13; C-14; C-15
28	-	182.8	-
29	0.91 3H (<i>s</i>)	33.0	C-21; C-30
30	0.93 3H (<i>s</i>)	23.6	C-20; C-21

31
Table-2: NMR Spectral Data of Compound-2 (β -sitosterol)

Position	δ_H (ppm) {m, <i>J</i> (Hz)}	δ_C (ppm)	HMBC
1	1.85 1H, (<i>m</i>)	37.3	C-2; C-5
	1.09 1H (<i>m</i>)		
2	1.98 1H (<i>m</i>)	31.9	-
	1.50 1H (<i>m</i>)		
3	3.53 1H (<i>m</i>)	71.8	-
4	2.24 1H (<i>m</i>)	42.3	C-3; C-5
	2.30 1H (<i>m</i>)		
5	-	140.8	-
6	5.36 1H (<i>m</i>)	121.7	C-4; C-8; C-10
7	1.85 1H (<i>m</i>)	31.6	-
	2.01 1H (<i>m</i>)		
8	1.98 1H (<i>m</i>)	31.9	-
9	0.93 1H (<i>m</i>)	50.2	C-10
10	-	36.5	-
11	1.50 2H (<i>m</i>)	21.1	-
12	2.01 1H (<i>m</i>)	39.8	C-11
	1.17 1H (<i>m</i>)		
13	-	42.3	-
14	0.97 1H (<i>m</i>)	56.8	-
15	1.09 1H (<i>m</i>)	24.3	C-14
	1.58 1H (<i>m</i>)		
16	1.85 1H (<i>m</i>)	28.3	C-17; C-20
	1.26 1H (<i>m</i>)		
17	1.09 1H (<i>m</i>)	56.1	-
18	0.69 3H (<i>s</i>)	11.9	C-12; C-13; C-17
19	1.01 3H (<i>s</i>)	19.4	C-1; C-9
20	1.36 1H (<i>m</i>)	36.2	-
21	0.93 3H (<i>m</i>)	18.8	C-17; C-22
22	1.33 1H (<i>m</i>)	33.9	C-20; C-21
	1.03 1H (<i>m</i>)		
23	1.17 2H (<i>m</i>)	26.1	-
24	0.93 1H (<i>m</i>)	45.8	-
25	1.67 1H (<i>m</i>)	29.2	C-24
26	0.82 3H (<i>d</i> , <i>J</i> = 6.8 Hz)	19.8	C-24; C-25; C-27
27	0.83 3H (<i>d</i> , <i>J</i> = 6.8 Hz)	19.0	C-26
28	1.28 1H (<i>m</i>)	23.1	C-23; C-24; C-29
	1.26 1H (<i>m</i>)		
29	0.85 3H (<i>m</i>)	11.8	C-28

In an anticancer assay, compound 1 was considered moderate active for both cancer cell lines. For compound 2, it was significantly toxic on the HeLa cell line. Its cytotoxicity was almost as toxic as the standard drug (doxorubicin), although it has moderate cytotoxicity against the T47D cell line. Anticancer activity of compound 1 and compound 2 was reported in Table-3, and the percentage of cell viability of these two compounds against HeLa cell and T47D were presented in Fig-3. The pure

compound is deemed as highly toxic when $IC_{50} \leq 4 \mu\text{g/mL}$ and toxic when $IC_{50} \leq 20 \mu\text{g/mL}$. Moderately toxic were considered when IC_{50} 20-100 $\mu\text{g/mL}$ and IC_{50} above 100 $\mu\text{g/mL}$ were suggested as non-toxic in literature.^{25-26.}

Table- 3: Anticancer Activity of Compound 1 and Compound 2 on HeLa and T47D Cell Lines

	$IC_{50} \pm SD$ ($\mu\text{g/mL}$)		
	Doxorubicin	Compound 1 (Oleanolic acid)	Compound 2 (β -sitosterol)
HeLa cell line	2.67 ± 0.25	64.38 ± 0.952	3.43 ± 0.002
T47D cell line	0.035 ± 0.012	20.15 ± 0.011	90.80 ± 5.710

*SD= Standard Deviation




Fig.-3: Percentage of Cell Viability of Compound 1 acid and Compound 2 on HeLa Cell and T47D

CONCLUSION

In this study, oleanolic acid and β -sitosterol were isolated from the stem bark of *S. aqueum*. The two compounds showed anticancer activity against both HeLa and T47D cell lines with MTT assay. Oleanolic acid was more active on the T47D cell line, but β -sitosterol was more active on the HeLa cell line.

ACKNOWLEDGMENT

The first author gratefully thanks Universitas Airlangga Indonesia for a Ph.D. scholarship (ADS). All of the authors are thankful to Indonesia Ministry, in Research, Technology, and Higher Education for the research grant (Penelitian Dasar, 2020), to Faculty of Pharmacy, Meijo University, Tempaku, Nagoya, Japan for NMR data and to Cancer Chemoprevention Research Center, Faculty of Pharmacy, Gajah Mada University, Indonesia for MTT Bioassay.

REFERENCES

1. W. Kooti, K. Servatyari, M. Behzadifar, M. Asadi-Samani, F. Sadeghi, B. Nouri, H. Z. Marzouni, *Journal of Evidence-Based Complementary & Alternative Medicine*, 1(2017), DOI:10.1177/2156587217696927
2. A. Nahata, *International Journal of Clinical Pharmacology & Toxicology*, 6(2), 250(2017), DOI:10.19070/2167-910X-1700042
3. A. B. S. Khumar, K. N. Kumar, C. Raja, M. R. Ezhilarasi, *Rasayan Journal of Chemistry*, 13(2), 1199(2020), DOI:10.31788/RJC.2020.1325495
4. T. Juwitaningsih, I. S. Jahro, I. Dumariris, E. Hermawati, Y. Rukayadi, *Rasayan Journal of Chemistry*, 13(2), 1096 (2020), DOI:10.31788/RJC.2020.1325614
5. N. Tyagi, G. N. Sharma, B. Shrivastava, P. Saxena, N. Kumar, *International Journal of Research and Development in Pharmacy & Life Science*, 6(7), 2732 (2017), DOI:10.21276/IJRDP.227 8-0238.2017
6. P. C. Situmorang, S. Ilyas, S. Hutahaean, R. Rosidah, D. Manurung, *Rasayan Journal of Chemistry*, 13(2), 780 (2020), DOI:10.31788/RJC.2020.1325621
7. A. Hemavathy, P. Shanthi, C. Sowndharya, S. T. Sundari, K. Priyadharshni, *International Journal of Pharmacognosy and Phytochemical Research*, 11(3), 199(2019).

8. A. H. Memon, Z. Ismail, F. S. R. Al-Suede, A. F. A. Aisha, M. S. R. Hamil, M. A. A. Saeed, M. Laghari, A. M. S. A. Majid, *Molecules*, **20**, 14212(2015), DOI:10.3390/molecules200814212
9. T. N. C. Vasconcelos, C. E. B. Proenca, B. Ahmad, D. S. Aguilar, R. Aguilar, B. S. Amorim, K. Campbell, I. R. Costa, P. S. De-Carvalho, J. E. Q Faria, A. Giaretta, P. W. Kooij, D. F Lima, F. F. Mazine, B. Peguero, G. Prenner, M. F. Santos, J. Soewarto, A. Wingyer, E. J. Lucas, *Molecular Phylogenetics and Evolution*, **109**, 113(2017), DOI:10.1016/j.ympev.2017.01.002
10. D. Mudiana, *Biodiversitas*, **17(2)**, 733(2016), DOI:10.13057/biodiv/d170248
11. B. Abera, L. Adane, F. Mamo, *Journal of Pharmacognosy and Phytochemistry*, **7(2)**, 3104(2018).
12. T. Manaham, S. Chakravarthi, A. K. Radhakrishnan, U. D. Palanisamy, *Toxicology Reports*, **1**, 718(2014), DOI:10.1016/j.toxrep.2014.09.006
13. H. Osman, A. A. Rahim, N. M. Isa, N. M. Bakhir, *Molecules*, **14**, 970(2009), DOI:10.3390/molecules14030970
14. M. Sobeh, M. F. Mahmoud, G. Petruk, S. Rezaq, M. L. Ashour, F. S. Youssel, A. M. El-Shazly, D. M. Monti, A. B. Abde-Naim, M. Wink, *Frontiers in Pharmacology*, **9**, 566(2018), DOI:10.3389/fphar.2018.00566
15. U. D. Palanisamy, L. T. Ling, T. Manaharan, V. Sivapalan, T. Subramaniam, M. H. Helme, T. Masilamani, *International Journal of Cosmetic Science*, **33**, 269(2011), DOI:10.1111/j.1468-2494.2010.00637.x
16. N. M. Saptarinli, I. E. Herawati, *Asian Journal of Pharmaceutical and Clinical Research*, **54**(2017), DOI:10.22159/ajpcr.2017.v10s2.19487
17. A. S. L. Lim, M. S. Rabeta, *Intentional Food Research Journal*, **20(2)**, 673(2013).
18. U. D. Palanisamy, T. Manaharan, *International Journal of Medicinal Plants and Natural Product*, **1098**, 13(2015), DOI:10.17660/ActaHortic.2015.1098.1
19. T. Manaharan, D. Appleton, H. M. Cheng and U. D. Palanisamy, *Food Chemistry*, **132**, 1802(2012), DOI:10.1016/j.foodchem.2011.11.147
20. G. Nonaka, Y. Aiko, K. Aritake and I. Nishioka, *Chemical and Pharmaceutical Bulletin*, **40(10)**, 2671(1992).
21. http://ccrc.farmasi.ugm.ac.id/wp-content/uploads/10_sop-uji-sitotoksik-metode-mtt.pdf, Cancer Chemoprevention Research Centre (CCRC), Prosedur Tetap Uji Sitotoksik Metode MTT, Faculty of Pharmacy Gadjah Mada University (UGM), Yogyakarta, (Text by Indonesia Language), (2014)
22. R. Mutiah, C. Indradmojo, H. H. Dwi, T. P. Griana, A. Listyana, R. Ramdhani, *Journal of Applied Pharmaceutical Science*, **7(10)**, 138(2017), DOI:10.7324/JAPS.2017.71020
23. D. Martins, L. L. Carrion, D. F. Ramos, K. S. Salome, P. E. A. Silva, A. Barison, C. V. Nunez1, *Hindawi Publishing Corporation Biomed Research International*, **2013**, 1(2013), DOI:10.1155/2013/605831
24. N. Kamal, C. Clements, A. I. Gray, R. Edrada-Ebel, *Journal of Applied Pharmaceutical Science*, **6(01)**, 102(2016), DOI:10.7324/JAPS.2016.600117
25. S. Sriwiryajan, T. Ninpesh, Y. Sukpondma, T. Nasomyon, P. Graidist, *Tropical Journal of Pharmaceutical Research*, **13(6)**, 192(2014), DOI: 10.4314/tjpr.v13i6.14
26. R. A. Fikroh, S. Matsjeh, C. Anwar, *Molekul*, **15(1)**, 34(2020), DOI:10.20884/1.jm.2020.15.1.558 [RJC-6106/2020]

ANTICANCER ACTIVITY OF ISOLATED COMPOUNDS FROM Syzygium aqueum STEM BARK

ORIGINALITY REPORT

20%
SIMILARITY INDEX

13%
INTERNET SOURCES

16%
PUBLICATIONS

2%
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to Universitas Airlangga **1%**
Student Paper

2 www.thieme-connect.com **1%**
Internet Source

3 G. Pasaribu, S. Hudiyono, E. Budianto, A.H. Cahyana, G. Pari. "In-vitro ANTICANCER ACTIVITY OF Saurauia vulcani LEAF EXTRACTS AGAINST HUMAN COLON WIDR AND HCT-116 CANCER CELLS", Rasayan Journal of Chemistry, 2021 **1%**
Publication

4 Marjon G Bolster, Ben J.M Jansen, Aede de Groot. "The synthesis of Ambra oxide related compounds starting from (+)-larixol. Part 3", Tetrahedron, 2002 **1%**
Publication

5 scholarworks.uark.edu **1%**
Internet Source

6

Toyooka, N.. "Stereodivergent synthesis of the 2,3,5,6-tetrasubstituted piperidine ring system: an application to the synthesis of alkaloids 223A and 205B from poison frogs", *Tetrahedron*, 20040712

Publication

1 %

7

academic-accelerator.com

Internet Source

1 %

8

mail.keosoe.com

Internet Source

1 %

9

Kartika Dwi Kurniasari, Ade Arsianti, Yully Astika Nugrahayning Aziza, Baiq Kirana Dyahningrum Mandasari et al.
"Phytochemical Analysis and Anticancer Activity of Seaweed *Gracilaria verrucosa* against Colorectal HCT-116 Cells", *Oriental Journal of Chemistry*, 2018

Publication

1 %

10

Harald Gross, Joachim Reitner, Gabriele M. König. "Isolation and structure elucidation of azoricasterol, a new sterol of the deepwater sponge *Macandrewia azorica*", *Naturwissenschaften*, 2004

Publication

1 %

11

bmccomplementmedtherapies.biomedcentral.com

Internet Source

1 %

cwww.intechopen.com

12

Internet Source

1 %

13

jnanobiotechnology.biomedcentral.com

Internet Source

1 %

14

rikkyo.repo.nii.ac.jp

Internet Source

1 %

15

journal.umpalangkaraya.ac.id

Internet Source

<1 %

16

Coleman, A.C.. "Radioactivity-Guided Isolation and Characterization of the Bicyclic Pseudopterodin Diterpene Cyclase Product from Pseudopterogorgia elisabethae", Tetrahedron, 20001201

Publication

<1 %

17

Gregory Domnic, Nelson Jeng-Yeou Chear, Siti Fairus Abdul Rahman, Surash Ramanathan et al. "Combinations of indole based alkaloids from Mitragyna speciosa (Kratom) and cisplatin inhibit cell proliferation and migration of nasopharyngeal carcinoma cell lines", Journal of Ethnopharmacology, 2021

Publication

<1 %

18

Mizuko Goto, Irie Miyoshi, Yusuke Ishii, Yukie Ogasawara et al. "Novel skeletal rearrangement of hydroindan derivatives into hydroazulenones via an alkoxy radical", Tetrahedron, 2002

<1 %

19

gcris.iyte.edu.tr

Internet Source

<1 %

20

patents.justia.com

Internet Source

<1 %

21

www.epibio.com

Internet Source

<1 %

22

Pravin L. Kotian, V. Satish Kumar, Tsu-Hsing Lin, Yahya El-Kattan et al. " An Efficient Synthesis Of Acyclic N - and N -Adenine Nucleosides Via Alkylation With Secondary Carbon Electrophiles to Introduce Versatile Functional Groups At the C-1 Position of Acyclic Moiety ", Nucleosides, Nucleotides and Nucleic Acids, 2006

Publication

<1 %

23

www.nature.com

Internet Source

<1 %

24

www.rasayanjournal.co.in

Internet Source

<1 %

25

www.tlsr.usm.my

Internet Source

<1 %

26

link.springer.com

Internet Source

<1 %

27

Sabrin R.M. Ibrahim, Gamal A. Mohamed, Mostafa A. Fouad, Ehab S. El-Khayat, Peter Proksch. " Iotrochotamides I and II: New ceramides from the Indonesian sponge ", Natural Product Research, 2009

Publication

<1 %

28

"World Congress on Medical Physics and Biomedical Engineering 2006", Springer Science and Business Media LLC, 2007

Publication

<1 %

29

K N W Tun, N S Aminah, A N Kristanti, R Ramadhan, Y Takaya. " Two Flavonoids From Stem Bark of and Their Antidiabetic and Antioxidant Activities ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

30

Timothy Newhouse, Alexandra Bodnar, Suzanne Szewczyk, Yang Sun, Yifeng Chen, Anson Huang. "A Comparative Mechanistic Analysis of Palladium- and Nickel-Catalyzed α,β -Dehydrogenation of Carbonyls via Organozinc Intermediates", American Chemical Society (ACS), 2022

Publication

<1 %

31

Yong-Qing Xiao. "A new compound from Gastrodia elata blume", Journal of Asian Natural Products Research, 1/1/2002

Publication

<1 %

32 Yu-Lin Li, Yu Zhang, Ping-Zhi Zhao, Zhan-Xing Hu, Yu-Cheng Gu, Jian Ye, Xiao-Jiang Hao. "Two new steroidal alkaloids from the rhizomes of *Veratrum nigrum* L. and their anti-TYLCV activity", *Fitoterapia*, 2020
Publication <1 %

33 ijrdpl.com
Internet Source <1 %

34 ir.lib.u-ryukyu.ac.jp
Internet Source <1 %

35 patents.google.com
Internet Source <1 %

36 pdfs.semanticscholar.org
Internet Source <1 %

37 repository.uin-malang.ac.id
Internet Source <1 %

38 uspto.report
Internet Source <1 %

39 www.authorstream.com
Internet Source <1 %

40 www.science.gov
Internet Source <1 %

41 www.tandfonline.com
Internet Source <1 %

www.teses.usp.br

42

<1 %

43

Paterson, I.. "A fully stereocontrolled total synthesis of (+)-leucascandrolide A", *Tetrahedron*, 20030825

Publication

<1 %

44

Rafael Pedrosa, Celia Andrés, Jesús M Iglesias, Manuel A Obeso. "Synthesis of enantiopure mono- and disubstituted tetrahydroisoquinolines by 6-exo radical cyclizations", *Tetrahedron*, 2001

Publication

<1 %

45

Zhao, Qingchun, Tayyab A. Mansoor, Jongki Hong, Chong-O. Lee, Kwang Sik Im, Dong Seok Lee, and Jee H. Jung. "New Lysophosphatidylcholines and Monoglycerides from the Marine Sponge *Stelletta* sp.", *Journal of Natural Products*, 2003.

Publication

<1 %

46

Minwan Wu, Yahya El-Kattan, Tsu-Hsing Lin, Ajit Ghosh, Satish Vadlakonda, Pravin L. Kotian, Yarlagadda S. Babu, Pooran Chand. "Synthesis of 9-[1-(Substituted)-3-(phosphonomethoxy)propyl]adenine Derivatives as Possible Antiviral Agents", *Nucleosides, Nucleotides and Nucleic Acids*, 2005

<1 %

47

N. S. Aminah, A. Yulvia, M. Tanjung. "Methyl-3,4-dihydroxybenzoate and 9-10-dihydrophenanthrene-2,4,7-triol two phenolic compounds from *Dioscorea alata* L. and their antioxidant activity", AIP Publishing, 2017

Publication

<1 %

48

Pichlmair, S.. "Evaluation of possible intramolecular [4+2] cycloaddition routes for assembling the central tetracyclic core of the potent marine antiinflammatory agent mangicol A", Tetrahedron, 20060529

Publication

<1 %

49

eprints.lancs.ac.uk
Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

ANTICANCER ACTIVITY OF ISOLATED COMPOUNDS FROM Syzygium aqueum STEM BARK

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
