

DE GRUYTER

2021 VOLUME 19

e-ISSN 2391-5420

# OPEN CHEMISTRY



 | OPEN ACCESS

DE  
G

[degruyter.com/chem](https://degruyter.com/chem)

Should you have **institutional access?** [Here's how to get it...](#)

[SUBJECTS](#) [SERVICES](#) [PUBLICATIONS](#) [ABOUT](#)



[Open Access](#) Published since March 1, 2003

# Open Chemistry

ISSN: 2391-5420

Impact Factor: 1.554

[OVERVIEW](#)

**[LATEST ISSUE](#)**

[ISSUES](#)

[RANKING](#)

[SUBMIT](#)

[EDITORIAL](#)

## Volume 20 Issue 1 January 2022

### Regular Articles

[Open Access](#) January 26, 2022

#### **[Photocatalytic degradation of Rhodamine B in aqueous phase by bimetallic metal-organic framework M/Fe-MOF \(M = Co, Cu, and Mg\)](#)**

*Thi Kim Ngan Tran, Huu Loc Ho, Huu Vinh Nguyen, Bich Thuy Tran, Trung Thanh Nguyen, Phuong Quynh Thi Bui, Long Giang Bach*

Page range: 52-60

[More ▾](#)

[Cite this](#)

[Download PDF](#)

[Open Access](#) February 8, 2022

#### **[Assessment of using electronic portal imaging device for analysing bolus material utilised in radiation therapy](#)**

*Alper Özseven*

Page range: 61-68

[More ▾](#)

[Cite this](#)

[Download PDF](#)

[Open Access](#) February 11, 2022

#### **[A detailed investigation on highly dense CuZr bulk metallic glasses for shielding purposes](#)**

*Huseyin Ozan Tekin, Ghada ALMisned, Gulfem Susoy, Hesham M. H. Zakaly, Shams A. M. Issa, Gokhan Kilic, Yasser Saad Rammah, Gandham Lakshminarayana, Antoaneta Ene*

Page range: 69-80

[More ▾](#)

[Cite this](#)

[Download PDF](#)

[Open Access](#) February 8, 2022

#### **[Simulation of gamma-ray shielding properties for materials of medical interest](#)**

*Mucize Sarihan*

Page range: 81-87

[More ▾](#)

[Cite this](#)

[Download PDF](#)

[Open Access](#) February 8, 2022

#### **[Environmental impact assesment regulation applications and their analysis in Turkey](#)**

*Hakan Ceylan*

Page range: 88-96

[More ▾](#)

[Cite this](#)

[Download PDF](#)

[Open Access](#) February 22, 2022

### **Sample age effect on parameters of dynamic nuclear polarization in certain difluorobenzen isomers/MC800 asphaltene suspensions**

Hüseyin Ovaliöglu

Page range: 97-104

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) March 10, 2022

### **Passenger demand forecasting for railway systems**

Melek Nar, Seher Arslankaya

Page range: 105-119

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) February 22, 2022

### **Design of a Robust sliding mode controller for bioreactor cultures in overflow metabolism via an interdisciplinary approach**

Pablo Antonio López-Peréz, Abraham Efraim Rodriguez-Mata, Omar Hernández-González, Leonel E. Amabilis-Sosa, Rogelio Baray-Arana, Jesus Leon-Borges

Page range: 120-129

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) March 2, 2022

### **Gamma, neutron, and heavy charged ion shielding properties of Er<sup>3+</sup>-doped and Sm<sup>3+</sup>-doped zinc borate glasses**

Huseyin Ozan Tekin, Ghada ALMisned, Hesham M. H. Zakaly, Abdallah Zamil, Dalia Khoucheich, Ghaida Bilal, Lubna Al-Sammaraie, Shams A. M. Issa, Mohammed Sultan Al-Buriah, Antoaneta Ene

Page range: 130-145

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) March 2, 2022

### **Bridging chiral de-tert-butylcalix[4]arenes: Optical resolution based on column chromatography and structural characterization**

Wen-Shan Liu, Hong-Ying Li, Hai-Xia Zhang, Xiu-Yuan Han, Xi-Chun Guo, Chuan-Hua Ding

Page range: 163-168

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) March 11, 2022

### **Petrology and geochemistry of multiphase post-granitic dikes: A case study from the Gabal Serbal area, Southwestern Sinai, Egypt**

Mohamed S. Kamar, Ibrahim A. Salem, Ibrahim E. El-Aassy, Abdu A. El-Sayed, Hamdy A. Awad, Huseyin Ozan Tekin, Abdullah M. Alzahrai, El Saeed R. Lasheen

Page range: 169-181

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) March 8, 2022

### **Comparison of the yield and purity of plasma exosomes extracted by ultracentrifugation, precipitation, and membrane-based approaches**

Wei-Jian Li, Hong Chen, Man-Li Tong, Jian-Jun Niu, Xiao-Zhen Zhu, Li-Rong Lin

Page range: 182-191

[More ▾](#)

[Cite this](#)

[Download PDF](#)

 [Open Access](#) March 10, 2022

### **Bioactive triterpenoids from Indonesian medicinal plant *Syzygium aqueum***

Alfinda Novi Kristanti, Ei Ei Aung, Nanik Siti Aminah, Yoshiaki Takaya, Hnin Thanda Aung, Rico Ramadhan

Page range: 204-211

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 15, 2022

### **Investigation of the effects of machining parameters on surface integrity in micromachining**

*Ayhan Etyemez, Umut Aydin*

Page range: 212-224

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 16, 2022

### **The mesoporous aluminosilicate application as support for bifunctional catalysts for *n*-hexadecane hydroconversion**

*Gulzira Vassilina, Kamilla Umbetkaliyeva, Albina Abdrassilova, Tursunay Vassilina, Zhassulan Zakirov*

Page range: 225-236

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 17, 2022

### **Gamma-ray shielding properties of Nd<sub>2</sub>O<sub>3</sub>-added iron-boron-phosphate-based composites**

*Bekir Oruncak*

Page range: 237-243

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 23, 2022

### **Numerical investigation on perforated sheet metals under tension loading**

*Cetin Karakaya*

Page range: 244-253

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 23, 2022

### **Statistical analysis on the radiological assessment and geochemical studies of granite rocks in the north of Um Taghir area, Eastern Desert, Egypt**

*Hamdy A. Awad, Ibrahim Abu El-Leil, Aleksey V. Nastavkin, Abdellah Tolba, Mostafa Kamel, Refaey M. El-Wardany, Abdalla Rabie, Antoaneta Ene, Huseyin O. Tekin, Shams A. M. Issa, Hesham M. H. Zakaly*

Page range: 254-266

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 24, 2022

### **Two new polypodane-type bicyclic triterpenoids from mastic**

*Xuerui An, Junlong Wang, Xuehui Yu, Hao Wu, Wei Liu*

Page range: 267-271

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 24, 2022

### **Structural, physical, and mechanical properties of the TiO<sub>2</sub> added hydroxyapatite composites**

*Ayhan Etyemez*

Page range: 272-276

[More ▼](#)[Cite this](#)[Download PDF](#)

 Open Access March 24, 2022

### **Tribological properties and characterization of borided Co-Mg alloys**

*Ismail Yıldız*

Page range: 277-286

[More ▾](#)[Cite this](#)[Download PDF](#)

## Review Articles

 [Open Access](#) January 20, 2022

### **Comprehensive review on synthesis, physicochemical properties, and application of activated carbon from the Arecaceae plants for enhanced wastewater treatment**

*Ngoc Bich Hoang, Thi Cam Quyen Ngo, Thi Kim Ngan Tran, Van Tan Lam*

Page range: 10–22

[More ▾](#)[Cite this](#)[Download PDF](#)

 [Open Access](#) January 26, 2022

### **Research progress on speciation analysis of arsenic in traditional Chinese medicine**

*Congnan Peng, Juntong Zhou, Chi Li, Yuxin Chen, Qing Huo, Fei Xie*

Page range: 23–39

[More ▾](#)[Cite this](#)[Download PDF](#)

## Special Issue on Applied Biochemistry and Biotechnology (ABB 2021)

 [Open Access](#) January 18, 2022

### **Study of solidification and stabilization of heavy metals by passivators in heavy metal-contaminated soil**

*Shenglan Ye, Luyao Wang, Tiancheng Liu*

Page range: 1–9

[More ▾](#)[Cite this](#)[Download PDF](#)

 [Open Access](#) March 8, 2022

### **Human health risk assessment and distribution of VOCs in a chemical site, Weinan, China**

*Yan Li, Bo Yan*

Page range: 192–203

[More ▾](#)[Cite this](#)[Download PDF](#)

## Special Issue on the Conference of Energy, Fuels, Environment 2020

 [Open Access](#) January 26, 2022

### **Improving the thermal performance of existing buildings in light of the requirements of the EU directive 2010/31/EU in Poland**

*Tomasz Sumera, Tadeusz Olkusi*

Page range: 40–51

[More ▾](#)[Cite this](#)[Download PDF](#)

## Special Issue on Ethnobotanical, Phytochemical and Biological Investigation of Medicinal Plants

 [Open Access](#) March 2, 2022

### **Study of plant resources with ethnomedicinal relevance from district Bagh, Azad Jammu and Kashmir, Pakistan**

*Tasneem Maqsood, Tayyaba Munawar, Yamin Bibi, Ahmad El Askary, Amal F. Gharib, Tariq E. Elmissbah, Basem H. Elesawy, Abdul Qayyum*

Page range: 146–162

[More ▾](#)[Cite this](#)[Download PDF](#)

[\(Deutsch\)](#)**Type:** Journal**Language:** English**Publisher:** De Gruyter Open Access**First published:** March 1, 2003**Publication Frequency:** 1 Issue per Year

## Subjects

[Architecture and Design](#)[Arts](#)[Asian and Pacific Studies](#)[Business and Economics](#)[Chemistry](#)[Classical and Ancient Near Eastern Studies](#)[Computer Sciences](#)[Cultural Studies](#)[Engineering](#)[General Interest](#)[Geosciences](#)[History](#)[Industrial Chemistry](#)[Islamic and Middle Eastern Studies](#)[Jewish Studies](#)[Law](#)[Library and Information Science, Book Studies](#)[Life Sciences](#)[Linguistics and Semiotics](#)[Literary Studies](#)[Materials Sciences](#)[Mathematics](#)[Medicine](#)[Music](#)[Pharmacy](#)[Philosophy](#)[Physics](#)[Social Sciences](#)[Sports and Recreation](#)[Theology and Religion](#)

## Services

[For journal authors](#)[For book authors](#)[For librarians](#)[Rights & Permissions](#)

## Publications

[Publication types](#)[Open Access](#)

## About

[Contact](#)[Career](#)[About De Gruyter](#)[Partnerships](#)[Press](#)[New website FAQs](#)[Help/FAQ](#)[Privacy policy](#)[Cookie Policy](#)[Accessibility](#)[Terms & Conditions](#)[Legal Notice](#)

© Walter de Gruyter GmbH 2022



Should you have **institutional access?** [Here's how to get it...](#)

[SUBJECTS](#)[SERVICES](#)[PUBLICATIONS](#)[ABOUT](#)

**Open Access** Published since March 1, 2003

# Open Chemistry

ISSN: 2391-5420

Impact Factor: 1.554

[OVERVIEW](#)[LATEST ISSUE](#)[ISSUES](#)[RANKING](#)[SUBMIT](#)[EDITORIAL](#)

## Editorial

### Editor-in-Chief

Snezana D. Zaric, University of Belgrade, Serbia

### Managing Editor

Karolina Kurtyka, Poland

### Editorial Advisory Board

Metin Hayri Acar, Istanbul Technical University, Turkey

Sergei Aldoshin, Russian Academy of Sciences, Russia

Roland Boese, University of Essen, Germany

David C. Clary, University of Oxford, UK

Graham Cooks, Purdue University, USA

Elias J. Corey, Harvard University, USA

Carlos Fernandez, Robert Gordon University, UK

[Boris Furtula](#), University of Kragujevac, Serbia

Jean-François Gérard, SGM INSA Lyon, CNRS, ECNP, France

Raquel P. Herrera, Isqch (Csic-Uz) Instituto De Síntesis Química Y Catálisis Homogénea, Spain

Janusz Jurczak, Warsaw University and Institute of Organic Chemistry, Poland

Alexei Khokhlov, Moscow State University and Nesmeyanov Institute of Organoelement Compounds, Russia

Alexander M. Klibanov, Massachusetts Institute of Technology, USA

Jacek Klinowski, University of Cambridge, UK

Shu Kobayashi, University of Tokyo, Japan

Pavel Kratochvíl, Academy of Sciences of the Czech Republic, Czech Republic

Janusz Lipkowski, Polish Academy of Sciences, Poland

Goverdhan Mehta, Indian Institute of Science, India

Achim Müller, University of Bielefeld, Germany

Stanislaw Penczek, Centre of Molecular and Macromolecular Studies, Poland

Chintamani Nagesa Ramachandra Rao, Jawaharlal Nehru Centre for Advanced Scientific Research, India

Thomas Rauchfuss, University of Illinois, USA

Vladimir Sklenar, Masaryk University, Czech Republic

Edward I. Solomon, Stanford University, USA

Barry Trost, Stanford University, USA

[Donald G. Truhlar](#), University of Minnesota, USA

Fosong Wang, Chinese Academy of Sciences, China

George Whitesides, Harvard University, USA

Frank Würthner, Institut für Organische Chemie & Center for Nanosystems Chemistry, Germany

Jung Woon Yang, Sungkyunkwan University, South Korea

### Editors

Mozhgan Afshari, Islamic Azad University, Shoushtar, Iran

Khuram Shahzad Ahmad, Fatima Jinnah Women University, Pakistan

Iskender Akkurt, Süleyman Demirel Üniv. Physics Dep. Nuclear Physics Div. Isparta, Turkey

Diego Alonso, Alicante University, Spain

Biljana Arsic, Department of Chemistry, Faculty of Sciences and Mathematics, University of Nis, Republic of Serbia

Aleksander Maria Astel, Pomeranian University in Słupsk, Poland



Maria Luisa Astolfi Sapienza University of Rome, Italy  
Ebaa Adnan Azooz, The Gifted Students` School in Najaf, Ministry of Education, Iraq  
Sezgin Bakirdere, Yıldız Technical University, Turkey  
Saikat Bala, Scientist II at Beam Therapeutics, Cambridge, Massachusetts, USA  
Csaba Balazsi, Centre for Energy Research, Centre of Excellence of Hungarian Academy of Sciences, Hungary  
Roya Boodaghi Malidarre, Payame Noor University, Tehran, Iran  
Arindam Bose, Harvard Medical School, USA  
Anthony J. Burke, University of Evora, Portugal  
Eugenijus Butkus, Vilnius University, Lithuania  
Sergio Carrasco, Universidad Rey Juan Carlos, Spain  
Domenico Cautela, Stazione Sperimentale per le Industrie delle Essenze e dei Derivati Dagli Agrumi (SSEA), Italy  
Paolo Censi, University of Palermo, Italy  
Christophoros Christophoridis, Aristotle University of Thessaloniki, Greece  
Łukasz Cieřła, University of Alabama, USA  
Dibyendu Dana, KemPharm Inc., USA  
Costel C. Darie, Clarkson University, USA  
Adi Darmawan, Faculty of Sciences and Mathematics, Diponegoro University, Indonesia  
Rajat Subhra Das, Omega Therapeutics, USA  
Joaquín R. Domínguez, Universidad de Extremadura, Spain  
Biswanath Dutta, University of Illinois Urbana Champaign, USA  
Chiara Fanali, Campu Bio-Medico University of Rome, Italy  
Is Fatimah, Universitas Islam Indonesia, Indonesia  
Huanhuan Feng, Harbin Institute of Technology (Shenzhen), China  
Robert Fraczkiewicz, Simulations Plus Inc., USA  
Iolanda Francolini, Sapienza University of Rome, Italy  
Ramesh L. Gardas, Indian Institute of Technology Madras, India  
Mazeyar Parvinzadeh Gashti, PRE Labs Inc, Canada  
Jose Gonzalez-Rodriguez, University of Lincoln, UK  
Sravanthi Devi Guggilapu, Nimble Therapeutics, USA  
Juan Luis Garcia Guirao, Technical University of Cartagena, Spain  
Oğuz Gürsoy, Burdur Mehmet Akif Ersoy University, Turkey  
Dariusz Guziejewski, University of Lodz, Poland  
Ahmed A. Hussein, Cape Peninsula University of Technology, South Africa  
Ahmed S. Ibrahim, Qatar University, Qatar  
Saravana Kumar Jaganathan, Universiti Teknologi Malaysia, Malaysia  
Agata Jakóbk-Kolon, Silesian University of Technology, Poland  
Paweł Jeźowski, Poznan University of Technology, Poland  
Hassan Karimi-Maleh, University of Electronic Science and Technology of China, China  
Yasar Nellyot Kavil, King Abdulaziz University, Saudi Arabia  
Przemysław Kowalczewski, Poznań University of Life Sciences, Poland  
Iryna Kravchenko, Odessa Polytechnic State University (Department of Organic and Pharmaceutical Technology), Ukraine  
Jerzy Langer, Adam Mickiewicz University, Poland  
Fei Li, Zhongnan University of Economics and Law, China  
Wenhui Li, Applied Materials Inc., USA  
Antonio Martin-Esteban, INIA-CSIC, Spain  
Zoran Mazej, Jozef Stefan Institute, Slovenia  
Mohsen Mhadhbi, National Institute of Research and Physical-chemical Analysis, Tunisia  
Christiana Mitsopoulou, National and Kapodistrian University of Athens, Greece  
Raj Mukherjee, Sanofi, USA  
Selvakumar Murugesan, University of Bayreuth, Germany  
Dmitry Murzin, Åbo Akademi, Finland  
Waqas Nazeer, GC University Lahore, Pakistan  
Awal Noor, King Faisal University, Saudi Arabia  
Mozaniel Santana de Oliveira, Adolpho Ducke Laboratory, Botany Coordination, Museu Paraense Emílio Goeldi, Brazil  
Shagufta Perveen, King Saud University, Saudi Arabia  
Christos Petrou, School of Sciences and Engineering, Cyprus  
Daniela Piazzase, University of Palermo, Italy  
Tanay Pramanik, University of Engineering and Management Kolkata, India  
María Mar Quesada-Moreno, University of Granada, Spain  
Fitria Rahmawati, Universitas Sebelas Maret, Indonesia  
Ponnadurai Ramasami, University of Mauritius, Mauritius

Abdul Rauf, University of Swabi, Pakistan  
Dominique Richon, MINES ParisTECH, France  
Juan Garcia Rodriguez, Complutense University, Spain  
Daily Rodriguez-Padron, Universidad de Cordoba, Spain  
Agnieszka Saeid, Wroclaw University of Science and Technology, Poland  
Suresh Sagadevan, FRSC, Nanotechnology & Catalysis Research Centre, University of Malaya, Malaysia  
Christian Schmitz, Hochschule Niederrhein, Germany  
Catinca Secuianu, Politehnica University of Bucharest, Romania  
Navpreet Kaur Sethi, Zhejiang University, China  
Belgin Sever, Anadolu University, Faculty of Pharmacy, Turkey  
Praveen Kumar Sharma, Lovely Professional University, India  
Francesco Siano, National Research Council, Italy  
Krishnamoorthy Sivakumar, SCSVMV University, India  
Gaweł Sołowski, Institute of Fluid Machinery, Poland  
Atul Srivastava, University of Chicago, USA  
Lakshmi Narayana Suvarapu, Yeungnam University, South Korea  
Jose Luis Toca-Herrera, BOKU, Austria  
Riaz Ullah, Department of Pharmacognosy, College of Pharmacy, King Saud University Riyadh Saudi Arabia  
Konstantin Volcho, Novosibirsk Institute of Organic Chemistry SB RAS, Russia  
Iveta Waczulikova, Comenius University, Fac of Math, Phys. and Inform., Slovakia  
Chunpeng Wan, Jiangxi Agricultural University, China  
Shin-ichi Yusa, Univ of Hyogo, Japan  
Constantinos K. Zacharis, Aristotle University of Thessaloniki, Greece  
Szczezan Zapotoczny, Jagiellonian University in Krakow, Poland  
Snezana Zaric, University of Belgrade, Serbia  
Zhien Zhang, Ohio State University, USA  
Tingting Zheng, Peking University Shenzhen Hospital, Shenzhen PKU-HKUST Meidical Center, Peking University, China  
Grigoris Zoidis, National and Kapodistrian University of Athens, Greece

**Publisher**

DE GRUYTER Poland  
Bogumiła Zuga 32A Str.  
01-811 Warsaw, Poland  
T: +48 22 701 50 15

**Editorial Contact**

[openchemistry@degruyter.com](mailto:openchemistry@degruyter.com)

**Assistant Managing Editor**

Deepan Selvaraj, Compuscript  
[AssistantManagingEditor@degruyter.com](mailto:AssistantManagingEditor@degruyter.com)

[\(Deutsch\)](#)

**Type:** Journal

**Language:** English

**Publisher:** De Gruyter Open Access

**First published:** March 1, 2003

**Publication Frequency:** 1 Issue per Year

---



## Subjects

---

Architecture and Design

Arts

Asian and Pacific Studies

Business and Economics

Chemistry

Classical and Ancient Near Eastern Studies

Computer Sciences

Cultural Studies

Engineering

General Interest

Geosciences

History

Industrial Chemistry

Islamic and Middle Eastern Studies

Jewish Studies

Law

Library and Information Science, Book Studies

Life Sciences

Linguistics and Semiotics

Literary Studies

Materials Sciences

Mathematics

Medicine

Music

Pharmacy

Philosophy

Physics

Social Sciences

Sports and Recreation

Theology and Religion

## Services

---

For journal authors

For book authors

For librarians

Rights & Permissions

## Publications

---

Publication types

Open Access

## About

---

Contact

Career

About De Gruyter

Partnerships

Press

New website FAQs

---

[Help/FAQ](#)

[Privacy policy](#)

[Cookie Policy](#)

[Accessibility](#)

[Terms & Conditions](#)

[Legal Notice](#)

---

© Walter de Gruyter GmbH 2022



SJR

Scimago Journal &amp; Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

# Open Chemistry


**COUNTRY**[Germany](#)Universities and research  
institutions in Germany**SUBJECT AREA AND CATEGORY**[Chemistry](#)[Chemistry \(miscellaneous\)](#)[Materials Science](#)[Materials Chemistry](#)**PUBLISHER**[Walter de](#)[Gruyter](#)[GmbH](#)**PUBLICATION TYPE**[Journals](#)**ISSN**

23915420

**COVERED PERIOD**2003,  
2015-  
2020

**SCOPE**

Open Chemistry is a peer-reviewed, open access journal that publishes original research, reviews and short communications in chemistry in an ongoing way. Our central goal is to provide a hub for researchers working across all subjects to present their work and to be a forum for the discussion of the important issues in the field. Our journal is the premier source for cutting edge research in chemistry and it provides high quality peer review services for its authors across the world. Moreover, it allows for libraries to avoid subscribing to multiple local publications, and to receive instead all the necessary chemistry research from a single source, serving the entire scientific community.







 Join the conversation about this journal




 Quartiles



**FIND SIMILAR JOURNALS** 

<p>1 <b>Journal of Chemistry</b> USA <b>67%</b> similarity</p>	<p>2 <b>Bulgarian Chemical Communications</b> BGR <b>57%</b> similarity</p>	<p>3 <b>Molecules</b> CHE <b>56%</b> similarity</p>	<p>4 <b>J S P</b></p>
--	---	---	---------------------------





**Open Chemistry**

Q3

Chemistry (miscellaneous)

best quartile

**SJR 2020**

0.27

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

**SCImago Graphica**

Explore, visually communicate and make sense of data with our **new free tool**.

Get it



Metrics based on Scopus® data as of April 2021

**Leave a comment**

Name

**Email**

(will not be published)

[Submit](#)

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

**EST MODUS IN REBUS**

Horatio (Satire 1,1,106)



## Research Article

Alfinda Novi Kristanti\*, Ei Ei Aung, Nanik Siti Aminah, Yoshiaki Takaya, Hnin Thanda Aung, Rico Ramadhan

# Bioactive triterpenoids from Indonesian medicinal plant *Syzygium aqueum*

<https://doi.org/10.1515/chem-2022-0138>

received December 23, 2021; accepted February 15, 2022

**Abstract:** A large common species, *Syzygium aqueum*, belonging to the genus *Syzygium* possesses numerous bioactive phytochemical constituents. Moreover, the different parts of this species have been used as folk medicine since centuries ago. In this study, a phytochemical exploration was carried out on the plant's stem bark. Isolation of the compounds was carried out through the extraction step with some organic solvents, followed by separation and purification using chromatography techniques until the two triterpenoids were isolated from nonpolar and semipolar extracts. Structure elucidation was done using spectroscopic methods. These compounds were identified as alphitolic acid and arjunolic acid. Subsequently, these two compounds were used in anti-cancer tests against human cancer cells HeLa, T47D, and A549 using colorimetric assay. The result showed that both compounds showed more inhibition of the growth of HeLa and T47D than A549 cancer cells, with the highest activity shown by arjunolic acid against HeLa cell lines.

\* **Corresponding author: Alfinda Novi Kristanti**, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Kampus C UNAIR, Jl. Mulyorejo-60115, Surabaya, Indonesia; Biotechnology of Tropical Medicinal Plants Research Group, Universitas Airlangga, Surabaya, Indonesia, e-mail: [alfinda-n-k@fst.unair.ac.id](mailto:alfinda-n-k@fst.unair.ac.id), tel: +62-31-5936501, fax: +62-31-5936502

**Ei Ei Aung:** Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Kampus C UNAIR, Jl. Mulyorejo-60115, Surabaya, Indonesia; Department of Chemistry, Yadanarbon University, Amarapura-05063, Mandalay, Myanmar

**Nanik Siti Aminah:** Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Kampus C UNAIR, Jl. Mulyorejo-60115, Surabaya, Indonesia; Biotechnology of Tropical Medicinal Plants Research Group, Universitas Airlangga, Surabaya, Indonesia

**Yoshiaki Takaya:** Faculty of Pharmacy, Meijo University, Tempaku-468-8502, Nagoya, Japan

**Hnin Thanda Aung:** Department of Chemistry, Mohnyin Degree College, Kachin state, Myitkyina, Myanmar

**Rico Ramadhan:** Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Kampus C UNAIR, Jl. Mulyorejo-60115, Surabaya, Indonesia

**Keywords:** *Syzygium aqueum*, alphitolic acid, arjunolic acid, MTT assay, XTT assay

## 1 Introduction

One of the probable medicament manufacturing countries, Indonesia, between the Pacific and Indian Oceans, is commonly known as a country with the second richest biodiversity of medicinal plants on the earth. Medicinal plants have therapeutic properties or provide beneficial pharmacological effects for the human body. Some examples are given here. The chemical composition of the essential oils extracted from *Elaeagnus umbellata* Thunb fruit demonstrated antioxidant, anticholinesterase, and antidiabetic activities that could be used as an alternative drug to treat oxidative stress-related diseases [1]. The essential oil of *Teucrium stocksianum* possesses a strong antinociceptive potential, which is further used as a topical analgesic [2]. The study of *Woodwardia unigemmata* (Makino) Nakai plant extracts, which were rich in polyphenolic compounds (total phenolic compounds and total flavonoid compounds) with efficient antioxidant activity, also showed remarkable antibacterial activity against plant and animal pathogenic bacteria [3]. The diverse Indonesian medicinal plants have been used as the sources of herbal medicines for the inhibition and treatment of human diseases for several thousand years [4].

The sixteenth largest genus of flowering plants in the Myrtaceae family, *Syzygium*, comprises about 1,200–1,800 species. Moreover, *Syzygium* species of tree and shrub types are distributed in tropical areas worldwide, especially Indonesia, Malaysia, Philippines, and Thailand. Some species of *Syzygium* are widely cultivated for economic importance, edible fruits, spices, flavoring agents, and pharmacological proprieties. *Syzygium* contains abundant diversity of bioactive phytochemical constituents such as terpenoids, steroids, chalcones, flavonoids, lignans, alkyl phloroglucinols, hydrolysable tannins, and chromone derivatives [5].

A medicinal species of *Syzygium* found in Indonesia, *Syzygium aqueum*, is a large common species in this genus found in Malaysia [6]. Fruits, leaves, and roots of this species have been used as folk medicine, especially for antibiotic properties, diabetic treatment, the treatment for a cracked tongue, relieving itches, stomach aches, dysentery, and reducing of swelling, respectively [7–11].

Some researchers were focused on bioactive phytochemical constituents based on the above medicinal values of this plant. Therefore, bioactive phytochemical constituents such as tannins, flavonoids, steroids, triterpenoids, and phenols from leaves and stem bark of this plant have already been reported in some research articles [12–17]. These articles also showed that some flavonoids from *S. aqueum* (leaves) strongly showed antidiabetic properties using  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibition assays and showed their cytotoxicity against MCF-cell line using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Moreover, some triterpenoids and steroids from *S. aqueum* (stem bark) were active with high cytotoxicity on HeLa, T47D, and A549 cancer cell lines using MTT and 2,3-bis-(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5-carboxanilide (XTT) assays.

Based on the literature study, the stem bark of *S. aqueum* was designated for this research. Primarily, some extracts of stem bark of *S. aqueum* were separated to collect bioactive constituents with vacuum liquid chromatography and column chromatography. The structure of isolated compounds was identified with spectroscopic methods, including UV, Fourier transform infrared (FTIR), nuclear magnetic resonance (NMR [1D and 2D]), and HR-DART-MS. After this, their anticancer activity was determined with colorimetric assay (MTT and XTT assays). The isolated bioactive constituents were triterpenoid compounds, first isolated from *S. aqueum*.

## 2 Materials and methods

### 2.1 Plant materials

The stem bark of *S. aqueum* was collected in December 2018 from Wage, Taman, Sidoarjo, East Java, Indonesia. The plant material was identified at the Department of Biology, Faculty of Science and Technology, Universitas Airlangga, and the voucher specimen (UA-MSa050918) was deposited at the Herbarium of Universitas Airlangga, Laboratory of Biosystematic, Department of Biology, Faculty of Science and Technology, Universitas Airlangga. The

collected stem bark was cleaned, chopped, and crushed into small pieces of coarse powder.

### 2.2 Instrumentations

A polarimeter with  $\varphi$  3.5 mm diameter (cell size)  $\times$  100 mm (cell length) was used to measure the specific rotation. For structure elucidation, some of the instruments used are a UV-vis 1800 spectrometer (Shimadzu), FTIR spectra on a Tracer-100 spectrophotometer (Shimadzu), NMR spectra on Bruker Avance III HD 600, and HR-DART-MS on an Exactive Plus Orbitrap DART Mass Spectrometer. Enzyme-linked immunosorbent assay (ELISA) reader was used for the anticancer test.

### 2.3 General procedure

For extraction, fractionation, isolation, and purification of a sample, organic solvents such as *n*-hexane (*n*-Hex), ethyl acetate (EtOAc), dichloromethane (DCM), and methanol (MeOH) were used. In analytical thin-layer chromatography (TLC), a precoated silica gel 60 F<sub>254</sub> (Merck) was used and an anisaldehyde-sulfuric acid reagent was sprayed on a TLC plate for visualization. Silica gel 60 (700–200 mesh ASTM) was applied for column chromatography. Then, vacuum liquid chromatography (VLC) was performed using Kieselgel 60 (F<sub>254</sub>, Merck). The melting point of pure compounds was measured on a Fisher-Johns melting point apparatus (Stuart SMP30). Spectroscopic methods, including UV-vis, IR, NMR, and MS, were used to determine the structure of isolated compounds. An anticancer test was carried out using MTT and XTT reagents (colorimetric assay), and the absorbance for an anticancer test was recorded on an ELISA reader.

### 2.4 Extraction and isolation

At room temperature, 2 kg of coarse powder of stem bark was extracted with a polar solvent (MeOH, 40 L) for 3  $\times$  24 h. This crude extract (450 g) was then mentioned to partition with a nonpolar solvent (*n*-hexane). It was separated into a nonpolar fraction and a methanol fraction. Then, 400 mL of aqueous was added to methanol fractions before partitioning with a semipolar solvent, EtOAc. After this, this mixture was separated into a semipolar fraction and an aqueous methanol residue. First, the

nonpolar fraction was chromatographed with gradient polarity solvents, *n*-Hex:EtOAc. Separation procedure was carried out for three times using column chromatography to obtain a pure compound. Compound 1 (30 mg) was acquired from a fraction eluted with *n*-Hex:EtOAc = 7:3. The semipolar fraction was separated by VLC using gradient solvent mixtures of *n*-Hex:DCM and DCM:EtOAc. Separation procedure using column chromatography was carried out twice to obtain a pure compound, Compound 2 (50 mg), that was eluted with DCM:EtOAc = 6:4.

## 2.5 Anticancer activity

### 2.5.1 Cell culture

Three cancer cell lines, HeLa (cervical cancer), T47D (breast cancer), and A459 (lung cancer), were obtained from American Type Culture Collection. Cells were cultured at 37°C in a CO<sub>2</sub> incubator for 24 h and 100% humidity in medium supplemented with 10% FBS, 1% L-glutamine, and 1% penicillin/streptomycin. The cell culture process was carried out in the Cancer Chemoprevention Research Center, Faculty of Pharmacy at Universitas Gadjah Mada, Indonesia and in Physic Laboratory, Faculty of Science of University Putra Malaysia (UPM), Malaysia.

### 2.5.2 MTT assay

An anticancer test of pure compounds was performed using MTT assay. Cells were cultured in 96 well plates at  $2.13 \times 10^4$  cell/well density and were incubated at 37°C in a CO<sub>2</sub> incubator for 24 h. The cells were then treated with 100 µL of the prepared sample with different concentration series (1.5625–100 µg/mL) at 37°C in a CO<sub>2</sub> incubator for 24 h. As much as 100 µL of the MTT reagent (50 mg in 10 mL of PBS) was filled in each well after incubation, and it was incubated again at 37°C in a CO<sub>2</sub> incubator for 2–4 h until purple formazan crystals are formed. After the crystal formation, cells were observed with a microscope, and 100 µL of 10% SDS stopper in 0.1 N HCl was added, and it was kept in a dark place overnight (room temperature). Next, the plate's absorbance value was read at a wavelength of 560 nm by using an ELISA reader [18,19].

### 2.5.3 XTT assay

The anticancer test was carried out using XTT assay. Cells were seeded ( $10^4$ – $10^5$  cells/well) into 96 well plates and

were incubated at 37°C in a CO<sub>2</sub> incubator for 24 h. As much as 100 µL of the prepared sample with various concentrations (1.5625–100 µg/mL) was inserted to treat cancer cells. The plate was incubated at 37°C in a CO<sub>2</sub> incubator for 24 h. And then, PMS solution (10 mM PMS solution and PBS (3 mg PMS + 1 mL PBS)) was prepared and 10 µL of this solution was added in 4 mL of XTT (4 mg of XTT dissolved in 4 mL of 37°C cells culture medium). After incubation, 25 µL of the XTT reagent (XTT/PMS solution) was filled in each well. After this, it was incubated for 2–4 h in a CO<sub>2</sub> incubator at 37°C until the changing of the orange formazan. The next step was to read the absorbance value at a wavelength of 450 nm using an ELISA reader [19,20]. Each experiment was carried out in triplicate, and the number of viable cells was calculated using the following formula:

$$\begin{aligned} & \% \text{ Cell viability} \\ &= \frac{\text{Absorbance of treatment} - \text{Absorbance of media}}{\text{Absorbance of negative control} - \text{Absorbance of media}} \\ & \times 100\%. \end{aligned}$$

## 3 Results and discussion

### 3.1 Isolation of compounds and structure elucidation

Nonpolar and semipolar fractions of *S. aqueum* (stem bark) gave one bioactive compound each, Compound 1 and Compound 2. Structures of these compounds were identified and elucidated using spectroscopic methods.

**Compound 1** was acquired as white powder,  $[\alpha]_D^{25}$  -9, m.p. 279–280°C. According to the HR-DART-MS spectrum, the molecular formula was C<sub>30</sub>H<sub>48</sub>O<sub>4</sub>, *m/z* 471.3462 [M–H]<sup>–</sup> (calc. mass for C<sub>30</sub>H<sub>48</sub>O<sub>4</sub>–H<sup>–</sup>, *m/z* 471.3474). The UV spectrum in MeOH was displayed at  $\lambda_{\text{max}}$  182 nm. In the FTIR spectrum (KBr), hydroxyl group (3,423 cm<sup>–1</sup>), symmetric and asymmetric sp<sup>3</sup> C–H stretching (2,941–1,870 cm<sup>–1</sup>), carbonyl acid group (1,699 cm<sup>–1</sup>), CH<sub>2</sub> bending (1,456 cm<sup>–1</sup>), and CH<sub>3</sub> bending (1,381 cm<sup>–1</sup>) were recorded. The proton signals, both  $\delta_{\text{H}}$ : 3.60 ppm (H-2 $\beta$ ) and  $\delta_{\text{H}}$ : 2.89 ppm (H-3 $\alpha$ ), were suggested as carbinol methine protons, which coupled each other according to the Double Quantum Filtered-COSY (DFQ-COSY) experiment and relative configuration indicated *trans* configuration due to the 9.6 Hz coupling constant of H-3 $\alpha$ . Nuclear Overhauser Effect Spectroscopy (NOESY) cross-peaks of H-3 $\alpha$  and H-5 $\alpha$  confirmed the presence of the *trans*-AB ring. Six methyl protons possessed some Heteronuclear Multiple Bond Correlation (HMBC) such as H-23 ( $\delta_{\text{H}}$ : 0.99 ppm) and C-24 ( $\delta_{\text{C}}$ : 15.8 ppm), H-24 ( $\delta_{\text{H}}$ : 0.78 ppm) and H-25 ( $\delta_{\text{H}}$ : 0.92 ppm) and C-5 ( $\delta_{\text{C}}$ : 55.2 ppm), H-26 ( $\delta_{\text{H}}$ : 0.97 ppm) and C-7 ( $\delta_{\text{C}}$ : 1.43 ppm), and H-27 ( $\delta_{\text{H}}$ : 1.01 ppm)




Figure 1: DQF-COSY, NOESY, and HMBC correlations of alphitolic acid from *S. aqueum*.


and C-14 ( $\delta_C$ : 42.3 ppm). Two vinylic protons (H-29) were observed at  $\delta_H$ : 4.60 ppm and  $\delta_H$ : 4.71 ppm. Their positions were confirmed by HMBC between these two H-29 protons with C-19 ( $\delta_C$ : 47.1 ppm) and C-30 ( $\delta_C$ : 18.2 ppm). Based on HSQC experiments, two carbinol methine carbons ( $R_2$ -CH-OH) gave correlations H-2 $\beta$ -C-2 ( $\delta_C$ : 68.4 ppm) and H-3 $\alpha$ -C-3 ( $\delta_C$ : 83.0 ppm). A vinylic group was appeared at  $\delta_C$ : 108.7 ppm (C-29) and  $\delta_C$ : 150.6 ppm (C-20) in the  $^{13}C$  NMR spectrum. The C-20 was supposed to be a quaternary carbon of the olefinic system. And then, the C-29 was suggested as methylene carbon of the same olefinic system ( $=CH_2$ ), which were connected by two protons [ $\delta_H$ : 4.60 ppm (H-29) and  $\delta_H$ : 4.71 ppm (H-29)] through the HSQC spectrum. Chemical shift of the carbonyl group appeared at  $\delta_C$ : 178.7 ppm (C-28) in the  $^{13}C$  NMR spectrum. In the HMBC spectrum, C-28 showed correlations with H-18 ( $\delta_H$ : 1.61 ppm) and H-22 ( $\delta_H$ : 1.43 ppm). Furthermore, the positions of quaternary carbons were verified based on the HMBC spectrum by observing the correlations of H-23 and H-24 to C-4 ( $\delta_C$ : 39.1 ppm), H-27 to C-8 ( $\delta_C$ : 40.6 ppm), H-1 to C-10 ( $\delta_C$ : 38.1 ppm), H-15 to C-14 ( $\delta_C$ : 42.3 ppm), and H-16 to C-17 ( $\delta_C$ : 56.1 ppm). The remaining peaks at  $\delta_C$ : 27.7 ppm (C-23), 15.8 ppm (C-24), 16.5 ppm (C-25), 15.3 ppm (C-26), 13.7 ppm (C-27), and 18.2 ppm (C-30) were suggested as primary carbons through Distortionless Enhancement by Polarization Transfer (DEPT) 135 information. Moreover, six methyl carbons, ten methylene carbons, seven methine carbons, and seven quaternary carbons were confirmed with the DEPT experiment (DEPT 90 and DEPT 135). Therefore, Compound 1 was verified as alphitolic acid according to the above data. DQF-COSY ( $^1H$ - $^1H$ ), NOESY ( $^1H$ - $^1H$ ), and HMBC ( $^1H$ - $^{13}C$ ) correlations of the structure are shown in Figure 1, as well as spectral data of both  $^1H$  NMR (600 MHz, methanol- $d_4$ ) with their coupling constant

and  $^{13}C$  NMR (150 MHz, methanol- $d_4$ ) were compared with the literature data in Table 1.

**Compound 2** was acquired as white powder, m.p. 338–340°C. The UV spectrum in MeOH appeared at  $\lambda_{max}$  220 nm. In FTIR spectra (KBr), hydroxyl group ( $3,371\text{ cm}^{-1}$ ), asymmetric and symmetric  $sp^3$  C-H stretching vibration ( $2,941\text{ cm}^{-1}$ ), carbonyl (COOH) group ( $1,693\text{ cm}^{-1}$ ),  $CH_2$  bending ( $1,462\text{ cm}^{-1}$ ) and  $CH_3$  bending ( $1,388$ – $1,365\text{ cm}^{-1}$ ), and C-O stretching ( $1,049\text{ cm}^{-1}$ ) were observed. In the  $^1H$  NMR spectrum, four carbinol protons were shown at  $\delta_H$ : 3.68 ppm (H-2 $\beta$ ),  $\delta_H$ : 3.35 ppm (H-3 $\alpha$ ),  $\delta_H$ : 3.27 ppm (H-23), and  $\delta_H$ : 3.50 ppm (H-23). The H-3 $\alpha$  and H-2 $\beta$  were coupled through the DQF-COSY spectrum. The *trans* configuration was confirmed due to the 10.2 Hz coupling constant of H-3 $\alpha$  and NOESY correlation of H-2 $\beta$  and H-24 $\beta$ . Thus, H-23 was correlated to C-3 ( $\delta_C$ : 76.8 ppm) according to the HMBC spectrum. The olefinic protons were shown at  $\delta_H$ : 5.25 ppm (H-12) and coupled to H-11 ( $\delta_H$ : 1.95 ppm) in the DQF-COSY spectrum. Moreover, six methyl protons were appeared at  $\delta_H$ : 0.70 ppm (H-24), 1.03 ppm (H-25), 0.82 ppm (H-26), 1.18 ppm (H-27), 0.91 ppm (H-29), and 0.94 ppm (H-30). These methyl protons have correlations observed on the HMBC spectrum, especially between H-24 and C-23 ( $\delta_C$ : 65.0 ppm), H-26 and C-8 ( $\delta_C$ : 39.2 ppm) and C-14 ( $\delta_C$ : 41.6 ppm), H-27 and C-8 ( $\delta_C$ : 39.2 ppm), C-14 ( $\delta_C$ : 41.6 ppm) and C-15 ( $\delta_C$ : 27.4 ppm), and H-29 and H-30 and C-20 ( $\delta_C$ : 30.2 ppm). On the  $^{13}C$  NMR spectrum, the three carbinol carbons were observed at  $\delta_C$ : 68.8 ppm (C-2), 76.8 ppm (C-3), and 65.0 ppm (C-23). The carbons of C-2 and C-3 were identified as methine alcohol ( $R_2$ -CH-OH), and C-23 was suggested as methylene alcohol ( $R$ - $CH_2$ -OH) according to DEPT (DEPT 135 and DEPT 90) experiments. Chemical shifts at  $\delta_C$ : 180.7 ppm in the  $^{13}C$  NMR spectrum indicated the presence of the carboxylic acid functional group (C-28), which was

**Table 1:** NMR spectra of arjunolic acid by comparing to literature

Position	DEPT	Experiment			Literature [21]		
		$\delta_C$ (ppm)	$\delta_H$ (ppm) {mult, J (Hz)}		$\delta_C$ (ppm)	$\delta_H$ (ppm) {mult, J (Hz)}	
1	CH <sub>2</sub>	46.9	0.82 (1H, m) 2.00 (1H, m)		46.5	—	—
2	CH	68.4	3.60 (1H, m)		68.6	3.46	(1H, dd, J = 4.0, 9.5 Hz)
3	CH	83.0	2.89 (1H, d, J = 9.6 Hz)		83.2	2.74	(1H, d, J = 9.5 Hz)
4	C	39.1	—		40.6	—	
5	CH	55.2	0.82 (1H, m)		55.2	—	
6	CH <sub>2</sub>	18.1	1.54 (1H, m) 1.43 (1H, m)		18.2	—	
7	CH <sub>2</sub>	34.1	1.43 (2H, m)		34.1	—	
8	C	40.6	—		39.2	—	
9	CH	50.6	1.39 (1H, m)		50.3	—	
10	C	38.1	—		38.2	—	
11	CH <sub>2</sub>	20.8	1.43 (2H, m)		20.9	—	
12	CH <sub>2</sub>	25.4	1.73 (1H, m) 1.08 (1H, m)		25.3	1.52 (1H, m) 1.64 (1H, m)	
13	CH	38.2	2.32 (1H, m)		38.1	—	
14	C	42.3	—		40.6	—	
15	CH <sub>2</sub>	29.4	1.30 (1H, m) 1.15 (1H, m)		30.4	—	
16	CH <sub>2</sub>	32.0	0.90 (2H, m)		32.1	—	
17	C	56.1	—		56.1	—	
18	CH	49.1	1.61 (1H, m)		48.1	—	
19	CH	47.1	3.03 (1H, m)		48.9	2.86	(1H, m)
20	C	150.6	—		150.7	—	
21	CH <sub>2</sub>	30.3	1.39 (2H, m)		29.5	—	
22	CH <sub>2</sub>	36.7	1.91 (1H, m) 1.43 (1H, m)		37.0	—	
23	CH <sub>3</sub>	27.7	0.99 (3H, s)		28.2	0.81	(3H, s)
24	CH <sub>3</sub>	15.8	0.78 (3H, s)		16.4	0.60	(3H, s)
25	CH <sub>3</sub>	16.5	0.92 (3H, s)		17.1	0.71	(3H, s)
26	CH <sub>3</sub>	15.3	0.97 (3H, s)		15.7	0.80	(3H, s)
27	CH <sub>3</sub>	13.7	1.01 (3H, s)		14.5	0.75	(3H, s)
28	COOH	178.7	—		179.3	—	
29	CH <sub>2</sub>	108.7	4.60 (1H, s) 4.71 (1H, s)		109.4	4.42 (1H, s) 4.54 (1H, s)	
30	CH <sub>3</sub>	18.2	1.70 (3H, s)		19.1	1.51	(3H, s)

**Figure 2:** DQF-COSY, NOESY, and HMBC correlations of arjunolic acid from *S. aqueum*.

correlated with H-18 ( $\delta_{\text{H}}$ : 2.86 ppm) on the HMBC spectrum. The double bond carbons at positions C-12 and C-13 were appeared at  $\delta_{\text{C}}$ : 121.9 ppm (methine carbon) and  $\delta_{\text{C}}$ : 144.1 ppm (quaternary carbon). The remaining quaternary carbon signals were shown at  $\delta_{\text{C}}$ : 42.7 ppm (C-4), 39.2 ppm (C-8), 41.6 ppm (C-14), 46.3 ppm (C-17), and 30.2 ppm (C-20). Carbon signals at  $\delta_{\text{C}}$ : 12.4 ppm (C-24), 16.2 ppm (C-25), 16.4 ppm (C-26), 25.1 ppm (C-27), 32.1 ppm (C-29), and 22.6 ppm (C-30) were suggested as primary carbons. The presence of eight quaternary carbons,

six methine carbons, ten methylene carbons, and six methyl carbons was also confirmed by the DEPT experiment. According to the above data, the structure of Compound 2 was verified as arjunolic acid, and its molecular formula was  $\text{C}_{30}\text{H}_{48}\text{O}_5$ . DQF-COSY ( $^1\text{H}$ - $^1\text{H}$ ), NOESY ( $^1\text{H}$ - $^1\text{H}$ ), and HMBC ( $^1\text{H}$ - $^{13}\text{C}$ ) correlations of the structure are demonstrated in Figure 2. And then, the spectral data of both  $^1\text{H}$  NMR (600 MHz, methanol- $d_4$ ) with their coupling constants and  $^{13}\text{C}$  NMR (150 MHz, methanol- $d_4$ ) are shown in Table 2, completed with the reported literature data.

**Table 2:** NMR spectra of arjunolic acid by comparing to literature

Position	DEPT	Experiment			Literature [22]		
		$\delta_{\text{C}}$ (ppm)	$\delta_{\text{H}}$ (ppm) (mult, $J$ in Hz)		$\delta_{\text{C}}$ (ppm)	$\delta_{\text{H}}$ (ppm) (mult, $J$ in Hz)	
1	CH <sub>2</sub>	46.5	0.89 (1H, m) 1.95 (1H, m)		47.8	—	
2	CH	68.8	3.68 (1H, m)		69.6	3.66 (1H, m)	
3	CH	76.8	3.35 (1H, d, $J$ = 10.2 Hz)		78.1	3.36 (1H, m)	
4	C	42.7	—		43.0	—	
5	CH	46.8	1.28 (1H, m)		48.3	—	
6	CH <sub>2</sub>	17.7	1.39 (1H, m) 1.48 (1H, m)		19.1	—	
7	CH <sub>2</sub>	32.0	1.28 (1H, m) 1.64 (1H, m)		33.8	—	
8	C	39.2	—		40.7	—	
9	CH	47.6	1.69 (1H, m)		48.1	—	
10	C	37.7	—		38.5	—	
11	CH <sub>2</sub>	23.2	1.95 (1H, m) 1.24 (1H, m)		24.0	—	
12	CH	121.9	5.25 (1H, m)		123.4	5.23 (1H, m)	
13	C	144.1	—		144.3	—	
14	C	41.6	—		42.7	—	
15	CH <sub>2</sub>	27.4	1.07 (1H, m) 1.78 (1H, m)		29.1	—	
16	CH <sub>2</sub>	22.6	1.59 (1H, m) 2.24 (1H, m)		24.4	—	
17	C	46.3	—		46.6	—	
18	CH	41.4	2.86 (1H, dd, $J$ = 3.9, 13.7 Hz)		43.3	—	
19	CH <sub>2</sub>	45.9	1.13 (2H, m)		44.1	—	
20	C	30.2	—		31.6	—	
21	CH <sub>2</sub>	33.5	1.39 (1H, m) 1.21 (1H, m)		34.8	—	
22	CH <sub>2</sub>	32.5	1.75 (1H, m) 1.54 (1H, m)		33.5	—	
23	CH <sub>2</sub>	65.0	3.50 (1H, d, $J$ = 11.0 Hz) 3.27 (1H, d, $J$ = 11.0 Hz)		66.2	3.50 (1H, s) 3.31 (1H, s)	
24	CH <sub>3</sub>	12.4	0.70 (3H, s)		13.9	0.69 (3H, s)	
25	CH <sub>3</sub>	16.2	1.03 (3H, s)		17.7	0.81 (3H, s)	
26	CH <sub>3</sub>	16.4	0.82 (3H, s)		17.5	0.91 (3H, s)	
27	CH <sub>3</sub>	25.1	1.18 (3H, s)		26.8	1.17 (3H, s)	
28	COOH	180.7	—		181.8	—	
29	CH <sub>3</sub>	32.1	0.91 (3H, s)		25.3	1.03 (3H, s)	
30	CH <sub>3</sub>	22.6	0.94 (3H, s)		24.6	0.94 (3H, s)	

**Table 3:** Anticancer activity against HeLa, T47D, and A549 cell lines

Compound name	MTT assay ( $IC_{50} \pm SD$ $\mu\text{g/mL}$ )		XTT assay ( $IC_{50} \pm SD$ $\mu\text{g/mL}$ )
	HeLa	T47D	A549
Alphitolic acid	16.12 $\pm 0.681$	7.37 $\pm 0.388$	$84.41 \pm 0.048$
Arjunolic acid	6.74 $\pm 0.015$	27.51 $\pm 0.036$	$168.22 \pm 0.090$
Doxorubicin	2.67 $\pm 0.25$	0.035 $\pm 0.012$	NT

The meaning of NT is not test, SD = standard deviation.

### 3.2 Anticancer activity

Isolated compounds in this study were evaluated their anticancer activity against the human cancer cells as HeLa (cervical cancer), T47D (breast cancer), and A549 (lung cancer) cell lines using MTT and XTT assays, which are the colorimetric measurement of cell viability. In the present anticancer activity test, doxorubicin was used as the standard. The pure compound is deemed highly toxic if the half-maximal inhibitory concentration ( $IC_{50}$ ) is  $\leq 4 \mu\text{g/mL}$  and toxic if the  $IC_{50}$  is  $\leq 20 \mu\text{g/mL}$ . The compound is moderately toxic if the  $IC_{50}$  is  $20\text{--}100 \text{ g/mL}$  and if  $IC_{50}$  above  $100 \text{ g/mL}$  is recommended as non-toxic [23,24].

Based on anticancer activity data obtained, alphitolic acid was suggested as a toxic compound against HeLa and T47D cell lines, with  $IC_{50}$  being  $16.12 \mu\text{g/mL}$  and  $7.37 \mu\text{g/mL}$ , respectively. However, its activity against the A549 cell line was presented as moderate inhibition ( $IC_{50} = 84.41 \mu\text{g/mL}$ ). Moreover, arjunolic acid with an  $IC_{50}$  of  $6.74 \mu\text{g/mL}$  was considered a toxic compound against HeLa cell lines. Arjunolic acid was a moderate compound against the T47D cancer cell line with an  $IC_{50}$  of  $27.15 \mu\text{g/mL}$ , although it was a nontoxic compound against the A549 cancer cell line. According to the data mentioned earlier, the two types of triterpenoids inhibited higher toxicity against HeLa and T47D cell lines than the A549 cell line, and their activities may improve to obtain the natural origin drug from this species. The  $IC_{50}$  values of the two compounds are listed in Table 3.

Many terpene compounds have interesting bioactivities, so it is necessary to carry out other activity tests such as agar-well diffusion methods to evaluate the minimum inhibitory concentrations, minimum bactericidal concentration,  $IC_{50}$ , and zone of inhibitions that could determine the potential antimicrobial and antifungal efficacy. The significant antioxidant potential could be evaluated through

1,1-diphenyl-2-picryl-hydrazyl and 2,2'-Azino-bis(3-ethyl-benzothiazoline-6-sulfonic acid) (ABTS) assays. The inhibiting of the  $\alpha$ -glucosidase represents an antidiabetic activity that also could be visualized through molecular docking simulations. The effective anti-inflammatory agent could be determined via the carrageenan-induced assay, while an appreciable analgesic activity could be observed through the acetic acid-induced writhing bioassay [25].

## 4 Conclusion

This study reports the success of isolating two bioactive triterpenoid compounds from stem bark of *S. aqueum*, an Indonesian medicinal plant, namely alphitolic acid and arjunolic acid. These two isolated triterpenoids were significantly toxic against cervical cancer and breast cancer, shown by small  $IC_{50}$  values. Based on this result, we recommended continuing exploring the potential of this plant so that it can be used as a source of secondary metabolite compounds with interesting bioactivity.

**Acknowledgments:** All authors express big thanks to the Indonesian Ministry of Education and Culture for the Research Grant (Penelitian Dasar, 2021), to the Faculty of Pharmacy, Meijo University, Tempaku, Nagoya, Japan, for NMR and MS data, to the Faculty of Pharmacy, Gajah Mada University, Indonesia, and the Faculty of Science, UPM, Malaysia for an anticancer test.

**Funding information:** This work was supported by the Indonesian Ministry of Education and Culture for the Research Grant (Penelitian Dasar, 2021).

**Author contributions:** Alfinda Novi Kristanti: conceptualization, formal analysis, funding acquisition, investigation, resources, supervision, and writing – review and editing; Ei Ei Aung: data curation, formal analysis, investigation, project administration, validation, visualization, and writing – original draft; Nanik Siti Aminah: conceptualization, formal analysis, funding acquisition, methodology, resources, and supervision; Yoshiaki Takaya: data curation, resources, software, validation, and visualization; Hnin Thanda Aung: data curation, formal analysis, methodology, and supervision; and Rico Ramadhan: data curation, formal analysis, software, and validation.

**Conflict of interest:** Authors state no conflict of interest.

**Ethical approval:** The conducted research is not related to either human or animal use.

**Data availability statement:** All data generated or analyzed during this study are included in this published article [and its supplementary information files].

## References

- [1] Nazir N, Zahoor M, Uddin F, Nisar M. Chemical composition, in vitro antioxidant, anticholinesterase, and antidiabetic potential of essential oil of *Elaeagnus umbellata* Thunb. *BMC Complement Med Ther.* 2021;21:73. doi: 10.1186/s12906-021-03228-y.
- [2] Shah SMM, Ullah F, Shah SMH, Zahoor M, Sadiq A. Analysis of chemical constituents and antinociceptive potential of essential oil of *Teucrium stocksianum* bioss collected from the North West of Pakistan. *BMC Compl Altern Med.* 2012;12:244. doi: 10.1186/1472-6882-12-244.
- [3] Takuli P, Khulbe K, Kumar P, Parki A, Syed A, Elgorban AM. Phytochemical profiling, antioxidant and antibacterial efficacy of a native Himalayan Fern: *Woodwardia unigemmata* (Makino) Nakai. *Saudi J Biol Sci.* 2020;27:1961–7. doi: 10.1016/j.sjbs.2020.06.006.
- [4] Elfahmi Y, Woerdenbag HJ, Kayser O. Jamu: Indonesia traditional herbal medicine towards rational phytopharmacological use. *J Herb Med.* 2016;4(2):51–73. doi: 10.1016/j.hermed.2014.01.002.
- [5] Memon AH, Ismail Z, Al-Suede FSR, Aisha AFA, Hamil MSR, Saeed MAA. Isolation, characterization, crystal structure elucidation of two flavanones and simultaneous RP-HPLC determination of five major compounds from *Syzygium campanulatum* Korth. *Molecules.* 2015;20:14212–33. doi: 10.3390/molecules200814212.
- [6] Abera B, Adane L, Mamo F. Phytochemical investigation the root extract of *Syzygium guineense* and isolation of 2,3,23-trihydroxy methyl oleanate. *J Pharmacogn Phytochem.* 2018;7(2):3104–11.
- [7] Sobeh M, Mahmoud MF, Petruk G, Rezaq S, Ashour ML, Youssef FS. *Syzygium aqueum*: a polyphenol-rich leaf extract exhibits antioxidant, hepatoprotective, pain-killing and anti-inflammatory activities in animal models. *Front Pharmacol.* 2018;9:1–14. doi: 10.3389/fphar.2018.00566.
- [8] Osman H, Rahim AA, Isa NM, Nornaemah M, Bakhir NM. Antioxidant activity and phenolic content of *Paederia foetida* and *Syzygium aqueum*. *Molecules.* 2009;14:970–8. doi: 10.3390/molecules14030970.
- [9] Sujanapal P, Sankaran KV. *Common plants of maldives.* Bangkok: FAO; 2016.
- [10] Costa JS, Cruz ENS, Setzer WN, Silva JKRS, Maia JGS, Figueiredo PLB. Essential oils from Brazilian *Eugenia* and *Syzygium* species and their biological activities. *Biomolecules.* 2020;10(1155):1–36. doi: 10.3390/biom10081155.
- [11] Palanisamy UD, Ling LT, Manaharan T, Sivapalan V, Subramaniam T, Helme MH. Standard extract of *Syzygium aqueum*: a safe cosmetic ingredient. *Int J Cosmet Sci.* 2011;33:269–75. doi: 10.1111/j.1468-2494.2010.00637.x.
- [12] Nonaka G, Aiko Y, Aritake K, Nishioka I, Samarangeni AB. novel proanthocyanidins with doubly bonded structures, from *Syzygium samarangense* and *Syzygium aqueum*. *Chem Pharm Bull.* 1992;40(10):2671–3.
- [13] Aung EE, Kristanti AN, Amina NS, Takaya Y, Ramadhan R. Isolation of 2,4-di-tert-butylphenol and butyrospermol 3- $\beta$ -O-palmitate from *Syzygium aqueum* stem bark. *MOJ Eco Environ Sci.* 2020;5(4):193–7. doi: 10.15406/mojes.2020.05.00193.
- [14] Manaharan T, Appleton D, Cheng HM, Palanisamy UD. Flavonoids isolated from *Syzygium aqueum* leaf extract as potential antihyperglycaemic agents. *Food Chem.* 2012;132:1802–7. doi: 10.1016/j.foodchem.2011.11.147.
- [15] Subarnas A, Diantini A, Abdulah R, Zuhrotun A, Hadisaputri YE, Puspitasari IM. Apoptosis induced in MCF-7 human breast cancer cells by 2',4'-gihydroxy-6-methoxy-3,5-dimethylchalcone isolated from *Eugenia aquea* burm f. leaves. *Oncol Lett.* 2015;9:2303–6. doi: 10.3892/ol.2015.2981.
- [16] Aung EE, Kristanti AN, Amina NS, Takaya Y, Ramadhan R. Cytotoxicity of butyrospermol and sitosterone from the stem bark of *Syzygium aqueum*. *Trop. J Nat Prod Res.* 2020;4(11):899–904. doi: 10.26538/tjnpr/v4i11.10.
- [17] Aung EE, Kristanti AN, Amina NS, Takaya Y, Ramadhan R, Aung HT. Anticancer activity of isolated compounds from *Syzygium aqueum* stem bark. *Rasayan J Chem.* 2021;14(1):312–8. doi: 10.31788/RJC.2021.1416106.
- [18] Cancer Chemoprevention Research Centre (CCRC); 2014. *Prosedur tetap uji sitotoksik metode MTT*, Faculty of Pharmacy Gadjah Mada University (UGM), Yogyakarta, Indonesia (in Indonesian). [http://ccrc.farmasi.ugm.ac.id/wp-content/uploads/10\\_sop-uji-sitotoksik-metode-mtt.pdf](http://ccrc.farmasi.ugm.ac.id/wp-content/uploads/10_sop-uji-sitotoksik-metode-mtt.pdf).
- [19] Mutiah R, Indradmojo C, Dwi HH, Griana TP, Listyana A, Ramdhani R. Induction of apoptosis and phase-cell cycle inhibition of G0-G1, S, G2-M of T47D breast cancer cells on treatment with ethyl acetate fraction of jackfruit parasite leaves (*Macrosolen cochinchensis*). *J Appl Pharm Sci.* 2017;7(10):138–43. doi: 10.7324/JAPS.2017.71020.
- [20] American Type Culture Collection (ATCC). *XTT cell proliferation assay Kit Instruction Manual*. <https://www.atcc.org/products/30-1011k>.
- [21] Marzouk AM. Hepatoprotective triterpenes from hairy root cultures of *Ocimum basilicum* L. *Z Naturforsch.* 2009;64c(3–4):201–9. doi: 10.1515/znc-2009-3-409.
- [22] Nono RN, Barboni L, Teponno RB, Quassinti L, Bramucci M, Vitali LA. Antimicrobial, antioxidant, anti-inflammatory activities and phytoconstituents of extracts from the roots of *Dissotis thollonii* Cogn. (Melastomataceae). *S Afr J Bot.* 2014;93:19–26.
- [23] Sriwiryajan S, Ninpesh T, Sukpondma Y, Nasomyon T, Graidist P. Cytotoxicity screening of plants of genus piper in breast cancer cell lines. *Trop J Pharm Res.* 2014;13(6):921–8. doi: 10.4314/tjpr.v13i6.14.
- [24] Fikroh RA, Matsjeh S, Anwar C. Synthesis and anticancer activity of (E)-2'-hydroxy-2-bromo-4,5-dimethoxychalcone against breast cancer (MCF-7) cell line. *Molekul.* 2020;15(1):34–9. doi: 10.20884/1.jm.2020.15.1.558.
- [25] Shah M, Murad W, Rehman NU, Mubin S, Al-Sabahi JN, Ahmad M, et al. GC-MS analysis and biomedical therapy of oil from n-hexane fraction of *Scutellaria edelbergii* Rech. f.: in vitro, in vivo, and in silico approach. *Molecules.* 2021;26:7676. doi: 10.3390/molecules26247676.