



purwo sri rejeki <purwo-s-r@fk.unair.ac.id>

Chiang Mai University Journal of Natural Sciences - Manuscript ID CMUJ-2021-00781 message

Priraya Rithaporn <onbehalf@manuscriptcentral.com>

Wed, Apr 14, 2021 at 1:16 PM

Reply-To: prirayar@hotmail.com

To: purwo-s-r@fk.unair.ac.id, purwo_faal@yahoo.com

14-Apr-2021

Dear Dr. Rejeki:

Your manuscript entitled "A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mus musculus" has been successfully submitted online and is presently being given full consideration for publication in the Chiang Mai University Journal of Natural Sciences.

Your manuscript ID is CMUJ-2021-0078.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your street address or e-mail address, please log in to ScholarOne Manuscripts at https://mc04.manuscriptcentral.com/cmuj_ns and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to https://mc04.manuscriptcentral.com/cmuj_ns.

Thank you for submitting your manuscript to the Chiang Mai University Journal of Natural Sciences.

Sincerely,
Chiang Mai University Journal of Natural Sciences Editorial Office

Chiang Mai University Journal of Natural Sciences - Decision on Manuscript ID CMUJ-2021-0078

1 message

Korakot Nganvongpanit <onbehalf@manuscriptcentral.com>

Tue, Jul 27, 2021 at 11:36 AM

Reply-To: korakot.n@cmu.ac.th

To: purwo-s-r@fk.unair.ac.id, purwo_faal@yahoo.com

27-Jul-2021

Dear Dr. Rejeki:

Manuscript ID CMUJ-2021-0078 entitled "A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mus musculus" which you submitted to the Chiang Mai University Journal of Natural Sciences, has been reviewed. The comments of the reviewer(s) are included at the bottom of this letter.

The reviewer(s) have recommended publication, but also suggest some major revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript.

To revise your manuscript, log into https://mc04.manuscriptcentral.com/cmuj_ns and enter your Author Center, where you will find your manuscript title listed under "Manuscripts with Decisions." Under "Actions," click on "Create a Revision." Your manuscript number has been appended to denote a revision.

You may also click the below link to start the revision process (or continue the process if you have already started your revision) for your manuscript. If you use the below link you will not be required to login to ScholarOne Manuscripts.

*** PLEASE NOTE: This is a two-step process. After clicking on the link, you will be directed to a webpage to confirm. ***

https://mc04.manuscriptcentral.com/cmuj_ns?URL_MASK=4a02fd6934594279b2f5a2dbcdf46865

You will be unable to make your revisions on the originally submitted version of the manuscript. Instead, revise your manuscript using a word processing program and save it on your computer. Please also highlight the changes to your manuscript within the document by using the track changes mode in MS Word or by using bold or colored text.

Once the revised manuscript is prepared, you can upload it and submit it through your Author Center.

When submitting your revised manuscript, you will be able to respond to the comments made by the reviewer(s) in the space provided. You can use this space to document any changes you make to the original manuscript. In order to expedite the processing of the revised manuscript, please be as specific as possible in your response to the reviewer(s). Also, please be sure to indicate by page and line number where the corresponding changes can be found in the revised version of your paper.

IMPORTANT: Your original files are available to you when you upload your revised manuscript. Please delete any redundant files before completing the submission.

Because we are trying to facilitate timely publication of manuscripts submitted to the Chiang Mai University Journal of Natural Sciences, your revised manuscript should be submitted by 26-Sep-2021. If it is not possible for you to submit your revision by this date, we may have to consider your paper as a new submission.

Once again, thank you for submitting your manuscript to the Chiang Mai University Journal of Natural Sciences and I look forward to receiving your revision.

Sincerely,
Dr. Korakot Nganvongpanit
Editor-in-Chief, Chiang Mai University Journal of Natural Sciences
korakot.n@cmu.ac.th

Note by editor:

This is an interesting study. In view of all this, it was found to require revision to complete the study. Please response all comments point by point. Making a highlight in all change of revision manuscript for easy visibility. Moreover, It

would be our thankful if you can cite some papers published in CMUJ Nat. Sci. because we published many papers related to your work.

Editor comments;

1 Figure 3 Author wrote “*Significant vs. positive control group” but in figure show * on K1 group. It is wrong, please revision.

2 We do not publish supplement data. All data (table and figure) in supplement are importance please move to main text.

Reviewer(s)' Comments to Author:

Comments to the Author

The title, I suggest:

A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mice (*Mus musculus*)
Mus musculus --> in italic

Abstract:

The statistical analysis performed using One Way-ANOVA and Least Significant Difference (LSD) post hoc test using Statistic Package for Social Science (SPSS) version 16 --> it can be skipped
please briefly explain: K1, K2, etc

Tables & figures:

explain what K1, K2, etc are, and write them below the tables and figures



purwo sri rejeki <purwo-s-r@fk.unair.ac.id>

Chiang Mai University Journal of Natural Sciences - Manuscript ID CMUJ-2021-0078.R1

1 message

Priraya Rithaporn <onbehalf@manuscriptcentral.com>
Reply-To: prirayar@hotmail.com
To: purwo-s-r@fk.unair.ac.id, purwo_faal@yahoo.com

Sat, Jul 31, 2021 at 11:29 AM

31-Jul-2021

Dear Dr. Rejeki:

Your manuscript entitled "A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mice (*Mus musculus*)" has been successfully submitted online and is presently being given full consideration for publication in the Chiang Mai University Journal of Natural Sciences.

Your manuscript ID is CMUJ-2021-0078.R1.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your street address or e-mail address, please log in to ScholarOne Manuscripts at https://mc04.manuscriptcentral.com/cmuj_ns and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to https://mc04.manuscriptcentral.com/cmuj_ns.

Thank you for submitting your manuscript to the Chiang Mai University Journal of Natural Sciences.

Sincerely,
Chiang Mai University Journal of Natural Sciences Editorial Office



purwo sri rejeki <purwo-s-r@fk.unair.ac.id>

Chiang Mai University Journal of Natural Sciences - Decision on Manuscript ID CMUJ-2021-0078.R1

2 messages

Korakot Nganvongpanit <onbehalf@manuscriptcentral.com>

Mon, Aug 9, 2021 at 3:13 PM

Reply-To: korakot.n@cmu.ac.th

To: purwo-s-r@fk.unair.ac.id, purwo_faal@yahoo.com

09-Aug-2021

Dear Dr. Rejeki:

It is a pleasure to accept your manuscript entitled "A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mice (*Mus musculus*)" in its current form for publication in the Chiang Mai University Journal of Natural Sciences. The comments of the reviewer(s) who reviewed your manuscript are included at the foot of this letter.

Thank you for your fine contribution. On behalf of the Editors of the Chiang Mai University Journal of Natural Sciences, we look forward to your continued contributions to the Journal.

Please be advised that you will be expected to assist the Journal with any editorial questions that may arise as part of the final editing and publication process, without which assistance your manuscript may potentially not be printed.

Sincerely,
Dr. Korakot Nganvongpanit
Editor-in-Chief, Chiang Mai University Journal of Natural Sciences
korakot.n@cmu.ac.th

Associate Editor Comments to Author:

Associate Editor
Comments to the Author:
(There are no comments.)

Reviewer(s)' Comments to Author:

purwo sri rejeki <purwo-s-r@fk.unair.ac.id>

Tue, Aug 10, 2021 at 8:07 PM

To: korakot.n@cmu.ac.th

Dear Dr. Korakot Nganvongpanit

Thank you for your letter. We appreciate and really glad to hear the decision of our article from you. We hope we will have other opportunities for publishing our other manuscripts in your journal in the future.

Sincerely
Purwo Sri Rejeki

[Quoted text hidden]



A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mice (*Mus musculus*)

Journal:	<i>Chiang Mai University Journal of Natural Sciences</i>
Manuscript ID	CMUJ-2021-0078.R1
Manuscript Type:	Original Article
Keywords:	Cancer, Diet, Inflammation

SCHOLARONE™
Manuscripts

1
2
3 **A high-fat diet decreases serum TNF-alpha and breast tumor area**
4 **on benzopyrene induced mice (*Mus musculus*)**
5
6
7
8
9

10
11 Purwo Sri Rejeki^{123*}, Dita Mega Utami⁴, Nabilah Izzatunnisa⁴, Adi Pranoto⁵, Devitya
12 Angielevi Sukarno⁶, and Minidian Fasitasari⁷
13
14
15
16
17

18
19 ¹ Department of Physiology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

20
21 ² Sport Health Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

22
23 ³ Basic Medical Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

24
25 ⁴ Medical Program, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

26
27 ⁵ Medical Science, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

28
29 ⁶ Laboratory of Physiology, Faculty of Medicine Universitas Surabaya, Surabaya, Indonesia

30
31 ⁷ Faculty of Medicine, Universitas Islam Sultan Agung Semarang, Semarang, Indonesia
32
33
34
35
36

37 *Correspondence Author:

38
39 Purwo Sri Rejeki

40
41 Department of Physiology

42
43 Faculty of Medicine, Universitas Airlangga

44
45 Prof. Dr. Moestopo No. 47 Street, Surabaya

46
47
48 Indonesia

49
50 +6282141559388

51
52 purwo-s-r@fk.unair.ac.id; purwo_faal@yahoo.com
53
54
55
56
57
58
59
60

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: PSR. Performed the experiments: PSR DMU NI. Analyzed the data: AP. Contributed reagents/materials/analysis tools: DAS MF. Wrote the paper: PSR DMU NI AP DAS MF.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

ACKNOWLEDGMENTS

This study is supported by Fundamental Research Program Ministry of Research, Technology and Higher Education, Indonesia, under Grant Number: 4/E1/KP.PTNBH/2019 and 543/UN3.14/LT/2019.

A high-fat diet decreases serum TNF-alpha and breast tumor area on benzopyrene induced mice (*Mus musculus*)

ABSTRACT

This study aims to analyze whether the high-fat diet decreases serum TNF- α and breast tumor area on benzopyrene induced mice (*Mus musculus*). This study was a true experimental with the randomized posttest-only control group design using 36 female mice (*Mus musculus*), 3-4 months age, 25 \pm 5 grams. Mice were induced with benzopyrene (BZP) subcutaneously with a dose of 0.3mg/20gBB/day for 14 days in the right breast area, then randomly divided into 6 groups, K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber). The high-fat diet was administered for 28 days. The mean of tumor area delta at K₁ (0.00 \pm 0.00) mm², K₂ (3.52 \pm 1.98) mm², K₃ (27.18 \pm 21.23) mm², K₄ (13.19 \pm 9.93) mm², K₅ (8.80 \pm 1.72) mm², K₆ (10.81 \pm 6.55) mm², and (p=0.001). The mean of TNF- α levels at K₁ (56.32 \pm 8.25) ng/mL, K₂ (65.99 \pm 2.82) ng/mL, K₃ (70.43 \pm 4.61) ng/mL, K₄ (58.05 \pm 5.80) ng/mL, K₅ (54.91 \pm 3.27) ng/mL, K₆ (59.67 \pm 3.63) ng/mL and (p=0.000). A high-fat diet lowers TNF- α levels and reduces the area of BZP-induced breast tumors. The lowest TNF- α levels and the lowest breast tumor area were found in groups with a fat : protein ratio = 60:30.

Keywords: High-fat diet, TNF- α , benzopyrene induced, breast tumor area

INTRODUCTIONS

The high-fat diet is a diet that produces blood ketones as an energy source. It is well known among people and is used for various purposes such as nonpharmacological therapy for cancer (Weber *et al.*, 2020; Zick *et al.*, 2018; Klement, 2017), weight loss (Paoli, 2014), type 2 diabetes mellitus (Saslow *et al.*, 2017), epilepsy and status epilepticus therapy in children and adults (Williams and Cervenka, 2017; Nabbout *et al.*, 2011) and also improve performance in competitive athletes (Taylor *et al.*, 2019). The high-fat diet provides a hostile environment with cancer cells, thereby causing a better prognosis (Lettieri-Barbato and Aquilano, 2018). The high-fat diet reduces the initiative, progression and metastasis of cancer cells (Zick *et al.*, 2018). But the explanation regarding the mechanism of the high-fat diet on the growth of cancer cells is still not known.

The high-fat diet is currently a new type of dietary trend in Indonesia and the world, but there are still many pros and cons of the effects of the high-fat diet on cancer. Study on the mechanism of the high-fat diet has not been done much, so the clarity is needed to monitor the benefits, side effects and complications caused by high-fat diet (Weber *et al.*, 2020; Klement, 2017). Currently, cancer is a major health problem in the world and Indonesia. GLOBOCAN data states that in 2018 there were 18.1 million new cases with a mortality rate of 9.6 million (GLOBOCAN, 2019). Cancer is the second leading cause of death in the world after cardiovascular disease. In Indonesia, the prevalence of cancer is quite high. Based on the results of the Basic Health Research in 2018, the prevalence of tumors/cancer in Indonesia reached 1.8 per 1000 population or around 480.000 people. This number increased compared to 2013, which was 1.4 per 1000 population or around 330.000 people (Riskesdas, 2018). According to GLOBOCAN

1
2
3
4 data, the incidence of cancer in Indonesia in 2018 (136.2/100.000 population) ranks 8th
5
6 in Southeast Asia, whereas in Asia ranks 23rd. The highest incidence rate in Indonesia
7
8 for women is breast cancer at 42.1 per 100.000 population with an average death of 17
9
10 per 100.000 population (GLOBOCAN, 2019). In addition to early detection and
11
12 treatment, more optimal management is needed, such as supporting therapies that can
13
14 improve the results of radiotherapy and chemotherapy.
15
16
17

18 According to the Warburg postulate increased glycolysis induces carcinogenesis.
19
20 Scientists today also have a similar hypothesis that genetic mutations cause cancer and
21
22 cancer cells metabolize glucose. Preclinical study data indicate that the insulin pathway
23
24 is related to the initiation and development of cancer. This pathway is regulated through
25
26 carbohydrate consumption (Erickson *et al.*, 2017). Interleukin-6 (IL-6), interleukin-8 (IL-
27
28 8), and serum tumor necrosis factor-alpha (TNF- α) levels can function as biological
29
30 markers of metastases and tumor invasion because studies have shown a correlation with
31
32 stage II or III carcinoma and metastases in lymph nodes (Ma *et al.*, 2017). According to
33
34 Wolczyk *et al.* (2016) high TNF- α levels are characteristic of many malignant cancers,
35
36 including breast cancer, and are often associated with cancer cell aggressiveness and poor
37
38 prognosis. Benzopyrene (BZP) is a kind of polycyclic aromatic hydrocarbon (PAH)
39
40 compound contained in cigarettes. This compound will undergo metabolism in the liver
41
42 by the cytochrome P4501A1 (CYP1A1) enzyme and produce carcinogenic benzopyrene-
43
44 7.8-dihydrodiol 9.10-epoxide (BPDE) metabolism which causes mutant cell formation
45
46 (Irmawati *et al.*, 2018). This compound can enter the body's tissues through the skin or
47
48 through the mouth by inhalation. Induction of BZP compounds with small doses has a
49
50 carcinogenic effect. According to some studies, induction of at least 0.24-0.3 mg of BZP
51
52 per 20 grams of body weight per day for 12-28 days can cause cancer in induced sites
53
54
55
56
57
58
59
60

(Irmawati *et al.*, 2018; Nansi *et al.*, 2015; Gao *et al.*, 2011; Anandakumar *et al.*, 2009).

The high-fat diet is a diet that uses a ratio of predominantly fat and low carbohydrate energy sources resulting in a state of ketosis in which the body will use the ketone bodies as the primary energy source (van Berkel *et al.*, 2018).

The high-fat diet is assumed to reduce blood glucose levels, so it is expected that there will be a decrease in the growth of cancer cells. The progression of cancer cells can be seen from serum TNF- α levels. This study aims to analyze whether the high-fat diet decreases serum TNF- α levels and breast tumor area on benzopyrene induced mice (*Mus musculus*). We hypothesized that the high-fat diet can decrease serum TNF- α levels and breast tumor area on benzopyrene induced mice (*Mus musculus*).

MATERIALS AND METHODS

Animals, diets, and experimental protocol

This study was a true experimental with the randomized posttest-only control group design using 36 female mice (*Mus musculus*), 3-4 months age, 25 \pm 5 grams. Mice were divided into 6 groups randomly. In K₁, BZP was not induced, while K₂-K₆ were induced with BZP subcutaneously as much as 0.3 mg/20gBB/day for 14 days in the right breast area. During D1-14 induction all groups were given a standard diet. D15-32 (28 days) dieting namely K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber). The study was conducted at the

Embryology Laboratory of the Faculty of Veterinary Medicine, Universitas Airlangga within a period of 5 months (July to November 2019). Experimental animals were placed at room temperature $25\pm 2^{\circ}\text{C}$ with a humidity of 50-60% and the lighting was regulated by a light-dark cycle with the regulation of 12 hours of light and 12 hours of darkness (Pinthong and Suanarunsawat, 2020; Wunjuntuk *et al.*, 2020; Santos *et al.*, 2019; Gao *et al.*, 2011; Anandakumar *et al.*, 2009). Mice cages measuring 30x45x20 cm, made of plastic covered with gauze equipped with a place to eat and drink bottles, each cage is filled with 1 group (6 mice). The feeding and high-fat diet were done at 09.00-10.00 a.m. All research procedures were approved by Faculty of Veterinary Universitas Airlangga Medicine Research Ethics Commission No. 2.KE142.07.2019. The present study followed animal welfare principles in experimental science published in the European Convention for the Protection of Vertebrate Animals.

Data collection

Body weight measurements were performed at the beginning and the end of the post-high-fat diet using a HL-3650 heles digital scale (0-5kg scale). Tumor area measurements were performed at the beginning and end of post high-fat diet using Vernier Caliper (Vernier Caliper Tube 150/0.02mm 536-161 Mitutoyo MT0003767 Include PPN). Measurement of blood ketone levels using Abbott Freestyle Optium Neo Blood Glucose and Ketones. Measurement of ketone levels requires a relatively small blood volume, i.e. 0.6-1.5 μL /test taken from the tail vein (Sussman *et al.*, 2013; Pinthong and Suanarunsawat, 2020). Measurement of ketone levels was carried out 24 hours after the last high-fat diet was given.

1
2
3
4 Blood was drawn from the left ventricle of experimental animals as much as 1 ml.
5
6 Blood sampling was performed 24 hours after the last high-fat diet is given. Blood was
7
8 centrifuged for 10 minutes at 3000 rpm. The serum was separated and stored at -80°C for
9
10 analysis of TNF- α levels the following day. Measurement of TNF- α levels using the
11
12 Enzyme-Linked Immunosorbent Assay (ELISA) BT-E0117Mo kit (Biossay Technology
13
14 Laboratory, Inc., Shanghai, China, 2019) with a standard curve range of 5-1000 ng/L and
15
16 TNF- α sensitivity level in the kit 2.49 ng/L.
17
18
19
20
21
22

23 **Statistical analysis**

24
25 The statistical analysis was using the Social Science (SPSS) Statistics for
26
27 Windows, version 16 (SPSS Inc., Chicago, IL, USA). The normality test uses the Shapiro-
28
29 Wilk test, while the homogeneity test uses the Levene test. Different test using One way-
30
31 ANOVA and continued with Least Significant Difference (LSD) post hoc test. Linear
32
33 correlation with Pearson product-moment model. All data is displayed with mean \pm
34
35 standard deviation (SD). All statistical analyzes used a significant level ($p < 0.05$).
36
37
38
39
40

41 **RESULTS**

42
43
44 The mean analysis results of initial and final body weight after the high-fat diet
45
46 can be seen in **Figure 1**.
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Mean initial, final and Δ (delta) body weight of mice on high-fat diet

Note: *Significant vs negative control group (K₁) $p < 0.05$. K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber).

Based on these data, it is found that the mean delta body weight of mice at K₁ is higher compared to K₂, K₃, K₄, K₅ and K₆. Based on One way-ANOVA test results showed that there were significant differences in the mean delta weight of mice ($p = 0.039$). Based on the LSD post hoc test results showed that there were significant differences in the weight difference between K₂ with K₁ ($p = 0.045$), K₃ with K₁ ($p = 0.004$) and K₄ with K₁ ($p = 0.005$), while K₁ with K₅ ($p = 0.164$), K₁ with K₆ ($p = 0.164$), K₂ with K₃ ($p = 0.322$), K₂ with K₄ ($p = 0.364$), K₂ with K₅ ($p = 0.507$), K₂ with K₆ ($p = 0.507$), K₃ with K₄ ($p = 0.934$), K₃ with K₅ ($p = 0.104$), K₃ with K₆ ($p = 0.104$), K₄ with K₅ ($p = 0.121$), K₄ with K₆ ($p = 0.121$) and K₅ with K₆ ($p = 1,000$) there was no significant difference ($p > 0.05$). The results of the average analysis of ketone levels after the high-fat diet can be seen in **Figure 2**.




Figure 2. Mean of ketone levels after high-fat diet treatment

Note: *Significant vs negative control group (K₁) $p < 0.05$. K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber).

Based on these data, it is found that the mean ketone levels at K₅ are higher compared to K₁, K₂, K₃, K₄ and K₆. Based on One way-ANOVA test results there were significant differences in the average ketone levels in mice ($p < 0.05$). The results of the analysis of the mean tumor area between the initial and final tumors area after the administration of the high-fat diet can be seen in **Table 1**.

Table 1. Mean of initial and final tumor areas after the treatment of the high-fat diet

Group	Tumor Areas (mm ²)			One way-ANOVA Δ (<i>p</i> -values)
	Initial	Final	delta (Δ)	
K ₁	0.00±0.00	0.00±0.00	0.00±0.00 ^a	0.001
K ₂	0.35±0.32	3.87±2.27	3.52±1.98 ^a	
K ₃	0.65±0.33	27.83±20.98	27.18±21.23 ^b	
K ₄	0.63±0.59	13.83±10.19	13.19±9.93 ^a	
K ₅	2.17±2.86	9.80±1.83	8.80±1.72 ^a	
K ₆	1.19±1.09	12.00±7.54	10.81±6.55 ^a	

Note: Different superscript shows significant differences in the LSD post hoc test with ($p < 0.05$). K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0%

carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber).

Based on these data, it is found that the mean delta area of mice tumors at K₃ is wider than K₁, K₂, K₄, K₅ and K₆. Based on the One way-ANOVA test results there were significant differences in the mean delta area of the tumor in mice ($p < 0.05$). The results of a comparative analysis of tumor area with body weight can be seen in **Table 2**.

Table 2. Comparison mean of final tumor area and final body weight

Group	Final Tumor Areas (mm ²)	Body Weight (gram)	TA/BW	One way-ANOVA TA/BW (<i>p-values</i>)
K ₁	0.00±0.00	30.00±2.61	0.00±0.00 ^a	0.000
K ₂	3.87±2.27	26.33±3.72	0.15±0.09 ^a	
K ₃	27.83±20.98	24.00±4.34	1.09±0.68 ^{bc}	
K ₄	13.83±10.19	23.00±4.19	0.62±0.48 ^b	
K ₅	9.80±1.83	26.67±1.21	0.36±0.07 ^a	
K ₆	12.00±7.54	27.00±3.95	0.46±0.28 ^b	

Note: Different superscript shows significant differences in the LSD post hoc test with ($p < 0.05$). TA: tumor area; BW: body weight. K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber).

Based on these data, it is found that the mean comparison of final tumor area with final body weight at K₃ is higher compared to K₁, K₂, K₄, K₅ and K₆. Based on the One way-ANOVA test results there were significant differences in the mean of the final tumor area with body weight in mice ($p < 0.05$). The results of the mean analysis of TNF- α levels after the administration of the high-fat diet can be seen in **Figure 3**.




Figure 3. Mean TNF- α levels in each group after the high-fat diet

Note: *Significant vs. positive control group, standard feed (K₂), and high-fat diet group with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber (K₃) $p < 0.05$. K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber).


Based on these data, it is found that the average level of TNF- α at K₃ is higher than K₁, K₂, K₄, K₅ and K₆. Based on One way-ANOVA test results there was a significant difference in the mean TNF- α levels ($p = 0.000$). Based on the LSD post hoc test results there were significant differences in the mean TNF- α levels between K₁ with K₂ ($p = 0.002$), K₄ with K₂ ($p = 0.011$), K₅ with K₂ ($p = 0.001$), K₆ with K₂ ($p = 0.039$), K₁ with K₃ ($p = 0.000$), K₄ with K₃ ($p = 0.000$), K₅ with K₃ ($p = 0.000$), K₆ with K₃ ($p = 0.001$), while K₁ with K₄ ($p = 0.559$), K₁ with K₅ ($p = 0.636$), K₁ with K₆ ($p = 0.261$), K₂ with K₃ ($p = 0.140$), K₄ with K₅ ($p = 0.294$), K₄ with K₆ ($p = 0.583$) and K₅ with K₆ ($p = 0.115$) did not show significant differences in mean TNF- α levels ($p > 0.05$). The results of the comparative analysis of TNF- α levels with body weight can be seen in **Table 3**.

Table 3. Comparison mean of TNF- α levels with final body weight

Group	TNF- α Levels (ng/mL)	Body Weight (gram)	TNF- α /BW	One way-ANOVA TNF- α /BW (<i>p</i> -values)
K ₁	0.00±0.00	30.00±2.61	1.89±0.35 ^a	0.004
K ₂	3.87±2.27	26.33±3.72	2.55±0.38 ^b	
K ₃	27.83±20.98	24.00±4.34	3.03±0.68 ^b	
K ₄	13.83±10.19	23.00±4.19	2.62±0.65 ^b	
K ₅	9.80±1.83	26.67±1.21	2.06±0.19 ^a	
K ₆	12.00±7.54	27.00±3.95	2.26±0.47 ^a	

Note: Different superscript shows significant differences in the LSD post hoc test with ($p < 0.05$). TNF- α : tumor necrosis factor alpha; BW: body weight. K₁ (negative control group, given standard feed), K₂ (positive control group, standard feed), K₃ (high-fat diet with a ratio of 60% protein, 0% carbohydrate, 30% fat, 10% fiber), K₄ (high-fat diet with a ratio of 45% protein, 0% carbohydrate, 45% fat, 10% fiber), K₅ (high-fat diet with a ratio of 30% protein, 0% carbohydrate, 60% fat, 10% fiber) and K₆ (high-fat diet on day 15 with a ratio of 15% protein, 0% carbohydrate, 75% fat, 10% fiber).

Based on these data, it is found that comparison mean of TNF- α with final body weight at K₃ is higher compared to K₁, K₂, K₄, K₅ and K₆. Based on One way-ANOVA test results there was a significant difference in mean placing TNF- α levels with final body weight ($p = 0.004$). The parametric analysis by Pearson's product-moment linear correlation shows that the tumor areas were significantly and negatively correlated with the body weight ($r = 0.482, p \leq 0.01$) and TNF- α levels ($r = 0.447, p \leq 0.01$). By contrast, the tumor areas were negatively associated with ketone levels ($r = -0.352, p \leq 0.05$) (Figure 4).






Figure 4. Tumor areas were positively associated with body weight, TNF- α levels. Tumor areas were negatively associated with keton levels. Pearson correlation coefficients (r) and p -values are shown in each graph.

DISCUSSION

The high-fat diet is a diet that produces ketones in the blood as an energy source and can be used as a nonpharmacological therapy for cancer (Zick *et al.*, 2018; Klement, 2017), lose weight (Paoli, 2014), type 2 diabetes mellitus (Saslow *et al.*, 2017), epilepsy and status epilepticus therapy in children and adults (Williams and Cervenka, 2017), also improve performance in competitive athletes (Taylor *et al.*, 2019). The administration of the high-fat diet has been shown to increase blood ketone levels in both rats and mice

(Roberts *et al.*, 2017; Nandivada *et al.*, 2016; Meidenbauer *et al.*, 2014). Based on the results of the present study showed that the mean ketone levels at K₅ were higher compared to K₁, K₂, K₃, K₄ and K₆. The One way-ANOVA test results showed that there were significant differences in the ketone levels mean in mice ($p < 0.05$). These results are in line with the results of a study conducted by Kephart *et al.* (2017) which concluded that the administration of the high-fat diet with a composition of 23% protein, 10% carbohydrate and 67% fat significantly increased serum β -ketones or β -hydroxybutyrate compared to the control group. Likewise, research conducted by Nandivada *et al.* (2016) concluded that the administration of the high-fat diet with a composition of 90.5% fat, 9.1% protein, and 0.4% carbohydrate significantly increased serum beta-hydroxybutyrate levels in mice. As well as the results of a study conducted by Sussman *et al.* (2013) which concluded that the administration of the high-fat diet with a composition of 67.4% fat, 0.6% carbohydrate, and 15.3% protein wt/wt for 30 days significantly increased ketone levels compared with controls. The study conducted by Roberts *et al.* (2017) also found similar results that the administration of the high-fat diet (89% kcal of fat) for 13 days and 26 days significantly increased β -hydroxybutyrate compared with the control group and the low-carbohydrate diets group. Increased levels of ketones in K₅ may be caused by the effect of the high-fat diet. The high-fat diet causes a decrease in blood glucose levels, thereby increasing lipolysis (Allen *et al.*, 2014). Increased lipolysis will continuously cause an increase in ketone production in the blood to be used as an energy source in all tissues (Weber *et al.*, 2020; van Berkel *et al.*, 2018).

During the high-fat diet the brain will look for alternative energy sources that come from the excess production of acetyl-CoA, the ketone bodies (Weber *et al.*, 2020; Soldati *et al.*, 2018). Ketone bodies consist of acetoacetate (AcAc), β -hydroxybutyrate

1
2
3
4 (βHB) and acetone (Weber *et al.*, 2020). Ketone bodies are produced in the liver by the
5 breakdown of fat or ketosis (Page *et al.*, 2009). Ketosis is a physiological mechanism that
6
7 can be an indicator of decreased body fat (Kephart *et al.*, 2017). Under the normal diet,
8
9 ketone bodies concentrations are less than 0.3 mmol/L and will not be used as an energy
10
11 source by the brain until this concentration reaches 4 mmol/L, the same as glucose
12
13 concentration. At this concentration, the ketone body will be used as an energy source by
14
15 all tissues. The maximum concentration of ketone bodies in individuals on a high-fat diet
16
17 will reach 7-8 mmol/L and without accompanied by changes in blood pH (pH=7.4).
18
19 Ketosis in the high-fat diet is still classified as physiological ketosis in contrast to diabetic
20
21 ketoacidosis which has a blood ketone level reaching 20 mmol/L and a decrease in blood
22
23 pH to <7.3 (Wisnu *et al.*, 2017; Azar *et al.*, 2016; Fukao *et al.*, 2004).
24
25
26
27
28

29
30 The high-fat diet causes a decrease in blood glucose, insulin, and insulin-like
31
32 growth factor (IGF-1) which has the potential to reduce cancer cell proliferation (Weber
33
34 *et al.*, 2018; Stock *et al.*, 2009) reduce the development of cancer cells and increase
35
36 normal cell resistance (Erickson *et al.*, 2017). The high-fat diet causes cancer cells to use
37
38 more glycolysis compared to oxidative phosphorylation to produce adenosine
39
40 triphosphate (ATP) (Weber *et al.*, 2018). The high-fat diet can cause Protein Kinase B
41
42 (Akt) to induce resistance to apoptosis, cause changes in cancer cell metabolism, decrease
43
44 beta-oxidation and increase lipid synthesis in the cytosol (Rose and Vona-Davis, 2012).
45
46 Therefore, the administration of the high-fat diet can provide a hostile environment with
47
48 cancer cells, which causes a better prognosis (Lettieri-Barbato and Aquilano, 2018), can
49
50 reduce initiation, progression and metastasis from cancer cells (Zick *et al.*, 2018).
51
52
53

54
55 Tumor necrosis factor alpha (TNF-α) is a pleiotropic cytokine (has various effects
56
57 on various cells) produced mainly by active macrophages and other immune cell types
58
59
60

1
2
3
4 such as T lymphocytes, NK cells and neutrophils (Zhou *et al.*, 2016). TNF- α can function
5
6 as a biomarker of metastases and tumor invasion, because from the results of previous
7
8 studies there was a correlation with stage II or III carcinoma and metastasis in lymph
9
10 nodes (Ma *et al.*, 2017). According to Wolczyk *et al.* (2016) high levels of TNF- α are
11
12 characteristic of many malignant cancers, including breast cancer and are often associated
13
14 with cancer cell aggressiveness and poor prognosis. Based on the results of the study
15
16 showed that the average levels of TNF- α at K₅ were lower than those of K₁, K₂, K₃, K₄
17
18 and K₆. Based on One way-ANOVA test results there were significant differences in the
19
20 mean TNF- α levels in mice ($p < 0.05$). These results are in line with the previous study
21
22 conducted by Dupuis *et al.* (2015) concluded that the administration of the high-fat diet
23
24 for 14 days significantly reduced TNF- α levels compared to the control group in mice.
25
26 As well as the results of a study conducted by Vizquete *et al.* (2013) which concluded that
27
28 the administration of the high-fat diet for 8 weeks reduced TNF- α levels compared to the
29
30 ketogenic group with increased omega-3 and control in rats. In the study conducted by
31
32 Nandivada *et al.* (2016) showed that the administration of a high-fat diet with a
33
34 composition of 90.5% fat, 9.1% protein and 0.4% carbohydrate reduced post-absorptive
35
36 serum TNF- α levels compared with controls (12.5% fat, 10% protein and 77.5%
37
38 carbohydrate) in mice. However, these results differ from the postprandial serum TNF- α
39
40 levels which showed a significant increase in TNF- α levels in the high-fat diet group
41
42 compared to the control group. The results of a study conducted by Santos *et al.* (2019)
43
44 also concluded that the administration of a high-fat diet with a composition of 24.2%
45
46 carbohydrate, 60.9% fat, and 14.9% protein for 6 weeks increased TNF- α levels compared
47
48 with controls in mice. The contrast result of this study may due to a different composition
49
50 of high-fat diet. In this study the lowest level of TNF- α reduction occurred in the high-
51
52
53
54
55
56
57
58
59
60

1
2
3
4 fat diet with a composition of 30% protein, 0% carbohydrate, 60% fat, 10% fiber. This is
5
6 because the high-fat diet provides a neuroprotective effect that can reduce cell injury and
7
8 modify inflammation in the brain. The high-fat diet triggers low levels of lymphocytes or
9
10 polymorphonuclear cells so the cells ability to synthesize proinflammatory cytokines is
11
12 also reduced. In addition, the high-fat diet also decreases the synthesis of arachidonic acid
13
14 and n-6 polyunsaturated fatty acids (PUFA) which are precursors for the synthesis of
15
16 proinflammatory eicosanoids thereby causing a decrease in TNF- α .
17
18
19

20
21 The inflammatory process has a major role in the pathogenesis of breast cancer.
22
23 The previous study conducted by Kamel *et al.* (2012) has proven that chronic
24
25 inflammation is associated with breast cancer recurrence and increased levels of
26
27 inflammatory biological markers associated with reduced ability of breast cancer patients
28
29 to survive. Kamel *et al.* (2012) states that TNF- α is the most widely expressed
30
31 inflammatory cytokine in breast cancer. Likewise, according to Malik *et al.* (2018) TNF- α
32
33 has been shown to increase breast cancer cell growth. The number of cells expressing
34
35 TNF- α in mammary carcinoma has been known to be related to tumor grade and plays a
36
37 role in metastasis in mammary carcinoma. A previous study by Malik *et al.* (2018)
38
39 obtained results if benzopyrene can activate the inflammatory pathway which includes
40
41 TNF- α , cyclooxygenase-2 (COX-2) and vascular endothelial growth factor A (VEGF-A)
42
43 in both types of mammary cell lines. TNF- α is known to increase the activity of protein
44
45 kinase C (PKC- α) and protein tyrosin kinase to activate nuclear factor- κ B (NF κ B) as a
46
47 COX-2 promoter which is a marker of aggressiveness and poor prognosis in breast cancer.
48
49 However, in the type of breast cancer cell michigan cancer foundation-7 (MCF-7), TNF- α
50
51 regulates COX-2 through a path that is free of NF κ B. The administration of the high-fat
52
53 diet causes the replacement of the main fuel cell metabolism from glucose to ketones.
54
55
56
57
58
59
60

1
2
3
4 Normal cells can adapt to this change, but cancer cells which are highly dependent on
5 glucose as the main fuel, are unable to metabolize ketones (Aponte *et al.*, 2018). If glucose
6 levels decrease, the inflammatory response caused by cancer cells also decreases which
7 marked by low TNF- α levels as a biological marker of cancer cell malignancy.
8
9

10
11
12
13 The results of the present study showed that after the induction of BZP at a dose
14 of 0.3mg/20gBB/day for 14 days subcutaneously on the right breast of mice, found a
15 tumor in the positive control group and the high-fat diet group, whereas in the negative
16 control group induced with *oleum olivarum* there is no tumor. This is in line with the
17 results of studies conducted by Nansi *et al.* (2015) and Liangan *et al.* (2015) which
18 concluded that induction of BZP at a dose of 0.3mg/20gBB/day for 14 days
19 subcutaneously can induce breast cancer in mice. Malik *et al.* (2018) states that exposure
20 to BZP has several effects, which can cause metabolic activation of deoxyribonucleic acid
21 (DNA)-damaging species and elicit various biological responses such as activation of aryl
22 hydrocarbon receptors (AhR) which cause changes in gene expression. These various
23 biological responses then initiate carcinogenesis. BZP is metabolized by Cytochrome
24 P450 Family 1 Subfamily A Polypeptide 1 (CYP1A1), Cytochrome P450 Family 1
25 Subfamily B Member 1 (CYP1B1) and Epoxide Hydrolase (EH) to Benzo(a)pyrene 7,8-
26 dihydrodiol-9,10-epoxide BPDE) which is a reactive carcinogen. Exposure to BZP
27 increases the risk of breast cancer because of the expression of CYP1A1 and CYP1B1 in
28 breast tissue (Akram, 2012).
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 Based on the present study the mean delta area of mice tumors at K₃ was wider
51 than K₁, K₂, K₄, K₅ and K₆. The One way-ANOVA test results showed that there were
52 significant differences in the mean delta area of the tumor in mice ($p < 0.05$). These results
53 are different from the results of research conducted by Allen *et al.* (2014) which
54
55
56
57
58
59
60

1
2
3
4 concluded that the administration of the high-fat diet reduced tumor growth and improved
5
6 the survival of experimental animal models with glioma malignancy, colon cancer, gastric
7
8 cancer and prostate cancer. The previous study by Zhou *et al.* (2007) also concluded that
9
10 mice with astrocytoma and glioma treated with a high-fat diet and calorie restriction for
11
12 45 days had smaller astrocytoma and glioma tumor sizes compared to the mice on the
13
14 standard diet without calorie restriction. The results showed that giving a high-fat diet for
15
16 4 weeks had not had a significant effect on reducing tumor area compared to the negative
17
18 control group and the positive control group who received standard feed. This difference
19
20 in results is probably also due to the fact that after benzopyrene induction, there was a
21
22 significant difference between the intervention group and the control group.
23
24
25
26
27
28
29

30 CONCLUSION

31
32 Based on the results of this study we concluded a high-fat diet lowers TNF- α
33
34 levels and reduces the breast tumor area on benzopyrene induced mice (*Mus musculus*).
35
36 The lowest TNF- α levels and the lowest breast tumor area were found in groups with a
37
38 fat : protein ratio = 60:30.
39
40
41
42
43

44 REFERENCES

- 45
46
47 1. Akram, M. 2012. Effects of Benzo(a)pyrene on Human Breast Cancer Cell Lines.
48
49 Thesis. University of Eastern Finland.
50
51 2. Allen, B.G., Bhatia, S.K., Anderson, C.M., Eichenberger-Gilmore, J.M., Sibenaller,
52
53 Z.A., Mapuskar, K.A., Schoenfeld, J.D., Buatti, J.M., Spitz, D.R. and Fath, M.A.
54
55 2014. Ketogenic diets as an adjuvant cancer therapy: History and potential
56
57 mechanism. Redox Biol, 2: 963-970. <https://doi.org/10.1016/j.redox.2014.08.002>.
58
59
60

- 1
2
3
4 3. Anandakumar, P., Kamaraj, S., Ramakrishnan, G., Jagan, S. and Devaki, T. 2009.
5
6 Chemopreventive Task of Capsaicin against Benzo(a)pyrene-induced Lung Cancer
7
8 in Swiss Albino Mice. *Basic & Clinical Pharmacology & Toxicology*, 104: 360-365.
9
10 <https://doi.org/10.1111/j.1742-7843.2009.00387.x>.
11
12
- 13 4. Aponte, Y.R.T., Calderon, V. and Gonzales, M.J. 2018. The Kotegetic Diet and
14
15 Cancer. *Archives of Oncology and Cancer Therapy*, 1(1): 48-53.
16
17
- 18 5. Azar, S.T., Beydoun, H.M. and Albadri, M.R. 2016. Benefits of Ketogenic Diet for
19
20 Management of Type Two Diabetes: A Review. *Obesity & Eating Disorders*, 2(22):
21
22 1-4. <https://doi.org/10.4172/2471-8203.100022>.
23
24
- 25 6. Basic Health Research (Riskesdas). 2018. National Report on Basic Health Research.
26
27 Jakarta: Ministry of Health Indonesia. Available at:
28
29 <http://www.kesmas.kemkes.go.id>.
30
31
- 32 7. Dupuis, N., Curatolo, N., Benoist, J-F. and Auvin, S. 2015. Ketogenic diet exhibits
33
34 anti-inflammatory properties. *Epilepsia*, 56(7): e95-e98.
35
36 <https://doi.org/10.1111/epi.13038>.
37
38
- 39 8. Erickson, N., Boscheri, A., Linke, B. and Huebner, J. 2017. Systematic review:
40
41 isocaloric ketogenic dietary regimes for cancer patients. *Med Oncol*, 34(5): 72.
42
43 <https://doi.org/10.1007/s12032-017-0930-5>.
44
45
- 46 9. Fukao, T., Lopaschuk, G.D. and Mitchell, G.A. 2004. Pathways and control of ketone
47
48 body metabolism: on the fringe of lipid biochemistry. *Prostaglandins Leukot Essent*
49
50 *Fatty Acids*, 70(3): 243-251. <https://doi.org/10.1016/j.plefa.2003.11.001>.
51
52
53
54
55
56
57
58
59
60

10. Gao, M., Li, Y., Sun, Y., Shah, W., Yang, S., Wang, Y. and Long, J. 2011. Benzo[a]pyrene Exposure Increases Toxic Biomarkers and Morphological Disorders in Mouse Cervix. *Basic & Clinical Pharmacology & Toxicology*, 109: 398-406. <https://doi.org/10.1111/j.1742-7843.2011.00755.x>.
11. GLOBOCAN. 2019. Indonesia Summary Statistics 2018 available from <http://gco.iarc.fr/today/data/factsheets/populations/360-indonesia-fact-sheets.pdf>.
12. Irmawati, A., Jasmin, N. and Sidarningsih. 2018b. The effect of moderate exercise on the elevation of Bax/Bcl-2 ratio in oral squamous epithelial cells induced by benzopyrene. *Veterinary world*, 11(2): 177-180. <https://doi.org/10.14202/vetworld.2018.177-180>.
13. Kamel, M., Shouman, S., El-Merzebany, M., Kilic, G., Veenstra, T., Saeed, M., Wagih, M., Diaz-Arrastia, C., Patel, D. and Salama, S. 2012. Effect of Tumour Necrosis Factor-Alpha on Estrogen Metabolic Pathways in Breast Cancer Cells. *Journal of Cancer*, 3: 310-321. <https://doi.org/10.7150/jca.4584>.
14. Kephart, W.C., Mumford, P.W., Mao, X., Romero, M.A., Hyatt, H.W., Zhang, Y., Mobley, C.B., Quindry, J.C., Young, K.C., Beck, D.T., Martin, J.S., McCullough, D.J., D'Agostino, D.P., Lowery, R.P., Wilson, J.M., Kavazis, A.N. and Roberts, M.D. 2017. The 1-Week and 8-Month Effects of a Ketogenic Diet or Ketone Salt Supplementation on Multi-Organ Markers of Oxidative Stress and Mitochondrial Function in Rats. *Nutrients*, 9(1019): 1-22. <https://doi.org/10.3390/nu9091019>.
15. Klement, R.J. 2017. Beneficial effects of ketogenic diets for cancer patients: a realist review with focus on evidence and confirmation. *Medical Oncology*, 34(8):132. <https://doi.org/10.1007/s12032-017-0991-5>.

16. Lettieri-Barbato, D. and Aquilano, K. 2018. Pushing the Limits of Cancer Therapy: The Nutrient Game. *Frontiers in oncology*, 8: 148. <https://doi.org/10.3389/fonc.2018.00148>.
17. Liangan, R., Kairupan, C. and Durry, M. 2015. The effect of giving galangal (*alpinia galanga*) extract on the histology of mice (*mus musculus*) breast induced by benzo(a)pyrene. *Jurnal e-Biomedik (eBm)*, 3(1): 480–485. <https://doi.org/10.35790/ebm.3.1.2015.7492>.
18. Ma, Y., Ren, Y., Dai, Z-J., Wu, C-J., Ji, Y-H. and Xu. J. 2017. IL-6, IL-8 and TNF- α levels correlate with disease stage in breast cancer patients. *Adv Clin Exp Med*, 26(3): 421-426. <https://doi.org/10.17219/acem/62120>.
19. Malik, D-S., David, R.M. and Gooderham, N. 2018. Mechanistic evidence that benzo[a]pyrene promotes an inflammatory microenvironment that drives the metastatic of human mammary cells. *Archives of Toxicology*, 92: 3223-3239. <https://doi.org/10.1007/s00204-018-2291-z>.
20. Meidenbauer, J.J., Ta, N. and Seyfried, T.N. 2014. Influence of a ketogenic diet, fish-oil, and calorie restriction on plasma metabolites and lipids in C57BL/6J mice. *Nutr Metab (Lond)*, 11:23. <https://doi.org/10.1186/1743-7075-11-23>.
21. Nabbout, R., Vezzani, A., Dulac, O. and Chiron, C. 2011. Acute encephalopathy with inflammation-mediated status epilepticus. *Lancet Neurol*, 10(1): 99-108. [https://doi.org/10.1016/S1474-4422\(10\)70214-3](https://doi.org/10.1016/S1474-4422(10)70214-3).
22. Nandivada, P., Fell, G.L., Pan, A.H., Nose, V., Ling, P-R., Bistrrian, B.R. and Puder, M. 2016. Eucaloric Ketogenic Diet Reduces Hypoglycemia and Inflammation in Mice with Endotoxemia. *Lipids*, 51(6): 703-714. <https://doi.org/10.1007/s11745-016-4156-7>.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
23. Nansi, E.M., Durry, M.F. and Kairupan, C. 2015. Histopathological description of mice (*mus musculus*) breast induced by benzo(a)pyrene and given with yellow extract (*curcuma longa* l.). *Jurnal e-Biomedik (eBm)*, 3(1): 510-515. <https://doi.org/10.35790/emb.3.1.2015.7504>.
24. Page, K.A., Williamson, A., Yu, N., McNay, E.C., Dzuira, J., McCrimmon, R.J. and Sherwin, R.S. 2009. Medium-chain Fatty Acids Improve Cognitive Function in Intensively Treated Type 1 Diabetic Patients and Support in Vitro Synaptic Transmission During Acute Hypoglycemia. *Diabetes*, 58(5): 1237-1244. <https://doi.org/10.2337/db08-1557>.
25. Paoli, A. 2014. Ketogenic diet for obesity: friend or foe?. *International journal of environmental research and public health*, 11(2): 2092-2107. <https://doi.org/10.3390/ijerph110202092>.
26. Pinthong, W. and Suanarunsawat, T. 2020. Tea Seed Oil Alleviates Metabolic Derangement and Oxidative Stress in Rats Fed with High Fat and High Fructose Diet. *CMU J. Nat. Sci.* 19(4): 665-683. <https://doi.org/10.12982/CMUJNS.2020.0043>.
27. Roberts, M.N., Wallace, M.A., Tomilov, A.A., Zhou, Z., Marcotte, G.R., Tran, D., Perez, G., Gutierrez-Casado, E., Koike, S., Knotts, T.A., Imai, D.M., Griffey, S.M., Kim, K., Hagopian, K., McMackin, M.Z., Haj, F.G., Baar, K., Cortopassi, G.A., Ramsey, J.J. and Lopez-Dominguez, J.A. 2017. A Ketogenic Diet Extends Longevity and Healthspan in Adult Mice. *Cell Metabolism*, 26: 539-546. <http://dx.doi.org/10.1016/j.cmet.2017.08.005>.
28. Rose, D.P. and Vano-Davis, L. 2012. The cellular and molecular mechanisms by which insulin influences breast cancer risk and progression. *Endocr Relat Cancer*, 19(6): R225-R241. <https://doi.org/10.1530/ERC-12-0203>.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
29. Santos, E.W., Oliveira, D.C., Hastreiter, A., Silva, G.B., Beltran, J.S.O., Rogero, M.M., Fock, R.A. and Borelli, P. 2019. Short-term high-fat diet affects macrophages inflammatory response, early signs of a long-term problem. *Brazilian Journal of Pharmaceutical Sciences*, 55: e17561. <http://dx.doi.org/10.1590/s2175-97902019000117561>.
30. Saslow, L.R., Mason, A.E., Kim, S., Goldman, V., Ploutz-Snyder, R., Bayandorian, H., Daubenmier, J., Hecht, F.M. and Moskowitz, J.T. 2017. An Online Intervention Comparing a Very Low-Carbohydrate Ketogenic Diet and Lifestyle Recommendations Versus a Plate Method Diet in Overweight Individuals With Type 2 Diabetes: A Randomized Controlled Trial. *Journal of medical Internet research*, 19(2): e36. <https://doi.org/10.2196/jmir.5806>.
31. Soldati, L., Di Renzo, L., Jirillo, E., Ascierio, P.A., Marincola, F.M. and De Lorenzo, A. 2018. The influence of diet on anti-cancer immune responsiveness. *Journal of Translational Medicine*, 16(75): 1-18. <https://doi.org/10.1186/s12967-018-1448-0>.
32. Sussman, D., van Eede, M., Wong, M.D., Adamson, X.L. and Henkelman, M. 2013. Effects of a ketogenic diet during pregnancy on embryonic growth in the mouse. *BMC Pregnancy and Childbirth*, 13:109. <http://www.biomedcentral.com/1471-2393/13/109>.
33. Taylor, M.K., Swerdlow, R.H., Burns, J.M. and Sullivan, D.K. 2019. An Experimental Ketogenic Diet for Alzheimer Disease Was Nutritionally Dense and Rich in Vegetables and Avocado. *Current developments in nutrition*, 3(4): nzz003. <https://doi.org/10.1093/cdn/nzz003>.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
34. van Berkel, A.A., Ijff, D.M. and Verkuyl, J.M. 2018. Cognitive benefits of the ketogenic diet in patients with epilepsy: A systematic overview. *Epilepsy Behav*, 87: 69-77. <https://doi.org/10.1016/j.yebeh.2018.06.004>.
35. Vizuete, A.F., de Souza, D.F., Guerra, M.C., Batassini, C., Dutra, M.F., Bernardi, C., Costa, A.P. and Gonçalves, C-A. 2013. Brain changes in BDNF and S100B induced by ketogenic diets in Wistar rats. *Life Sciences*, 92(16-17): 923–928. <http://dx.doi.org/10.1016/j.lfs.2013.03.004>.
36. Weber, D.D., Aminazdeh-Gohari, S. and Kofler, B. 2018. Ketogenic diet in cancer therapy. *Aging*, 10(2): 164-165. <https://doi.org/10.18632/aging.101382>.
37. Weber, D.D., Aminzadeh-Gohari, S., Tulipan, J., Catalano, L., Feichtinger, R.G. and Kofler, B. 2020. Ketogenic diet in the treatment of cancer e Where do we stand?. *Molecular Metabolism*, 33: 102-121. <https://doi.org/10.1016/j.molmet.2019.06.026>.
38. Williams, T.J. and Cervenka, M.C. 2017. The role for ketogenic diets in epilepsy and status epilepticus in adults. *Clinical neurophysiology practice*, 2: 154-160. <https://doi.org/10.1016/j.cnp.2017.06.001>.
39. Wisnu, G.N.P.P., Khairun N.B. and Riyan W. 2017. Ketogenic diet: a promising non-pharmacological therapy for epilepsy. *Jurnal Kedokteran Universitas Negeri Lampung*, 7(1): 118-122.
40. Wolczyk, D., Zaremba-Czogalla, M., Hryniewicz-Jankowska, A., Tabola, R., Grabowski, K., Sikorski, A. F. and Augoff, K. 2016. TNF- α promotes breast cancer cell migration and enhances the concentration of membrane-associated proteases in lipid rafts. *Cellular oncology (Dordrecht)*, 39(4): 353-363. <https://doi.org/10.1007/s13402-016-0280-x>.

- 1
2
3
4 41. Wunjuntuk, K., Chareonkiatkul, S. and Kettawan, A. 2020. Parboiled Germinated
5 Brown Rice Decreases Inflammatory Cytokines Levels in Various Regions of Brain
6
7
8
9 in Rats Induced by Carbon Tetrachloride. *CMU J. Nat. Sci.* 19(2): 265-279.
10
11 <https://doi.org/10.12982/CMUJNS.2020.0018>.
12
- 13 42. Zhou, S., Kachhap, S., Sun, W., Wu, G., Chuang, A., Poeta, L., Grumbine, L.,
14
15 Mithani, S.K., Chatterjee, A., Koch, W., Westra W.H., Maitra, A., Glazer, C.,
16
17 Carducci, M., Sidransky, D., McFate, T., Verma, A. and Califano, J.A. 2007.
18
19 Frequency and phenotypic implications of mitochondrial DNA mutations in human
20
21 squamous cell cancers of the head and neck. *Proceedings of the National Academy*
22
23 *of Science of the United States of America*, 104(18): 7540-7545.
24
25
26
27 <https://doi.org/10.1073/pnas.0610818104>.
28
- 29 43. Zhou, J., Liu, B., Liang, C., Li, Y. and Song, Y-H. 2016. Cytokine Signaling in
30
31 Skeletal Muscle Wasting. *Trends in Endocrinology & Metabolism*, 27(5): 335-347.
32
33
34 <https://doi.org/10.1016/j.tem.2016.03.002>.
35
- 36 44. Zick, S.M., Snyder, D. and Abrams, D.I. 2018. Pros and Cons of Dietary Strategies
37
38 Popular Among Cancer Patients. *Oncology (Williston Park)*, 32(11): 542-547.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60