

Periodontitis Affects Skeletal Muscle Metabolism Through an Increase in Proinflammatory Cytokines

by Risma Aprinda Kristanti

Submission date: 23-Mar-2022 11:55AM (UTC+0800)

Submission ID: 1790677532

File name: Metabolism_Through_an_Increase_in_Proinflammatory_Cytokines.pdf (181.12K)

Word count: 4993

Character count: 27721

Periodontitis Affects Skeletal Muscle Metabolism Through an Increase in Proinflammatory Cytokines

Risma Aprinda Kristanti¹, Taufan Bramantoro², Pratiwi Soesilawati^{3*},
Erni Maduratna Setiawati⁴, Bambang Purwanto⁵

1. Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.
2. Departement of Dental Public Health, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.
3. Departement of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.
4. Departement of Periodontology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya Indonesia.
5. Department of Physiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

Abstract

Periodontitis is an inflammatory disease of the oral cavity which attacks the gingival and periodontal tissue. One of the results of immune response activity in periodontitis is an increase in proinflammatory cytokines both locally and systematically. An increase in proinflammatory cytokines will affect tissue metabolism, including skeletal muscle tissue. The mechanism of periodontitis in affecting skeletal muscle is still being discussed and studied, this review will discuss the mechanism of periodontitis in affecting skeletal muscle metabolism. The interaction between bacteria and the immune system in periodontal tissues will activate the immune system and can cause an increase in proinflammatory cytokines such as IL-6, IL-1 β , and TNF- α both locally and systemically. The presence of bacteria in the body's circulation and an enhancement of proinflammatory cytokines systemically can affect other tissues in the body, one of which is skeletal muscle. Several researchers have proven that periodontitis affects the regeneration process, glucose uptake, and recovery in skeletal muscle. An increase in IL-6, IL-1 β , and TNF- α caused by periodontitis can reduce the process of muscle regeneration, reduce glucose uptake in skeletal muscle, and decreases the level of muscle responsiveness to training loads.

Review (J Int Dent Med Res 2021; 14(4): 1623-1628)

Keywords: Cytokine, periodontitis, skeletal muscle, immunology.

Received date: 01 July 2021

Accept date: 26 October 2021

Introduction

Periodontitis is an inflammatory disease characterized by inflammation of the soft tissues and the periodontal ligament and also alveolar bone damage.¹ Periodontal disease is mainly caused by infection from gram-negative bacteria such as *Porphyromonas gingivalis* (*P.gingivalis*).² Inflammation will triggered by the interaction between these bacteria and the host's immune systems, causing damage to collagen tissue as one of the tooth supporting tissue structures.^{3,4} The process of recognizing bacterial components by the host immune system in the periodontal tissue will cause various immune cells to be

activated in the inflamed area.⁵ This causes an increase in the production of inflammatory mediators, such as chemokines, ROS, and proinflammatory cytokines where they can be the sign of disease progression level.^{6,7,8,9} The activation of inflammatory mediators in periodontitis has a significant effect on the host response to systemic inflammation.⁸ There are several cytokines such as IL-6, IL-1, IL-8, and TNF- α , prostaglandins, and MMP produced from periodontitis.^{10,11} Bacteria and cytokines in periodontal tissues can enter the body's systemic circulation and cause a response from several tissues.^{12,13,14} A study conducted in a systematic review has proven that periodontitis can affect physical fitness through its effects on muscles.¹⁵ The increased level of cytokines can cause muscle wasting.¹⁶ Proinflammatory cytokines such as IL-6, IL-1 β , and TNF- α can trigger muscle damage.^{17,18}

The anabolism and catabolism processes

*Corresponding author:

Pratiwi Soesilawati,
Universitas Airlangga, JL. Mayjend. Prof. Dr. Moestopo No.47,
Surabaya, Indonesia.
E-mail: pratiwi.soesilawati@gmail.com

are such important factors that they can maintain muscle mass. Anabolic factors consist of physical and chemical factors, while catabolic factors consist of muscle that is rarely used, oxidative stress, proinflammatory cytokines, acidosis, and glucocorticoid hormones.¹⁹ Structural changes such as a decrease in muscle mass and fiber can cause muscle weakness, reducing the muscles strength and endurance.^{20,21} Immunological factors such as cytokines and myeloid cells also influence muscle metabolism. An increase in proinflammatory cytokines can cause an increase of lysis in muscle fibers and affect the process of differentiation of muscle cells.²²

Until now, the mechanism of periodontitis in influencing muscle metabolism has not been thoroughly explained. The purpose of this article is to review the effect of periodontitis on the regeneration process, glucose uptake, and muscle recovery process seen from the aspect of increasing proinflammatory cytokines.

Review

In periodontitis, there is an increased level of IL-6 and TNF- α cytokines. The plasma levels of IL-6 and TNF- α are higher in people with periodontitis than people without it.^{23,24} The increased levels of IL-6 experienced by patients with periodontitis occur in both plasma and saliva.²⁵ A study on experimented animals showed that rats induced by periodontitis using *P. gingivalis* bacteria experienced an enhancement in serum concentration of the proinflammatory cytokine IL-6 on day 15 which indicated a association between serum IL-6 levels and disease activity in periodontitis.²⁶ In periodontitis, there is also an increase in IL-1 β , which can be detected through the gingival cervical fluid (GCF) and saliva, periodontitis patients show higher IL-1 β concentrations than people without it.^{27,28} A study on an experimented rat in periodontitis model induced by *P. gingivalis* showed an increase in serum IL-1 β .²⁶ This suggests that periodontitis can cause a systemic increase in IL-1 β . In addition to IL-6 and IL-1 β , there is also an increase in TNF- α concentration in patients with periodontitis. The study showed the results of an enhancement in TNF- α levels in the saliva of periodontitis patients.²⁹ An increase in TNF- α concentration also occurs in the serum of patients with periodontitis.³⁰

There was a relationship between

P. gingivalis-induced on experimented animals and sarcopenia.³¹ The animals induced by *P. gingivalis* using doses adjusted to systemic *P. gingivalis* levels in periodontitis patients resulted in a decrease in skeletal muscle mass, especially in fast-type skeletal muscle.³¹ A study by Visser *et al.* proved that there was a reduce in muscle mass and strength as IL-6 and TNF- α concentration increased.³² Meanwhile, another study conducted by Sullivan indicates that IL-1 β plays a role in suppressing myoblast fusion to become muscle fibers, resulting in a decrease in muscle regeneration.³³

P. gingivalis bacterial infection in periodontal tissue can reduce skeletal muscle glucose uptake.³⁴ There is a possibility that the decrease in skeletal muscle glucose uptake due to periodontitis is caused by an increase in the IL-6 cytokine in response to an increase in the amount of bacteria. A study in experimental animals has proven that chronic exposure to IL-6 can cause insulin resistance by decreasing the translocation process from glucose transporter type 4 (GLUT4) to muscle cell membranes. The GLUT4 is needed when taking glucose from the blood circulation which then will be used as a source of energy formation for the muscles to contract and relax.^{35,36}

The study conducted by De Souza *et al.* proved that periodontitis can affect the recovery process in rats' skeletal muscles which is given a treadmill exercise. Periodontal disease is able to change the leukocyte count and systemic interleukin levels so that it will contribute to the training load and have an indirect effect on muscle catabolism.³⁷

Systemic increase in chronic proinflammatory cytokines can cause skeletal muscle atrophy.³⁸ Periodontitis induces an increase in TNF- α and IL-6 but in rats which were given treadmill exercise and induced periodontitis did not change the concentration of these cytokines.³⁷ This suggests that periodontitis may indirectly impair muscle responsiveness to training loads due to the presence of local TNF- α . The presence of an increase in local TNF- α triggers the catabolism process in muscle fibers in response to training loads, and this effect is very visible in type 2 muscle fibers because the gastrocnemius muscle shows a decrease in fiber hypertrophy in rats given training loads and induced periodontitis.³⁷

Discussion

An Increase in Proinflammatory Cytokines in Periodontitis

In periodontitis, the process of recognizing the bacterial cell wall is carried out by Toll-like Receptors (TLRs).³⁹ Toll-like Receptor 4 is the main receptor for the LPS signaling process.⁴⁰ Lipopolysaccharide (LPS) which binds to TLR4 in gingival fibroblasts activates several second messengers and NF- κ B which play a role in the expression of inflammatory cytokines including IL-6.^{40,41} Meanwhile, the increase in IL-1 β production is triggered by the interaction between the host and microbiota and is widely induced by the expansion and activation of Th1 and Th2 cells.⁴² LPS from bacteria will induce monocytes and macrophages to produce TNF- α and IL-1 β .⁴³ Oral polymorphonuclear leukocytes (PMN) have mRNA for both TNF- α and IL-1 β which are able to produce both types of cytokines in sufficient quantities without stimulation.^{44,45}

Periodontitis - Skeletal muscle regeneration process

Interleukin-6 and IL-1 β can affect the process of muscle regeneration through their action on IGF-1. In sufficiently low concentrations, IL-1 β can suppress protein synthesis, myogenin expression and myoblast differentiation significantly. This will inhibit the work of IGF-1 in muscle to increase protein synthesis.⁴⁶ A study conducted by Li (2009) using rats proved that exposure to IL-1 β for 2 days can reduce myotube width by 13%. In addition, there was a decrease in actin synthesis by 13% to 14%.⁴⁷ Interleukin 1 β is able to inhibit Growth Hormone (GH) responses to the Serine protease inhibitor 2.1 (Spi 2.1). At 10 ng/ml, IL-1 β has been shown to inhibit GH from expressing IGF-1 and Spi 2.1 mRNA in hepatocyte cultures.^{48,49} A study by Bergad *et al.* even showed that at lower concentrations, ie 0.25 or 0.5 ng/ml, IL-1 β could inhibit the expression of Spi 2.1 mRNA and its activity.⁵⁰ Growth hormone stimulation of IGF-1 and Spi 2.1 mRNA was inhibited by IL-1 by suppressing the JAK2/STAT5 pathway.⁵¹ There was a decrease in the activation ability of STAT 5 which was phosphorylated and bound to DNA.⁵² The target of IL-1 inhibition on STAT5 is histone acetyl transferase which plays a role in chromatin remodeling and it is a coactivator for the regulation of transcriptional activity of STATs.⁵³

Interleukin-6 can cause a decrease in circulating IGF-1 while the production of IGF-1 in the liver and the number of acid-labeled subunits in the serum do not change, but the circulating levels of IGFBP-3 decrease along with the increase in proteolysis.⁵⁴ The binding between IGF-1 and its receptor will trigger phosphoinositide-3-kinase (P13-K) to produce phosphatidylinositol (4,5)-biphosphate (PIP2) which in turn stimulates the production of phosphatidylinositol 3,4,5-triphosphate (PIP3).^{55,56,57} Phosphatidylinositol 3,4,5-triphosphate will bind to phosphoinositide-dependent kinase-1 (PDK1) which will then bind to the pleckstrin homology (PH) domain of Akt.^{55,57} Akt activates mammalian target of rapamycin (mTOR).⁵⁷ The phosphorylation process of mTORC1 in mTOR²⁴⁴⁸ and mTOR²⁴⁸¹ will cause muscle protein synthesis.^{58,59} Therefore, a decrease in IGF-1 in the systemic circulation caused by an increase in IL-6 will inhibit muscle protein synthesis.

Several studies have shown that TNF- α triggers the breakdown of myosin heavy chain protein (MHCP).^{60,61,62} Exposure to TNF- α for a long time can cause inhibition of cell differentiation and muscle regeneration.⁶³ Studies in muscle cells owned by humans, pigs and rats have shown that exposure to TNF- α and IL-1 β prevents IGF-1 from stimulating protein synthesis.^{64,65}

Periodontitis - Skeletal muscle glucose uptake

Chronic exposure to IL-6 attenuates insulin stimulation of glucose uptake and GLUT4 translocation at C2C12 and neonatal myotubes. Insulin activity in stimulating IRS-1 and AKT phosphorylation was inhibited by IL-6. The absence of AMPK and AS160 phosphorylation further highlights the negative effects of insulin signaling due to chronic IL-6 exposure.^{66,67}

Insulin can increase glucose transport to peripheral tissues including muscle by increasing translocation of GLUT4 from intracellular to plasma membrane where this process is mediated by activation of PI3K, protein kinase B (AKT), and several isoforms of protein kinase C.⁶⁸ Skeletal muscle has a mechanism to increase glucose transport by activating AMPK which can be triggered in conditions of hypoxia, ischemia, or exercise, and it is an insulin independent mechanism.⁶⁹ AKT substrate 160 (AS160) is a 160 kDa protein which is an effector

for glucose transport and triggers GLUT4 translocation.⁷⁰ Interleukin-6 increases glucose uptake by activating the serine/threonine protein kinase 11 (LKB1)/AMP-activated protein kinase/protein kinase B pathway substrate of 160 kDa (AS160). The opposite effect occurred when miotube and rat experimental animals were exposed to IL-6 for a long (chronic) period where IL-6 decreased GLUT4 translocation to the plasma membrane, resulting in a decrease in insulin signaling on insulin receptor substrate (IRS-1). The three mechanisms involved include: activation of c-jun NH2-terminal kinase (JAK1/2), accumulation of suppressor of cytokine signaling 3 (SOCS3) mRNA, and increased activity of PTP1B.³⁵

Periodontitis - Skeletal muscle recovery

When inflammation occurs in which TNF- α increases systemically, the skeletal muscle sarcolemma can function as receptors for TNF- α .^{71,72} TNF- α can activate NF- κ B which is a transcription factor and activation of NF- κ B causing the degradation of skeletal muscle protein.^{73,74} Nuclear Factor Kappa Beta is in the form of inactive homodimer or heterodimer complexes in the cell cytoplasm, when activated by cytokines, NF- κ B dimers will translocate to the nucleus and increase the expression of several genes encoding proteins, including proteins that trigger muscle atrophy.⁷⁵

The effect of muscle damage by TNF can occur through the activation of Janus Kinase. The activated Janus Kinase can inhibit the expression of IGF-1 mRNA, during inflammation and increase in cytokine IGF-1 signaling mechanisms and the interaction between TNF and IGF-1 can cause muscular dystrophy.⁶⁴

TNF- α which binds to its receptor will activate JAK which then, as a signaling molecule, will inhibit the activation of Insulin Receptor Substrate 1 (IRS-1). Inhibited IRS-1 will not be able to activate Akt, causing inhibition of the muscle protein synthesis process.⁷⁶

Conclusions

An increase in cytokines IL-6, IL-1 β , and TNF- α in periodontitis can reduce the process of muscle regeneration by affecting the activity of IGF-1. IL-6 can decrease IGF-1 levels in the systemic circulation and IL-1 β can reduce the ability of hepatocyte cells to express IGF-1. IL-1 β

and TNF- α are able to prevent IGF-1 from stimulating muscle protein synthesis. In addition, IL-6 can also reduce glucose uptake in skeletal muscle where the glucose is needed as a source of contraction for muscles. Periodontitis also indirectly affects the process of muscle recovery if given a training load, in which the effect is thought to be through TNF- α which reduces the level of muscle responsiveness to training loads.

Acknowledgements

All authors have the same contribution to this manuscript.

Declaration of Interest

The authors report no conflict of interest.

References

1. Hajshengallis G. Periodontitis: from microbial immune subversion to systemic inflammation. *Nature Reviews Immunology* 2015; 15(1): 30-44.
2. Xu W, Zhou W, Wang H, Liang S. Roles of Porphyromonas gingivalis and its virulence factors in periodontitis. *Advances in Protein Chemistry and Structural Biology* 2020; 120: 46.
3. Pozo P, Valenzuela MA, Melej C, et al. Longitudinal analysis of metalloproteinases, tissue inhibitors of metalloproteinases and clinical parameters in gingival crevicular fluid from periodontitis-affected patients. *J Periodontol Res* 2005; 40: 199-207.
4. De Carlo AA Jr., Windsor LJ, Bodden MK, et al. Activation and novel processing of matrix metalloproteinases by a thiol-proteinase from the oral anaerobe Porphyromonas gingivalis. *J Dent Res* 1997; 76: 1260-70.
5. Craig RG, Pemat AM, Pemat, Pecoits-Filho R, Levin NW, and Kotanko P. Periodontal Diseases and Systemic Inflammation. *Seminars in Dialysis* 2013; 26 (1): 16-39.
6. Gariet GP, Aranha AMF, Silveira EM, Vieira AE, Queiroz-Junior CM, Madeira MFM, Fukada SY, and Silva TA. The Role of Chemokines and Cytokines in The Pathogenesis of Periodontal And Periapical Lesions: Current Concepts. In *Cell and Molecular Biology, Immunology and Clinical Basis*, ed. In Tech. 2012: 219-64.
7. Khalaf H., Lonn J, and Bengtsson T. Cytokines and Chemokines Are Differentially Expressed in Patients with Periodontitis: Possible Role For TGF-B1 As A Marker for Disease Progression. *Cytokine* 2014; 67(1): 29-35.
8. Leishman SJ, Seymour GJ, and Ford PJ. Local and Systemic Inflammatory Responses to Experimentally Induced Gingivitis. *Disease Markers* 2013; 35(5): 543-49.
9. Ramseier CA, Kinney JS, Herr AE, Braun T, Sugai JV, Shelburne CA, Rayburn LA, Tran HM, Singh AK, and Giannobile WV. Identification of Pathogen and Host-Response Markers Correlated with Periodontal Disease. *Journal of Periodontology* 2009; 80 (3): 436-46.
10. Fentoglu O, Koroglu BK, Hıçyılmaz H, et al. Pro-inflammatory cytokine levels in association between periodontal disease and hyperlipidaemia. *J Clin Periodontol* 2011; 38: 8-16.
11. Vernal R, Dezerega A, Dutzan N, et al. RANKL in human periapical granuloma: possible involvement in periapical bone destruction. *Oral Dis* 2006; 12: 283-289.
12. Rahmadi A, Yulianti A, Soesilawati P. The Role of Macrophags In Pregnant Rats with Chronic Periodontitis as A Risk of Preeclampsia. *Biochem Cell Arch* 2020; 20(2): 4865-69.

13. Ardalan MR, Ghabili K, Pourabbas R, Shoja MM. A causative link between periodontal disease and glomerulonephritis: a preliminary study. *Therapeutics and Clinical Risk Management* 2011; (7): 93-8.
14. Han P, Dianxing Sun D, Yang J. Interaction between periodontitis and liver diseases (Review). *Biomedical Reports* 2016; 5: 267-76.
15. Bramantoro T, Hariyani N, Setyowati D, Purwanto B, Zulfiana AA, Irmalia WR. The impact of oral health on physical fitness: A systematic review. *Heliyon* 2020;6(4).
16. Matsukawa S, Kai S, Seo H, Suzuki K, Fukuda K. Activation of the β -adrenergic receptor exacerbates lipopolysaccharide-induced wasting of skeletal muscle cells by increasing interleukin-6 production. *Plos One* 2021: 1-19.
17. Corbin CB. A multidimensional hierarchical model of physical fitness: a basis for integration and collaboration. *Quest* 1991; 43(3): 296-306.
18. Cannon JG, Fielding RA, Fiatarone MA, Orencole SF, Dinarello CA, Evans WJ. Increased interleukin 1 beta in human skeletal muscle after exercise. *Am J Physiol Integr Comp Physiol* 1989; 257(2): 451-55.
19. Tidball JG and Villalta SA. Regulatory Interactions Between Muscle and The Immune System During Muscle Regeneration. *Am J Physiol Regul Integr Comp Physiol* 2010;298(5): 1173-87.
20. Krogh-Madsen, R., Plomgaard, P., Moller, K., Mittendorfer, B., & Pedersen, B.K. Influence of TNF α and IL-6 infusions on insulin sensitivity and expression of IL-18 in humans. *Am J Physiol Endocrinol Metab* 2006; 291(1): 108-14.
21. Munoz-Canoves P, Scheele C, Pedersen BK, Serrano AL. Interleukin-6 Myokine Signaling in Skeletal Muscle: A Double-Edged Sword? *FEBS J* 2013; 280(17): 4131-48.
22. Villalta SA, Rinaldi C, Deng B, Liu G, Fedor B, Tidball JG. Interleukin-10 reduces the pathology of mdx muscular dystrophy by deactivating M1 macrophages and modulating macrophage phenotype. *Hum Mol Genet* 2011; 20(4): 790-805.
23. Noh MK, Jung M, Kim SH, Lee SR, Park KH, Kim DH, Kim HH, Park YG. Assessment of IL-6, IL-8 and TNF- α Levels in The Gingival Tissue of Patients with Periodontitis. *Experimental and Therapeutic Medicine* 2013;6.
24. Medara N, Lenzo JC, Walsh KA, Darby IB, Simpson NMO, Reynolds AC. T helper 17 cell-related cytokines in serum and saliva during management of periodontitis. *Cytokine* 2020; 134: 1-12.
25. Costa PP, Trevisan GL, Macedo GO, Palioto DB, Souza SLS, Grisi MFM, Novaes AB, Mario Taba M. Salivary Interleukin-6, Matrix Metalloproteinase-8, and Osteoprotegerin in Patients with Periodontitis and Diabetes. *J Periodontol* 2010; 85(3).
26. Saadi-Thiers K, Huck O, Simonis P, Tilly P, Fabre JE, Tenenbaum H, Davideau JL. Periodontal and Systemic Responses in Various Mice Models of Experimental Periodontitis: Respective Roles of Inflammation Duration and Porphyromonas gingivalis Infection. *J Periodontol* 2013; 84(3).
27. Rangbulla V, Nirola A, Gupta M, Batra P. Salivary IgA, Interleukin-1beta and MMP-8 As Salivary Biomarkers in Chronic Periodontitis Patients. *Chin J Dent Res* 2017; 20: 43-51.
28. Kinney JS et al. Crevicular Fluid Biomarkers and Periodontal Disease Progression. *J Clin Periodontol* 2014; 41: 113-20.
29. Singh P, Gupta ND, Bey A, and Khan S. Salivary TNF-alpha: A Potential Marker Of Periodontal Destruction. *Indian Society Periodontology* 2014; 18(3).
30. Gokul K. Estimation of the level of tumor necrosis factor- α in gingival crevicular fluid and serum in periodontal health and disease: A biochemical study. *Indian Journal of Dental Research* 2012; 23(3).
31. Kawamura N, Ohnuki Y, Matsuo I, Suita K, Ishikawa M, Mototani Y, Shiozawa K, Ito A, Yagisawa Y, Hayakawa Y, Nariyama M, Umeki D, Ujije Y, Gomi K, Okumura S. Effects of chronic Porphyromonas gingivalis lipopolysaccharide infusion on skeletal muscles in mice. *The Journal of Physiological Sciences* 2019; 69: 503-11.
32. Visser M, Pahor M, Taaffe DR, Goodpaster BH, Simonsick EM, Newman AB, Nevitt M, Harris TB. Relationship of Interleukin-6 and Tumor Necrosis Factor- α With Muscle Mass and Muscle Strength in Elderly Men and Women: The Health ABC Study. *Journal of Gerontology* 2002; 57A(5).
33. Sullivan BE. Interleukin 1-beta Decreases Myoblast Fusion In Vitro. Honors Scholar Theses. University of Connecticut. 2015
34. Watanabe K, Katagiri S, Takahashi H, Sasaki N, Maekawa S, Komazaki R, Hatasa M, Kitajima Y, Maruyama Y, Shiba T, Komatsu K, Ohsugi Y, Tanaka K, Matsuzawa A, Hirota T, Tohara H. Porphyromonas gingivalis impairs glucose uptake in skeletal muscle associated with altering gut microbiota. *The FASEB Journal* 2020; 00: 1-16.
35. Nieto-Vazquez I, Fernandez-Veledo S, de Alvaro C, and Lorenzo M. Dual Role of Interleukin-6 in Regulating Insulin Sensitivity in Murine Skeletal Muscle. *Diabetes* 2008; 57.
36. Sherwood L. *Human Physiology: From Cell to Systems*. Jakarta: EGC. 2016.
37. De Souza BC, Ribas ME, Lopes AL, Teixeira BC, Lames ML. Periodontal Disease Influences The Recovery Processes In The Skeletal Muscles In Trained Mice. *Journal of Physical Education and Sport* 2017; 17(2): 572 - 81.
38. Li YP and Reid MB. NF-kappaB Mediates The Protein Loss Induced by TNF-alpha In Differentiated Skeletal Muscle Myotubes. *Am J Physiol Regul Integr Comp Physiol* 2000; 279(4): 1165-70.
39. Hoshino, K., Takeuchi, O., Kawai, T., Sanjo, H., Ogawa, T., Takeda, Y., Takeda, K., & Akira, S. Pillars Article: Cutting Edge: Toll-Like Receptor 4 (TLR4)-Deficient Mice Are Hyporesponsive to Lipopolysaccharide: Evidence for TLR4 as the Lps Gene Product. *J Immunol* 1999 2016; 162: 3749-52.
40. Wang PL, Azuma Y, Shinohara M, Ohura K. Toll-like receptor 4-mediated signal pathway induced by Porphyromonas gingivalis lipopolysaccharide in human gingival fibroblasts. *Biochemical and Biophysical Research Communications* 2000; 273(3): 1161-67.
41. Tomchuck SL, Zvezdaryk KJ, Coffelt SB, Waterman RS., Danka ES, Scandurro AB. Toll-Like Receptors on Human Mesenchymal Stem Cells Drive Their Migration and Immunomodulating Responses. *Stem Cells* 2008; 26(1): 99-107.
42. Ben-Sasson SZ et al. IL-1 Acts Directly on CD4 T Cells to Enhance Their Anti Gen Driven Expansion and Differentiation. *Proc Natl Acad Sci* 2009; 7119-24.
43. Nishimura F, Iwamoto Y, Mineshima J, Shimizu A, Soga Y, Murayama Y. Periodontal disease and diabetes mellitus: the role of tumor necrosis factor alpha in a "two-way relationship". *Journal of Periodontology* 2003; 74: 97-102.
44. Hendley TM, Steed RB, Galbraith GMP. Interleukin-1 β geneexpression in human oral polymorphonuclearleukocytes. *J Periodontology* 1995; 66: 761-65.
45. Galbraith GMP, Hagan C, Steed RB, Sanders JJ, Javed T. Cytokine production by oral and peripheral blood neutrophils in adult Periodontitis. *J Periodontology* 1997; 68: 832-38.
46. Spate U and Schulze PC. Pro-inflammatory Cytokines and Skeletal Muscle. *Curr Opin Clin Nutr Metab Care* 2004; 7: 265-69.
47. Li W, Moylan JS, Chambers MA, Smith J, Reid MB. Interleukin-1 Stimulates Catabolism in C2C12 Myotubes. *Am J Physiol Cell Physiol* 2009; 297(3): 706-14.
48. Simar-Blanchet AE, Legraverend C, Thissen JP, Cam AL. Transcription of the Rat Serine Protease Inhibitor 2.1 Gene *in Vivo*: Correlation with GAGA Box Promoter Occupancy and Mechanism of Cytokine-Mediated Down-Regulation. *Molecular Endocrinology* 1998; 12(3): 391-404.
49. Thissen JP and Josiane Verniers J. Inhibition by Interleukin-1 β and Tumor Necrosis Factor- α of the Insulin-Like Growth Factor I Messenger Ribonucleic Acid Response to Growth Hormone in Rat Hepatocyte Primary Culture. *Endocrinology* 1997; 138(3): 1078-84.

50. Bergad EL, Schwarzenberg SJ, Humbert JT, Morrison M, Amarasinghe S, Towle HC, Berry SA. Inhibition of Growth Hormone Action in Models of Inflammation. *Am J Physiol Cell Physiol* 2000; 279: 1906-17.
51. Shumate ML, Yumet G, Ahmed TA, Cooney RN. Interleukin-1 Inhibits The Induction Of Insulin-Like Growth Factor-1 By Growth Hormone In CWSV-1 Hepatocytes. *Am J Physiol Gastrointest Liver Physiol* 2005; 289: 227-39.
52. Groner B, Fritsche M, Stocklin E, Berchtold S, Merkle C, Moriggi R, and Pfizner E. Regulation of the trans-activation potential of STAT5 through its DNA-binding activity and interactions with heterologous transcription factors. *Growth Hormone IGF Res* 2000; 15-20.
53. Rasche A and Lees E. Chromatin acetylation and remodeling at the cis promoter during STAT5-induced transcription. *Nucleic Acids Res* 2003; 31: 6882-90.
54. De Benedetti F, Meazza C, Oliveri M, Pignatti P, Vivarelli M, Alonzi T, Fattori E, Garrone S, Barreca A & Martini A. Effect of IL-6 on IGF binding protein-3: a study in IL-6 transgenic mice and in patients with systemic juvenile idiopathic arthritis. *Endocrinology* 2001; 142: 4818-26.
55. Stitt TN, Drujan D, Clarke BA, Panaro F, Timofeyeva Y, Kline WO, et al. The IGF-1/PI3K/akt pathway prevents expression of muscle atrophy-induced ubiquitin ligases by inhibiting FOXO transcription factors. *Mol Cell* 2004; (14): 395-403.
56. O'Neill BT, Lauritzen HP, Hirshman MF, Smyth G, Goodyear LJ, Kahn CR. Differential role of insulin/IGF-1 receptor signaling in muscle growth and glucose homeostasis. *Cell Rep* 2015; 11: 1220-35.
57. Schiaffino S and Mammucari C. Regulation of skeletal muscle growth by the IGF1-akt/PKB pathway: Insights from genetic models. *Skelet Muscle* 2011;1(1):4.
58. Deldicque L, Theisen D, Francaux M. Regulation of mTOR by amino acids and resistance exercise in skeletal muscle. *Eur J Appl Physiol* 2005; 94:1-10.
59. Bolster DR, Kubica N, Crozier SJ, Williamson DL, Farrell PA, Kimball SR, et al. Immediate response of mammalian target of rapamycin (mTOR)-mediated signalling following acute resistance exercise in rat skeletal muscle. *J Physiol* 2003; 553: 213-20.
60. Teles RP, Likhari V, Socransky SS, Haffajee AD. Salivary Cytokine Levels in Chronic Periodontitis and Periodontally Healthy Subjects. A cross-sectional Study. *J Periodontol Res* 2009; 44(3): 411-17.
61. Hodgetts S, Radley H, Davies M, Grounds MD. Reduced necrosis of dystrophic muscle by depletion of host neutrophils, or blocking TNF alpha function with Etanercept in mdx mice. *Neuromuscul Disord* 2006; 16(9-10): 591-602.
62. Clark, I.A. How TNF was recognized as a key mechanism of disease. *Cytokine Growth Factor Rev* 2007; 18(3-4): 335-43.
63. Dargelos E, Renaud V, Decossas M, Bure C, Lambert O, Poussard S. Caveolae-mediated effects of TNF- α on human skeletal muscle cells. *Experimental Cell Research* 2018; 370(2): 623-31.
64. Frost RA, Nystrom GJ, Lang CH. Tumor Necrosis Factor- α Decreases Insulin Like Growth Factor-I Messenger Ribonucleic Acid Expression in C2C12 Myoblast Via a Jun N-terminal Kinase Pathway. *Endocrinology* 2003; 144(5): 1770-79.
65. Broussard SR, Mc Cusker RH, Novakofski JE, Strle K, Shen, WH, Johnson RW, et al. Cytokine-hormone interactions: tumor necrosis factor alpha impairs biologic activity and downstream activation signals of the insulin-like growth factor I receptor in myoblasts. *Endocrinology* 2003; 144: 2988-96.
66. Kern PA, Ranganathan S, Li C, Wood L, Ranganathan G. Adipose tissue tumor necrosis factor and interleukin-6 expression in human obesity and insulin resistance. *Am J Physiol Endocrinol Metab* 2001; 280: 745-51.
67. Bastard JP, Maachi M, Van Nhieu JT, Jardel C, Bruckert E, Grimaldi A, Robert JJ, Capeau J, Hainque B. Adipose tissue IL-6 content correlates with resistance to insulin activation of glucose uptake both in vivo and in vitro. *J Clin Endocrinol Metab* 2002; 87(5).
68. Huang and Czech MP. The GLUT4 glucose transporter. *Cell Metab.* 2007; 5:237-52
69. Fujii N, Jessen N, Goodyear LJ. AMP-activated protein kinase and the regulation of glucose transport. *Am J Physiol Endocrinol Metab* 2006; 291: 867-77.
70. Kramer HF, Witzack CA, Fujii N, Jessen N, Taylor, EB, Arnolds DE, Sakamoto K, Hirshman MF, Goodyear LJ. Distinct signals regulate AS160 phosphorylation in response to insulin, AICAR, and contraction in mouse skeletal muscle. *Diabetes* 2006; 55:2067-76.
71. Winkelmann, C. Inactivity and Inflammation: Selected Cytokines as Biologic Mediators in Muscle Dysfunction During Critical Illness. *AACN Clinical Issues* 2004; 15(1): 74-82.
72. Reid MB and Li YP. Tumor necrosis factor-alpha and muscle wasting: a cellular perspective. *J Clin Endocrinol Metab* 2001; 2(5): 269-72.
73. Sulistio E, Masulili SL, Lessang R, Auerkari EI. The Influence of Smoking on IL-17 Cytokine in Chronic Periodontitis Patients. *Journal of International Dental and Medical Research* 2019;12(1):199-202.
74. Li H, Malhotra S, Kumar A. Nuclear factor-kappa B signaling in skeletal muscle atrophy. *Journal of Molecular Medicine* 2008; 86(10): 1113-26.
75. Amin MF, Meidyawati R, Djamil MS, Latief BS. Inflammatory cytokine serum levels in sockets following extraction of teeth with apical periodontitis. *Journal of International Dental and Medical Research* 2019;12(1):129-32.
76. Grounds, M. D., Radley, H. G., Gebiski, B. L., Bogoyevitch, M. A, & Shavlakadze, T. Implications of cross-talk between tumour necrosis factor and insulin-like growth factor-1 signalling in skeletal muscle. *Clinical and Experimental Pharmacology and Physiology* 2008; 35(7): 846-51.

Periodontitis Affects Skeletal Muscle Metabolism Through an Increase in Proinflammatory Cytokines

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Davidson Fróis Madureira, Izabella Lucas De Abreu Lima, Giselle Cabral Costa, Elizabeth Maria Bastos Lages et al. "Tumor Necrosis Factor-alpha in Gingival Crevicular Fluid as a Diagnostic Marker for Periodontal Diseases: A Systematic Review", *Journal of Evidence Based Dental Practice*, 2018 1%

Publication

- 2** Leandro Chambrone, Adriana M. Foz, Mariana R. Guglielmetti, Cláudio M. Pannuti et al. "Periodontitis and chronic kidney disease: a systematic review of the association of diseases and the effect of periodontal treatment on estimated glomerular filtration rate", *Journal of Clinical Periodontology*, 2013 1%

Publication

- 3** Nouf Alshibani. "Resolvins as a Treatment Modality in Experimental Periodontitis: A Systematic Review of Preclinical Studies", *Cureus*, 2022 1%

Publication

4	"Free Communications Sessions 17-32 Thursday, August 29, 2013", International Dental Journal, 2013. Publication	1 %
5	link.springer.com Internet Source	1 %
6	www.myukk.org Internet Source	<1 %
7	www.wjgnet.com Internet Source	<1 %
8	Anilei Hoare, Cristopher Soto, Victoria Rojas-Celis, Denisse Bravo. "Chronic Inflammation as a Link between Periodontitis and Carcinogenesis", Mediators of Inflammation, 2019 Publication	<1 %
9	Makoto Matsushita, Kazutoshi Fujita, Koji Hatano, Marco A. De Velasco, Hirotsugu Uemura, Norio Nonomura. "Connecting the Dots Between the Gut-IGF-1-Prostate Axis: A Role of IGF-1 in Prostate Carcinogenesis", Frontiers in Endocrinology, 2022 Publication	<1 %
10	etheses.bham.ac.uk Internet Source	<1 %
11	A. Ginion. "Inhibition of the mTOR/p70S6K pathway is not involved in the insulin-	<1 %

sensitizing effect of AMPK on cardiac glucose uptake", AJP Heart and Circulatory Physiology, 05/20/2011

Publication

12

Pediatric Inflammatory Bowel Disease, 2013.

Publication

<1 %

13

e-journal.unair.ac.id

Internet Source

<1 %

14

pdfs.semanticscholar.org

Internet Source

<1 %

15

www.apps.org.au

Internet Source

<1 %

16

www.atsjournals.org

Internet Source

<1 %

17

www.nature.com

Internet Source

<1 %

18

Akhmedov, Dmitry, and Rebecca Berdeaux. "The effects of obesity on skeletal muscle regeneration", Frontiers in Physiology, 2013.

Publication

<1 %

19

Bing Han. "Insulin-like growth factor-1 (IGF-1) and leucine activate pig myogenic satellite cells through mammalian target of rapamycin (mTOR) pathway", Molecular Reproduction and Development, 05/2008

Publication

<1 %

20

Gillian M. P. Galbraith, R. Britt Steed, John J. Sanders, Janardan P. Pandey. "Tumor Necrosis Factor Alpha Production by Oral Leukocytes: Influence of Tumor Necrosis Factor Genotype", Journal of Periodontology, 1998

Publication

<1 %

21

Nora Silva. "Characterization of progressive periodontal lesions in chronic periodontitis patients: levels of chemokines, cytokines, matrix metalloproteinase-13, periodontal pathogens and inflammatory cells", Journal Of Clinical Periodontology, 3/2008

Publication

<1 %

22

Tae Ho Kim, Sung E Choi, Eun Suk Ha, Jong Gab Jung, Seung Jin Han, Hae Jin Kim, Dae Jung Kim, Yup Kang, Kwan Woo Lee. "IL-6 induction of TLR-4 gene expression via STAT3 has an effect on insulin resistance in human skeletal muscle", Acta Diabetologica, 2011

Publication

<1 %

23

external.doyma.es

Internet Source

<1 %

24

mafiadoc.com

Internet Source

<1 %

25

www.mdpi.com

Internet Source

<1 %

26

"The Plasticity of Skeletal Muscle", Springer Science and Business Media LLC, 2017

Publication

<1 %

27

Bárbara Capitanio de Souza, Bibiana Franzen Matte, André Luiz Lopes, Bruno Costa Teixeira, Marcelo Lazzaron Lamers. "Periodontal Disease Impairs Muscle Recovery by Modulating the Recruitment of Leukocytes", Inflammation, 2019

Publication

<1 %

28

Leslie E Stefanyk. "The interaction between adipokines, diet and exercise on muscle insulin sensitivity :", Current Opinion in Clinical Nutrition & Metabolic Care, 05/2010

Publication

<1 %

29

Otis, Jeffrey S., Sarah Niccoli, Nicole Hawdon, Jessica L. Sarvas, Melinda A. Frye, Adam J. Chicco, and Simon J. Lees. "Pro-Inflammatory Mediation of Myoblast Proliferation", PLoS ONE, 2014.

Publication

<1 %

30

Yue Hou, Xin Wang, Cong-Xiao Zhang, Yu-Dan Wei, Li-Li Jiang, Xiao-Yu Zhu, Yu-Jun Du. "Risk factors of periodontal disease in maintenance hemodialysis patients", Medicine, 2017

Publication

<1 %

31

atm.amegroups.com

Internet Source

<1 %

32	biolres.biomedcentral.com Internet Source	<1 %
33	citeseerx.ist.psu.edu Internet Source	<1 %
34	ijirms.in Internet Source	<1 %
35	ir.lib.uth.gr Internet Source	<1 %
36	mspace.lib.umanitoba.ca Internet Source	<1 %
37	pure.uva.nl Internet Source	<1 %
38	scholar.qsensei.com Internet Source	<1 %
39	vuir.vu.edu.au Internet Source	<1 %
40	www.intechopen.com Internet Source	<1 %
41	www.karger.com Internet Source	<1 %
42	"Salivary Bioscience", Springer Science and Business Media LLC, 2020 Publication	<1 %

43 C. A. Witczak. "CaMKII regulates contraction- but not insulin-induced glucose uptake in mouse skeletal muscle", AJP Endocrinology and Metabolism, 06/01/2010 <1 %

Publication

44 Hong Li. "Nuclear factor-kappa B signaling in skeletal muscle atrophy", Journal of Molecular Medicine, 10/2008 <1 %

Publication

45 Ji-Youn Kim, Ki-Rim Kim, Han-Na Kim. "The Potential Impact of Salivary IL-1 on the Diagnosis of Periodontal Disease: A Pilot Study", Healthcare, 2021 <1 %

Publication

46 N/A. "Current World Literature", Current Opinion in Clinical Nutrition and Metabolic Care, 05/2004 <1 %

Publication

47 Ram R. Miller. "Association Between Interleukin-6 and Lower Extremity Function After Hip Fracture—The Role of Muscle Mass and Strength", Journal of the American Geriatrics Society, 4/11/2008 <1 %

Publication

48 Young Park. "Assessment of IL-6, IL-8 and TNF- α levels in the gingival tissue of patients <1 %

with periodontitis", Experimental and
Therapeutic Medicine, 2013

Publication

49

J. P. Scarth. "Modulation of the growth hormone–insulin-like growth factor (GH–IGF) axis by pharmaceutical, nutraceutical and environmental xenobiotics: An emerging role for xenobiotic-metabolizing enzymes and the transcription factors regulating their expression. A review", Xenobiotica, 2008

Publication

<1 %

50

Katy Thouvenot, Teva Turpin, Janice Tailé, Karine Clément, Olivier Meilhac, Marie-Paule Gonthier. "Links between Insulin Resistance and Periodontal Bacteria: Insights on Molecular Players and Therapeutic Potential of Polyphenols", Biomolecules, 2022

Publication

<1 %

51

Oral Infections and General Health, 2016.

Publication

<1 %

52

P. J. Klover, T. A. Zimmers, L. G. Koniaris, R. A. Mooney. "Chronic Exposure to Interleukin-6 Causes Hepatic Insulin Resistance in Mice", Diabetes, 2003

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Periodontitis Affects Skeletal Muscle Metabolism Through an Increase in Proinflammatory Cytokines

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
