



purwo sri rejeki &lt;purwo-s-r@fk.unair.ac.id&gt;

---

**Your manuscript ID JBCPP.2021.0393.R1 - submission confirmation**

1 message

---

**Journal of Basic and Clinical Physiology and Pharmacology**

Sat, Feb 5, 2022 at 10:05

&lt;onbehalf@manuscriptcentral.com&gt;

AM

Reply-To: alberto\_marra@hotmail.it

To: purwo-s-r@fk.unair.ac.id, purwo\_faal@yahoo.com

04-Feb-2022

Dear Dr. Rejeki,

The revision of your manuscript entitled "Moderate-Intensity Exercise Decreases the Circulating Level of Betatrophin and Its Correlation among Markers of Obesity in Women" has been successfully submitted online and is presently being given full consideration for publication in Journal of Basic and Clinical Physiology and Pharmacology (JBCPP).

Your manuscript ID is JBCPP.2021.0393.R1.

Please mention the above manuscript ID in all future correspondence or when calling the office for questions. If there are any changes in your affiliation, street address or e-mail address, please log in to ScholarOne Manuscripts at <https://mc.manuscriptcentral.com/jbcpp> and edit your user information as appropriate.

You can also view the status of your manuscript at any time by checking your Author Center after logging in to <https://mc.manuscriptcentral.com/jbcpp>.

Thank you for submitting your manuscript to JBCPP.

Kind regards

Mr. Deepan Selvaraj

Journal of Basic and Clinical Physiology and Pharmacology

[journal.editor@degruyter.com](mailto:journal.editor@degruyter.com)

For news highlights from this journal and other De Gruyter publications, check out our blog De Gruyter Conversations at <https://blog.degruyter.com/>



purwo sri rejeki &lt;purwo-s-r@fk.unair.ac.id&gt;

---

**JBCPP.2021.0393.R1 - DecisionAccept**

1 message

---

**Journal of Basic and Clinical Physiology and Pharmacology**

Wed, Feb 16, 2022 at 1:49

&lt;onbehalf@manuscriptcentral.com&gt;

AM

Reply-To: journal.editor@degruyter.com

To: purwo-s-r@fk.unair.ac.id, purwo\_faal@yahoo.com

15-Feb-2022

Dear Dr. Rejeki:

I would like to thank you for submitting your manuscript entitled "Moderate-Intensity Exercise Decreases the Circulating Level of Betatrophin and Its Correlation among Markers of Obesity in Women" to Journal of Basic and Clinical Physiology and Pharmacology (JBCPP). Your manuscript has been reviewed, and it is a pleasure to accept it for publication in JBCPP. The comments of the reviewer(s) are included at the bottom of this letter.

The JBCPP production office will contact you for proofreading in the near future. Your article will be published ahead of print as soon as possible, and assigned to an online issue at a later time.

Thank you for your fine contribution. On behalf of the Editors of Journal of Basic and Clinical Physiology and Pharmacology we look forward to your continued contributions to the Journal.

Kind regards

Dr. Mariarosaria De Luca

Associate Editor, Journal of Basic and Clinical Physiology and Pharmacology

Reviewer(s)' Comments to Author:

Reviewer: 1

Comments to the Author

Thanks for the add e good luck for your paper!

For news highlights from this journal and other De Gruyter publications, check out our blog De Gruyter Conversations at <https://blog.degruyter.com/>

## Publish Open Access with De Gruyter!

1 message

no-reply@copyright.com <no-reply@copyright.com>  
To: purwo-s-r@fk.unair.ac.id

Sat, Feb 19, 2022 at 6:23 AM



DE GRUYTER

### *Publish Open Access with De Gruyter!*

Dear Purwo Sri Rejeki,

Congratulations on being accepted for publication in *Journal of Basic and Clinical Physiology and Pharmacology* for the following manuscript:

Manuscript DOI: 10.1515/JBCPP-2021-0393

Manuscript ID: JBCPP.2021.0393.R1

Manuscript Title: Moderate-Intensity Exercise Decreases the Circulating Level of Betatrophin and Its Correlation among Markers of Obesity in Women

Published by: Walter De Gruyter GmbH

Your article has been accepted in a hybrid open access journal.

Your institution may have agreed to pay open access charges on your behalf. Click the link below to opt into open access or to determine whether your institution has negotiated free or reduced-cost publication on your behalf. However, you are not required to publish open access.

We cooperate with the Copyright Clearance Center (CCC) to guide you through the entire process via their RightsLink® e-commerce solution.

[I want to publish Open Access!](#)

To publish open access and determine whether your institution has agreed to pay fees, please [click here](#).

To complete a secure transaction, you will need a [RightsLink account](#). If you do not have one already, you will be prompted to register as you are checking out your author charges. This is a very quick process; the majority of your registration form will be pre-populated automatically with information we have already supplied to RightsLink.

If you have any questions about these charges, please contact CCC [Customer Service](#)

using the information below.

Sincerely,  
Walter De Gruyter GmbH

Tel.: +1-877-622-5543 / +1-978-646-2777  
[publicationservices@copyright.com](mailto:publicationservices@copyright.com)  
[www.copyright.com](http://www.copyright.com)



This message (including attachments) is confidential, unless marked otherwise. It is intended for the addressee(s) only. If you are not an intended recipient, please delete it without further distribution and reply to the sender that you have received the message in error.

---

## Overdue - Check your proof pdf: JBCPP.2021.0393.R1

---

noreply@degruyter.com <noreply@degruyter.com>

Fri, Mar 11, 2022 at 7:35 AM

To: purwo-s-r@fk.unair.ac.id

Cc: dean.maharani.widiatmaja-2018@fk.unair.ac.id, pertreesia@tnq.co.in

11 March 2022

Dear Purwo Sri Rejeki,

This task is overdue now.

MS-ID: JBCPP.2021.0393.R1

Purwo Sri Rejeki: Moderate-Intensity Exercise Decreases the Circulating Level of Betatrophin and Its Correlation among Markers of Obesity in Women

Please find below the link to access your galley proof. Please check it carefully and upload your corrections as soon as possible.

The attached Users Guide provides step-by-step instructions for submitting your corrections.

[Click here to access your galley proof](#)

For Online Proofs:

Please use Firefox, Google Chrome or Safari to access the proof. The browser Microsoft Edge is not being supported.

For PDF Proofs:

The attached User's Guide provides step-by-step instructions for inserting your corrections.

When making corrections, please always adhere to the requirements below.

- Mark your corrections directly within the PDF file or on a printed copy.
- DO NOT modify any source files (.doc, .tex, etc.) or the text in the PDF!
- Confirm that all authors' names and affiliations are listed correctly.
- Check all hyperlinks within your article (text, footnotes, references) for accuracy and functionality.

If you want to use KUDOS ([www.growkudos.com](http://www.growkudos.com)) for your publication, please click **here** to enter your "Plain Language Summary".

Once you submit your galley proof PDF any corrections will be implemented and your article will be published online at [www.degruyter.com](http://www.degruyter.com).

Sincerely,

TNQ Vendor

[dg-wms@tnq.co.in](mailto:dg-wms@tnq.co.in)

Production Department de Gruyter:

Pertreesia Thomas

[pertreesia@tnq.co.in](mailto:pertreesia@tnq.co.in)

PLEASE NOTE: *Please do not reply directly to this message.* It was sent from an automated mailbox that cannot accept incoming email. For questions, please refer to the contact(s) mentioned above.



DG\_Authors\_guide\_on\_viewing\_and\_annotating\_PDF\_files.pdf

451K

# Author's guide on viewing and annotating PDF files

---

Version 1.0

A PDF file can be used to review and annotate galley proofs on-screen and to transfer them between the author/editor and De Gruyter. It allows the user to add comments easily on-screen. **Advantages:** Reduction of delivery time and cost, elimination of errors caused by illegible or confusing handwritten annotations, easy to access and store (do not restrict by a password!), less use of paper and therefore more environmentally friendly.

The basics of commenting are shown in a short video provided on the official Adobe website:

<http://tv.adobe.com/watch/learn-acrobat-x/getting-started-the-basics-of-commenting/>


(Copyright © 2012 Adobe Systems Incorporated).

## 1 How to use the annotation tools

Please use the freely available program Adobe Acrobat Reader (version XI or higher) to view and annotate the galley proofs of your article or book. It can be downloaded for Windows PC, Mac and other platforms at:

<http://get.adobe.com/reader/>.

- Take care to annotate your corrections as simply, as short and as clear as possible. General comments should be written in an e-mail.
- Do not change the actual text in the pdf file, and use the Comment tool bar to annotate your changes to be incorporated (see below).




## 2 Standard tools in Adobe Acrobat Reader XI (or higher versions)

Open the PDF file and click on 'Comments' (if this option is not available you will need to upgrade your current version of Adobe Reader to a higher version; see the link above).

This opens a task pane and, below that, a list of all possible 'Annotation' tools you are able to use:


### 2.1 Insert Text

Select the icon 'Insert Text' from the Comment tool bar. Then click to set the cursor location in the text and simply start typing. The text will appear in a comment box. You may also copy-and-paste text from another file into the comment box.




### 2.2 Replace Text

Select the icon 'Replace Text' from the Comment tool bar. To highlight the text to be replaced, click and drag the cursor over the text. Then simply type in the replacement text, which will appear in a comment box. You may also copy-and-paste text from another file into this box. To replace formatted text (e.g. an equation) please attach an authorized file (see below 'Attach a file').



### 2.3 Remove Text

Select the icon 'Strikethrough' from the Comment tool bar. Then click and drag over the text to be deleted and it will be struck through.




### 2.4 Highlight Text

Select the icon 'Highlight Text' from the Comment tool bar. Then click and drag over the text (e.g. to highlight an index entry). To make a comment, double click on the highlighted text and simply start typing (see below 'Add note to text').




## 2.5 Add Note to Text

Select the icon 'Add note to text' from the Comment tool bar. Then click and drag over the text and a comment box will open automatically to simply start typing.




## 2.6 Attach a File

Select the icon 'Attach File' from the Comment tool bar. Then click on the element to be replaced (fig., tab. or text passage). A window will automatically open allowing you to attach the file and a graphic (graph, paperclip, attachment, tag) will appear at the chosen spot indicating the insertion of a file.



## 2.7 Leave a Note

Select the icon 'Add Sticky Note' from the Comment tool bar. Then click to set the placement of the note on the document and simply start typing. **Attention:** Do not use this feature to make text edits!




## 2.8 Drawing Markups

The 'Drawing Markups' may be used for corrections that other tools may not be able to express (e.g. alignment or spacing comments). Use the comment box by double clicking on the drawing markup (or right click and select 'Open Pop-Up Note') and typing in your annotation.




## 2.9 Review (Comments List)

Your changes will appear automatically in a list below the Comment tool bar. You can navigate by clicking on a comment in the list. Alternatively, double click on any comment to open the comment box.



## 2.10 Undo Comment / Delete Comment

To undo any comments made, click on 'Edit' in the main Adobe menu and then 'Undo'. To delete comments, use the right click and select 'Delete' (or choose the comment to be deleted in the Comments List and click 'Del' on your keyboard).





### 3 Author queries and margin notes

Please ensure that you have answered any questions raised in author queries and margin notes in the galley proofs (see examples below) and that you have inserted all corrections. Please note that later inclusions of any subsequent corrections cannot be guaranteed.

Query No	Page No	Query
Q1	1	Please indicate corresponding author and provide email also.
Q2	1	Please provide the missing city and country for affiliations [1 and 2].
Q3	3	Please check if the inserted short title is OK as typeset.

any  $u \in \mathbb{R}^d$ . Hence,  $b|s| < n$  implies  $|A(u)s| < n$  and  $|A(u)s| < n$  implies  $a|s| < n$ . Therefore,

$$\begin{aligned} \int_{\mathbb{R}^d} \int_{a|s| < n} p_T(u, s) ds du - \int_{\mathbb{R}^d} \int_{|A(u)s| < n} p_T(u, s) ds du \\ \leq \int_{\mathbb{R}^d} \int_{a|s| < n < b|s|} |p_T(u, s)| ds du. \end{aligned}$$

**CE Note 2:**  
Please check  
the “|” in  
formula

Purwo Sri Rejeki\*, Pradika Gita Baskara, Lilik Herawati, Adi Pranoto, Hayuris Kinandita Setiawan, Ronny Lesmana and Shariff Halim

# Moderate-intensity exercise decreases the circulating level of betatrophin and its correlation among markers of obesity in women

<https://doi.org/10.1515/jbcpp-2021-0393>

Received December 22, 2021; accepted February 15, 2022; published online March 15, 2022

## Abstract

**Objectives:** Positive energy homeostasis due to over-nutrition and a sedentary lifestyle triggers obesity. Obesity has a close relationship with elevated levels of betatrophin and may increase the risk of developing metabolic syndrome. Therefore, lifestyle modification through a non-pharmacological approach based on physical exercise is the right strategy in lowering betatrophin levels. This study aimed to analyze the effect of moderate-intensity interval and continuous exercises on decreased betatrophin levels and the association between betatrophin levels and obesity markers in women.

**Methods:** A total of 30 women aged 20–24 years old were randomly divided into three groups. Measurement of betatrophin levels using Enzyme-Linked Immunosorbent Assay (ELISA). Data analysis techniques used were one-way ANOVA and parametric linear correlation.

**Results:** The results showed that the average levels of betatrophin pre-exercise were  $200.40 \pm 11.03$  pg/mL at CON,  $203.07 \pm 42.48$  pg/mL at MIE,  $196.62 \pm 21.29$  pg/mL at MCE, and  $p=0.978$ . Average levels of betatrophin post-exercise were  $226.65 \pm 18.96$  pg/mL at CON,  $109.31 \pm 11.23$  pg/mL at

MIE,  $52.38 \pm 8.18$  pg/mL at MCE, and  $p=0.000$ . Pre-exercise betatrophin levels were positively correlated with age, BMI, FM, WHR, FBG, and PBF ( $p \leq 0.001$ ).

**Conclusions:** Our study showed that betatrophin levels are decreased by 10 min post-MIE and post-MCE. However, moderate-intensity continuous exercise is more effective in lowering betatrophin levels than moderate-intensity interval exercise. In addition, pre-exercise betatrophin levels also have a positive correlation with obesity markers.

**Keywords:** betatrophin; moderate-intensity exercise; myokine; obesity.

## Introduction

Obesity is a metabolic disease that is the third leading cause of death and has become a global epidemic [1, 2]. Obesity is already considered an epidemic and has now become a pandemic [3]. This is due to the prevalence rate of obesity experiencing a continuous increase in both developed countries and developing countries [1, 4]. It was estimated that 1.9 billion people over the age of 18 were overweight, and 650 million of who were obese comprising 11% of men and 15% of women [5], meaning that more than one-third of adults globally were obese [6]. If every year, the prevalence of obesity continues to increase, it is estimated that by 2025, the prevalence of obesity will be 18% of men and 21% of women [7]. Based on Basic Health Research (Riskesdas) in 2018 showed that the prevalence of obesity in Indonesia people aged over 18 years old had increased, which was 21.8%, the number was higher than in 2013 (14.8%) and in 2007 (10.5%) [8]. Meanwhile, the prevalence of obesity in children in recent decades has also increased worldwide [9, 10]. In 2016 estimated that the prevalence of obesity in the world in children and adolescents aged 5–19 years is 124 million people and 213 million people are overweight [11]. Based on previous data, it was also reported that more than 70% of adolescents worldwide have low levels of physical activity [12]. Physiologically, changes in lifestyle behavior, such as increased time sitting, lying down [13], and reduced desire to exercise are

\*Corresponding author: Purwo Sri Rejeki, Department of Medical Physiology and Biochemistry, Physiology Division, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, E-mail: purwo-s-r@fk.unair.ac.id, Phone: +62 821 4155 9388

Pradika Gita Baskara, Sport Health Science, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

Lilik Herawati and Hayuris Kinandita Setiawan, Department of Medical Physiology and Biochemistry, Physiology Division, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

Adi Pranoto, Doctoral Program of Medical Science, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

Ronny Lesmana, Department of Biomedical Science, Physiology Division, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

Shariff Halim, Clinical Research Centre, Management and Science University, Shah Alam, Malaysia

the causes of increased body weight and risk of obesity [14, 15].

Obesity increases the risk of developing metabolic syndrome, as obesity is associated with increased levels of total cholesterol, increased triglyceride and low-density lipoprotein (LDL) levels, and decreased levels of high-density lipoprotein (HDL) [16], so that obese individuals are susceptible to type 2 diabetes mellitus [17], hypertension [18], hypercholesterolemia [19], several types of cancer [20, 21], nonalcoholic fatty liver disease (NAFLD), and dyslipidemia [22]. In addition, obesity also causes the heart workload to increase, thus increasing the risk of cardiovascular disease [23, 24]. In the condition of obesity, there is also impaired insulin secretion by pancreatic  $\beta$ -cells caused by increased levels of betatrophin [25–28]. Elevated levels of betatrophin can also have an effect on the kidneys and female organs, resulting in kidney problems [29] and polycystic ovarian syndrome (PCOS) [30]. Excessive levels of betatrophin can also inhibit the performance of the lipoprotein lipase (LPL) enzymes [31, 32]. The stunted performance of LPL enzymes causes fat to be irreversible into an energy source [33], resulting in increased fat storage in adipocytes [34]. Increased fat storage in adipocytes is one of the causes of obesity [35, 36]. Therefore, the discovery of new solutions for obesity prevention is becoming a very important target to do and potential therapies with high visibility at a low cost for obesity prevention in the global community are urgently needed.

Over the past decade, several interventions have been made on how to tackle obesity and boost metabolism without increasing the activity of internal stressors in the body [37, 38]. Recent findings suggest that exercise may lower betatrophin levels in obese people [26, 27]. Physical activity is organized in exercise programs oriented to strength development, low-intensity fitness, or a combination of these two [39]. The design of a program must follow specific rules with regard to the exercise to be undertaken, as well as its intensity, duration, and frequency, in order to obtain benefits [39]. A research conducted by Abu-Farha et al. [26] reported that the combination of both moderate-intensity aerobic (30 min) and resistance exercises using either a treadmill or cycling (10 min) lowered betatrophin levels in obesity. Likewise, the study of Susanto et al. [37] proved that moderate-intensity exercise lowered betatrophin levels in nonprofessional athletes. However, a research conducted by Enteshary et al. [40] found different results in which betatrophin levels increased post-moderate-intensity exercise in women with Type 2 Diabetes (T2D). Thus, the fundamental impact of moderate-intensity exercise on decreasing betatrophin levels remains controversial. Therefore, the aim of the present paper was to elaborate the

potential regulation of moderate-intensity exercise decreases the circulating level of betatrophin and its correlation among markers of obesity in women.

## Materials and methods

### Experimental design

This research is a real experiment with a research design of randomized pretest–posttest control group design. The total subjects participating in the study were 30 women aged  $21.27 \pm 0.25$  years old with body mass index (BMI) of  $28.82 \pm 0.30$  kg/m<sup>2</sup>, blood pressure (systolic blood pressure  $113.33 \pm 1.30$  mmHg; diastolic blood pressure  $76.00 \pm 1.23$  mmHg), resting heart rate (RHR) of  $75.57 \pm 2.23$  bpm, fasting blood glucose (FBG) of  $90.23 \pm 1.33$  mg/dL, hemoglobin (Hb) of  $14.99 \pm 0.24$  g/dL, and maximum oxygen volume ( $VO_{2max}$ ) of  $28.08 \pm 0.66$  mL/kg/min examined using astrand 6 min cycle test method. All subjects were examined both physically and psychologically. All subjects obtained information orally or in writing about the research. The subjects filled out and signed informed consent before participating in the research. All of our research procedures comply with the Declaration of the World Medical Association of Helsinki regarding the ethical conduct of research involving human subjects.

### Exercise protocol

The exercise program was applied and supervised by professional officers from the Fitness Center of Malang Health Office. The subjects were randomly divided into three groups: CON (n=10, control without intervention), MIE (n=10, moderate-intensity interval exercise), and MCE (n=10, moderate-intensity continuous exercise). The MIE intervention was carried out by employing the subjects running on a treadmill with moderate intensity of 60–70%  $HR_{max}$  for 45 min with details of 5 min of warming up (50–60%  $HR_{max}$ ), 35 min of core exercise (5 min of work (60–70%  $HR_{max}$ ) interspersed with active recovery on the treadmill for 2.5 min (50–60%  $HR_{max}$ ) performed 5 times a while, and 5 min of cooling (50–60%  $HR_{max}$ ) [41–44]. The intervention was conducted at 07.00–09.00 am [45, 46] using a treadmill (Pulsar 4.0 HP Cosmos Sports & Medical, Nussdorf-Traunstein, Germany) [47]. Heart rate monitoring during exercise used a polar heart rate monitor (Polar H10 Heart Rate Sensor, Inc., USA). The research environment had a room temperature of  $26 \pm 1$  °C and a room humidity level of 50–70% [48, 49].

### Blood samples

Blood samples were collected from a 4 mL cubital vein after a 12 h overnight fasting [50, 51]. At the time of the blood draw, the subjects were in a lying position. Blood samples were taken at 30 min pre-exercise and 10 min post-exercise. At the time of pre-exercise and post-exercise blood draws, the subjects remained in a fasting condition but they were allowed to drink mineral water without calories to prevent dehydration. The blood was centrifuged for 15 min at a speed of 3,000 rpm. The serum was then separated and stored at  $-80$  °C for analysis of betatrophin levels on the next day [51–53].

## Blood analysis

Examination of betatrophin levels was conducted at the Laboratory of Physiology of the Faculty of Medicine, Universitas Brawijaya, Malang, using the enzyme-linked immunosorbent assay (ELISA) kit (Cat No. E11644h; EIAab Science Co., Wuhan) with a standard curve range of 78–5,000 pg/mL and betatrophin sensitivity levels in a kit of 32 pg/mL. FBG examination was carried out using Accu-Chek Performa (Roche, Mannheim, Germany) with mg/dL concentration unit, while Hb examination used Easy Touch (Easy Touch, Hsinchu, Taiwan) with g/dL concentration unit.

## Anthropometric measurements and physical fitness

Body height measurements were conducted using a stadiometer (SECA, Chino, CA, USA). Measurements of obesity markers included body weight, body mass index (BMI), body fat percentage (PBF), and fat mass (FM) using TANITA Body Composition Analyzer DC3607601(2)-1604 FA (TANITA Corporation of America, Inc., Arlington Heights, IL, USA). Measurement of the waist to hip ratio (WHR) was conducted by calculating waist circumference (WC) divided by hip circumference (HC). Measurement of maximal oxygen volume ( $VO_{2max}$ ) by the Astrand 6 min cycle test method was performed using a Monark 828 E Version 1010 ergo cycle (Monark, Vansbro, Sweden). Measurement of resting heart rate (RHR) was conducted using Beurer Pulse Oximeter (PO 30 Pulse Oximeter, Beurer North America LP, Hallandale Beach, FL, USA). Blood pressure was measured using an automated device OMRON (OMRON Model HEM-7130 L, Omron Co., Osaka, Japan) at the non-dominant arm 3 times consecutively with a 1–2 min interval between two measurements while the participants were in a seated position.

## Statistical analysis

Statistical analysis was conducted using SPSS software version 17 (SPSS Inc., Chicago, IL, USA). The normality test was conducted using

the Shapiro-Wilk test, while the homogeneity test used the Levene test. Different tests were conducted using paired sample T-Test, one-way ANOVA, and then followed by Tukey's honestly significant difference (HSD) post hoc test and linear correlation with Pearson product-moment model. Data were presented as the mean  $\pm$  standard error of the mean (SEM). Significant will be considered with p value  $\leq 0.05$ .

## Results


The basic profiles of the samples are displayed in Table 1. Table 1 shows that the average characteristic data of the study subjects showed no significant differences in all parameters of each group ( $p \geq 0.05$ ). The results of the analysis of betatrophin levels between pre-exercise and post-exercise in each group are presented in Figure 1.

Figure 1 shows that betatrophin levels between pre-exercise and post-exercise in the CON group had no change and tended to be the same, while in the MIE and MCE groups, there was a change and the betatrophin levels tended to decrease. Paired sample T-Test results in the CON group showed that there was no significant difference in average betatrophin levels between pre-exercise and post-exercise ( $200.40 \pm 11.03$  vs.  $226.65 \pm 18.96$  pg/mL, (p-value=0.269)) (Figure 1). However, the MIE group showed significant differences in average betatrophin levels between pre-exercise and post-exercise ( $203.08 \pm 42.48$  vs.  $109.31 \pm 11.23$  pg/mL, (p-value=0.046)) (Figure 1). Likewise, the MCE group showed significant differences in average levels of betatrophin between pre-exercise and post-exercise ( $196.62 \pm 21.29$  vs.  $52.38 \pm 8.18$  pg/mL, (p-value=0.000)) (Figure 1). The results

**Table 1:** The basic profiles of the samples.

Parameter	Group			One way-ANOVA p-Value
	CON (n=10)	MIE (n=10)	MCE (n=10)	
Age, yrs	21.30 $\pm$ 0.30	21.50 $\pm$ 0.64	21.00 $\pm$ 0.33	0.734
Body height, m	1.57 $\pm$ 0.01	1.59 $\pm$ 0.02	1.55 $\pm$ 0.01	0.060
Body weight, kg	73.30 $\pm$ 1.80	73.33 $\pm$ 2.93	67.63 $\pm$ 1.56	0.123
BMI, kg/m <sup>2</sup>	29.42 $\pm$ 0.40	28.81 $\pm$ 0.70	28.23 $\pm$ 0.38	0.281
PBF (%)	45.17 $\pm$ 1.14	42.40 $\pm$ 1.43	42.98 $\pm$ 0.61	0.199
FM, kg	35.66 $\pm$ 1.36	34.32 $\pm$ 2.49	31.87 $\pm$ 1.82	0.391
WHR	0.81 $\pm$ 0.01	0.81 $\pm$ 0.01	0.79 $\pm$ 0.01	0.420
Hb, g/dL	15.09 $\pm$ 0.57	14.97 $\pm$ 0.29	14.93 $\pm$ 0.39	0.964
FBG, mg/dL	91.90 $\pm$ 1.82	90.50 $\pm$ 2.49	88.30 $\pm$ 2.59	0.552
SBP, mmHg	115.00 $\pm$ 2.69	111.00 $\pm$ 1.79	114.00 $\pm$ 2.21	0.439
DBP, mmHg	78.00 $\pm$ 2.00	73.00 $\pm$ 2.13	77.00 $\pm$ 2.13	0.220
RHR, bpm	73.70 $\pm$ 3.73	77.30 $\pm$ 4.92	75.70 $\pm$ 3.05	0.815
$VO_{2max}$ , mL/kg/min	27.40 $\pm$ 0.61	27.59 $\pm$ 1.29	29.24 $\pm$ 1.37	0.467

BMI, Body mass index; PBF, Percentage of body fat; FM, Fat mass; WHR, Waist to hip ratio; Hb, Hemoglobin; FBG, Fasting blood glucose; SBP, Systolic blood pressure; DBP, Diastolic blood pressure; RHR, Resting heart rate;  $VO_{2max}$ , Maximum oxygen volume; CON, Control group; MIE, Moderate-intensity interval exercise group; MCE, Moderate-intensity continuous exercise group. One way-ANOVA. Data are presented as mean  $\pm$  SEM.



**Figure 1:** Pre-exercise vs. post-exercise betatrophin levels. CON, Control group; MIE, Moderate-intensity interval exercise group; MCE, Moderate-intensity continuous exercise group. Data are presented as mean  $\pm$  SEM. p-Value was obtained using paired sample T-Test to compare post-exercise and pre-exercise betatrophin levels.

of the analysis of betatrophin levels in each group based on the time of blood collection can be seen in Table 2.

Table 2 of the One way-ANOVA test results shows that there was no significant difference in average betatrophin levels based on the timing of pre-exercise blood collection in each group ( $p \geq 0.05$ ), while in post-exercise and delta ( $\Delta$ ) (Post–Pre), the results showed significant differences in average levels of betatrophin ( $p \leq 0.001$ ). Tukey’s HSD post hoc test results showed that there was a significant difference in average levels of post-exercise betatrophin between MIE and CON ( $p \leq 0.001$ ), MCE and CON ( $p \leq 0.001$ ), and MCE and MIE ( $p \leq 0.05$ ). Likewise, delta ( $\Delta$ ) (Post–Pre) showed significant differences in average levels of betatrophin between MIE and CON ( $p \leq 0.05$ ) and MCE and CON ( $p \leq 0.001$ ), while MCE and MIE showed no significant difference ( $p \geq 0.05$ ). The results of the correlation of pre-exercise betatrophin levels with obesity markers and characteristic parameters of the study subjects are shown in Figure 2.

Preliminary research results found a significant association of pre-exercise betatrophin levels with obesity markers and the characteristic parameters of study subjects including age and FBG. Pearson product-moment

linear correlation parametric analysis showed that pre-exercise betatrophin levels were positively correlated with age ( $r=0.557$ ,  $p \leq 0.05$ ), BMI ( $r=0.660$ ,  $p \leq 0.001$ ), FM ( $r=0.385$ ,  $p \leq 0.05$ ), WHR ( $r=0.543$ ,  $p \leq 0.05$ ), FBG ( $r=0.536$ ,  $p \leq 0.05$ ), and PBF ( $r=0.698$ ,  $p \leq 0.001$ ).


## Discussions

Obesity is a growing health problem in society globally and is associated with an increased risk of premature death [54]. Betatrophin is one of the newly identified proteins, mainly expressed in white adipose tissue (WAT), brown adipose tissue (BAT), and liver [55]. Evidence currently developing suggests that betatrophin plays an important role in metabolism, including the synthesis and degradation of lipids in cells, adipocyte differentiation [54], and energy balance [27]. In addition, several interventions have been done on how to overcome obesity and increase metabolism without increasing the activity of internal stressors in the body. However, the fundamental impact of moderate-intensity exercises on decreasing betatrophin levels in individuals with obesity remains controversial.

**Table 2:** Betatrophin levels in each group based on the time of blood collection.

Time	Betatrophin, pg/mL			One Way-ANOVA p-Value
	CON (n=10)	MIE (n=10)	MCE (n=10)	
Pre-exercise	200.40 $\pm$ 11.03	203.07 $\pm$ 42.48	196.62 $\pm$ 21.29	0.978
Post-exercise	226.65 $\pm$ 18.96	109.31 $\pm$ 11.23*	52.38 $\pm$ 8.18*†	0.000
Delta ( $\Delta$ ) post–Pre	26.25 $\pm$ 22.31	–93.76 $\pm$ 40.64*	–144.24 $\pm$ 22.33*	0.001

One way-ANOVA, followed by Tukey’s HSD post hoc test, was used to compare the differences among groups. Data are presented as mean  $\pm$  SEM. \*Significant vs. control group (CON) ( $p \leq 0.001$ ). †Significant vs. moderate-intensity interval exercise group (MIE) ( $p \leq 0.05$ ).



**Figure 2:** Correlation between pre-exercise betatrophin levels and obesity markers and the characteristic parameters of the study subjects. The significant linear correlation between parameters was visualized in the plot model ( $p \leq 0.05$ ). \*Significant with  $p \leq 0.05$  by Pearson's product-moment correlation test. \*\*Significant with  $p \leq 0.001$  by Pearson's product-moment correlation test.

Therefore, we evaluated the effects of moderate-intensity interval exercise and moderate-intensity continuous exercise on decreasing betatrophin levels in obese women.

Our study demonstrated that there was no difference in betatrophin levels between pre-exercise vs. post-exercise on CON, while MIE and MIC showed significant differences in betatrophin levels between pre-exercise and post-exercise (Figure 1). The results were in line with the results of a study

by Abu-Farha et al. [26] reporting that a combination of moderate-intensity aerobic exercise (30 min) and endurance exercise using a treadmill or cycling (10 min) significantly lowered betatrophin levels in obese subjects. Likewise, a research by Susanto et al. [37] reported that moderate-intensity exercise significantly lowered betatrophin levels in non-professional athlete subjects. The decrease in betatrophin levels is likely due to intervention factors (MIE and

MICE). At the time of intervention, there is an increase in the need for energy obtained from blood glucose, thus causing blood glucose levels to decrease. A research by Zheng et al. [56] reported that moderate-intensity aerobic exercise lowered blood glucose levels and improved glycemic control. Likewise, a study by van Dijk et al. [57] using obese subjects who were given moderate-intensity acute exercise interventions concluded that moderate-intensity acute exercise significantly lowered blood glucose levels. Decreased blood glucose levels can cause betatrophin secretion to decrease [58]. Zhang [59] reported that high blood glucose levels in obesity increase betatrophin secretion in the liver. Gusarova et al. [60] revealed that glycemic control in addition to avoiding hypoglycemia and hyperglycemia could also decrease excessive betatrophin expression.

Obesity is a condition where there is an excessive accumulation of body fat [21]. Excess fat in the body leads to increased triglyceride levels (hypertriglyceridemia) [28]. Increase of triglyceride levels could stimulate betatrophin levels, so that obese individuals experience an increase in betatrophin levels [26, 27]. Elevated levels of betatrophin can inhibit the performance of the lipoprotein lipase (LPL) enzymes [31, 32, 59]. Inhibition of LPL enzymes makes triglycerides difficult to convert into energy [33]. In addition to having high levels of betatrophin, obese people also have high blood glucose levels [27, 61]. Meanwhile, the administration of acute exercise intervention with moderate-intensity requires glucose (glucose uptake) to be converted into energy (ATP) from carbohydrates, fats, and proteins [62]. Increased glucose uptake to be converted into energy makes glucose levels in the blood and betatrophin secretion decrease [58], so that acute exercise with moderate intensity can respond to the decrease in betatrophin levels in the body faster [26, 27].

The literature states that exercise can be used as a useful tool for the prevention, treatment, and rehabilitation of several diseases, such as diabetes mellitus and cardiovascular disease [39]. In diabetic patients exercise can be recommended and prescribed to manage blood glucose and improve the quality of general good health [13]. However, this recommendation must also take into account the lifestyle and habits of diabetics, so DM treatment should be carried out by a multidisciplinary team specializing in pharmacological aspects and physical exercise to increase the effectiveness of therapy [13]. In addition, an exercise that is prescribed and performed properly and correctly has also been shown to be safe and effective for people with cardiovascular disease [39].

Betatrophin is a newly discovered adipocytokine and is believed to play an important role in the body's metabolism [54]. Betatrophin is one of the main focuses in

obesity research and relevant research results have been published continuously [26, 63]. Several epidemiological studies have shown that there is an independent relationship between circulating betatrophin levels and obesity [64–66], while other studies reported no association [63, 67–69]. Inconsistent results are likely due to too few subject counts and also influenced by many factors [54]. Therefore, we conducted a re-study to further evaluate the relationship between circulating betatrophin levels and obesity.

Our data present that betatrophin levels have a positive relationship with age, body mass index (BMI), fat mass (FM), waist-to-hip ratio (WHR), percentage of body fat (PBF), and fasting blood glucose (FBG) (Figure 2). The results confirm previous findings that reported that betatrophin levels were positively correlated with age, BMI, FBG [26, 33, 61, 70–72], WHR [34], FM [65], and percentage of body fat (FAT %) [70]. This association is likely due to an increase in obese individuals with elevated levels of betatrophin [27, 28, 63, 64, 67]. It is as evidenced by the results of a meta-analysis conducted by Ye et al. [54] which reported that betatrophin levels were higher in obese individuals compared to those in healthy individuals. Elevated levels of betatrophin can inhibit the performance of the LPL enzymes [31, 32]. The stunted performance of LPL enzymes causes fat to be irreversible into an energy source [70], resulting in increased fat storage in adipocytes [34].

Adipose tissue, known as an important endocrine organ, has attracted a lot of attention because it has many effects on several metabolic processes, such as glucose homeostasis, lipid metabolism, inflammation, and blood pressure [54]. Betatrophin is known as angiopoietin-like protein 8 (ANGPTL8), refeeding induced fat and liver (RIFL), chromosome 19 open reading frame 80 (C19ORF80), hepatocellular carcinoma-associated protein TD26, and lipasin, which is a new adipokine secreted from adipose and liver tissue [55]. Previous studies using mouse models implied an important role of betatrophin in some metabolic-related pathways, such as lipid metabolism and energy balance [26]. Based on a meta-analysis of 9 observational studies, individuals with obesity have higher levels of betatrophin than normal-weight individuals [54]. There is emerging evidence supporting a positive association between betatrophin levels and obesity [26, 54, 65, 66]. *In vitro* studies conducted by Ren et al. [73] showed that betatrophin expression increased more than 100-fold during cell adipogenesis 3T3-L1. Its knockout in 3T3-L1 cells during adipogenesis and its knockdown lead to a decrease in adipogenesis. Furthermore, betatrophin expression increased about 8 times in white adipose tissue (WAT) in obese model mice compared

to wild mice [74]. Likewise, betatrophin mRNA is expressed very highly in humans with obesity [26]. High concentrations of betatrophin in circulation can directly increase the risk of obesity in adults [54]. Therefore, betatrophin can serve as a viable therapeutic target in the fight against obesity in adults.

There are some limitations to this study. First, the number of subjects who participated in this study was still relatively small who met the inclusion criteria and participated in the training program. Thus, a large number of samples may be needed to further analyze the effect of moderate-intensity exercise on the modulation of betatrophin levels in obese women. Second, the type of exercise intervention for this study is a single intervention (acute exercise), several possible types and intensity of exercise can affect the regulation of betatrophin. Therefore, further research is needed on the type and intensity of exercise. Third, circulating betatrophin levels can be influenced by gender and lipid profile, so further research can add to the examination of body fat profiles, such as total cholesterol (TC), triglyceride (TG), low-density lipoprotein (LDL), high-density lipoprotein (HDL), and comparing betatrophin levels between men and women.

## Conclusions

Taken together, the results of this study showed that moderate-intensity interval exercise and moderate-intensity continuous exercise performed for 30–35 min lowered betatrophin levels. However, moderate-intensity continuous exercise is more effective in lowering betatrophin levels compared to moderate-intensity interval exercise. Therefore, moderate-intensity continuous exercise can be a major modality in lowering betatrophin levels and betatrophin can serve as a promising therapeutic target for obesity in adults. Further research is needed to explore in more detail the physiological mechanisms of exercise in lowering betatrophin levels in obesity.

**Acknowledgments:** We would like to express our gratitude to the Faculty of Sport Science, State University of Malang that has provided facilities in the screening process of a prospective research subject and the Fitness Center of Health Ministry of Malang that has provided facilities well. Also, we greatly appreciate and wish to thank *Palang Merah Indonesia* (PMI) Blood Transfusion Unit (UTD) of Malang that has assisted the blood sampling and blood centrifuge processes. This includes but is not limited to all parties of Physiology Laboratory, Faculty of Medicine,

Universitas Brawijaya Malang who have helped the analysis process of betatrophin levels and all volunteers who have participated in this study.

**Research funding:** This study is supported by the Fundamental Research Program of the Ministry of Research, Technology and Higher Education, Indonesia, under Grant Number: 4/AMD/E1/KP.PTNBH/2020 and 580/UN3.14/PT/2020.

**Author contribution:** All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

**Competing interests:** Authors state no conflict of interest.

**Informed consent:** Informed consent was obtained from all individuals included in this study.

**Ethical approval:** Based on the ethical standards of the Declaration of Helsinki 1975, this study has been approved by the Health Research Ethics Commission of the Faculty of Medicine, Universitas Airlangga, Surabaya with registration number: 309/EC/KEPK/FKUA/2019.

## References

- Aladeniyi I, Adeniyi OV, Fawole O, Adeolu M, Goon DT, Ajayi AI, et al. Pattern and correlates of obesity among public service workers in Ondo State, Nigeria: a cross-sectional study. *S Afr Fam Pract* 2017;59:195–200.
- Aktar N, Qureshi NK, Ferdous HS. Obesity: a review of pathogenesis and management strategies in adult. *Delta Med Coll J* 2017;5:35–48.
- Chooi YC, Ding C, Magkos F. The epidemiology of obesity. *Metab Clin Exp* 2019;92:6–10.
- Windarti N, Hlaing SW, Kakinaka M. Obesity Kuznets curve: international evidence. *Publ Health* 2019;169:26–35.
- World Health Organization (WHO). Obesity and overweight. Geneva: WHO; 2015. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/>.
- Kibria GM. Prevalence and factors affecting underweight, overweight and obesity using Asian and World Health Organization cutoffs among adults in Nepal: analysis of the Demographic and Health Survey 2016. *Obes Res Clin Pract* 2019; 13:129–36.
- NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 2016;387:1377–96.
- Basic Health Research (Risksdas). National Report on Basic Health Research. Jakarta: Ministry of Health Indonesia; 2018. Available at: <http://www.kesmas.kemkes.go.id>.
- Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes* 2006;1:11–25.
- Masuet-Aumatell C, Ramon-Torrell JM, Banqu -Navarro M, D valos-Gamboa Mdel R, Monta o-Rodr guez SL. Prevalence of overweight and obesity in children and adolescents from Cochabamba (Bolivia); a cross-sectional study. *Nutr Hosp* 2013; 28:1884–91.



11. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* 2017;390:2627–42.
12. Guthold R, Stevens GA, Riley LM, Bull FC. Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 16 million participants. *Lancet Child Adolesc Health* 2020;4:23–35.
13. Palermi S, Iacono O, Sirico F, Modestino M, Ruosi C, Spera R, et al. The complex relationship between physical activity and diabetes: an overview. *J Basic Clin Physiol Pharmacol* 2021;1–13. <https://doi.org/10.1515/jbcp-2021-0279>.
14. Shen CC, Tseng YH, Shen MS, Lin HH. Effects of Sports massage on the physiological and mental health of college students participating in a 7-week intermittent exercises program. *Int J Environ Res Publ Health* 2021;18:5013.
15. Korczak DJ, Madigan S, Colasanto M. Children’s physical activity and depression: a meta-analysis. *Pediatrics* 2017;139:e20162266.
16. Klop B, Elte JWF, Cabezas MC. Dyslipidemia in obesity: mechanisms and potential targets. *Nutrients* 2013;4:1218–40.
17. Boden G. 45Obesity, insulin resistance and free fatty acids. *Curr Opin Endocrinol Diabetes Obes* 2011;18:139–43.
18. Midha T, Krishna V, Nath B, Kumari R, Rao YK, Pandey U, et al. Cut-off of body mass index and waist circumference to predict hypertension in Indian adults. *World J Clin Cases* 2014;2:272–8.
19. Félix-Redondo FJ, Grau M, Baena-Díez JM, Décano IR, De León AC, Guembe MJ. Prevalence of obesity and associated cardiovascular risk: the DARIOS study. *BMC Public Health* 2013;13:1–10.
20. Nimpitsch K, Konigorski S, Pischon T. Diagnosis of obesity and use of obesity biomarkers in science and clinical medicine. *Metab Clin Exp* 2019;92:61–70.
21. Hruby A, Hu FB. The epidemiology of obesity: a big picture, department of nutrition. *Pharmacoeconomics* 2015;33:673–89.
22. Fruh SM. Obesity: risk factors, complications, and strategies for sustainable long-term weight management. *J Am Assoc Nurse Pract* 2017;29:S3–14.
23. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global regional and national prevalence of overweight and obesity in children and adults 1980–2013: a systematic analysis. *Lancet* 2014;384:766–81.
24. Segula D. Complications of obesity in adults: a short review of the literature. *Malawi Med J* 2014;26:20–4.
25. Kusminski CM, Bickel PE, Scherer PE. Targeting adipose tissue in the treatment of obesity-associated diabetes. *Nat Rev Drug Discov* 2016;15:639–60.
26. Abu-Farha M, Sriraman D, Cherian P, AlKhairi I, Elkum N, Behbehani K, et al. Circulating ANGPTL8/betatrophin is increased in obesity and reduced after exercise training. *PLoS One* 2016;11: e0147367.
27. Crujeiras AB, Zulet MA, Abete I, Amil M, Carreira MC, Martínez JA, et al. Interplay of atherogenic factors, protein intake and betatrophin levels in obese–metabolic syndrome patients treated with hypocaloric diets. *Int J Obes* 2016;40:403–10.
28. Fu Z, Abou-Samra AB, Zhang R. An explanation for recent discrepancies in levels of human circulating betatrophin. *Diabetologia* 2014;57:2232–4.
29. Maurer L, Schwarz F, Fischer-Rosinsky A, Schlueter N, Brachs S, Möhlig M, et al. Renal function is independently associated with circulating betatrophin. *PLoS One* 2017;12:1–10.
30. Qu Q, Zhao D, Zhang F, Bao H, Yang Q. Serum betatrophin levels are increased and associated with insulin resistance in patients with polycystic ovary syndrome. *J Int Med Res* 2017;45:193–202.
31. Chi X, Britt EC, Shows HW, Hjelmaas AJ, Shetty SK, Cushing EM, et al. ANGPTL8 promotes the ability of ANGPTL3 to bind and inhibit lipoprotein lipase. *Mol Metabol* 2017;6:1137–49.
32. Bace E, Karpisek M, Stejskal D, Kotolova H, Ruzicka V, Hlozankova M. Betatrophin – a new insight into lipid homeostasis. *Atherosclerosis* 2017;263:e111–282.
33. Xie X, Gao T, Yang M, Chen P, Jin H, Yang L, et al. Associations of betatrophin levels with irisin in Chinese women with normal glucose tolerance. *Diabetol Metab Syndrome* 2015;7:1–8.
34. Zheng J, Liu J, Hong BS, Ke W, Huang M, Li Y. Circulating betatrophin/ANGPTL8 levels correlate with body fat distribution in individuals with normal glucose tolerance but not those with glucose disorders. *BMC Endocr Disord* 2020;20:1–9.
35. Longo M, Zatterale F, Naderi J, Parrillo L, Formisano P, Raciti GA, et al. Adipose tissue dysfunction as determinant of obesity-associated metabolic complications. *Int J Mol Sci* 2019;20:1–23.
36. Gómez-Hernández A, Beneit N, Díaz-Castroverde S, Escribano Ó. Differential role of adipose tissues in obesity and related metabolic and vascular complications. *Int J Endocrinol* 2016; 2016. <https://doi.org/10.1155/2016/1216783>.
37. Susanto H, Taufiq A, Sugiharto, Merawati D, Badu KM, Purnomo JD, et al. Moderate-intensity exercise and musical Co-treatment decreased the circulating level of betatrophin. *Int J Endocrinol* 2020; 2020:1–9.
38. Palermi S, Sacco AM, Belviso I, Romano V, Montesano P, Corrado B, et al. Guidelines for physical activity-A cross-sectional study to assess their application in the general population. Have we achieved our goal? *Int J Environ Res Publ Health* 2020;17:3980.
39. Palermi S, Bragazzi NL, Cular D, Ardigò LP, Padulo J. How chest press-based exercises can alleviate the burden of cardiovascular diseases. *Hum Mov* 2022;23. <https://doi.org/10.5114/hm.2021.106911>.
40. Enteshary M, Esfarjani F, Reisi J. Comparison of the effects of two different intensities of combined training on irisin, betatrophin, and insulin levels in women with type 2 diabetes. *Asian J Sports Med* 2019;10: e68943.
41. Rejeki PS, Pranoto A, Prasetya RE, Sugiharto. Irisin serum increasing pattern is higher at moderate-intensity continuous exercise than at moderate-intensity interval exercise in obese females. *Comp Exerc Physiol* 2021;17:475–84.
42. Dias KA, Ingul CB, Tjonna AE, Keating SE, Gomersall SR, Follstad T, et al. Effect of high-intensity interval training on fitness, fat mass and cardiometabolic biomarkers in children with obesity: a randomised controlled trial. *Sports Med* 2017;48:733–46.
43. Wewege M, van den Berg R, Ward RE, Keech A. The effects of high-intensity interval training vs. moderate-intensity continuous training on body composition in overweight and obese adults: a systematic review and meta-analysis. *Obes Rev* 2017;18:635–46.
44. Norton K, Norton L, Sadgrove D. Position statement on physical activity and exercise intensity terminology. *J Sci Med Sport* 2010; 13:496–502.
45. Nygaard H, Slettalokken G, Vegge G, Hollan I, Whist JE, Strand T, et al. Irisin in blood increases transiently after single sessions of

- intense endurance exercise and heavy strength training. *PLoS One* 2015;10:1–12.
46. Kraemer RR, Shockett P, Webb ND, Shah U, Castracane VD. A transient elevated irisin blood concentration in response to prolonged, moderate aerobic exercise in young men and women. *Horm Metab Res* 2014;46:150–4.
  47. Nedić O, Šunderić M, Miljuš G, Valdevit Z, Jakovljević V, Glibetić M, et al. Preparatory training attenuates drastic response of the insulin-like growth factor binding protein 1 at the point of maximal oxygen consumption in handball players. *J Sport Health Sci* 2017;6:372–7.
  48. Qiu S, Bosnyák E, Treff G, Steinacker JM, Nieß AM, Krüger K, et al. Acute exercise-induced irisin release in healthy adults: associations with training status and exercise mode. *Eur J Sport Sci* 2018;18:1226–33.
  49. Huh JY, Siopi A, Mougios V, Park KH, Mantzoros CS. Irisin in response to exercise in humans with and without metabolic syndrome. *J Clin Endocrinol Metab* 2015;100:E453–7.
  50. Elizondo-Montemayor L, Silva-Platas C, Torres-Quintanilla A, Rodríguez-López C, Ruiz-Esparza GU, Reyes-Mendoza E, et al. Association of irisin plasma levels with anthropometric parameters in children with underweight, normal weight, overweight, and obesity. *BioMed Res Int* 2017;2017:1–11.
  51. Daskalopoulou SS, Cooke AB, Gomez YH, Mutter AF, Filippaios A, Mesfum ET, et al. Plasma irisin levels progressively increase in response to increasing exercise workloads in young, healthy, active subjects. *Eur J Endocrinol* 2014;171:343–52.
  52. Tsuchiya Y, Ando D, Goto K, Kiuchi M, Yamakita M, Koyama K. High-intensity exercise causes greater irisin response compared with low-intensity exercise under similar energy consumption. *Tohoku J Exp Med* 2014;233:135–40.
  53. Tsuchiya Y, Ando D, Takamatsu K, Goto K. Resistance exercise induces a greater irisin response than endurance exercise. *Metab Clin Exp* 2015;64:1042–50.
  54. Ye J, Qin Y, Wang D, Yang L, Yuan G. The relationship between circulating ANGPTL8/betatrophin concentrations and adult obesity: a meta-analysis. *Dis Markers* 2019;2019:5096860.
  55. Kong FJ, Ma LL, Li G, Chen YX, Zhou JQ. Circulating betatrophin levels and gestational diabetes mellitus: a systematic review and meta-analysis. *PLoS One* 2017;12: e0169941.
  56. Zheng X, Qi Y, Bi L, Shi W, Zhang Y, Zhao D, et al. Effects of exercise on blood glucose and glycemic variability in type 2 diabetic patients with dawn phenomenon. *BioMed Res Int* 2020;2020:6408724.
  57. van Dijk JW, Venema M, van Mechelen W, Stehouwer CD, Hartgens F, van Loon LJ. Effect of moderate-intensity exercise versus activities of daily living on 24-hour blood glucose homeostasis in male patients with type 2 diabetes. *Diabetes Care* 2013;36:3448–53.
  58. Fu Z, Berhane F, Fite A, Seyoum B, Abou-Samra AB, Zhang R. Elevated circulating lipasin/betatrophin in human type 2 diabetes and obesity. *Sci Rep* 2014;4:1–5.
  59. Zhang R. Lipasin, a novel nutritionally-regulated liver-enriched factor that regulates serum triglyceride levels. *Biochem Biophys Res Commun* 2012;424:786–92.
  60. Gusarova V, Alexa CA, Na E, Stevis PE, Xin Y, Bonner-Weir S, et al. ANGPTL8/betatrophin does not control pancreatic beta cell expansion. *Cell* 2014;159:691–6.
  61. Wang H, Lai Y, Han C, Liu A, Fan C, Wang H, et al. The effects of serum ANGPTL8/betatrophin on the risk of developing the metabolic syndrome - a prospective study. *Sci Rep* 2016;6:28431.
  62. Hall JE. *Guyton and Hall Textbook of Medical Physiology*, 13th ed. United States of America: Elsevier; 2016.
  63. Guo K, Lu J, Yu H, Zhao F, Pan P, Zhang L, et al. Serum betatrophin concentrations are significantly increased in overweight but not in obese or type 2 diabetic individuals. *Obesity* 2015;23:793–7.
  64. Gómez-Ambrosi J, Pascual E, Catalán V, Rodríguez A, Ramírez B, Silva C, et al. Circulating betatrophin concentrations are decreased in human obesity and type 2 diabetes. *J Clin Endocrinol Metab* 2014;99:E2004–9.
  65. Barja-Fernández S, Folgueira C, Seoane LM, Casanueva FF, Dieguez C, Castela C, et al. Circulating betatrophin levels are increased in anorexia and decreased in morbidly obese women. *J Clin Endocrinol Metab* 2015;100:E1188–96.
  66. Wu S, Gao H, Ma Y, Fu L, Zhang C, Luo X. Characterisation of betatrophin concentrations in childhood and adolescent obesity and insulin resistance. *Pediatr Diabetes* 2016;17:53–60.
  67. Fenzl A, Itariu BK, Kosi L, Fritzer-Szekeres M, Kautzky-Willer A, Stulnig TM, et al. Circulating betatrophin correlates with atherogenic lipid profiles but not with glucose and insulin levels in insulin-resistant individuals. *Diabetologia* 2014;57:1204–8.
  68. Roth CL, Elfers C, Lass N, Reinehr T. Betatrophin: no relation to glucose metabolism or weight status in obese children before and after lifestyle intervention. *Pediatr Diabetes* 2017;18:485–91.
  69. Battal F, Türkön H, Aylanç N, Aylanç H, Yıldırım Ş, Kaymaz N, et al. Investigation of blood betatrophin levels in obese children with non-alcoholic fatty liver disease. *Pediatr Gastroenterol, Hepatol Nutr* 2018;21:111–7.
  70. Pu D, Li L, Yin J, Liu R, Yang G, Liao Y, et al. Circulating ANGPTL8 is associated with the presence of metabolic syndrome and insulin resistance in polycystic ovary syndrome young women. *Mediat Inflamm* 2019;2019:6321427.
  71. Abu-Farha M, Abubaker J, Noronha F, Al-Khairi I, Cherian P, Alarouj M, et al. Lack of associations between betatrophin/ANGPTL8 level and C-peptide in type 2 diabetic subjects. *Cardiovasc Diabetol* 2015;14:112.
  72. Maurer L, Brachs S, Decker AM, Brachs M, Leupelt V, von Schwartzberg RJ, et al. Weight loss partially restores glucose-driven betatrophin response in humans. *J Clin Endocrinol Metab* 2016;101:4014–20.
  73. Ren G, Kim JY, Smas CM. Identification of RIFL, a novel adipocyte-enriched insulin target gene with a role in lipid metabolism. *Am J Physiol Endocrinol Metab* 2012;303:E334–51.
  74. Abu-Farha M, Abubaker J, Tuomilehto J. ANGPTL8 (betatrophin) role in diabetes and metabolic diseases. *Diabetes/Metab Res Rev* 2017;33:e2919.