

Negative Correlation between Serum Brain-derived Neurotrophic Factor Levels and Obesity Predictor Markers and Inflammation Levels in Females with Obesity

by Slamet Raharjo

Submission date: 22-Jun-2022 12:58PM (UTC+0800)

Submission ID: 1861131630

File name: ain-derived_Neurotrophic_Factor_Levels_and_Obesity_Predictor.pdf (495.51K)

Word count: 4366

Character count: 23894



Negative Correlation between Serum Brain-derived Neurotrophic Factor Levels and Obesity Predictor Markers and Inflammation Levels in Females with Obesity

Slamet Raharjo¹, Adi Pranoto², Purwo Sri Rejeki³, A. S. M. Harisman¹, Yualita Putri Pamungkas⁴, Olivia Andiana¹

¹Department of Sports Science, Faculty of Sports Science, State University of Malang, Malang, Indonesia; ²Department of Medical Science, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; ³Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia; ⁴Department of Public Health, Faculty of Sports Science, State University of Malang, Malang, Indonesia

Abstract

Edited by: Branislav Filipović
Citation: Raharjo S, Pranoto A, Rejeki PS, Harisman ASM, Pamungkas YP, Andiana O. Negative Correlation between Serum Brain-Derived Neurotrophic Factor Levels and Obesity Predictor Markers and Inflammation Levels in Females with Obesity. Open Access Maced J Med Sci. 2021 Sep 30; 9(B):1021-1026. https://doi.org/10.3889/oamjms.2021.6840

Keywords: Obesity; Brain-derived neurotrophic factor; Body mass index; Waist hip ratio; Interleukin-6; Tumor necrosis factor- α

Correspondence: Purwo Sri Rejeki, Department of Physiology and Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. E-mail: purwo-s-r@fkk.unair.ac.id

Received: 13-Jul-2021
Revised: 19-Aug-2021
Accepted: 20-Sep-2021

Copyright: © 2021 Slamet Raharjo, Adi Pranoto, Purwo Sri Rejeki, A. S. M. Harisman, Yualita Putri Pamungkas, Olivia Andiana

Funding: The authors used their personal cost as the resource of research funding.
Competing interests: The authors declare that they have no competing interests.

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: Obesity has been widely associated with structural and functional changes in brain, whereas inflammation is one of the potential mechanisms involved in these changes.

AIM: This study aims to prove the relationship between serum brain-derived neurotrophic factor (BDNF) levels and obesity predictor markers (body mass index [BMI] and waist to hip ratio [WHR]) and inflammation (interleukin-6 [IL-6] and tumor necrosis factor-alpha [TNF- α]) levels of in females with obesity.

METHODS: This study used a cross-sectional study method using 33 female with obesity aged 19–23 years, BMI >27.5 kg/m², normal blood pressure, normal resting heart rate, normal hemoglobin, and fasting blood glucose \leq 100 mg/dL. The examination of serum BDNF, IL-6, and TNF- α levels using the Enzyme-Linked Immunosorbent Assay method. The data were analyzed using Pearson product-moment test with a significant levels $p < 0.05$.

RESULTS: The results indicated that there is a negative correlation between serum BDNF levels and BMI ($r = -0.759$; $p < 0.001$), WHR ($r = -0.675$; $p < 0.001$), IL-6 levels ($r = -0.530$; $p < 0.001$), and TNF- α levels ($r = -0.561$; $p < 0.001$).

CONCLUSION: Based on the results of the study, there is a negative correlation between serum BDNF levels and BMI, WHR, and inflammation (IL-6 and TNF- α) levels in females with obesity. Further studies are needed to confirm the present findings.

Introduction

The worldwide prevalence of obesity nearly tripled between 1975 and 2016 [1]. Based on data from the WHO [1] reports, around 13% of the world's population aged 18 years in 2016 were categorized as obese. Obesity is regarded as a condition of excessive fat accumulation, and thus it can affect individual health [2]. According to Wang *et al.* [3], individuals with obesity have a lower life expectancy and have a higher risk of premature death than those with normal weight [4]. Today's obesity is also a major contributor to public health problems since it increases the risk factors for chronic non-communicable diseases, such as cardiovascular disease, non-alcoholic fatty liver disease, diabetes mellitus type II, and several types of cancer [5], [6].

A study conducted by Nota *et al.* [7] reported that obesity has also been associated with structural

and functional changes in the brain. Potential mechanisms in these changes include inflammation and vascular and metabolic changes [8]. Recently, obesity has also been associated with a low-grade inflammatory state [9], [10], [11]. This condition triggers systemic inflammation which becomes an important factor for microglial activation and neuroinflammation in neurodegeneration [12]. This is caused by excess macronutrients in adipose tissue stimulating the release of inflammatory mediators, such as tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6), and decreasing the production of anti-inflammatory agents, such as adiponectin [13]. The results of a study conducted by Calabrese *et al.* [13] indicated that the production of pro-inflammatory cytokines significantly reduces the expression of brain derived neurotrophic factor (BDNF). The decrease in BDNF expression has been widely associated with cognitive dysfunction and dementia [14], [15], [16], [17], [18], [19], [20], [21], [22].

BDNF dysregulation is not only associated with the pathogenesis of neurodegenerative and psychiatric disorders, but also inflammation, metabolism, and cardiovascular disease [23].

BDNF is the most expressed neurotrophin in the brain and plays a key role in learning and memory, especially in cognitive function [23]. A current study have revealed that activation of tropomyosin receptor kinase B (TrkB) by BDNF can stimulate the signaling pathways of mitogen-activated protein kinase/ERK, phosphatidylinositol 3-kinase/Akt, and phospholipase C-gamma [24]. Intensive investigations into this signaling had been performed since BDNF/TrkB-mediated intracellular signaling is involved in many neuronal aspects [25]. The BDNF regulates brain function and acts as a neuroprotective through the mechanism of increasing neurogenesis, neuronal survival, axonal growth, dendritic growth, synaptic plasticity, neuronal development, and maintenance in the central nervous system neurons [24]. The BDNF is also associated with the pathophysiology of obesity and metabolic syndrome [26]. However, the relationship of BDNF and obesity is still not clearly revealed. The results of the meta-analysis conducted by Sandrini *et al.* [27] showed that obesity is not associated with a decrease in circulating BDNF levels. However, the study conducted by of Goltz *et al.* [28] reported that BDNF levels is correlated with central obesity. In addition, the relationship between BDNF levels and inflammation levels is still controversial. A study conducted by Patas *et al.* [29] reported that BDNF levels is correlated with IL-6 levels and has no correlation with TNF- α levels. However, the results of a study conducted by Zhang *et al.* [30] showed that the BDNF levels are not correlated with IL-6 and TNF- α levels. On this basis, the objective of this study is to analyze the relationship between serum BDNF levels and predictor markers of obesity (body mass index (BMI) and waist to hip ratio [WHR]) and inflammation levels (IL-6 and TNF- α) in adolescent girls with obesity. The authors hypothesized that there is a significant correlation between serum BDNF levels and predictor markers of obesity and inflammation levels in adolescent girls with obesity.

Methods

Research design

The authors used a cross-sectional study method by employing 33 female with obesity aged 19–23 years with BMI >27.5 kg/m², normal blood pressure, normal resting heart rate (RHR), hemoglobin (Hb) normal, and fasting blood glucose (FBG) ≤ 100 mg/dL. All respondents received information orally and in writing about the research. They were

required to fill out and sign an informed consent before participating in the study.

Ethical clearance

All research procedures have been approved by the Health Research Ethics Commission, Faculty of Medicine, Universitas Brawijaya Malang number 81/EC/KEPK-S1/04/2020.

Analysis of obesity predictor markers

The height of the respondents was measured using a stadiometer (SECA, Chino, CA, USA), while the body weight was measured using an electronic scale (Tech 05[®], China). The BMI was measured by calculating body weight (kg) divided by height in m² [31]. The blood pressure was measured using an OMRON digital blood pressure meter (OMRON Model HEM-7130 L, Omron Co., Osaka, Japan). The waist circumference (WC) was measured by circling the anthropometric tape measure on the halfway between the lower rib and iliac bones parallel to the midaxillary straight line. The hip circumference (HC) was measured by circling the anthropometric tape measure right on the great trochanters, while the waist to WHR measurement was performed by calculating its WC divided by HC.

Analysis of BDNF and inflammatory levels

The blood was taken from the cubital vein amounted to 4 ml after undergoing overnight fasting for 12 h. The respondents were asked to do sleeping position while the blood was taken. The collected blood was centrifuged for 15 min at 3000 rpm. The serum was separated and stored at -30°C for obtaining an analysis of BDNF, IL-6, and TNF- α levels in the next day. The blood sampling was performed at 07.00–08.00 a.m. (Jakarta time). The measurement of serum BDNF levels was performed using the Enzyme-Linked Immunosorbent Assay (ELISA) kit method (Catalog No. E-EL-H0010; Elabscience, Inc., China) with a standard curve range of 31.25–2000 pg/mL and the sensitivity levels of BDNF in the kit 18.75 pg/mL. The IL-6 levels were measured using ELISA kit method (Catalog No. E-EL-H0102; Elabscience, Inc., China) with a standard curve range of 7.81–500 pg/mL and the sensitivity levels of IL-6 in the kit was 4.69 pg/mL. The TNF- α levels were measured using the ELISA kit method (Catalog No. E-EL-H0109; Elabscience, Inc., China) with a standard curve range of 7.81–500 pg/mL and sensitivity 4.69 pg/mL. Blood taking for examination of FBG and Hb levels was carried out in the capillaries located at the tip of the middle finger. The FBG examination was performed using Accu-Chek Performance (Roche, Mannheim, Germany) with the levels of mg/dL, while the Hb examination was performed using Easy Touch GCHb (Easy Touch, Hsinchu, Taiwan) with the levels of g/dL.

Statistical analysis

The statistical analysis technique was performed using Statistical Package for the Social Science (SPSS) software version 21 (SPSS Inc., Chicago, IL, USA). The normality test was made using Kolmogorov–Smirnov test. Data with normal distribution were tested using Pearson product-moment with significant levels of $p < 0.05$. All data were then displayed with the mean \pm standard error of the mean.

Results

The analysis on the characteristics of respondents showed several results, that is, the mean age (20.76 ± 0.19 years), anthropometric parameters of height (1.57 ± 0.01 m), body weight (73.02 ± 1.48 kg), BMI (29.55 ± 0.52 kg/m²), WC (87.61 ± 2.03 cm), HC (107.58 ± 1.27 cm), WHR (0.80 ± 0.01), SBP (113.82 ± 0.68 mmHg), DBP (76.06 ± 0.85 mmHg), RHR (73.18 ± 1.08 bpm), FBG (88.00 ± 1.04 mg/dL), and Hb (14.65 ± 0.14 g/dL). The mean serum BDNF levels showed (341.33 ± 18.97 pg/mL) and the mean inflammation levels showed IL-6 (10.03 ± 0.79 pg/mL) and TNF- α (17.91 ± 0.40 pg/mL). The relationship between serum BDNF levels and BMI, WHR, IL-6, TNF- α is presented in Figure 1.

Based on Figure 1, the results of the Pearson product-moment linear correlation parametric analysis indicates that there is a negative correlation between serum BDNF levels and BMI ($r = -0.759$; $p < 0.001$), WHR ($r = -0.675$; $p < 0.001$), IL-6 levels ($r = -0.530$; $p < 0.001$), and TNF- α levels ($r = -0.561$; $p < 0.001$).




Figure 1: Negative correlation between serum brain-derived neurotrophic factor levels and body mass index, waist to hip ratio, interleukin-6, and tumor necrosis factor- α . *Significant with $p < 0.05$ by Pearson's product-moment correlation test

Discussion

The results of the analysis showed that there was a negative correlation between serum BDNF levels and BMI and WHR (Figure 1). These results are in line with the results of a study conducted by Si *et al.* [32] that there is a negative correlation between BMI and BDNF levels in male and female students. A study conducted by Yang *et al.* [33] also reported that BDNF levels have a significant correlation with BMI in female. Furthermore, a study conducted by Jung *et al.* [34] confirmed that BDNF levels have a correlation with BMI. Zhang *et al.* [35], through his study, also reported that BDNF levels are associated with BMI. Goltz *et al.* [28] in his research also reported that BDNF levels correlated with WHR in obese individuals. These findings confirm that there is a relationship between BDNF levels and predictor markers of obesity. This relationship may be underlined by a significant reduction in BDNF levels in individuals with obesity [36]. The study conducted by El-Alamey *et al.* [26] proved that children with obesity have a significant decrease in serum BDNF levels. Besides, the study conducted by El-Gharbawy *et al.* [37] also reported that serum BDNF is lower in children and adolescents with obesity than children and adolescents with normal or ideal weight. The decrease in BDNF levels can inhibit the activation of tropomyosin kinase B (TrkB) receptors and primary BDNF receptors, resulting in hyperphagia and obesity [38]. The inhibition in activation between BDNF and TrkB triggers a series of reactions that inhibit an eager to eat and weight gain [38]. This shows that BDNF has a central role in inhibiting appetite, thereby increasing negative energy balance [36]. Being overweight and obese in early adulthood are associated with a higher risk of cardiovascular disease in future life, and it also has a negative impact on academic performance and executive control which is characterized by accelerated cognitive decline and brain atrophy in later years [32].

Based on the results of statistical analysis, there is a negative correlation between serum BDNF levels and inflammation levels (IL-6 and TNF- α) in female with obesity (Figure 1). These results are in line with the results of study conducted by Patas *et al.* [29] that BDNF levels have a correlation with IL-6 levels, but the BDNF levels do not have a relationship with TNF- α levels in patients with major depressive disorder (MDD). These results are also different from that of the study conducted by Zhang *et al.* [30] that BDNF levels do not have correlation with IL-6 and TNF- α levels in patients with chronic schizophrenia. The difference may be caused by the fact that this study used female with obesity as the respondents, whereas the previous studies used chronic schizophrenia patients and MDD patients. It has recently been reported that obesity is associated with a low-grade inflammatory state [9], [10], [27]. Such a condition triggers systemic inflammation which

is an important factor for microglial activation and neuroinflammation in neurodegeneration [11]. This may happen due to excess macronutrients in adipose tissue that stimulate the release of pro-inflammatory mediators, such as TNF- α and IL-6 and reduce the production of anti-inflammatory agents, such as adiponectin [12]. The results of a study conducted by Calabrese *et al.* [13] revealed that an increase in production of pro-inflammatory cytokines significantly reduces the BDNF levels.

Based on the results of a review conducted by Sui and Pasco [39], obesity also has a relationship with a decrease in cognitive function, neuronal plasticity, brain volume, and brain structure. Besides, the changes in BDNF levels are affected by genetic factors, but other evidence also showed that BDNF levels are associated with the changes in lifestyle and dietary habit [39]. An excessive calorie intake as compared to calorie expenditure in the body causes an increase in body weight which is characterized by an increase in white adipose tissue and can also lead to a decrease in the BDNF levels [27]. Having proven by the results of a study by Lee *et al.* [40], the baseline BDNF levels in individuals with obesity are lower than that of in the individuals without obesity. Furthermore, a study conducted by Si *et al.* [32] reported that was associated with the decrease in BDNF levels in adolescents aged 18–20 years. Obesity can reduce serum neurotrophic factor levels and cause blood-brain barrier (BBB) dysfunction [41]. This leads to peripheral pro-inflammatory cytokines to pass through the BBB, causing central inflammation that occurs in the brain and causes damage to the nervous system [42].

Conclusion

Based on the results of the study, there is a negative correlation between serum BDNF levels and BMI, WHR, and inflammation levels (IL-6 and TNF- α) in female with obesity. The authors suggest that future researchers can reveal the physiological mechanism of correlation between serum BDNF and BMI, WHR, and inflammation levels (IL-6 and TNF- α) in female with obesity more detail. Further studies are needed to confirm the present findings.

References

- World Health Organization. Obesity and overweight. Geneva: World Health Organization; 2020. Available from: <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>. <https://doi.org/10.1787/a47d0cd2-en> [Last accessed on 2021 May 15].
- Maria P, Evagelia S. Obesity disease. *Health Sci J*. 2009;3(3):132-8.
- Wang C, Chan JS, Ren L, Yan JH. Obesity reduces cognitive and motor functions across the lifespan. *Neural Plasticity*. 2016;2016:2473081. <https://doi.org/10.1155/2016/2473081> PMID:26881095
- Peeters A, Barendregt JJ, Willekens F, Mackenbach JP, Al Mamun A, Bonneux L, *et al.* Obesity in adulthood and its consequences for life expectancy: a life-table analysis. *Ann Intern Med*. 2003;138(1):24-32. <https://doi.org/10.7326/0003-4819-138-1-200301070-00008>.
- Goossens GH. The metabolic phenotype in obesity: Fat mass, body fat distribution, and adipose tissue function. *Obes Facts*. 2017;10(3):207-15. <https://doi.org/10.1159/000471488>.
- Kopelman PG. Obesity as a medical problem. *Nature*. 2000;404(6778):635-43. <https://doi.org/10.1038/35007508> PMID:10766250
- Nota M, Vreeken D, Wiesmann M, Aarts EO, Hazebroek EJ, Kiliaan AJ. Obesity affects brain structure and function- rescue by bariatric surgery? *Neurosci Biobehav Rev*. 2020;108:646-57. <https://doi.org/10.1016/j.neubiorev.2019.11.025> PMID:31794778
- Nguyen JC, Killcross AS, Jenkins TA. Obesity and cognitive decline: Role of inflammation and vascular changes. *Front Neurosci*. 2014;8:375. <https://doi.org/10.3389/fnins.2014.00375> PMID:25477778
- Donath MY, Shoelson SE. Type 2 diabetes as an inflammatory disease. *Nat Rev Immunology*. 2011;11(2):98-107. <https://doi.org/10.1038/nri2925> PMID:21233852
- Monteiro R, Azevedo I. Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm*. 2010;2010:289645. <https://doi.org/10.1155/2010/289645> PMID:20706689
- Ugalde-Muñiz P, Fetter-Pruneda I, Navarro L, García E, Chavarria A. Chronic systemic inflammation exacerbates neurotoxicity in a Parkinson's disease model. *Oxid Med Cell Longev*. 2020;2020:4807179. <https://doi.org/10.1155/2020/4807179> PMID:32015787
- Ellulu MS, Patimah I, Khaza'ai H, Rahmat A, Abed Y. Obesity and inflammation: The linking mechanism and the complications. *Arch Med Sci*. 2017;13(4):851-63. <https://doi.org/10.5114/aoms.2016.58928> PMID:28721154
- Calabrese F, Rossetti AC, Racagni G, Gass P, Riva MA, Molteni R. Brain-derived neurotrophic factor: A bridge between inflammation and neuroplasticity. *Front Cell Neurosci*. 2014;8:430. <https://doi.org/10.3389/fncel.2014.00430> PMID:25565964
- Pérez LM, Pareja-Galeano H, Sanchis-Gomar F, Emanuele E, Lucia A, Gálvez BG. Adipaging: Ageing and obesity share biological hallmarks related to a dysfunctional adipose tissue. *J Physiol*. 2016;594(12):3187-207. <https://doi.org/10.1113/JP271691> PMID:26926488
- Weinstein G, Beiser AS, Choi SH, Preis SR, Chen TC, Vorges D, *et al.* Serum brain-derived neurotrophic factor and the risk for dementia: The Framingham heart study. *JAMA Neurol*. 2014;71(1):55-61. <https://doi.org/10.1001/jamaneurol.2013.4781> PMID:24276217
- Autry AE, Monteggia LM. Brain-derived neurotrophic factor and neuropsychiatric disorders. *Pharmacol Rev*. 2012;64(2):238-58.

- <https://doi.org/10.1124/pr.111.005108>
PMid:22407616
17. Cunha C, Brambilla R, Thomas KL. A simple role for BDNF in learning and memory?. *Front Mol Neurosci.* 2010;3:1. <https://doi.org/10.3389/fnro.2010.001.2010>
PMid:20162032
 18. Komulainen P, Pedersen M, Hänninen T, Bruunsgaard H, Lakka TA, Kivipelto M, *et al.* BDNF is a novel marker of cognitive function in ageing female: The DR's EXTRA study. *Neurobiol Learn Memory.* 2008;90(4):596-603. <https://doi.org/10.1016/j.nlm.2008.07.014>
PMid:18707012
 19. Binder DK, Scharfman HE. Brain-derived neurotrophic factor. *Growth Factors (Chur, Switzerland).* 2004;22(3):123-31. <https://doi.org/10.1080/08977190410001723308>
PMid:15518235
 20. Yamada K, Mizuno M, Nabeshima T. Role for brain-derived neurotrophic factor in learning and memory. *Life Sci.* 2002;70(7):735-44. [https://doi.org/10.1016/s0024-3205\(01\)01461-8](https://doi.org/10.1016/s0024-3205(01)01461-8)
PMid:11833737
 21. Holsinger RM, Schnarr J, Henry P, Castelo VT, Fahnstock M. Quantitation of BDNF mRNA in human parietal cortex by competitive reverse transcription-polymerase chain reaction: decreased levels in Alzheimer's disease. *Brain research.* *Mol Brain Res.* 2000;76(2):347-54. [https://doi.org/10.1016/s0169-328x\(00\)00023-1](https://doi.org/10.1016/s0169-328x(00)00023-1)
PMid:10762711
 22. Phillips HS, Hains JM, Armanini M, Laramée GR, Johnson SA, Winslow JW. BDNF mRNA is decreased in the hippocampus of individuals with Alzheimer's disease. *Neuron.* 1991;7(5):695-702. [https://doi.org/10.1016/0896-6273\(91\)90273-3](https://doi.org/10.1016/0896-6273(91)90273-3)
PMid:1742020
 23. Ieraci A, Beggiano S, Ferraro L, Barbieri SS, Popoli M. Kynurenine pathway is altered in BDNF Val66Met knock-in mice: Effect of physical exercise. *Brain Behav Immunity.* 2020;89:440-50. <https://doi.org/10.1016/j.bbi.2020.07.031>
 24. Numakawa T, Odaka H, Adachi N. Actions of brain-derived neurotrophin factor in the neurogenesis and neuronal function, and its involvement in the pathophysiology of brain diseases. *Int J Mol Sci.* 2018;19(11):3650. <https://doi.org/10.3390/ijms19113650>
PMid:30463271
 25. Konishi K, Cherkerzian S, Aroner S, Jacobs EG, Rentz DM, Remington A, *et al.* Impact of BDNF and sex on maintaining intact memory function in early midlife. *Neurobiol Aging.* 2020;88:137-49. <https://doi.org/10.1016/j.neurobiolaging.2019.12.014>
 26. El-Alameey IR, Ahmed HH, Abushady MM. Role of lifestyle intervention program in regulating brain derived neurotrophic factor in obese children with metabolic syndrome components. *Biomed Pharmacol J.* 2019;12(3):1317-28.
 27. Sandrini L, Di Minno A, Amadio P, Ieraci A, Tremoli E, Barbieri SS. Association between obesity and circulating brain-derived neurotrophic factor (BDNF) levels: Systematic review of literature and meta-analysis. *Int J Mol Sci.* 2018;19(8):2281. <https://doi.org/10.3390/ijms19082281>
PMid:30081509
 28. Goltz A, Janowitz D, Hannemann A, Nauck M, Hoffmann J, Seyfart T, *et al.* Association of brain-derived neurotrophic factor and Vitamin D with depression and obesity: A population-based study. *Neuropsychobiology.* 2017;76(4):171-81. <https://doi.org/10.1159/000489864>
PMid:29920493
 29. Patas K, Penninx BW, Bus BA, Vogelzangs N, Molendijk ML, Elzinga BM, *et al.* Association between serum brain-derived neurotrophic factor and plasma interleukin-6 in major depressive disorder with melancholic features. *Brain Behav Immunity.* 2014;36:71-9. <https://doi.org/10.1016/j.bbi.2013.10.007>
PMid:24140302
 30. Zhang XY, Tan YL, Chen DC, Tan SP, Yang FD, Wu HE, *et al.* Interaction of BDNF with cytokines in chronic schizophrenia. *Brain Behav Immunity.* 2016;51:169-75. <https://doi.org/10.1016/j.bbi.2015.09.014>
 31. Nimptsch K, Konigorski S, Pischon T. Diagnosis of obesity and use of obesity biomarkers in science and clinical medicine. *Metabolism.* 2019;92:61-70. <https://doi.org/10.1016/j.metabol.2018.12.006>
PMid:30586573
 32. Si J, Zhang H, Zhu L, Chen A. The relationship between overweight/obesity and executive control in college students: The mediating effect of BDNF and 5-HT. *Life (Basel, Switzerland).* 2021;11(4):313. <https://doi.org/10.3390/life11040313>
PMid:33916706
 33. Yang F, Wang K, Du X, Deng H, Wu HE, Yin G, *et al.* Sex difference in the association of body mass index and BDNF levels in Chinese patients with chronic schizophrenia. *Psychopharmacology.* 2019;236(2):753-62. <https://doi.org/10.1007/s00213-018-5107-1>
 34. Jung SH, Kim J, Davis JM, Blair SN, Cho HC. Association among basal serum BDNF, cardiorespiratory fitness and cardiovascular disease risk factors in untrained healthy Korean men. *Eur J Appl Physiol.* 2011;111(2):303-11. <https://doi.org/10.1007/s00421-010-1658-5>
PMid:20878177
 35. Zhang XY, Zhou DF, Wu GY, Cao LY, Tan YL, Haile CN, *et al.* BDNF levels and genotype are associated with antipsychotic-induced weight gain in patients with chronic schizophrenia. *Neuropsychopharmacology.* 2008;33(9):2200-5. <https://doi.org/10.1038/sj.npp.1301619>
PMid:17987059
 36. Alomari MA, Khabour OF, Alawneh K, Alzoubi KH, Maikano AB. The importance of physical fitness for the relationship of BDNF with obesity measures in young normal-weight adults. *Heliyon.* 2020;6(3):e03490. <https://doi.org/10.1016/j.heliyon.2020.e03490>
PMid:32154423
 37. El-Gharbawy AH, Adler-Wailles DC, Mirch MC, Theim KR, Ranzenhofer L, Tanofsky-Kraff M, *et al.* Serum brain-derived neurotrophic factor concentrations in lean and overweight children and adolescents. *J Clin Endocrinol Metab.* 2006;91(9):3548-52. <https://doi.org/10.1210/jc.2006-0658>
PMid:16787984
 38. Xu B, Goulding EH, Zang K, Cepoi D, Cone RD, Jones KR, *et al.* Brain-derived neurotrophic factor regulates energy balance downstream of melanocortin-4 receptor. *Nat Neurosci.* 2003;6(7):736-42. <https://doi.org/10.1038/nn1073>
PMid:12796784
 39. Sui SX, Pasco JA. Obesity and brain function: The brain-body crosstalk. *Medicina (Kaunas, Lithuania).* 2020;56(10):499. <https://doi.org/10.3390/medicina56100499>
PMid:32987813
 40. Lee SS, Yoo JH, Kang S, Woo JH, Shin KO, Kim KB, *et al.* The effects of 12 weeks regular aerobic exercise on brain-derived neurotrophic factor and inflammatory factors in juvenile obesity and Type 2 diabetes mellitus. *J Phys Ther Sci.* 2014;26(8):1199-204. <https://doi.org/10.1589/jpts.26.1199>
PMid:25202180
 41. Roh HT, So WY. The effects of aerobic exercise training on

oxidant-antioxidant balance, neurotrophic factor levels, and blood-brain barrier function in obese and non-obese men. *J Sport Health Sci.* 2017;6(4):447-53. <https://doi.org/10.1016/j.jshs.2016.07.006>
PMid:30356625

42. Pugazhenti S, Qin L, Reddy PH. Common neurodegenerative pathways in obesity, diabetes, and Alzheimer's disease. *Biochim BiophysActa. Molecular basis of disease*, 2017;1863(5):1037-45. <https://doi.org/10.1016/j.bbadis.2016.04.017>
PMid:27156888

Negative Correlation between Serum Brain-derived Neurotrophic Factor Levels and Obesity Predictor Markers and Inflammation Levels in Females with Obesity

ORIGINALITY REPORT

20%

SIMILARITY INDEX

15%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Yazgaldi Nazari, Masoud Nikbakht, Abdolhamid Habibi, Saeed Shakeryan. "Acute and Chronic Effects of Combined Training on Brain-Derived Neurotrophic Factor Levels and Its Association with Anthropometric Variables in Overweight Men", *Annals of Military and Health Sciences Research*, 2016
Publication 1%
- 2 www.mdpi.com
Internet Source 1%
- 3 Yasmin Karimah Ikhsan, Soebagijo Adi Soelistijo, Johanes Nugroho Eko Putranto. "Profile of cardiovascular disease risk in type 2 diabetes mellitus patients receiving statin therapy: A cross-sectional study", *Annals of Medicine and Surgery*, 2022
Publication 1%
- 4 Masatoshi Kudo. "Combination Immunotherapy with Anti-PD-1/PD-L1 1%

Antibody plus Anti-VEGF Antibody May Promote Cytotoxic T Lymphocyte Infiltration in Hepatocellular Carcinoma, Including in the Noninflamed Subclass", Liver Cancer, 2022

Publication

5

Ritter, Cristiane, Aline S Miranda, Vinícius Giombelli, Cristiane D Tomasi, Clarissa M Comim, Antonio Teixeira, João Quevedo, and Felipe Dal-Pizzol. "Brain-derived neurotrophic factor plasma levels are associated with mortality in critically ill patients even in the absence of brain injury", Critical Care, 2012.

Publication

1 %

6

Wenlei Jiang. "Direct evidence for central proinflammatory mechanisms in rats with experimental acute liver failure: protective effect of hypothermia", Journal of Cerebral Blood Flow & Metabolism, 03/04/2009

Publication

1 %

7

Yu-Hui Liu, Shu-Sheng Jiao, Ye-Ran Wang, Xian-Le Bu et al. "Associations Between ApoE ϵ 4 Carrier Status and Serum BDNF Levels—New Insights into the Molecular Mechanism of ApoE ϵ 4 Actions in Alzheimer's Disease", Molecular Neurobiology, 2014

Publication

1 %

8

Novi Silvia Hardiany, Yohana Yohana, Septelia Inawati Wanandi. "TNFR, TRAF2, NF- κ B mRNA

1 %

Levels of Glioblastoma Multiforme Cells Treated by Conditioned Medium of Umbilical Cord-derived Mesenchymal Stem Cells", The Indonesian Biomedical Journal, 2019

Publication

9

Sajjad Rezaei, Mahdieh Molanouri Shamsi, Mehdi Mahdavi, Azadeh Jamali et al.

"Endurance exercise training decreased serum levels of surfactant protein D and improved aerobic fitness of obese women with type-2 diabetes", Diabetology & Metabolic Syndrome, 2017

Publication

1 %

10

Boby Pratama Putra, Felix Nugraha Putra.

"MO494: The Prospective Roles of Urinary Dickkopf-3 for Detecting Renal Function Deterioration and Monitoring Chronic Kidney Disease Progression: Systematic Review of Cohort Studies", Nephrology Dialysis Transplantation, 2022

Publication

1 %

11

Giselli Scaini, Tássia Tonon, Carolina F. Moura de Souza, Patricia F. Schuk et al. "Serum Markers of Neurodegeneration in Maple Syrup Urine Disease", Molecular Neurobiology, 2016

Publication

<1 %

12

[Research-Repository.griffith.edu.au](https://research-repository.griffith.edu.au)

Internet Source

<1 %

13

Sahil Malhotra, Shweta Dumoga, Parul Sirohi, Neetu Singh. "Red Blood Cells-Derived Vesicles for Delivery of Lipophilic Drug Camptothecin", ACS Applied Materials & Interfaces, 2019

Publication

<1 %

14

Yinghui Hu. "BDNF and the diseased nervous system: a delicate balance between adaptive and pathological processes of gene regulation", Journal of Neurochemistry, 2/28/2008

Publication

<1 %

15

labanimres.biomedcentral.com

Internet Source

<1 %

16

medicopublication.com

Internet Source

<1 %

17

research.birmingham.ac.uk

Internet Source

<1 %

18

singaporeanjbem.com

Internet Source

<1 %

19

www.ssph-journal.org

Internet Source

<1 %

20

cdn.easo.org

Internet Source

<1 %

21

doi.org
Internet Source

<1 %

22

dr.library.brocku.ca
Internet Source

<1 %

23

pureadmin.qub.ac.uk
Internet Source

<1 %

24

www.ijmr.org.in
Internet Source

<1 %

25

Despina A. Tata, Evgenia Dandi, Evangelia Spandou. "Expression of synaptophysin and BDNF in the medial prefrontal cortex following early life stress and neonatal hypoxia - ischemia", *Developmental Psychobiology*, 2020
Publication

<1 %

26

Fernández-Sánchez, Alba, Eduardo Madrigal-Santillán, Mirandeli Bautista, Jaime Esquivel-Soto, Ángel Morales-González, Cesar Esquivel-Chirino, Irene Durante-Montiel, Graciela Sánchez-Rivera, Carmen Valadez-Vega, and José A. Morales-González. "Inflammation, Oxidative Stress, and Obesity", *International Journal of Molecular Sciences*, 2011.
Publication

<1 %

27

Juan Wu, Ping Niu, Yueqiang Zhao, Yanyang Cheng, Weiping Chen, Lan Lin, Jingmei Lu, Xue Cheng, Zhiliang Xu. "Impact of miR-223-3p and

<1 %

miR-2909 on inflammatory factors IL-6, IL-1 β , and TNF- α , and the TLR4/TLR2/NF- κ B/STAT3 signaling pathway induced by lipopolysaccharide in human adipose stem cells", PLOS ONE, 2019

Publication

28

Lamb, Yvette N., Nicole S. McKay, Christopher S. Thompson, Jeffrey P. Hamm, Karen E. Waldie, and Ian J. Kirk. "Brain-derived neurotrophic factor Val66Met polymorphism, human memory, and synaptic neuroplasticity : BDNF, human memory, and synaptic plasticity", Wiley Interdisciplinary Reviews Cognitive Science, 2014.

Publication

29

Stephanie von Holstein-Rathlou, Nicolas Caesar Petersen, Maiken Nedergaard. "Voluntary running enhances glymphatic influx in awake behaving, young mice", Neuroscience Letters, 2018

Publication

30

cris.maastrichtuniversity.nl

Internet Source

31

ijms.sums.ac.ir

Internet Source

32

journals.plos.org

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

33

www.eneuro.org

Internet Source

<1 %

34

www.jneurosci.org

Internet Source

<1 %

35

www.researchsquare.com

Internet Source

<1 %

36

Neeti Bhat, Narayan Bahadur Mahotra, Lava Shrestha, Vibina Aryal, Nirmala Rayamajhi, Sanyukta Gurung. "Relationship between body mass index and simple visual reaction time in students of a medical college of Nepal", Journal of Physiological Society of Nepal, 2020

Publication

<1 %

37

Usir.salford.ac.uk

Internet Source

<1 %

38

bmcgeriatr.biomedcentral.com

Internet Source

<1 %

39

digibug.ugr.es

Internet Source

<1 %

40

jissn.biomedcentral.com

Internet Source

<1 %

41

physoc.onlinelibrary.wiley.com

Internet Source

<1 %

42

web.archive.org

Internet Source

<1 %

43

www.frontiersin.org

Internet Source

<1 %

44

www.researchpublish.com

Internet Source

<1 %

45

Takashi Hoshikawa, Naomichi Okamoto, Tomoya Natsuyama, Rintaro Fujii et al. "Associations of Serum Cytokines, Growth Factors, and High-Sensitivity C-Reactive Protein Levels in Patients with Major Depression with and without Type 2 Diabetes Mellitus: An Explanatory Investigation", *Neuropsychiatric Disease and Treatment*, 2022

Publication

<1 %

46

Farbod Raiszadeh, Mark I. Travin. "Are people with normal radionuclide perfusion imaging studies better-off if they are obese?", *Journal of Nuclear Cardiology*, 2010

Publication

<1 %

47

Orestes Vicente Forlenza, Breno Satler Diniz, Antonio Lucio Teixeira, Elida Benquique Ojopi et al. "Effect of brain-derived neurotrophic factor Val66Met polymorphism and serum levels on the progression of mild cognitive

<1 %

impairment", The World Journal of Biological Psychiatry, 2010

Publication

48

Robin M. Voigt, Shohreh Raeisi, Jingyun Yang, Sue Leurgans et al. "Systemic brain derived neurotrophic factor but not intestinal barrier integrity is associated with cognitive decline and incident Alzheimer's disease", PLOS ONE, 2021

Publication

<1 %

49

Sung Soo Lee, Jae Ho Yoo, Sung Kang, Jin Hee Woo, Ki Ok Shin, Kwi Beak Kim, Su Youn Cho, Hee Tae Roh, Young Il Kim. "The Effects of 12 Weeks Regular Aerobic Exercise on Brain-derived Neurotrophic Factor and Inflammatory Factors in Juvenile Obesity and Type 2 Diabetes Mellitus", Journal of Physical Therapy Science, 2014

Publication

<1 %

50

Yin Ting, Jia-xing Zhang, Yi Zhao, Xiao-xia Li, Xiu-ying Liu, Nan Li, Yu-hong Zhang. "Analysis and Comparison of Different Obesity Evaluation Indices and Their Effects on Hypertension, Diabetes, and Dyslipidaemia", Research Square, 2020

Publication

<1 %

51

Zhang, Xiang Yang, Yun-Long Tan, Da-Chun Chen, Shu-Ping Tan, Fu-De Yang, Hanjing

<1 %

Emily Wu, Giovana B. Zunta-Soares, Xu-Feng Huang, Thomas R. Kosten, and Jair C. Soares. "Interaction of BDNF with cytokines in chronic schizophrenia", Brain Behavior and Immunity, 2016.

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Negative Correlation between Serum Brain-derived Neurotrophic Factor Levels and Obesity Predictor Markers and Inflammation Levels in Females with Obesity

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
