

# Potassium intake is associated with nutritional quality and actual diet cost a study at formulating a low sodium high potassium (LSHP) healthy diet

*by Farapti Farapti*

---

**Submission date:** 22-Jun-2022 01:02PM (UTC+0800)

**Submission ID:** 1861133645

**File name:** is\_associated\_with\_nutritional\_quality\_and\_actual\_diet\_cost.pdf (377.5K)

**Word count:** 8141

**Character count:** 42151

### RESEARCH ARTICLE

## Potassium intake is associated with nutritional quality and actual diet cost: a study at formulating a low sodium high potassium (LSHP) healthy diet

Farapti Farapti<sup>1,2\*</sup>, Annas Buanasita<sup>3</sup>, Dominikus R. Atmaka<sup>1</sup>, Stefania W. Setyaningtyas<sup>1</sup>, Merryana Adriani<sup>1</sup>, Purwo S. Rejeki<sup>4</sup>, Yoshio Yamaoka<sup>5,6,7</sup> and Muhammad Miftahussurur<sup>7,8</sup>

<sup>1</sup>Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya 60115, Indonesia

<sup>2</sup>Post Graduate Doctoral Program, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

<sup>3</sup>Nutrition Department, Surabaya Health Polytechnic, Surabaya 60282, Indonesia

<sup>4</sup>Department of Physiology, Faculty of Medicine, Universitas Airlangga, Surabaya 60132, Indonesia

<sup>5</sup>Department of Environmental and Preventive Medicine, Oita University Faculty of Medicine, Yufu 879-5593, Japan

<sup>6</sup>Department of Medicine, Gastroenterology and Hepatology Section, Baylor College of Medicine, Houston, TX, USA

<sup>7</sup>Gastroentero-Hepatology Division, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60286, Indonesia

<sup>8</sup>Institute of Tropical Disease, Universitas Airlangga, Surabaya 60115, Indonesia

(Received 6 February 2021 – Final revision received 30 November 2021 – Accepted 3 December 2021)

Journal of Nutritional Science (2022), vol. 11, e11, page 1 of 9

doi:10.1017/jns.2021.104

#### Abstract

Increasing potassium and reducing sodium intake have been identified as a priority intervention to reduce non-communicable diseases. A low sodium high potassium (LSHP) healthy diet can be a predictor of overall dietary quality and is associated with higher diet costs. The present study was a randomised controlled-feeding trial, formulating menus of low sodium and potassium-rich healthy diet and comparing with usual diet (a control diet based on typical Indonesian diet) to assess the association of potassium intake in the menus with other nutritional contents and diet cost. Totally seventy menus, which consisted of LSHP diets and the usual diets for a 7-d cycle, were composed from the analysis of the Indonesian food composition database. The correlation coefficient of the potassium content of all menus with nutritional quality and diet cost was analysed using the Pearson test. Multiple linear regression analysis was performed to determine the most important nutrient in determining diet cost. A comparison of nutrition quality and diet cost from the two menus was analysed using the independent *t*-test. LSHP diet had significantly higher carbohydrate, protein, fibre, magnesium, calcium, vitamin C, potassium density and diet cost but lower total fat, saturated fat and energy density than the usual diet ( $P < 0.05$ ). Furthermore, there was a strong positive correlation between fibre, potassium and diet cost (coefficient correlation of  $>0.8$ ). Potassium is a nutrient that is closely related to diet quality although the cost of the diet often may inhibit its intake. A targeted and effective strategy is required to provide affordable food for achieving a sustainable nutrient-rich diet.

**Key words:** Diet cost: Healthy diet: Non-communicable disease: Nutritional assessment: Potassium: Sodium

#### Introduction

The global disease burden shows that non-communicable diseases (NCDs) are the leading cause of mortality and morbidity globally. There is increasing evidence that lifestyle and poor

diet are the most important modifiable risk factors for NCDs<sup>(1,2)</sup>. Low potassium intake and high sodium intake have also been associated with a number of NCDs, including hypertension<sup>(3)</sup>, cardiovascular diseases<sup>(4)</sup>, obesity<sup>(5)</sup>, osteoporosis<sup>(6)</sup>,

\* Corresponding author: Farapti Farapti; fax +6231-592-0948, email: farapti@fkm.umair.ac.id

Abbreviations: DASH, dietary approaches to stop hypertension; LSHP, low sodium high potassium; NCD, non-communicable diseases

© The Author(s), 2022. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



stroke<sup>(7)</sup> and all-cause mortality<sup>(8)</sup>. In 2012, the World Health Organization (WHO) published guidelines regarding sodium and potassium intakes for adults and children and recommended the adoption of a healthy eating pattern to prevent the onset of diet-related diseases<sup>(9,10)</sup>.

As part of the salt reduction programme, WHO Member States agreed to reduce the global population's dietary salt intake by 30 % by 2025<sup>(11,12)</sup>. A recent systematic review reported that a total of seventy-five countries now have a national salt reduction strategy, twelve countries have reported reductions in population salt intake, nineteen reduced salt content in foods and six improvements in consumer knowledge, attitudes or behaviours relating to salt<sup>(13)</sup>. The successful decrease in sodium intake is based on reports that low sodium diet is likely to be feasible when considering such factors as cost, the need to meet other nutritional recommendations and having familiar meals to enhance acceptability<sup>(14,15)</sup>. In addition, the role of public health programme also contributed to increased awareness for reducing salt intake in the population<sup>(16)</sup>.

In contrast to the salt programme, programmes for increasing potassium intake at the population level are still rare and not progressive. However, it is suggested that a daily intake of 4700 mg of potassium intake has been identified as a priority intervention to reduce NCDs<sup>(10,17,18)</sup>. In reality, most populations around the world consume less than the recommended levels of potassium. The WHO suggests a potassium intake of at least 90 mmol/d (3510 mg/d) for adults; however, the WHO report revealed that potassium intake levels varied among studies and ranged from 45 to 100 mmol/d, with a median value of 73 mmol/d<sup>(10)</sup>. Other population studies showed wide ranges of potassium intake; some consumed approximately 35 mmol/d in the lowest group and 65 mmol/d in the highest group<sup>(10,18)</sup>. The newest meta-analysis showed potassium intake is less than half of the recommended intake and most people do not get enough potassium<sup>(19)</sup>.

Several randomised clinical studies and systematic reviews provided evidence and strengthened the conclusion that increasing potassium has beneficial effects on health. The intervention to increase potassium included supplementation and potassium source food; however, it is recommended that potassium should be consumed through food because of the safety and no upper limit<sup>(10,20–22)</sup>. The WHO and the Dietary Guidelines for Americans Advisory Committee recommendations for increased potassium consumption are found in many foods, including vegetables, fruits, beans and milk products<sup>(10,20)</sup>. The WHO also included potassium as a component of their healthy diet indicator (HDI)<sup>(23)</sup>. In addition, a previous study reported a potassium-rich diet or a potassium-dense diet was associated with higher diet costs<sup>(15)</sup>. This is relevant to many epidemiological studies that observed diets of better quality are costlier, healthy diets tend to be expensive and low-cost energy-dense diets also tend to be nutrient-poor<sup>(24–26)</sup>. However, most people who do not get enough potassium might indicate they consume low-quality diet, and this might lead to the global barrier to healthy eating particularly for low-income households<sup>(19,20)</sup>.

Many countries have applied dietary patterns and developed guidelines to maintain healthy eating in their population.

Furthermore, the WHO has recommended the adoption of a healthy eating pattern to prevent the onset of diet-related diseases. These healthy diets, such as dietary approaches to stop hypertension (DASH) diet, Mediterranean diet and balanced diet, are not only focusing on sodium and potassium, but also essential nutrients categorised as nutrient-rich foods or nutrient-dense foods. These nutrients include potassium, magnesium, calcium, vitamin A (as carotenoids), vitamin C, vitamin E and fibre that are generally consumed in small amounts (less than 60 % of the recommended dietary allowance (RDA), especially for elderly people that appear to lack several nutrients in their diets<sup>(27,28)</sup>).

Since it is plausible that potassium intake can be a predictor of overall dietary quality and can lead to beneficial effects on human health, it is urgently needed to achieve the recommended potassium intake and nutrient-rich diet<sup>(29–31)</sup>. However, food cost affects how people eat healthy diets tend to be more expensive. This might be one reason why many people did not meet their nutrient needs, particularly the recommended potassium intake<sup>(15)</sup>. Therefore, the present study aimed to assess the association between potassium intake and other nutritional factors and diet cost to understand if potassium is a predictor of overall dietary quality and confirm that diet cost might be a barrier limiting many people from meeting the recommended potassium intake.

## 81 Materials and methods

### Study design

The low sodium high potassium (LSHP) study was randomised, controlled-feeding trial comparing the effects of two dietary patterns on blood pressure and other metabolic markers among older people. The present study was ancillary to the LSHP diet trial which was conducted among adult participants in Surabaya, Indonesia. The two dietary patterns were tested in the present study which included the LSHP diet and the usual diet (control diet). We formulated the diet using a 7-d cycle menu, and the nutrient targets of two experimental diets used in the present study were similar to diets used in the DASH-sodium trial, the diet that recommended as a healthy dietary standard by several countries due to its nutritional and metabolic benefits<sup>(20,32–35)</sup>. The menu cycle consisted of three meals (breakfast, lunch and dinner) and two snacks (mid-morning and afternoon). So, totally two dietary patterns provided seventy menus, each LSHP diet and usual diet consisted of thirty-five menus. The menus were developed based on the traditional Indonesian menu, the foods most consumed by the Indonesian population and culturally acceptable.

### Food composition database compilation

22  
Two dietitians with long-standing experience in nutritional fieldwork developed food compositions high in potassium and sodium. The food composition data for calculating nutrients were based on the two Indonesian food composition databases (Tabel Komposisi Pangan Indonesia/TKPI 2017



and Data Komposisi Bahan Makanan/DKBM 2017) (available at [www.panganku.org](http://www.panganku.org)) as the primary nutrient data sources<sup>(36)</sup>. A total of 1151 food items from TKPI and 1375 food items from DKBM were available for analysis. From the food list, we deleted similar food items, resulting in 1146 food items after sorting based on food names. Thereafter, we sorted the food list based on potassium content per 100 g and sodium content per 100 g. Then, we classified the food list differently for both potassium and sodium into four categories: (1) very high ( $\geq 1000$  mg/100 g), (2) high (500–999 mg/100 g), (3) medium (100–499 mg/100 g) and (4) low ( $< 100$  mg/100 g). After classifying the food list, we ranked each category based on potassium and sodium content. Food lists that have higher potassium and sodium ratio were prioritised for use in the menu development.

#### LSHP menu development

From the food database, we developed menus that fulfil the recommendation of high potassium and low sodium intakes. We called the LSHP diet. We formulated the diet using a 7-d cycle menu to develop the two experimental dietary patterns, the LSHP diet and the usual diet. All menus were developed at energy levels of 1800 kcal, potassium 4700 mg and sodium 1500 mg for the LSHP diet and potassium 2000 mg and sodium 2300 mg for the usual diet. We prepared the energy diet for the elderly age group based on the Indonesian RDA at approximately 1800 mg/d both for the LSHP diet and the usual diet<sup>(37)</sup>.

The nutritional composition of the foods or nutrient content of the diets was calculated using the Nutri survey software (version 2007). All foods used in the daily menu had no missing nutrient values in the database. All foods and ingredients were weighed on an electronic scale and balanced to the nearest 0.5 g, except for salt, oils and foods weighing less than 10 g which were weighed to the nearest 0.1 g. To ensure uniformity and consistency of nutrients, menus were developed with similar foods and recipes. For setting sodium content, we applied less salt to regular salted foods or recipes for the LSHP diet, and more salt was added to foods or recipes for the usual diet.

We prepared similar menus to the DASH diet. The diet is recommended as a healthy dietary standard and applied by several countries in the world. The DASH diet is rich in fruits and vegetables and limits the consumption of sugar, salt/sodium and total fat. It also provides increased amounts of potassium, calcium, magnesium and fibre. Furthermore, it parallels the healthy diet menus based on adherence to WHO's 2015 nutrition guidelines. The HDI assesses seven components which include  $\geq 400$  g of fruits and vegetables per day, total fat of  $< 30$  % for total energy,  $< 10$  % saturated fatty acid for total energy and  $\geq 25$  g/d dietary fibre<sup>(23)</sup>. Specifically, for potassium intake, we adjusted the 2019 Indonesian RDA to 4700 mg/d and sodium 1500 mg/d<sup>(34)</sup>.

#### Usual diet

We compiled a control diet as a comparison to the LSHP diet which is propositional to energy intake. A control diet was

made based on regular standards of what many Indonesians eat, and called it as usual diet. The usual menu has the same energy values as the LSHP menu. However, the main difference of those menus is potassium and sodium content: potassium 4700 mg and sodium 1500 mg for the LSHP diet and potassium 2000 mg and sodium 2300 mg for the usual diet. Similar to the LSHP diet, the menu cycle of the usual diet consisted of three meals and two snacks. In the applied menus, for example for snack, we chose cheese bread for the usual diet rather than fresh salad, juice and smoothies in the LSHP diet. In other cycle menus, two groups got the same menus such as juice; however, we chose lower potassium source foods from the fruits group for the usual diet.

#### Nutritional quality

The assessment of nutritional quality was based on the mean of energy-dense diets (energy density), nutrient-dense foods (potassium density), the amount (g) of fruit and vegetable per day, the amount (g) of dietary fibre per day and nutrient content of macronutrient and micronutrient such as calcium, magnesium and vitamin C. All nutrition compositions were obtained from a 7-d cycle menu.

Dietary energy density (kcal/g) was calculated by dividing total dietary energy by the edible weight of foods. Potassium density is the potassium consumption per calorie intake (mg/kcal/d). In the present study, the association between the potassium content in the diet and nutritional quality was measured using the Pearson correlation. Furthermore, we used the independent *t*-test to compare the nutritional quality between the LSHP diet and the usual diet.

#### Actual diet cost

Actual diet cost was the mean cost of the total 7-d cycle menu that was calculated per 1800 kcal of food purchased by using actual expenditures and a standard food price database. To calculate the cost of the diet, we collected regular prices from some major grocery retailers in Surabaya and food price databases. A standard food price database was obtained by estimating the standard price of each of the food items from the information centre of national strategic food price (PIHPS/hargapangan.id), the information system of availability and price development of food staples in East Java (siskaperbapo.com) and prices of food staples in Surabaya's traditional market. After adjustment for preparation and waste, food prices in the databases were expressed in rupiah (IDR) per 100 g of edible portion. All prices were collected between September and November 2019.

#### Statistical analysis

Univariate analysis was performed in the table to show the classification of potassium food sources. The content of potassium from source foods (per 100 g) was divided into three categories in each food group, that is,  $> 1000$ , 500–1000 and 100–500 mg based on two Indonesian food composition databases (Tabel Komposisi Pangan Indonesia/TKPI 2017 and Data



Komposisi Bahan Makanan/DKBM 2017). Furthermore, the classification based on the food types was also presented.

Bivariate analysis was used to measure the correlation between the LSHP diet and diet quality indicators, in terms of nutrient value and actual diet cost. Statistical analysis was performed using the BMI SPSS software. The statistical analysis started by performing the Kolmogorov–Smirnov test to assess the normality of data distribution. Since all data were distributed normally, correlation analysis of potassium intake of the 7-d cycle menu and actual diet cost was performed using the Pearson correlation test and the difference of actual diet cost between the LSHP diet and the usual diet was assessed using the independent *t*-test. Linear regression analysis was performed to determine the correlation between diet cost as a dependent variable and other nutrients as independent variables. The result is considered statistically significant if  $P < 0.05$ .

## Results

The food list containing high potassium was identified through sorting of the food list based on potassium content per 100 g and classifying based on the food group source of carbohydrate, protein, fat, fruits, vegetables and dairy food. Supplementary Table S1 shows that skim milk and flour fortified with dairy were sources of dairy food containing potassium  $>1000$  mg/100 g food weight. Plant protein from the bean group and some kinds of fish were categorised as high potassium food. Fruits and vegetables as sources of potassium generally contained potassium content of 100–1000 mg/100 g. Carbohydrates and proteins, on the other hand, had low potassium content.

The difference in the nutrient content between the LSHP diet and the usual diet can be seen in Table 1. It showed that carbohydrate, protein, fibre, magnesium, calcium, vitamin C and potassium density were significantly higher in the LSHP diet compared to the usual diet. On the other hand, the LSHP

diet contained lower total fat, saturated fat and energy density than the usual diet. Furthermore, the diet cost of LSHP menus was more expensive than that of the usual diet.

The Pearson test was used to assess the correlation between potassium content and other nutrients in all menus (Table 2). The result showed a positive relationship between potassium content and the nutrients in all menus: (1) fibre, calcium, fruits, potassium density and diet cost correlated strongly with potassium (coefficient correlation of  $>0.8$ ); (2) potassium correlated moderately with carbohydrate, protein, magnesium, vitamin C, total fat and saturated fat and (3) a weak correlation was observed between energy and potassium. Further analysed by linear regression, the result of the study showed a significant correlation between diet cost and energy, carbohydrate, fibre, fruits, magnesium and potassium. Moreover, fibre and potassium were the most important nutrients in determining cost (Table 3).

## Discussion

To the best of our knowledge, this is the first study focusing on the formulation of daily menus with recommended potassium content and analyses of potassium content and actual diet cost in the menus. Population studies demonstrated that globally, most people of all ages consume less than the recommended potassium intake<sup>(10)</sup>. Since potassium intake was associated with metabolic syndrome and cardiovascular disease, it is important to fulfil the daily recommended potassium intake. The present study formulated a chronic disease-related diet, with a focus on LSHP foods that are palatable and mostly consumed by the Indonesian population. Furthermore, we adopted the DASH diet, a healthy dietary standard approved by several countries due to its benefit in terms of metabolic and nutritional status<sup>(32–34)</sup>.

Before formulating the daily menus of high recommended potassium healthy diet, it is foremost important to know the potassium food sources and their characteristics. Secondly,

**Table 1.** Macronutrient and micronutrient of the LSHP and usual diets

	High potassium diet (n 35 menus/7-d cycle)	Usual potassium diet (n 35 menus/7-d cycle)	<i>P</i> (significancy)
Energy	1836.27 ± 33.85	1811.14 ± 50.87	0.3
Carbohydrate	299.7 ± 17.34 (65.19 %)	276.63 ± 11.17 (61.13 %)	0.012*
Fat	46.54 ± 6.41 (22.73 %)	54.04 ± 6.01 (26.8 %)	0.043*
Protein	73.44 ± 4.65 (16 %)	66.77 ± 4.77 (14.92 %)	0.021*
Fibre (g)	30.87 ± 7.08	16.53 ± 3.01	<0.001*
Potassium	4577.74 ± 119.29	2127.11 ± 75.86	<0.001*
Sodium	1449.3 ± 136.77	2387.94 ± 172.26	<0.001*
Magnesium	291.33 ± 43.42	235.07 ± 29.79	0.015*
Calcium	840.17 ± 146.10	524.74 ± 83.4	<0.001*
Vitamin C	328.12 ± 266.75	97.14 ± 21.01	0.04*
Vitamin E	4.46 ± 3.11	6.01 ± 1.67	0.27
Vitamin A	2196.16 ± 2050.9	1613.24 ± 806.66	0.5
Saturated fat	5.86 ± 4.32	12.2 ± 3.34	0.01*
Cholesterol	235.89 ± 81.99	293.37 ± 70.61	0.185
Energy density	1.29 ± 0.89	2.02 ± 0.14	<0.001*
Potassium density	2.49 ± 0.99	1.18 ± 0.54	<0.001*
Fruits (g)	737.14 ± 115.10	150 ± 0.00	<0.001*
Vegetables (g)	223.57 ± 75.09	179.29 ± 24.91	0.18
Diet cost (IDR)	35 192.10 ± 3318.27	26 205.71 ± 3678.74	<0.001*

\* Significant  $P < 0.05$ , independent *t*-test.

**Table 2.** Correlation between potassium and other nutrients

	Total		LSHP diet		Usual diet	
	Coefficient correlation	P value	Coefficient correlation	P value	Coefficient correlation	P value
Energy (kcal/d)	0.279 <sup>a</sup>	0.334*	-0.58	0.17	-0.001	0.99
Carbohydrate (g)	0.613	0.02*	-0.57	0.18	-0.66	0.11
Fat (g)	-0.518	0.06	0.65	0.11	0.07	0.88
Protein (g)	0.602	0.023*	0.29	0.53	0.28	0.54
Fibre (g)	0.787	0.001*	0.58	0.01*	0.06	0.39
Sodium (g)	0.201	0.49	-0.38	0.41	0.24	0.61
Magnesium (g)	0.62	0.018*	-0.37	0.41	0.19	0.69
Calcium (g)	0.832	<0.001*	0.29	0.54	0.5	0.25
Vitamin C (mg)	0.551	0.041*	0.07	0.88	-0.39	0.39
Vitamin E (mg)	-0.289	0.316	0.3	0.51	0.72	0.07
Vitamin A (µg)	0.229	0.43	0.64	0.12	-0.34	0.46
Saturated fat	-0.635	0.015*	0.42	0.35	0.6	0.15
Cholesterol	-0.38	0.18	-0.29	0.53	0.31	0.49
Energy density	-0.956	<0.001*	-0.18	0.7	0.34	0.45
Potassium density	0.998	<0.001*	0.92	0.003*	0.78	0.04*
Fruits (g)	0.975	<0.001*	0.58	0.18	0.61	0.15
Vegetables (g)	0.358	0.209	-0.52	0.23	-0.46	0.29
Diet cost (IDF)	0.809	<0.001*	0.45	0.31	0.68	0.09

LSHP, low sodium high potassium.

<sup>a</sup> All nutrition compositions were obtained from a 7-d cycle menu.

\* Significant  $P < 0.05$ , Pearson's test.

the obstacle to arranging the menus of a healthy diet should be identified. In the present study, food sources containing more than 500 mg potassium per 100 g were fruits, vegetables and plant protein, especially nut and dairy food. Conversely, most of the carbohydrate and fat food sources contained low potassium. Our result is similar to a previous study that reported that highly refined wheat flour contains less than half of the potassium level of complete flour and refined fats or sugars have very low potassium content and their presence or addition in foods results in a lowering of potassium concentration as well as nutritional density<sup>(20,21,35)</sup>. Finally, we chose foods that are low-cost sources of potassium such as potatoes, bananas and carrots to meet targeted dietary recommendations. This is similar to a previous study that highlighted that it is important to identify and promote lower-cost food as an effective approach to improve potassium intake<sup>(15)</sup>.

Potassium is an essential element and is found in all foods. However, previous studies demonstrated that food sources of potassium come from particularly fruits and vegetables<sup>(20,30)</sup>. Several large surveys have been published indicating that fruits and vegetables are the richest food sources of potassium among children<sup>(38)</sup>, adult<sup>(39)</sup> and the elderly<sup>(28)</sup>. Many fresh

fruits and vegetables are rich in potassium. In the present study, however, fruits had a strong correlation with potassium ( $r = 0.975$ ,  $P = 0.000$ ), but not vegetables. Dietary guidelines also suggested the consumption of adequate fruits and vegetables ranging from 400 to 600 mg/d<sup>(40)</sup>. Available empirical evidence have also found that fruits and vegetables are substantially more nutrient-dense with low energy content than other food groups, as evident by their highest nutrient density score<sup>(24,25,27)</sup>. Therefore, adequate intakes of fruits and vegetables, which are the major sources of potassium, are highly recommended. Besides the potassium benefit, fruit and vegetable intakes were associated with reduced risk of cardiovascular disease, cancer and all-cause mortality<sup>(29-31)</sup>. Unfortunately, despite these established benefits, most people consume less fruits and vegetables than recommended<sup>(41-45)</sup>.

The difficulty encountered in formulating the LSHP diet is related to energy content. The present study showed that the higher the potassium content in the menus serving, the higher the macronutrient intake (energy, protein and carbohydrate). It was particularly challenging to arrange the menus containing calorie restriction with high potassium and low sodium content, particularly for obesity treatment. Consistently, the previous studies showed that sodium and potassium intakes were positively correlated with energy intake<sup>(46,47)</sup>. The same trend was observed by a previous research which reported that potassium intake was correlated with total calories but further stated that potassium alone, which is not more effective than the DASH diet, improves endothelial function<sup>(48)</sup>. On the contrary, other studies confirmed that high potassium intake was associated with lower body mass index (BMI)<sup>(30,49,50)</sup>. The increase in dietary potassium consumption was a strong predictor of a decline in BMI, which might have been caused by the potassium density<sup>(49)</sup>. The study showed that participants who achieved a higher loss in BMI also had higher potassium density. The present study revealed that potassium

**Table 3.** Linear regression analysis between diet cost and other nutrients

	Standardised coefficients ( $\beta$ )	P value
Energy (kcal/d)	0.46	0.007*
Carbohydrate (g)	-0.49	0.02*
Potassium	0.53	0.004
Magnesium	-0.48	0.023*
Fibre (g)	1.22	0.04*
Vegetables (g)	-0.39	0.013

\* Significant  $P < 0.05$ .



intake had a strong positive correlation with potassium density, while it had a negative correlation with energy density. The present findings also revealed that potassium density was significantly higher in the LSHP diet compared to the usual diet. Thus, it is likely that potassium density, defined as potassium consumption per calorie intake, might become a useful marker to assess nutritional quality.

To ensure the consumption of a healthy and nutritious diet, identifying foods that are rich in nutrients is paramount. Drewnowski developed a nutrient-rich food index that focused on consuming fewer calories but with more beneficial nutrients<sup>(51)</sup>. Potassium is one of the beneficial nutrients with nutrient-rich density. A study conducted in 2019 developed a nutrient-rich food score that is useful to assess the nutrient density of diets among the elderly. The tool consists of seven nutrients that should be encouraged (protein, dietary fibre, folate, vitamin D, calcium, magnesium and potassium) and three nutrients to limit (saturated fat, sodium and mono- and disaccharides)<sup>(28)</sup>. This 2019 study compares well with the findings of the present study which showed that potassium intake and potassium density were strongly correlated with other nutrients such as magnesium, calcium, fibre and vitamin C. Compared to the usual potassium diet, the LSHP diet showed significantly higher potassium, calcium, magnesium, fibre, vitamin C, protein and carbohydrate contents. On the other hand, the total fat and sodium content was lower.

The formulation of the LSHP diet in the present study is similar to the DASH which is high in potassium, magnesium, calcium and fibre content, and low in sodium. The DASH diet was originally proposed for the prevention and treatment of hypertension and is currently recommended as a healthy dietary standard by several countries due to its metabolic and nutritional benefits<sup>(32–34)</sup>. The benefits of this diet rich in fruits and vegetables are attributable to its micronutrient composition which provides increased amounts of potassium, magnesium, calcium and fibre. The diet was originally developed to contain foods that increase potassium, magnesium and calcium based on established links to lower blood pressure and successful demonstration of a clinically meaningful blood pressure-lowering effect<sup>(32–34,52)</sup>. Interactions between these nutrients can also have blood pressure lowering effects, such as the interaction between potassium and calcium and the ability to increase sodium excretion by the kidneys. Potassium and calcium have previously been demonstrated to interact with the renin–angiotensin system by affecting plasma renin activity, and potassium can assist with sodium balances and has also been demonstrated to potentially lower blood pressure through endothelium-dependent vascular effects<sup>(3,53,54)</sup>. Population studies suggested that high fruits and vegetables in DASH lower blood pressure and improve endothelial function in obese hypertensive subjects<sup>(48)</sup>. Moreover, the umbrella review of the systematic review concluded that the DASH dietary pattern is associated with decreased incidence of cardiovascular disease and improves blood pressure with evidence of other cardiometabolic advantages in people with and without diabetes<sup>(34)</sup>. In the future, the LSHP diet in the present study can be applied to increase

potassium intake in healthy populations or people with NCDs.

One of the problems about a healthy diet, particularly high potassium intake, is the high cost. The present study showed that the diet cost of LSHP is significantly more expensive than the usual diet. It means that a nutrient-rich healthy diet is expensive. The daily mean cost of the LSHP diet was IDR 35 192.10 ± 3318.27 per person per day. It was categorised as high food expenditure based on daily food cost for Indonesian civil servants<sup>(55)</sup>. A similar categorisation was demonstrated by an epidemiological study which found that lower household expenditure and shorter education were associated with lower potassium intake<sup>(56)</sup>. The same trend was observed in previous studies which found that healthy diets tend to be expensive and low-cost energy-dense diets also tend to be nutrient-poor<sup>(24–26)</sup>. Foods that supply relatively more nutrients than calories are defined as nutrient-dense. In the present study, the LSHP diet is a more nutrient-dense and nutrient-rich food than the usual diet. It is well documented that nutrient density is an accurate marker of healthy diets, distinguishing between diets that are energy-dense and those that are nutrient-rich<sup>(57)</sup>. It was observed that besides potassium, the components of carbohydrate, protein, fibre, magnesium, calcium, vitamin C, fruit and potassium density were significantly higher in the LSHP diet compared to the usual diet. Adherence to healthy diet indicator (HDI) in a study in Japan is reported to be poor (6.6 %) and higher adherence to HDI was significantly associated with diets higher in protein, polyunsaturated fatty acids (PUFA), dietary fibre, potassium and vitamins<sup>(23)</sup>. The LSHP diet in the present study met the healthy diet recommendation by the WHO. It contained ≥400 g of fruits and vegetables per day, ≥25 g/d dietary fibre, ≥3500 mg/d potassium and total fat of <30 % for total energy. Furthermore, higher-quality diets were not only more costly but were also consumed by persons of higher educational levels. In contrast, foods with lower nutritional values and lower-quality diets generally cost less and tended to be consumed by groups of lower socio-economic status<sup>(58)</sup>. This finding is strengthened by other studies which found that nutrient density was negatively associated with energy density and positively associated with cost<sup>(59,60)</sup>.

The linear regression analysis in the present study showed that fibre and potassium were the most important nutrients in determining cost. This is similar to a previous study that reported a potassium-rich diet or a potassium-dense diet which was associated with higher diet costs<sup>(15)</sup>. In contrast, it differed from cost analysis of essential nutrients in the National Health and Nutrition Examination Survey from 2011 to 2014 that shows protein food was one of the most expensive food categories<sup>(26)</sup>. Interestingly, the present study proved fibre was the most nutrient costly. It might be still related to potassium since source food fibre is similar to potassium such as fruits and vegetables. Fruits and vegetables provided key nutrients at a reasonable cost when compared with other foods<sup>(24)</sup>. It was also proven by other countries particularly the low-income countries where healthy foods, including fruits and vegetables, were categorised as expensive food<sup>(59,60)</sup>. Fruits, vegetables and complex carbohydrates



together contributed to high fibre content in the menus, and they may have contributed to the high cost of fibre.

Several population studies showed that the consumption of high potassium healthy diets appears to still be a problem. There is a nutrient gap due to economic constraints in acquiring a nutritious diet. A study on diet cost in Indonesia revealed that affording a nutritious diet depended on household expenditure and geographic area. More people in the urban area can afford to consume high-quality diets<sup>(61)</sup>. A study by Temple and Steyn revealed that a healthy diet is unaffordable for most South Africans<sup>(62)</sup>. Another study showed that children from food-insecure households were more likely to consume low-quality diets, and higher parental income was associated with higher diet costs affordability<sup>(63)</sup>. The same trends were observed by Marty *et al.* when they found that among low-income population, actual diet cost was positively correlated with nutritional quality<sup>(64)</sup>. In addition, the study by Monsivais *et al.* demonstrated that higher-quality diets were costly and consumed by persons of higher educational level<sup>(59)</sup>. Similar findings were also reported in a previous study that a potassium-rich diet or a potassium-dense diet was associated with higher diet costs<sup>(15)</sup>.

The notable strength of the present study is that we identified food sources with high potassium and low sodium and utilised these directly in the formulation of healthy diet menus. Specifically, the optimal nutrients in the menus were developed based on the traditional Indonesian menu and the suitable Indonesian balance diet guideline. Therefore, the formulation of the LSHP diet was culturally acceptable, palatable and nutrient-rich. One important application of diet models is achieving a nutrient-rich diet at an affordable cost. Although we tried to arrange menus at an affordable cost, the limitation of the present study is that the LSHP diet is only applicable to the specific population and may limit generalisability to other populations. Since a large number of epidemiological studies showed that a healthy diet and nutrient-rich food are expensive, it is not easy to identify and providing healthy menus with optimal nutrients at an affordable cost is an uphill task.

## Conclusions

The main findings of the present study are that the LSHP diet had significantly higher essential nutrients than the usual diet. Potassium is one of the essential nutrients of a healthy diet, so it is important to consume the recommended high potassium healthy diet intake for maintaining health and preventing NCDs. In the present study, there was a strong positive correlation between fibre, potassium and diet cost. Since foods rich in potassium and fibre were relatively costly, cost may be a barrier to adopting a healthier diet and improving diet quality in populations. It may require a more effective and progressive strategy and price intervention from government to provide affordable and nutrient-rich food for achieving a sustainable healthy eating pattern. In addition, the menus were developed based on the traditional Indonesian menu, so they may not be applied in other ethnic groups and can not be extrapolated to a larger population. It is suggested to evaluate the menus based

on different ethnicity and to a larger population to gain more comprehensive results and adaptability.

## Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/jns.2021.104>.

## Acknowledgments

We would like to thank all investigators who analysed the present study's data and provided feedback during the preparation of this manuscript.

All authors contributed to the review of articles. F.F. involved in the conception and design, data analysis, draft manuscript, and review and editing. A.B., D.R.A. and S.W.A. helped to formulate menus, compiled data to be processed and analysed the data. M.A., P.S.R., M.M. and Y.Y. validated and supervised the study. Y.Y. provided review and editing. Finally, all authors read and approved the final manuscript.

This work was supported by Universitas Airlangga.

The authors declare that they have no conflicts of interest.

## References

- Eckel RH, Jakicic JM & Ard JD (2014) AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* **129**, S76–S99.
- World Health Organization (2016) *Assessing National Capacity for the Prevention and Control of Noncommunicable Diseases: Report of the 2015 Global Survey*. Geneva: World Health Organization.
- Adrogué HJ & Madias NE (2007) Sodium and potassium in the pathogenesis of hypertension. *N Engl J Med* **356**, 1966–1978.
- Cook NR, Obarzanek E, Cutler JA, *et al.* (2009) Joint effects of sodium and potassium intake on subsequent cardiovascular disease: the Trials of Hypertension Prevention follow-up study. *Arch Intern Med* **169**, 32–40.
- Jain N, Minhajuddin AT, Neeland IJ, *et al.* (2014) Association of urinary sodium-to-potassium ratio with obesity in a multiethnic cohort. *Am J Clin Nutr* **99**, 992–998.
- Zhu K, Devine A & Prince RL (2009) The effects of high potassium consumption on bone mineral density in a prospective cohort study of elderly postmenopausal women. *Osteoporos Int* **20**, 335–340.
- Jayedi A, Ghomashi F, Zargar MS, *et al.* (2019) Dietary sodium, sodium-to-potassium ratio, and risk of stroke: a systematic review and nonlinear dose-response meta-analysis. *Clin Nutr* **38**, 1092–1100.
- Yang Q, Liu T, Kuklina EV, *et al.* (2011) Sodium and potassium intake and mortality among US adults: prospective data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* **171**, 1183–1191.
- World Health Organization (2012) *Guideline: Sodium Intake for Adults and Children*. Geneva: World Health Organization.
- World Health Organization (2012) *Guideline: Potassium Intake for Adults and Children*. Geneva: World Health Organization.
- World Health Organization (2007) *Reducing Salt Intake in Populations: Report of a WHO Forum and Technical Meeting*. France: World Health Organization.
- Webster J, Trieu K, Dunford E, *et al.* (2014) Target salt 2025: a global overview of national programs to encourage the food industry to reduce salt in foods. *Nutrients* **6**, 3274–3287.





13. Trieu K, Neal B, Hawkes C, *et al.* (2015) Salt reduction initiatives around the world: a systematic review of progress towards the global target. *PLoS ONE* **10**, 1–22.
14. Wilson N, Nghiem N & Foster RH (2013) The feasibility of achieving low-sodium intake in diets that are also nutritious, low-cost, and have familiar meal components. *PLoS ONE* **8**, e58539.
15. Drewnowski A, Rehm CD, Maillot M, *et al.* (2015) The relation of potassium and sodium intakes to diet cost among U.S. adults. *J Hum Hypertens* **29**, 14–21.
16. Farapti F, Fatimah AD, Astutik E, *et al.* (2020) Awareness of salt intake among community-dwelling elderly at coastal area: the role of public health access program. *J Nutr Metab.* doi: 10.1155/2020/8793869.
17. McDonough AA & Youn JH (2017) Potassium homeostasis: the knowns, the unknowns, and the health benefits. *Physiology (Bethesda)* **32**, 100–111.
18. Burnier M (2019) Should we eat more potassium to better control blood pressure in hypertension? *Nephrol Dial Transplant* **34**, 184–193.
19. Tan M, He FJ, Wang C, *et al.* (2019) Twenty-four-hour urinary sodium and potassium excretion in China: a systematic review and meta-analysis. *J Am Heart Assoc* **8**, 1–12.
20. US Department of Health and Human Services (2006) *Lowering Your Blood Pressure with DASH*. Bethesda: NH Publication.
21. Demigné C, Sabbah H, Rémésy C, *et al.* (2004) Protective effects of high dietary potassium: nutritional and metabolic aspects. *J Nutr* **134**, 2903–2906.
22. Palmer BF & Clegg DJ (2016) Achieving the benefits of a high-potassium, paleolithic diet, without the toxicity. *Mayo Clin Proc* **91**, 496–508.
23. Kanauchi M & Kanauchi K (2018) The World Health Organization's healthy diet indicator and its associated factors: a cross-sectional study in central Kinki, Japan. *Prev Med reports* **12**, 198–202.
24. Darmon N, Darmon M, Maillot M, *et al.* (2005) A nutrient density standard for vegetables and fruits: nutrients per calorie and nutrients per unit cost. *J Am Diet Assoc* **105**, 1881–1887.
25. Drewnowski A & Specter SE (2004) Poverty and obesity: the role of energy density and energy costs. *Am J Clin Nutr* **79**, 6–16.
26. Hess JM, Cifelli CJ, Agarwal S, *et al.* (2019) Comparing the cost of essential nutrients from different food sources in the American diet using NHANES 2011–2014. *Nutr J* **18**, 68.
27. Kennedy E, Racsa P, Dallal G, *et al.* (2008) Alternative approaches to the calculation of nutrient density. *Nutr Rev* **66**, 703–709.
28. Berendsen AAM, Kramer CS & de Groot LCPGM (2019) The newly developed elderly nutrient-rich food score is a useful tool to assess nutrient density in European older adults. *Front Nutr* **6**, 119.
29. Aune D, Giovannucci E, Boffetta P, *et al.* (2017) Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality: a systematic review and dose-response meta-analysis of prospective studies. *Int J Epidemiol* **46**, 1029–1056.
30. Cai X, Li X, Fan W, *et al.* (2016) Potassium and obesity/metabolic syndrome: a systematic review and meta-analysis of the epidemiological evidence. *Nutrients* **8**, 183.
31. Blanch N, Clifton PM & Keogh JB (2015) A systematic review of vascular and endothelial function: effects of fruit, vegetable and potassium intake. *Nutr Metab Cardiovasc Dis* **25**, 253–266.
32. Lin P-H, Allen JD, Li Y-J, *et al.* (2013) Blood pressure-lowering mechanisms of the DASH dietary pattern. *J Nutr Metab.* doi:10.1155/2012/472396.
33. Jenkins DJA, Jones PJ, Frohlich J, *et al.* (2015) The effect of a dietary portfolio compared to a DASH-type diet on blood pressure. *Nutr Metab Cardiovasc Dis* **25**, 1132–1139.
34. Chiavaroli L, Vigliouk E, Nishi SK, *et al.* (2019) DASH dietary pattern and cardiometabolic outcomes: an umbrella review of systematic reviews and meta-analyses. *Nutrients* **11**, 338.
35. Lin P-H, Aickin M, Champagne C, *et al.* (2003) Food group sources of nutrients in the dietary patterns of the DASH-sodium trial. *J Am Diet Assoc* **103**, 488–496.
36. Ministry of Health (2017) *Tabel Komposisi Pangan Indonesia/TKPI 2017 and Data Komposisi Bahan Makanan/DKBM 2017*. Jakarta: Ministry of Health Republic Indonesia.
37. Ministry of Health (2019) *Dietary Allowance*. Jakarta: Ministry of Health Republic Indonesia.
38. Huybrechts I, De Keyzer W, Lin Y, *et al.* (2012) Food sources and correlates of sodium and potassium intakes in Flemish pre-school children. *Public Health Nutr* **15**, 1039–1046.
39. Meneton P, Lafay L, Tard A, *et al.* (2009) Dietary sources and correlates of sodium and potassium intakes in the French general population. *Eur J Clin Nutr* **63**, 1169–1175.
40. Health Department of Republic Indonesia (2014) *Balanced Diet Guideline*. Jakarta: Health Department of Republic Indonesia.
41. Salehi L, Eftekhar H, Mohammad K, *et al.* (2010) Consumption of fruit and vegetables among elderly people: a cross sectional study from Iran. *Nutr J* **13**, 2.
42. Kamphuis CBM, Giskes K, de Bruijn G-J, *et al.* (2006) Environmental determinants of fruit and vegetable consumption among adults: a systematic review. *Br J Nutr* **96**, 620–635.
43. Di Noia J & Byrd-Bredbenner C (2014) Determinants of fruit and vegetable intake in low-income children and adolescents. *Nutr Rev* **72**, 575–590.
44. Trude ACB, Kharmats AY, Hurley KM, *et al.* (2016) Household, psychosocial, and individual-level factors associated with fruit, vegetable, and fiber intake among low-income urban African American youth. *BMC Public Health* **16**, 872.
45. Msambichaka B, Eze IC, Abdul R, *et al.* (2018) Insufficient fruit and vegetable intake in a low- and middle-income setting: a population-based survey in semi-urban Tanzania. *Nutrients* **10**, 222.
46. Athanasatou A, Kandyliari A, Malisova O, *et al.* (2018) Sodium and potassium intake from food diaries and 24-h urine collections from 7 days in a sample of healthy Greek adults. *Front Nutr* **5**, 13.
47. Farapti F, Sulistyowati M, Artanti KD, *et al.* (2019) Highlighting of urinary sodium and potassium among Indonesian schoolchildren aged 9–12 years: the contribution of school food. *J Nutr Metab.* doi:10.1155/2019/1028672.
48. Al-Solaiman Y, Jesri A, Mountford WK, *et al.* (2010) DASH lowers blood pressure in obese hypertensives beyond potassium, magnesium and fibre. *J Hum Hypertens* **24**, 237–246.
49. Tal B, Sack J, Yaron M, *et al.* (2019) Increment in dietary potassium predicts weight loss in the treatment of the metabolic syndrome. *Nutrients* **11**, 1256.
50. Elfassy T, Mossavar-Rahmani Y, Van Horn L, *et al.* (2018) Associations of sodium and potassium with obesity measures among diverse US Hispanic/Latino adults: results from the Hispanic community health study/study of Latinos. *Obesity (Silver Spring)* **26**, 442–450.
51. Drewnowski A (2010) The Nutrient Rich Foods Index helps to identify healthy, affordable foods. *Am J Clin Nutr* **91**, 1095S–1101S.
52. Sacks FM, Svetkey LP, Vollmer WM, *et al.* (2001) Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med* **344**, 3–10.
53. Karppanen H, Karppanen P & Mervaala E (2005) Why and how to implement sodium, potassium, calcium, and magnesium changes in food items and diets? *J Hum Hypertens* **19**, S10–S19.
54. Houston MC & Harper KJ (2008) Potassium, magnesium, and calcium: their role in both the cause and treatment of hypertension. *J Clin Hypertens (Greenwich)* **10**, 3–11.
55. Finance Minister Republic Indonesia (2020) *Budget Cost Standards of 2020*. Available at: [www.jdih.kemkeu.go.id](http://www.jdih.kemkeu.go.id).
56. Miyagawa N, Okuda N, Nakagawa H, *et al.* (2018) Socioeconomic status associated with urinary sodium and potassium excretion in Japan: NIPPON DATA 2010. *J Epidemiol* **28**, S29–S34.
57. Drewnowski A & Fulgoni VL (2014) Nutrient density: principles and evaluation tools. *Am J Clin Nutr* **99**, 1223S–1228S.
58. Darmon N & Drewnowski A (2015) Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev* **73**, 643–660.



59. Monsivais P, McLain J & Drewnowski A (2010) The rising disparity in the price of healthful foods: 2004–2008. *Food Policy* **35**, 514–520.
60. Drewnowski A, Monsivais P, Maillot M, *et al.* (2007) Low-energy-density diets are associated with higher diet quality and higher diet costs in French adults. *J Am Diet Assoc* **107**, 1028–1032.
61. Baldi G, Martini E, Catharina M, *et al.* (2013) Cost of the diet (CoD) tool: first results from Indonesia and applications for policy discussion on food and nutrition security. *Food Nutr Bull* **34**, S35–S42.
62. Temple NJ & Steyn NP (2011) The cost of a healthy diet: a South African perspective. *Nutrition* **27**, 505–508.
63. Bukambu E, Lieffers JRL, Ekwaru JP, *et al.* (2020) The association between the cost and quality of diets of children in Canada. *Can J Public Health* **111**, 269–277.
64. Marty L, Dubois C, Gaubard MS, *et al.* (2015) Higher nutritional quality at no additional cost among low-income households: insights from food purchases of 'positive deviants'. *Am J Clin Nutr* **102**, 190–198.

# Potassium intake is associated with nutritional quality and actual diet cost a study at formulating a low sodium high potassium (LSHP) healthy diet

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

10%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Silvia Bel-Serrat, Antje von der Schulenburg, Amy Mullee, Celine Murrin. "Determinants of vegetable intake among urban socio-economically disadvantaged adolescents: a systematic review of quantitative studies", *Public Health Nutrition*, 2021  
Publication 1%
- 2 Marthe-Lise Næss-Andresen, Anne Karen Jenum, Jens Petter Berg, Ragnhild Sørum Falk, Line Sletner. "Prevalence of postpartum anaemia and iron deficiency by serum ferritin, soluble transferrin receptor and total body iron, and associations with ethnicity and clinical factors: a Norwegian population-based cohort study", *Journal of Nutritional Science*, 2022  
Publication 1%
- 3 Christian Demigné, Houda Sabboh, Christian Rémésy, Pierre Meneton. "Protective Effects of High Dietary Potassium: Nutritional and 1%

# Metabolic Aspects", The Journal of Nutrition, 2004

Publication

---

4	Masao Kanauchi, Kimiko Kanauchi. "The World Health Organization's Healthy Diet Indicator and its associated factors: A cross-sectional study in central Kinki, Japan", Preventive Medicine Reports, 2018 Publication	1 %
5	"Nutrition in Lifestyle Medicine", Springer Science and Business Media LLC, 2017 Publication	<1 %
6	Drewnowski, A, C D Rehm, M Maillot, and P Monsivais. "The relation of potassium and sodium intakes to diet cost among US adults", Journal of Human Hypertension, 2015. Publication	<1 %
7	<a href="http://bmcpediatr.biomedcentral.com">bmcpediatr.biomedcentral.com</a> Internet Source	<1 %
8	Monsivais, P.. "Lower-Energy-Density Diets Are Associated with Higher Monetary Costs per Kilocalorie and Are Consumed by Women of Higher Socioeconomic Status", Journal of the American Dietetic Association, 200905 Publication	<1 %
9	<a href="http://www.fao.org">www.fao.org</a> Internet Source	<1 %

---

10

[www.nutritionsocty.org](http://www.nutritionsocty.org)

Internet Source

&lt;1 %

11

Kathy Trieu, Bruce Neal, Corinna Hawkes, Elizabeth Dunford, Norm Campbell, Rodrigo Rodriguez-Fernandez, Branka Legetic, Lindsay McLaren, Amanda Barberio, Jacqui Webster. "Salt Reduction Initiatives around the World – A Systematic Review of Progress towards the Global Target", PLOS ONE, 2015

Publication

&lt;1 %

12

[repo.lib.tokushima-u.ac.jp](http://repo.lib.tokushima-u.ac.jp)

Internet Source

&lt;1 %

13

Thein Myint, Muhammad Miftahussurur, Ratha-korn Vilaichone, New Ni et al. "Characterizing Helicobacter pylori cagA in Myanmar", Gut and Liver, 2018

Publication

&lt;1 %

14

Natalie Marinho Dantas. "Avaliação sensorial de alimentos com redução de sódio: perspectivas de consumidores universitários", Universidade de Sao Paulo, Agencia USP de Gestao da Informacao Academica (AGUIA), 2021

Publication

&lt;1 %

15

[nutritionj.biomedcentral.com](http://nutritionj.biomedcentral.com)

Internet Source

&lt;1 %

16

Suzana Shahar, Yee Xing You, NurZetty Sofia Zainuddin, Viola Michael et al. "Sodium content in sauces—a major contributor of sodium intake in Malaysia: a cross-sectional survey", *BMJ Open*, 2019

Publication

&lt;1 %

17

Giulia Baldi, Elviyanti Martini, Maria Catharina, Siti Muslimatun et al. "Cost of the Diet (CoD) Tool: First Results from Indonesia and Applications for Policy Discussion on Food and Nutrition Security", *Food and Nutrition Bulletin*, 2013

Publication

&lt;1 %

18

*Metal Ion in Stroke*, 2012.

Publication

&lt;1 %

19

Putri Dwi Kurnianingsih Darely, Muhammad Iqbal. "Perbandingan efektivitas penggunaan database makanan berbasis website, dekstop dan buku pada mahasiswa gizi di Indonesia", *AcTion: Aceh Nutrition Journal*, 2019

Publication

&lt;1 %

20

Reham Alaini, Roslee Rajikan, Siti Masitah Elias. "Diet optimization using linear programming to develop low cost cancer prevention food plan for selected adults in Kuala Lumpur, Malaysia", *BMC Public Health*, 2019

Publication

&lt;1 %

21 Mokhtari, Z., S. Hosseini, R. Miri, A. R. Baghestani, M. Zahedirad, M. Rismanchi, and J. Nasrollahzadeh. "Relationship between Dietary Approaches to Stop Hypertension score and Alternative Healthy Eating Index score with plasma asymmetrical dimethylarginine levels in patients referring for coronary angiography", Journal of Human Nutrition and Dietetics, 2015.  
Publication

<1 %

22 Willem De Keyzer, Yi Lin, Carine Vereecken, Lea Maes et al. "Dietary sources of energy and macronutrient intakes among Flemish preschoolers", Archives of Public Health, 2011  
Publication

<1 %

23 kidney360.asnjournals.org  
Internet Source

<1 %

24 link.springer.com  
Internet Source

<1 %

25 lpi.oregonstate.edu  
Internet Source

<1 %

26 Francesca Biandolino, Isabella Parlapiano, Lucia Spada, Antonella Di Leo et al. " Occurrence and patterns of nutritional traits and polycyclic aromatic hydrocarbons (PAHs) in sea cucumber ( ) tissues: benefits and risk

<1 %

for human health ", Food Quality and Safety,  
2022

Publication

---

27

Ayyanathan, . "Author Notes", Cancer Cell  
Signaling, 2014.

Publication

---

<1 %

28

Bénédicte Tremblay, Frédéric Guénard, Benoît  
Lamarche, Louis Pérusse, Marie-Claude Vohl.  
"Genetic and Common Environmental  
Contributions to Familial Resemblances in  
Plasma Carotenoid Concentrations in Healthy  
Families", Nutrients, 2018

Publication

---

<1 %

29

Swapnil Hiremath. "Towards better blood  
pressure: Do non-pharmacological strategies  
provide the right path?", The Journal of  
Clinical Hypertension, 2018

Publication

---

<1 %

30

N. Unwin, K. G. M. M. Alberti. "Chronic non-  
communicable diseases", Annals of Tropical  
Medicine & Parasitology, 2013

Publication

---

<1 %

31

Colin D. Rehm, Adam Drewnowski. "Replacing  
Dairy Fat With Polyunsaturated and  
Monounsaturated Fatty Acids: A Food-Level  
Modeling Study of Dietary Nutrient Density  
and Diet Quality Using the 2013–16 National

<1 %



Health and Nutrition Examination Survey",  
Frontiers in Nutrition, 2019

Publication

32

Corné van Dooren, Annely Douma, Harry Aiking, Pier Vellinga. "Proposing a Novel Index Reflecting Both Climate Impact and Nutritional Impact of Food Products", Ecological Economics, 2017

Publication

<1 %

33

"Integrative and Functional Medical Nutrition Therapy", Springer Science and Business Media LLC, 2020

Publication

<1 %

34

[WWW.MDPI.COM](http://WWW.MDPI.COM)

Internet Source

<1 %

35

Yume Nohara - Shitama, Hisashi Adachi, Mika Enomoto, Ako Fukami et al. "Twenty - four - Hour Urinary Potassium Excretion, But Not Sodium Excretion, Is Associated With All - Cause Mortality in a General Population", Journal of the American Heart Association, 2018

Publication

<1 %

36

[cdn.doctoronly.co.il](http://cdn.doctoronly.co.il)

Internet Source

<1 %

37

[edepot.wur.nl](http://edepot.wur.nl)

Internet Source

<1 %

38	<a href="http://kclpure.kcl.ac.uk">kclpure.kcl.ac.uk</a> Internet Source	<1 %
39	<a href="http://www.wrc.org.za">www.wrc.org.za</a> Internet Source	<1 %
40	"OS10: Asthma 2 : Oral Sessions", <i>Respirology</i> , 2013. Publication	<1 %
41	Adelais Athanasatou, Aikaterini Kandyliari, Olga Malisova, Alex Pepa, Maria Kapsokefalou. "Sodium and Potassium Intake from Food Diaries and 24-h Urine Collections from 7 Days in a Sample of Healthy Greek Adults", <i>Frontiers in Nutrition</i> , 2018 Publication	<1 %
42	Darmon, N.. "A Nutrient Density Standard for Vegetables and Fruits: Nutrients per Calorie and Nutrients per Unit Cost", <i>Journal of the American Dietetic Association</i> , 200512 Publication	<1 %
43	Katelyn Bowden, Nicholas A Gray, Elizabeth Swanepoel, Hattie H Wright. "A Mediterranean lifestyle is associated with favourable cardiometabolic markers in people with non-dialysis dependent chronic kidney disease", <i>Journal of Nutritional Science</i> , 2021 Publication	<1 %

44

Internet Source

&lt;1 %

45

[open.library.ubc.ca](https://open.library.ubc.ca)

Internet Source

&lt;1 %

46

[www.jpmpm.org](http://www.jpmpm.org)

Internet Source

&lt;1 %

47

[www.rombio.eu](http://www.rombio.eu)

Internet Source

&lt;1 %

48

[zombiedoc.com](http://zombiedoc.com)

Internet Source

&lt;1 %

49

Katherine J Overwyk, Zerleen S Quader, Joyce Maalouf, Marlana Bates et al. "Dietary Sodium Intake and Health Indicators: A Systematic Review of Published Literature between January 2015 and December 2019", *Advances in Nutrition*, 2020

Publication

&lt;1 %

50

Matthieu Maillot, Florent Vieux, Fabien Delaere, Anne Lluch, Nicole Darmon. "Dietary changes needed to reach nutritional adequacy without increasing diet cost according to income: An analysis among French adults", *PLOS ONE*, 2017

Publication

&lt;1 %

51

S. Susan Hedayati, Abu T. Minhajuddin, Adeel Ijaz, Orson W. Moe, Essam F. Elsayed, Robert

&lt;1 %

F. Reilly, Chou-Long Huang. "Association of Urinary Sodium/Potassium Ratio with Blood Pressure: Sex and Racial Differences", Clinical Journal of the American Society of Nephrology, 2012

Publication

---

52

Sigrid Denver, Tove Christensen, Jonas Nordström. "Consumer preferences for low-salt foods: a Danish case study based on a comprehensive supermarket intervention", Public Health Nutrition, 2021

Publication

---

<1 %

53

Sigrunn Hernes, Rona N. Cabo, Mohammad Azam Mansoor, Margaretha Haugen. "Eating patterns are associated with biomarkers in a selected population of university students and employees", Journal of Nutritional Science, 2012

Publication

---

<1 %

54

T.L. Felicio, E.A. Esmerino, A.G. Cruz, L.C. Nogueira, R.S.L. Raices, R. Deliza, H.M.A. Bolini, M.A.R. Pollonio. "Cheese. What is its contribution to the sodium intake of Brazilians?", Appetite, 2013

Publication

---

<1 %

55

Temple, N.J.. "The cost of a healthy diet: A South African perspective", Nutrition, 201105

Publication

---

<1 %

56	lib.ugent.be Internet Source	<1 %
57	sumsdbweb.shiga-med.ac.jp Internet Source	<1 %
58	"Nutrition Guide for Physicians and Related Healthcare Professions", Springer Science and Business Media LLC, 2022 Publication	<1 %
59	"Science and Technology of Fibers in Food Systems", Springer Science and Business Media LLC, 2020 Publication	<1 %
60	Celia A Prentice, Claire Smith, Rachael M McLean. "Sodium in commonly consumed fast foods in New Zealand: a public health opportunity", Public Health Nutrition, 2015 Publication	<1 %
61	Jang H. Youn, Young Taek Oh, Stefania Gili, Alicia A. McDonough, John Higgins. "Estimating In Vivo Potassium Distribution and Fluxes with Stable Potassium Isotopes", American Journal of Physiology-Cell Physiology, 2022 Publication	<1 %
62	Katherine M Livingstone, Gavin Abbott, Steven J Bowe, Joey Ward, Catherine Milte, Sarah A McNaughton. "Diet quality indices,	<1 %

genetic risk and risk of cardiovascular disease and mortality: a longitudinal analysis of 77 004 UK Biobank participants", *BMJ Open*, 2021

Publication

---

63

Ling Li, Zhenjuan Zhao, Yini Wang, Xueqin Gao, Guojie Liu, Bo Yu, Ping Lin. "Association Between Dietary Intakes and Plaque Vulnerability Measured by Optical Coherence Tomography in Patients With Coronary Heart Disease: A Mediation Analysis of Inflammatory Factors", *Frontiers in Nutrition*, 2022

Publication

---

<1 %

64

Marieke AH Hendriksen, Joop MA van Raaij, Johanna M Geleijnse, Caroline Wilson-van den Hooven et al. "Monitoring salt and iodine intakes in Dutch adults between 2006 and 2010 using 24 h urinary sodium and iodine excretions", *Public Health Nutrition*, 2013

Publication

---

<1 %

65

Qinyu Yang, Weihong Jiang, Yongmei He, Lin Yang, Congke Zhao, Lijun Li, Pingting Yang, Lu Yin, Xiaohui Li, Xin Huang, Ying Li. "The association of arterial stiffness with estimated excretion levels of urinary sodium and potassium and their ratio in Chinese adults", *Journal of Human Hypertension*, 2022

Publication

---

<1 %

66

S. C. Francisco, L. F. Araújo, R. H. Griep, D. Chor et al. "Adherence to the Dietary Approaches to Stop Hypertension (DASH) and hypertension risk: results of the Longitudinal Study of Adult Health (ELSA-Brasil)", *British Journal of Nutrition*, 2020

Publication

&lt;1 %

67

Samuel J. Dicken, John Joseph Mitchell, Jessica Newberry Le Vay, Emma Beard, Dimitra Kale, Aleksandra Herbec, Lion Shahab. "Impact of COVID-19 pandemic on diet behaviour among UK adults: a longitudinal analysis of the HEBECO study", *Cold Spring Harbor Laboratory*, 2021

Publication

&lt;1 %

68

Sangah Shin, Hyojee Joung. "A dairy and fruit dietary pattern is associated with a reduced likelihood of osteoporosis in Korean postmenopausal women", *British Journal of Nutrition*, 2013

Publication

&lt;1 %

69

Sheikh Mohammed Shariful Islam, Tina Dannemann Purnat, Nguyen Thi Anh Phuong, Upendo Mwingira et al. "Non - Communicable Diseases (NCDs) in developing countries: a symposium report", *Globalization and Health*, 2014

Publication

&lt;1 %

70	Y Al-Solaiman. "Low-sodium DASH reduces oxidative stress and improves vascular function in salt-sensitive humans", Journal of Human Hypertension, 04/30/2009 Publication	<1 %
71	eprints.gla.ac.uk Internet Source	<1 %
72	theses.dur.ac.uk Internet Source	<1 %
73	gedos.usal.es Internet Source	<1 %
74	lib.unisayogya.ac.id Internet Source	<1 %
75	mattioli1885journals.com Internet Source	<1 %
76	pdffox.com Internet Source	<1 %
77	www.icomst2009.dk Internet Source	<1 %
78	www.plosmedicine.org Internet Source	<1 %
79	Alfonso Mendoza, Ana E Pérez, Anju Aggarwal, Adam Drewnowski. "Energy density of foods and diets in Mexico and their monetary cost by socioeconomic strata:	<1 %



analyses of ENSANUT data 2012", Journal of Epidemiology and Community Health, 2017

Publication

80

[library.wur.nl](http://library.wur.nl)

Internet Source

<1 %

81

Donald J. Naismith, Alessandro Braschi. "An investigation into the bioaccessibility of potassium in unprocessed fruits and vegetables", International Journal of Food Sciences and Nutrition, 2009

Publication

<1 %

82

Julie M. Hess, Christopher J. Cifelli, Sanjiv Agarwal, Victor L. Fulgoni. "Comparing the cost of essential nutrients from different food sources in the American diet using NHANES 2011–2014", Nutrition Journal, 2019

Publication

<1 %

83

L. Azadbakht. "The Dietary Approaches to Stop Hypertension Eating Plan Affects C-Reactive Protein, Coagulation Abnormalities, and Hepatic Function Tests among Type 2 Diabetic Patients", Journal of Nutrition, 06/01/2011

Publication

<1 %

84

Salehi-Abargouei, Amin, Zahra Maghsoudi, Fatemeh Shirani, and Leila Azadbakht. "Effects of Dietary Approaches to Stop Hypertension (DASH)-style diet on fatal or

<1 %

nonfatal cardiovascular diseases—Incidence: A systematic review and meta-analysis on observational prospective studies", Nutrition, 2013.

Publication

---

85

"Proceedings of the Andalus International Public Health Conference 2017", BMC Public Health, 2017

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On

# Potassium intake is associated with nutritional quality and actual diet cost a study at formulating a low sodium high potassium (LSHP) healthy diet

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---