



1







4. If you find mistakes in the attached pdf document of the article,

submit them to us by email ... secretary4@jced.es
You have to describe, in the email, the line and page
of the pdf
document where you find the mistakes. Do not try to
add the
modifications in the pdf.

Please inform us within the next 5 days of receipt. We will not admit any change after those 5 days.

If you have any doubt let us know - E-mail to: secretary4@jced.es

Yours sincerely.

Norma Vidal Editorial Office

## **AUTHORS**

- Initials of first Name of 1 author: A.

THE SEC OF THE SEC OF S

Last

Name of 1

author: Rizqiawan

- Initials of first Name of 2 author: A.

Last

Name of 2

author: Zahratur-Rasyida

- Initials of first Name of 3 author: I.

Last

Name of 3

author: Mulyawan

## CORRESPONDENCE:

Department of Oral and Maxillofacial Surgery Faculty of Dental Medicine, Universitas Airlangga Jl. Mayjend Prof. Dr. Moestopo No. 47 Surabaya 60132, Indonesia



III



- Initials of first Name of 3 author: I. Last

Name of 3

author: Mulyawan

## CORRESPONDENCE:

Department of Oral and Maxillofacial Surgery Faculty of Dental Medicine, Universitas Airlangga Jl. Mayjend Prof. Dr. Moestopo No. 47 Surabaya 60132, Indonesia

TITLE: Transoral surgical approach to solitary fibrous tumors in buccal space with infratemporal tumor extension: A case report

## ABSTRACT:

A solitary fibrous tumour (SFT) is a rare spindle-cell neoplasm of

mesenchymal origin usually located in the pleura. It has been recently

described as occurring in various head and neck sites, including the

oral cavity. The purpose of this article is to report a case of SFT

originating in the buccal space and extending into the infratemporal

space treated by means of transoral approach surgery. A 25-year-old

female patient reported to the Department of Oral and Maxillofacial

Surgery, Universitas Airlangga Hospital, chiefly complaining of a

painless lump in the left cheek which had been present for nine months.

The diagnosis was arrived at on the basis of a combination of clinical

investigation, imaging studies and histopathological examination

(biopsy). The surgical approach involved transoral incision through the

