

Bisindole alkaloids from Voacanga grandifolia leaves

by Idha Kusumawati

Submission date: 19-Apr-2021 10:49AM (UTC+0800)

Submission ID: 1563045223

File name: Jurnal_C-1.pdf (1.11M)

Word count: 4085

Character count: 18768



Bisindole alkaloids from *Voacanga grandifolia* leaves


Alfarius Eko Nugroho¹ · Yurika Ono¹ · Eunji Jin¹ · Yusuke Hirasawa¹ · Toshio Kaneda¹ · Abdul Rahman¹ · Idha Kusumawati² · Takahiro Tougan³ · Toshihiro Horii⁴ · Noor Cholies Zaini² · Hiroshi Morita¹

Received: 26 October 2020 / Accepted: 7 December 2020 / Published online: 19 January 2021
© The Japanese Society of Pharmacognosy 2021

Abstract

Two new bisindole alkaloids, 12'-*O*-demethyl-vobtusine-5-lactam and isovobtusine-*N*-oxide (**1** and **2**), were isolated from the leaves of *Voacanga grandifolia*, together with two known bisindole alkaloids. Their structures were elucidated on the basis of 1D and 2D NMR data. **1** and **2** showed potent antimalarial activity against *Plasmodium falciparum* 3D7 and very low cytotoxic activity against a human cell line, HepG2 cells.

Graphical abstract



Keywords *Voacanga grandifolia* · Bisindole alkaloids · Apocynaceae · Antimalarial activity

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11418-020-01475-w>.

✉ Hiroshi Morita
moritah@hoshi.ac.jp

- ¹ Faculty of Pharmaceutical Sciences, Hoshi University, Ebara 2-4-41 Shinagawa-ku, Tokyo 142-8501, Japan
- ² Faculty of Pharmacy, Airlangga University, Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia
- ³ Research Center for Infectious Disease Control, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamadaoka, Suita, Osaka 565-0871, Japan
- ⁴ Department of Molecular Protozoology, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamadaoka, Suita, Osaka 565-0871, Japan

Introduction

Voacanga is a small genus of the Apocynaceae family consisting of 12 species. Species of this genus are distributed mainly in the tropical Africa and Malaysia, and have been reported to contain vobasine, eburnane, iboga, and aspidoasperma type of monoterpene indole alkaloids [1]. Various activities have been reported for monoterpene indole alkaloids, such as cytotoxic activity [2–4], anti-melanogenesis [5], anti-plasmodial [6], and vasorelaxant activities [7]. In the search for new bioactive compounds from tropical plants [8–15], alkaloid constituents of *V. grandifolia* leaves were investigated and two new bisindole alkaloids, 12'-*O*-demethyl-vobtusine-5-lactam and isovobtusine-*N*-oxide (**1** and **2**,

Fig. 1), were isolated together with vobtusine-*N*-oxide [16], and vobtusine-3-lactam [17]. The isolation and structure elucidation of **1** and **2** consisting of bis-aspidosperma-type skeleton, from the leaves of *Voacanga grandifolia*, and their antimalarial activities are reported herein.

Results and discussions

Compound **1** $\{[\alpha]_D^{24} - 117.6^\circ (c 1.0, \text{MeOH})\}$ was isolated as an optically active, yellow amorphous solid. The molecular formula was determined by HRESIMS (m/z 719.3453 $[\text{M} + \text{H}]^+$, $\Delta + 0.8$ mmu) as $\text{C}_{42}\text{H}_{46}\text{N}_4\text{O}_7$, and the IR absorption at 3340 cm^{-1} and 1715 cm^{-1} indicated the presence of hydroxy and carbonyl groups, respectively. Analysis of the ^1H and ^{13}C NMR data (Table 1) and the HSQC spectrum of **1** revealed the presence of six sp^3 quaternary carbons, five sp^3 methines, fourteen sp^3 methylenes, one methyl, seven sp^2 methines, and nine sp^2 quaternary carbons. Among them, two sp^3 methines ($\delta_{\text{C}} 63.6$; $\delta_{\text{H}} 2.60$, $\delta_{\text{C}} 64.9$; $\delta_{\text{H}} 3.97$) and four sp^3 methylenes ($\delta_{\text{C}} 44.5$; $\delta_{\text{H}} 2.85$ and 4.16 , $\delta_{\text{C}} 44.6$; $\delta_{\text{H}} 3.13$ and 4.82 , $\delta_{\text{C}} 48.8$; $\delta_{\text{H}} 2.29$ and 2.84 , and $\delta_{\text{C}} 52.0$; $\delta_{\text{H}} 2.21$ and 3.02) ascribed to those attached to a nitrogen atom.

The gross structure of **1** was elucidated by analysis of 2D NMR data including the ^1H - ^1H COSY, TOCSY, HSQC, and HMBC spectra in CDCl_3 (Fig. 2). The ^1H - ^1H COSY spectrum revealed the connectivity of seven partial structures **a** (C-9' ~ C-11'), **b** (C-5' ~ C-6'), **c** (C-3', C-14' ~ C-15'), **d** (C-16' ~ C-17', C-22'), **e** (C-18' ~ C-19'), **f** (C-18 ~ C-19), and **g** (C-9 ~ C-12) as shown in Fig. 2. These partial structures were classified into two units, **a** ~ **e** and **f** ~ **g**.

The connectivity of partial structures **b** and **c** and C-21' through a nitrogen atom was revealed by the HMBC

correlations of H-5'b and H-21' to C-3' ($\delta_{\text{C}} 48.8$). The HMBC correlations from H-18'a and H-19'a to C-15' ($\delta_{\text{C}} 80.5$), H-14'a to C-20' ($\delta_{\text{C}} 44.0$), and H-19'b to C-17' ($\delta_{\text{C}} 32.6$) and C-21' ($\delta_{\text{C}} 63.6$) established the connections among C-15', C-17', C-19', and C-21' through C-20'.

In addition, HMBC cross peaks of H-6'b to C-8' ($\delta_{\text{C}} 134.9$), H-9' to C-21', H-21' to C-2' ($\delta_{\text{C}} 93.5$) and C-6' ($\delta_{\text{C}} 31.2$), and H-16' to C-7' ($\delta_{\text{C}} 55.9$) revealed the connectivity between C-6', C-16', C-21', and an indoline ring. Then, the HMBC correlations of H-23'b and OH-12' to C-13' ($\delta_{\text{C}} 138.2$), OH-12' to C-12' ($\delta_{\text{C}} 139.8$), and H-22'a and H-23'a to C-2' suggested the presence of hydroxyl groups at C-2' and C-12' in the indoline ring. These NMR data were characteristic of an aspidosperma-type skeleton.

On the other hand, the similar aspidosperma-type skeleton with a methoxy carbonyl at C-16 and a double bond between C-2 and C-16 was suggested by the HMBC correlations in the units **f** and **g** as shown in Fig. 2. The connection of two indole alkaloids through C-14 was revealed by the HMBC correlations from H-23'a to C-14 ($\delta_{\text{C}} 38.1$) and C-22' ($\delta_{\text{C}} 35.2$) and from H-23'b to C-3 ($\delta_{\text{C}} 44.5$) and C-15 ($\delta_{\text{C}} 87.8$). In particular, the HMBC correlations of H-3a, H₂-6 and H-21 to C-5 ($\delta_{\text{C}} 172.6$) confirmed the presence of a carbonyl at C-5. These data suggested the structure of **1** as 12'-*O*-demethyl-vobtusine-5-lactam.

The relative configuration of each monoterpene indole unit in **1** was assigned by NOESY correlations as shown in computer-generated 3D drawing (Fig. 3). In **1a** (partial structures **a** ~ **e**), the NOESY correlations of H-21'/H-18'a and H-9', and H-15'/H-18'b and H-17'b suggested that both C-21' and C-15' should be *S** and C-7' and C-20' should be *R**. The relative configuration of C-2' ($\delta_{\text{C}} 93.5$) was deduced to be *R** based on the similarity with the chemical shift ($\delta_{\text{C}} 93.7$) of vobtusine [17]. Stereochemistry of H-16' was elucidated to be α -oriented by $^3J_{\text{H-16'/H-17'a}}$ (13.0 Hz) and $^3J_{\text{H-16'/H-22'b}}$ (13.4 Hz), and the NOESY correlations in **1a**. In **1b** (partial structures **f** ~ **g**), H-15 is β -oriented and C-20 should be *R** by the NOESY correlations of H-21/H-18 and H-19a. Stereochemistry of a spiro carbon (C-14) in **1c** should be assigned as *R** by the NOESY correlations, suggesting that the diazaspiro [5.5] undecane ring took a chair conformation, as shown in Fig. 3. Finally, the NOESY correlations and ^1H - ^1H coupling constant values suggested the relative configuration of **1** to be, as shown in Fig. 1, the same as vobtusine-3-lactam.

Compound **2** $\{[\alpha]_D^{24} - 123.6^\circ (c 1.0, \text{MeOH})\}$ was isolated as an optically active, yellow amorphous solid. The molecular formula was determined by HRESIMS (m/z 735.3765 $[\text{M} + \text{H}]^+$, $\Delta 0.7$ mmu) as $\text{C}_{43}\text{H}_{50}\text{N}_4\text{O}_7$, and the IR absorption at 3340 cm^{-1} and 1685 cm^{-1} indicated the presence of hydroxy and carbonyl groups, respectively. In addition, the ^{13}C NMR data of **2** (Table 1) were similar to those of vobtusine-*N*-oxide and isovobtusine [16, 17]. The





Fig. 1 Structures of isolated compounds (**1** and **2**) from *Voacanga grandifolia*

Table 1. ^1H (600 MHz) and ^{13}C (150 MHz) NMR data of **1** and **2** in CDCl_3

	1		2	
	δ_{H} (J, Hz)	δ_{C}	δ_{H} (J, Hz)	δ_{C}
2		164.4		167.5
3a	4.16 (1H, d, 13.0)	44.5	2.68 (1H, d, 10.8)	58.2
3b	2.85 (1H, d, 13.0)		2.64 (1H, d, 10.8)	
5a		172.6	2.89 (1H, m)	51.5
5b			2.74 (1H, m)	
6a	2.97 (1H, d, 18.0)	48.3	2.01 (1H, m)	44.4
6b	2.65 (1H, d, 18.0)		1.70 (1H, m)	
7		47.4		55.1
8		135.7		138.1
9	7.12 (1H, d, 7.5)	121.2	7.20 (1H, d, 7.2)	121.7
10	6.94 (1H, dd, 7.6, 7.5)	121.8	6.87 (1H, dd, 7.6, 7.2)	120.7
11	7.21 (1H, dd, 7.8, 7.6)	129	7.12 (1H, dd, 7.7, 7.6)	127.7
12	6.86 (1H, d, 7.8)	109.8	6.77 (1H, d, 7.7)	109.3
13		142.9		143
14		38.1		39.2
15	3.42 (1H, s)	87.8	3.17 (1H, s)	81.8
16		94		94.5
17a	2.52 (1H, s)	26.1	2.81 (1H, d, 13.8)	28.5
17b	2.52 (1H, s)		2.20 (1H, d, 13.8)	
18a	3.72 (1H, m)	63.9	3.65 (1H, m)	63.6
18b	3.72 (1H, m)		3.30 (1H, m)	
19a	1.56 (1H, m)	35.9	1.26 (1H, m)	35
19b	1.39 (1H, ddd, 13.0, 7.9, 4.8)		1.17 (1H, m)	
20		46.5		47.9
21	3.97 (1H, s)	64.9	2.83 (1H, s)	70
CO		168		168.4
OMe	3.80 (3H, s)	51.4	3.71 (3H, s)	51
2'		93.5		92.3
3'a	2.84 (1H, m)	48.8	3.45 (1H, m)	58.7
3'b	2.29 (1H, m)		3.45 (1H, m)	
5'a	3.02 (1H, m)	52	3.52 (1H, m)	67.2
5'b	2.21 (1H, m)		3.32 (1H, m)	
6'a	2.79 (1H, m)	31.2	3.58 (1H, m)	30.7
6'b	1.24 (1H, m)		1.36 (1H, m)	
7'		55.9		55.6
8'		134.9		133.6
9'	6.74 (1H, d, 7.3)	116	6.64 (1H, d, 7.6)	111.1
10'	6.61 (1H, dd, 7.3, 7.2)	119.1	6.62 (1H, dd, 7.6, 7.6)	118.1
11'	6.72 (1H, d, 7.2)	121.6	6.59 (1H, d, 7.6)	112.5
12'		139.8		146.3
13'		138.2		137.7
14'a	1.95 (1H, br d, 12.0)	25.8	2.92 (1H, m)	22.8
14'b	1.86 (1H, m)		2.01 (1H, m)	
15'	3.44 (1H, br s)	80.5	3.66 (1H, dd, 2.9, 2.9)	78.7
16'	1.87 (1H, m)	31.6	2.03 (1H, m)	33.9
17'a	1.78 (1H, dd, 13.0, 13.0)	32.6	2.56 (1H, dd, 12.5, 12.5)	33.7
17'b	0.86 (1H, br d, 13.0)		0.91 (1H, br d, 12.5)	
18'a	4.05 (1H, m)	65.4	3.98 (1H, m)	65
18'b	3.90 (1H, m)		3.90 (1H, m)	

Table 1. (continued)

	1		2	
	δ_{H} (J, Hz)	δ_{C}	δ_{H} (J, Hz)	δ_{C}
19'a	2.35 (1H, m)	36.5	2.48 (1H, ddd, 12.5, 8.1, 3.9)	39.7
19'b	1.52 (1H, m)		1.69 (1H, m)	
20'		44.0		45.4
21'	2.60 (1H, s)	63.6	3.56 (1H, s)	75.3
22'a	1.48 (1H, br d, 13.6)	35.2	2.00 (1H, m)	35
22'b	2.13 (1H, dd, 13.6, 13.4)		2.00 (1H, m)	
23'a	4.82 (1H, d, 14.5)	44.6	4.69 (1H, d, 14.1)	46.7
23'b	3.13 (1H, d, 14.5)		3.02 (1H, d, 14.1)	
12'-OH	6.22 (1H, s)			
12'-OMe			3.75 (3H, s)	54.9

Fig. 2 Selected 2D NMR correlations of **1**

chemical shift differences between **2** and vobtusine-*N*-oxide (i.e. C-3, C-14, C-15, C-18, C-16', C-22' and C-23') suggested **2** to be the 14-*epi* derivative of vobtusine-*N*-oxide, isovobtusine-*N*-oxide, and the downfield shifts of signals ascribed to C-3', C-5' and C-21' in **2** relative to isovobtusine further supported this conclusion.

Biological activity of **1** and **2**

The antimalarial activity and cytotoxic activity of **1** and **2** were evaluated in vitro. Against *P. falciparum* 3D7 strain, **1** and **2** showed potent antimalarial activity [Table 2: half-maximal (50%) inhibitory concentration (IC_{50}) = 5.1 and 3.3 μM , respectively]. Chloroquine was used as a positive control (IC_{50} = 0.026 μM). In contrast, the cytotoxic activity of both **1** and **2** was low [Table 2: half-maximal (50%) cytotoxic concentration (CC_{50}) \geq 50 μM] against the human cell line, HepG2 cells.

Experimental section

General experimental procedures

Optical rotations were measured on a JASCO DIP-1000 polarimeter, UV spectra on a Shimadzu UVmini-1240 spectrophotometer, and IR spectra on a JASCO FT/IR-4100 spectrophotometer. High-resolution ESI MS were obtained on a JMS-T100LP (JEOL). ^1H and 2D NMR spectra were measured on a 600 MHz spectrometer at 300 K, and ^{13}C NMR spectra on a 150 MHz spectrometer. Residual solvent chemical shifts were used as internal standard, δ_{H} 7.26 and δ_{C} 77.0 for CHCl_3 . Standard pulse sequences were used for the 2D NMR experiments.

Material

The leaves of *V. grandiflora* were collected at the Purwodadi Botanical Garden in March 2008. The botanical identification was made by Ms. Sri Wuryanti, Purwodadi Botanical Garden. A voucher specimen has been deposited in the herbarium at Purwodadi Botanical Garden, Pasuruan, Indonesia.

Extraction and isolation

Finely powdered leaves of *V. grandiflora* (400 g) were extracted with MeOH (1L \times 3) at room temperature. The combined extracts were evaporated under reduced pressure to give a residue (30 g). The extract was dissolved in 3% aqueous tartaric acid (pH 2) and then partitioned with EtOAc. The aqueous layer was treated with Na_2CO_3 to pH 9 and was partitioned successively by CHCl_3 and *n*-BuOH. The CHCl_3 soluble materials (3.9 g) was subjected to an LH-20 column ($\text{CHCl}_3/\text{MeOH}$, 1:1) to obtain ten fractions.




Fig. 3 Selected NOESY correlations of **1**

Table 2 Antimalarial and cytotoxic activities of **1** and **2** against *Plasmodium falciparum* 3D7

Compound	Antimalarial activity		Cytotoxic activity
	IC ₅₀ (μM)	CC ₅₀ (μM)	
1	5.1	> 50	
2	3.3	> 50	

Fraction 7 (360 mg) was fractionated by silica gel column chromatography (CHCl₃/MeOH, 0:1 → 1:0), ODS column chromatography (MeOH/H₂O) and RP-HPLC (Nacalai tesque Cholester, 10 × 250 mm; 24% MeCN in 0.1% aqueous HCO₂H; flow rate 3.5 mL/min; UV detection at 254 nm) to obtain **1** (1.5 mg, 0.0004%), **2** (1.5 mg, 0.0004%), vobtusine-*N*-oxide (1.9 mg, 0.0005%) [16], and vobtusine-3-lactam (0.8 mg, 0.0002%) [17].

12'-demethyl-vobtusine-5-lactam (**1**): Yellow amorphous solid; [α] -117.6 (c 1.0, MeOH); IR (film) ν_{\max} 3340 and 1715 cm⁻¹; UV/Vis λ_{\max} (MeOH) nm (log ε): 306 (3.57), 273 (3.84), 227 (4.41) and 206 (4.41) nm; ESIMS (pos.) *m/z* 719 (M + H)⁺; HRMS-ESI: *m/z* [M + H]⁺ calcd for C₄₂H₄₇N₄O₇: 719.3445; found 719.3453.

Isovobtusine-*N*-oxide (**2**): Yellow amorphous solid; [α] -123.6 (c 1.0, MeOH); IR (film) ν_{\max} 3370 and 1685 cm⁻¹; UV/Vis λ_{\max} (MeOH) nm (log ε): 328 (4.22), 299 (4.16), 259 (3.93) and 220 (4.47) nm; ESIMS (pos.) *m/z* 735 (M + H)⁺; HRMS-ESI: *m/z* [M + H]⁺ calcd for C₄₃H₅₁N₄O₇: 735.3758; found 735.3765.

Parasite culture

P. falciparum laboratory strain 3D7 was obtained from Prof. Masatsugu Kimura (Osaka City University, Osaka, Japan). For the assessment of antimalarial activity of the compounds in vitro, the parasites were cultured in Roswell Park

Memorial Institute (RPMI) 1640 medium supplemented with 0.5 g/L L-glutamine, 5.96 g/L HEPES, 2 g/L sodium bicarbonate (NaHCO₃), 50 mg/L hypoxanthine, 10 mg/L gentamicin, 10% heat-inactivated human serum, and red blood cells (RBCs) at a 3% hematocrit in an atmosphere of 5% CO₂, 5% O₂, and 90% N₂ at 37 °C as previously described [18]. Ring-form-infected RBCs were collected using the sorbitol synchronization technique [19]. Briefly, the cultured cells were collected by centrifugation at 840 g for 5 min at room temperature, suspended in a fivefold volume of 5% D-sorbitol (Nacalai Tesque, Kyoto, Japan) for 10 min at room temperature, and then they were washed twice with RPMI 1640 medium to remove the D-sorbitol. The utilization of blood samples of healthy Japanese volunteers for the parasite culture was approved by the institutional review committee of the Research Institute for Microbial Diseases (RIMD), Osaka University (approval number: 22–3).

Antimalarial activity

Ring-form-synchronized parasites were cultured with compounds **1** and **2** at sequentially decreasing concentrations (50, 15, 5, 1.5, 0.5, 0.15, 0.05, and 0.015 μM) for 48 h for the flow cytometric analysis using an automated hematology analyzer, XN-30. The XN-30 analyzer was equipped with a prototype algorithm for cultured falciparum parasites [prototype; software version: 01–03, (build 16)] and used specific reagents (CELLPACK DCL, SULFOLYSER, Lysercell M, and Fluorocell M) (Sysmex, Kobe, Japan) [20,21]. Approximately 100 μL of the culture suspension diluted with 100 μL phosphate-buffered saline was added to a BD Microtainer MAP Microtube for Automated Process K2 EDTA 1.0 mg tube (Becton Dickinson and Co., Franklin Lakes, NJ, USA) and loaded onto the XN-30 analyzer with an auto-sampler as described in the instrument manual (Sysmex). The parasitemia shown as MI-RBC% was automatically reported [20]. Then 0.5% DMSO alone or containing 5 μM artemisinin used as the negative and positive controls, respectively. The growth inhibition (GI) rate was calculated from the MI-RBC% according to the following equation:

$$GI(\%) = 100 - \frac{(\text{test sample} - \text{positive control})}{(\text{negative control} - \text{positive control})} \times 100$$

The IC₅₀ was calculated from GI (%) using GraphPad Prism version 5.0 (GraphPad Prism Software, San Diego, CA, USA) [22].

Cytotoxic activity

HepG2 (JCRB1054) cell line was obtained from the Japanese Collection of Research Bioresources (JCRB, Osaka,

Japan). The cells were cultured in Dulbecco's modified Eagle's medium [DMEM (1.0 g/L glucose) with L-glutamine and sodium pyruvate; Nacalai Tesque, Kyoto, Japan] supplemented with 10% (v/v) fetal bovine serum (FBS; Gibco-BRL, Grand Island, NY, USA) in a humidified incubator with 5% CO₂ at 37 °C. For the assay, the cells (5 × 10³/well) were seeded in a 96-well plate. **1** and **2** at gradually decreasing concentrations (50, 15, 5, 1.5, 0.5, 0.15, 0.05, and 0.015 μM) were added to the cell culture after 24 h and the cells were subsequently cultured for 48 h. Cell viability was measured using a Cell Counting Kit-8 (Dojindo, Kumamoto, Japan) according to the manufacturer's instructions. Briefly, 10 μL CCK-8 reagent was added to each well containing culture medium and incubated for 2 h under standard culture conditions. The absorbance of the sample was measured at 450 nm using a PowerWave HT microplate spectrophotometer (BioTek Instruments, Winooski, VT, USA). The cell viability was expressed as a percentage of the absorbance of the untreated control cells after subtracting the appropriate background intensity. The CC₅₀ was calculated from the cell viability using GraphPad Prism version 5.0.

Acknowledgment We thank Prof. Masatsugu Kimura (Osaka City University, Osaka, Japan) for the kind gift of the 3D7 strain. We also thank Mr. Yuji Toya and Dr. Kinya Uchihashi (Sysmex, Kobe, Japan) for the setting of the XN-30 analyzer and Ms. Toshie Ishisaka and Ms. Sawako Itagaki for their technical assistance. This work was supported by JSPS KAKENHI Grant Number JP19K07152 to MH and partially supported by Sysmex Corporation to TT.

References

- Macabeo APG, Alejandro GJD, Hallare AV, Vidar WS, Villaflores OB (2009) Phytochemical Survey and Pharmacological Activities of the Indole Alkaloids in the Genus *Voacanga* Thouars (Apocynaceae) - An Update. *Phcog Rev* 3:132–142
- Hirasawa Y, Shoji T, Arai T, Nugroho AE, Deguchi J, Hosoya T, Uchiyama N, Goda Y, Awang K, Hadi AHA, Shiro M, Morita H (2010) Bisleuconothine a, an eburnane-aspidosperma bisindole alkaloid from *Leuconotis griffithii*. *Bioorg Med Chem Lett* 20:2021–2024
- Haseo A, Nugroho AE, Hirasawa Y, Kaneda T, Shirota O, Rachman A, Kusumawati I, Zaini NC, Morita H (2015) A new indole alkaloids from *Voacanga grandifolia*. *Heterocycles* 90:601–606
- Wong CP, Seki A, Horiguchi K, Shoji T, Arai T, Nugroho AE, Hirasawa Y, Sato F, Kaneda T, Morita H (2015) Bisleuconothine a induces autophagosome formation by interfering with AKT-mTOR signaling pathway. *J Nat Prod* 78:1656–1662
- Koyama K, Hirasawa Y, Hosoya T, Hoe TC, Chan K-L, Morita H (2010) Alpneumines A-H, new anti-melanogenic indole alkaloids from *Alstonia pneumatophora*. *Bioorg Med Chem* 18:4415–4421
- Nugroho AE, Sugai M, Hirasawa Y, Hosoya T, Awang K, Hadi AHA, Ekasari W, Widyawaruyanti A, Morita H (2011) New anti-plasmodial indole alkaloids from *Hunteria zeylanica*. *Bioorg Med Chem Lett* 21:3417–3419
- Hirasawa Y, Hara M, Nugroho AE, Sugai M, Zaima K, Kawahara N, Goda Y, Awang K, Hadi AHA, Litaudon M, Morita H (2010) Bisnicalaterines B and C, atropisomeric bisindole alkaloids from

- hunteria zeylanica*, showing vasorelaxant activity. *J Org Chem* 75:4218–4223
8. Nugroho AE, Hashimoto A, Wong CP, Yokoe H, Tsubuki M, Kaneda T, Hadi AHA, Morita H (2018) Ceramicines M-P from *Chisocheton ceramicus*: isolation and structure–activity relationship study. *J Nat Med* 72:64–72
 9. Nugroho AE, Inoue D, Wong CP, Hirasawa Y, Kaneda T, Shirota O, Hadi AHA, Morita H (2018) Reinereins A and B, new onocerane triterpenoids from *Reinwardtiidendron cinereum*. *J Nat Med* 72:588–592
 10. Kaneda T, Matsumoto M, Sotozono Y, Fukami S, Nugroho AE, Hirasawa Y, Hamid AHA, Morita H (2019) Cycloartane triterpenoid (23R, 24E)-23-acetoxymangiferonic acid inhibited proliferation and migration in B16–F10 melanoma via MITF downregulation caused by inhibition of both β -catenin and c-Raf–MEK1–ERK signaling axis. *J Nat Med* 73:47–58
 11. Tang Y, Nugroho AE, Hirasawa Y, Tougan T, Horii T, Hamid AHA, Morita H (2019) Leucophyllinines A and B, Bisindole alkaloids from *Leuconotis eugenifolia*. *J Nat Med* 73:533–540
 12. Amelia P, Nugroho AE, Hirasawa Y, Kaneda T, Tougan T, Horii T, Morita H (2019) Indole alkaloids from *Tabernaemontana macrocarpa* Jack. *J Nat Med* 73:820–825
 13. Hirasawa Y, Dai X, Deguchi J, Hatano S, Ohtsuka R, Nugroho AE, Kaneda T, Morita H (2019) New Vasorelaxant Indole Alkaloids, Taberniacins A and B, from *Tabernaemontana divaricata*. *J Nat Med* 73:627–632
 14. Prema WCP, Awouafack MD, Nugroho AE, Win YY, Win NN, Ngwe H, Morita H, Morita H (2019) Two new quassinoids and other constituents from the *Picrasma javanica* wood and their biological activities. *J Nat Med* 73:589–596
 15. Prema WCP, Kodama T, Nugroho AE, El-Desoky AH, Awouafack MD, Win YY, Ngwe H, Abe I, Morita H, Morita H (2020) Three new quassinoids isolated from the wood of *Picrasma javanica* and their anti-Vpr activities. *J Nat Med* 74:571–578
 16. Agwada VC, Naranjo J, Hesse M, Schmid H, Rolland Y, Kunesch N, Poisson J, Chatterjee A (1977) Die struktur des bisindolalkaloides amatain (= grandifolin, subessilin). *Helv Chim Acta* 60:2830–2853
 17. Rolland Y, Kunesch N, Poisson J, Hagaman EW, Schell FM, Wenkert E (1976) *Voacanga* alkaloids. 16. Carbon-13 nuclear magnetic resonance spectroscopy of naturally occurring substances. 43. Carbon-13 nuclear magnetic resonance analysis of bis-indoline alkaloids of two *Voacanga* species. *J Org Chem* 41:3270–3275
 18. Trager W, Jensen JB (1976) Human malaria parasites in continuous culture. *Science* 193:673–675
 19. Lambros C, Vanderberg JP (1979) Synchronization of *Plasmodium falciparum* erythrocytic stages in culture. *J Parasitol* 65:418–420
 20. Tougan T, Suzuki Y, Itagaki S, Izuka M, Toya Y, Uchihashi K, Horii T (2018) An automated haematology analyzer XN-30 distinguishes developmental stages of *falciparum* malaria parasite cultured in vitro. *Malar J* 17:59
 21. Toya T, Tougan T, Horii T, Uchihashi K (2021) Lysercell M enhances the detection of stage-specific *Plasmodium*-infected red blood cells in the automated hematology analyzer XN-31 prototype. *Parasitol Int* 80:102206. <https://pubmed.ncbi.nlm.nih.gov/33049417/>
 22. Tougan T, Toya Y, Uchihashi K, Horii T (2019) Application of the automated haematology analyzer XN-30 for discovery and development of anti-malarial drugs. *Malar J* 18:8

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Bisindole alkaloids from Voacanga grandifolia leaves

ORIGINALITY REPORT

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Jin-Biao Xu, Yuan Lin, Shi-Hui Dong, Fei Wang, Jian-Min Yue. " Trichinenlides A-T, Mexicanolide-Type Limonoids from ", Journal of Natural Products, 2013
Publication 1%
- 2 Hosoya, T.. "Five novel flavonoids from Wasabia japonica", Tetrahedron, 20050718
Publication 1%
- 3 www.mdpi.com
Internet Source 1%
- 4 crossmark.crossref.org
Internet Source 1%
- 5 Fei Liu, Xing-De Wu, Juan He, Xu Deng, Li-Yan Peng, Huai-Rong Luo, Qin-Shi Zhao. "Casuarines A and B, Lycopodium alkaloids from Lycopodium casuarinoides", Tetrahedron Letters, 2013
Publication 1%
- 6 test.dovepress.com
Internet Source 1%

7	www.reproduction-online.org Internet Source	1 %
8	www.spandidos-publications.com Internet Source	1 %
9	Jing-Yu Su. "New tetracyclic diterpenoid and new ceramides from the soft coral <i>Sinularia conferta</i> ", <i>Journal of Asian Natural Products Research</i> , 4/1/2005 Publication	1 %
10	tcr.amegroups.com Internet Source	1 %
11	www.readbag.com Internet Source	1 %
12	mafiadoc.com Internet Source	<1 %
13	www.thieme-connect.com Internet Source	<1 %
14	Fukuyama, Y.. "Neurotrophic sesquiterpene-neolignans from <i>magnoliaobovata</i> : structure and neurotrophic activity", <i>Tetrahedron</i> , 1992 Publication	<1 %
15	Shagufta Perveen, Sher Bahadar Khan, Abdul Malik, Rasool Bakhsh Tareen, Sarfraz Ahmad Nawaz, Muhammad Iqbal Choudhary. "Phenolic constituents from ", <i>Natural Product Research</i> , 2006	<1 %

16

[e-sciencecentral.org](https://www.e-sciencecentral.org)

Internet Source

<1 %

17

[genecelltissue.com](https://www.genecelltissue.com)

Internet Source

<1 %

18

Wen-Fang Yi, Xiao Ding, Yuan-Zhi Chen, Tiwalade A. Adelakun, Yu Zhang, Xiao-Jiang Hao. " Tabernaesines A–I, Cytotoxic Aspidosperma–Aspidosperma-Type Bisindole Alkaloids from ", Journal of Natural Products, 2020

Publication

<1 %

19

data.mendeley.com

Internet Source

<1 %

20

Fivelman, Q.L.. "Improved synchronous production of Plasmodium falciparum gametocytes in vitro", Molecular & Biochemical Parasitology, 200707

Publication

<1 %

21

Hitotsuyanagi, Y.. "A structure-activity relationship study of brusatol, an antitumor quassinoid", Tetrahedron, 20060424

Publication

<1 %

22

Zhang, Z.. "Cytotoxic triterpenoid saponins from the fruits of Aesculus pavia L.", Phytochemistry, 200708

Publication

<1 %

23

jpet.aspetjournals.org

Internet Source

<1 %

24

www.tandfonline.com

Internet Source

<1 %

25

Xiaofeng Shi, Quhuan Ma, Zhenyu Yao, Junmin Zhang, Dongyan Liu, Wei Shen, Bin Fan, Xindi Wang. "Further iridoids from the roots of *Patrinia scabra*", *Phytochemistry Letters*, 2015

Publication

<1 %

26

Ai-Qun Lin. "iso- α -Cyclopiazonic acid, a new natural product isolated from the marine-derived fungus *Aspergillus flavus* C-F-3", *Chemistry of Natural Compounds*, 11/17/2009

Publication

<1 %

27

Honzawa, S.. "The 2@-a-(3-hydroxypropyl) group as an active motif in vitamin D³ analogues as agonists of the mutant vitamin D receptor (Arg274Leu)", *Bioorganic & Medicinal Chemistry*, 20080315

Publication

<1 %

28

JINYANG YUAN. "Peroxisome proliferator-activated receptor gamma is frequently underexpressed in renal cell carcinoma", *International Journal of Urology*, 3/2006

Publication

<1 %

29

Meng, Y.. "Flavones and flavone glycosides from *Halophila johnsonii*", *Phytochemistry*,

<1 %

30

Uttpal Anand, Samapika Nandy, Avinash Mundhra, Neela Das, Devendra Kumar Pandey, Abhijit Dey. "A review on antimicrobial botanicals, phytochemicals and natural resistance modifying agents from Apocynaceae family: Possible therapeutic approaches against multidrug resistance in pathogenic microorganisms", Drug Resistance Updates, 2020

Publication

<1 %

31

[Www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

32

univoak.eu

Internet Source

<1 %

33

www.frontiersin.org

Internet Source

<1 %

34

Sunita Gamre, Mrityunjay Tyagi, Sucheta Chatterjee, Birija S. Patro, Subrata Chattopadhyay, Dibakar Goswami. " Synthesis of Bioactive Diarylheptanoids from and Their Mechanism of Action for Anticancer Properties in Breast Cancer Cells ", Journal of Natural Products, 2021

Publication

<1 %

35

Boyle, S.. "Rational design of high affinity tachykinin NK² receptor antagonists", Bioorganic & Medicinal Chemistry, 199402

Publication

<1 %

36

Keiko Nishizawa, Kenta Teruya, Ayumi Oguma, Yuji Sakasegawa, Hermann Schätzl, Sabine Gilch, Katsumi Doh-ura. "Preparation and Characterization of Cellulose Ether Liposomes for the Inhibition of Prion Formation in Prion-Infected Cells", Journal of Pharmaceutical Sciences, 2019

Publication

<1 %

37

M. Ishibashi. "Isolation of Bioactive Natural Products from Myxomycetes", Medicinal Chemistry, 2005

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Bisindole alkaloids from Voacanga grandifolia leaves

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
