

## Editorial Team

### Editor in Chief

#### Prof R K Sharma

Former Head, Department of Forensic medicine & Toxicology

All-India Institute of Medical Sciences, New Delhi, India

E-mail: [editor.ijfmt@gmail.com](mailto:editor.ijfmt@gmail.com)

### Editor

#### Prof. Adarsh Kumar

Forensic Medicine & Toxicology,

All-India Institute of Medical Sciences, New Delhi, India

### EDITORIAL ADVISORY BOARD

1. Prof Sudhir K Gupta, Head, FMT. AIIMS, New Delhi , India
2. Prof Mete Gulmen ,Cukurova University, TURKEY
3. Prof. Leandro Duarte De Carvalho , Minas Gerais, Belo Horizonte, BRAZIL
4. Dr. Valery Gunas, National Pirogov Memorial Medical University,Vinnytsya, UKRAINE
5. Prof T. Nataraja Moorthy , Forensic Science, Faculty of Health & Life Sciences ,Management & Science University, MALAYSIA
5. Dr. Rahul Pathak Forensic Science, Dept of Life Sciences ,Anglia Ruskin University, Cambridge, UNITED KINGDOM
6. Prof Emilo Nuzzalese, University of Turin , Italy
7. Dr Noha A. Magdy Elrafie, Forensic Toxicology, Ain Shams University, Cairo, EGYPT
8. Dr Rituja Sharma, Associate Prof, Law, Banasthali Vidyapeeth Jaipur
9. Dr Shankar Bakkanwar (*Associate Professor*) Forensic Medicine, Kasturba Medical College, Manipal, Karnataka
10. Dr K. Ravikumar , Raksha Shakti University, Ahmedabad, Gujrat.
11. Dr C. Behera (*Addl. Prof*) Dept of FMT, AIIMS, New Delhi
12. Dr. Kanak Lata Verma, Deputy Director, Toxicology ,RFSL, Chanakyapuri New Delhi
13. Dr. Asha Srivastava (*Senior Scientific Officer*) Forensic Psychology,Central Forensic Science Laboratory, CBI, Delhi
14. Dr. Raghvendra Kumar Vidua, (Associate Prof), FMT, AIIMS Bhopal
15. Dr. Vaibhav Saran (*Asst.Prof.*) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences,Allahabad
16. Ms Aparna R. Asst. Prof. Forensic Serology & Biology, Jain University, Bengaluru

17. Dr. Deepali Jain, Asst Prof, Forensic Science ,BB Ambedkar University, Lucknow
18. Prof. SK Dhatarwal, Forensic Medicine, PGIMS, Rohtak, Haryana
19. Prof. NK Aggrawal Forensic Medicine, UCMS, Delhi
20. Prof. Manoj Kumar Mohanty, Forensic Medicine, AIIMS, Bhubneshwar, Odisha
21. Prof. Amar Jyoti Patowary, Forensic Medicine, NEIGRIHMS, Shillong, Meghalaya
22. Prof S. Venkata Raghava , Forensic Medicine, Bangalore Medical College, Bengaluru
23. Prof. Shalini Gupta Oral Pathology and Microbiology, Faculty of Dental Sciences, King George Medical University, Lucknow
24. Prof. Virender Kumar Chhoker Forensic Medicine, Santosh Medical College, Ghaziabad, UP
25. Prof. Dayanand G Gannur , Forensic Medicine , Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
26. Prof Praveen Arora, Forensic Medicine, SAIMS, Indore, MP
27. Prof Barkha Gupta , Saraswathi Institute of Medical Sciences Hapur, Uttar Pradesh India
28. Prof M Prashant Apollo Medical College Hyderabad
29. Prof Dimple Patel , Anatomay, AMC MET Medical College , Ahmedabad , Gujarat
30. Dr Mohammed Nader Shalaby, Associate Professor of Biological Sciences and Sports Health Department, Faculty of Physical Education, Suez Canal University, Egypt

## Current Issue

ATOM 1.0

RSS 2.0

RSS 1.0

[Make a Submission](#)

## Browse

Open Journal Systems

## Information

[For Readers](#)

[For Authors](#)

Platform &  
workflow by  
**OJS / PKP**

## Vol. 15 No. 3 (2021): Indian Journal of Forensic Medicine & Toxicology

DOI: <https://doi.org/10.37506/ijfmt.v15i3>

Published: 2021-06-02

### Articles

#### Assess the Knowledge, Utilization & Barrier of Non-Utilization Regarding Maternal Health Services

Sonu, Akoijam Mamata Devi, Anu Grover, Deepak

1-6

 pdf

#### A Research Study on Medicolegal Autopsies Conducted at Mortuary of Government Medical College, Ongole, Andhra Pradesh from 1st January to 31st December 2020

Venati Jayashankar, Chaitanya. R, G.B. Raja Kumar

7-10

 pdf

#### Dietary and Nutritional Strategy for the Prevention and Management of Hyperuricemia

A. Priyadharshini, Manali Sudhir, Anirudh Manikandan, Ahalya S.P

11-15

 pdf

#### A Medico Legal Study of Fetal Deaths in and Around Guntur

A.Krishnajaneyulu, N. Devchand, Raja Sekhar Uppu, T.T.K.Reddy

16-22

 pdf

#### Exploring the Depth- Influence by Female Sex Hormones Seen in the Periodontium

Aishwarya Dham, Deepa G Kamath

23-31

**Basic Dental Age Estimation Using Radicular Dentine Translucency in Indian (Himachal Pradesh) Population-A Digital Method**

Akshay Sujith, Neeta Sharma, R Guru Prasad

32-39

 Pdf

**An Epidemiological Study of Snake Envenomation in a Tertiary Care Hospital of North India**

Amit Kumar, Shweta Tanwar, Rajesh Chetiwal, Rajesh Patnaik Thonangi

60-65

 Pdf

**Learning Styles: An Overview**

Amita Sood, Jyoti Sarin

66-71

 pdf

**A Novel Method to Estimate Height of an Individual with Facial Index**

Arunashri, Lalitha C., Venkateshu K.V.

40-43

 Pdf

**Evaluation of Snake Bite Cases Admitted To Hospital in Tertiary Care**

Avinash H. Waghmode, Pravir Bodkha, Suvarna N.Patil, A.J.Ajankar, Sonali Kotkar

44-48

 Pdf

**Current Trends of Poisoning Cases Admitted at Tertiary Care Hospital**

Avinash H. Waghmode, Prakash Mohite, Sudhir Ninave, Pravir Bodkha, Sonali Kotkar

49-53

 pdf

**Analysis of Questionnaire based Response of Indian Medical Students about the Distinguishing Features of Relevant Toxic Plants, their Systemic Effects and Legal Provisions Under Narcotic Drugs and Psychotropic Substance Act, 1985**

Amit Kumar Singh, Samiul Haque, Jaun Jeb Rizvi

54-59

 pdf

**Knowledge, Attitude and Perception Regarding the Procedure of Autopsy in COVID-19**

## **Issues among Doctors Doing Post Mortem Examination**

Anil Mangeshkar, Ashok Najan, Seema Sutay

72-77

 pdf

## **A Critical Appraisal of Ethical issues in E-Learning**

Anjankar A J, Mohite P M, Waghmode A, Patond S, Ninave S

78-81

 pdf

## **Do Nursing Students Have High Social Intelligence? Findings from Selected Nursing College, South India**

Anju Philip.T, Sunitha P.P, Lakshmi S Babu, Vrinda R

82-87

 pdf

## **An Observational Study of the Effect of Dexmedetomidine on Haemodynamic and Airway Responses During Extubation of Trachea Following Major Surgery Under General Anaesthesia**

Anuja Agrawal, Riddhi Shah, Kirti Patel, Pooja Shah, Dinesh Chauhan


88-95

 Pdf

## **Determination of Sex by various Morphometric Traits of Clavicle in the population of Central India (Bhopal region)**

Anuradha Singh, Rajendra Baraw, Jayanthi Yadav


96-103

 pdf

## **The Relationship between the Severity of Obstructive Sleep Apnea (OSA) and Tension-Type Headache Frequency**

Aprilia, Herlina Suryawati, M.I. Widiastuti, Aris Catur Bintoro, EndangKustiowati, Jimmy Eko Budi Hartono

104-109

 pdf


## **Need for Psychological Check-ups and Medical Treatment to the witnesses of Heinous Crimes**

Archit Panwar, Komal Vig

110-115


## **Prevalence and therapeutical Study of Gastrointestinal Nematodes in Local Goats in Northern of Iraq**

Alaa Kamil Mahmood, Zeyad Taha Hussein, Omar-Althani S. Saed, Maher S. Owain  
1985-1990




## **Histological Effect of Onion, Tea Against Lead Acetate in Some Organs of Sprague-Dawley Rats**

Ammar Abdulhasan Aldhalemi, Ali Joodi shuna Al-Fatlawi, Zinah Talib Alsallami, Ashraf Raof  
Mohammed Ali, Murtadha Abdulhasan Aldhalemi  
1991-1996




## **The Relationship of Antral Follicle Count (AFC) with Levels of Estradiol Serum and Mature Ovarian Follicles in Women Undergoing the IVF Program at the Graha Amerta Fertility Clinic of General Hospital of Dr. Soetomo in the Period 2018**

Anak Agung Renjani Kanya Kumari, RenyI'tishom, Hendy Hendarto, Sri Ratna Dwiningsih  
1997-2003




## **Facial Reconstruction after Surgical Excision of Basal Cell Carcinoma**

Anas H. Abed, Mohammed Kh. Abduljalel, Kamal Turki Aftan  
2004-2010




## **Azithromycin, Cefixime, Vitamin C, Vitamin D, and Zinc to Treat Mild Covid-19: A Case Report**

Anas Khaleel, Haidar AL Shammari, Mohammad Niazi, Talal Salem Al-Qaisi, Rafat Abutaleb, Amneh H. Tarkhan  
2011-2014



## **Antibacterial Activity Test of *Jatropha multifida* L. sap against *Staphylococcus aureus* and Methicillin Resistant *Staphylococcus aureus* (MRSA) in vitro**

Anastassya Yuniar Setyo Putri, Marijam Purwanta, DantiNur Indiasuti, Arthur Pohan Kawilarang  
2015-2020



## **Development of Information System-Based Policy for COVID-19 Affected Students in the Semi-Arid Area of Indonesia**

# The Relationship of Antral Follicle Count (AFC) with Levels of Estradiol Serum and Mature Ovarian Follicles in Women Undergoing the IVF Program at the Graha Amerta Fertility Clinic of General Hospital of Dr. Soetomo in the Period 2018

Anak Agung Renjani Kanya Kumari<sup>1</sup>, RenyI'tishom<sup>2</sup>, Heady Hendarto<sup>3</sup>, Sri Ratna Dwiningsih<sup>4</sup>

<sup>1</sup>Researcher, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, <sup>2</sup>Associate Professor, Department of Medical Biology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, <sup>3</sup>Professor, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, <sup>4</sup>Associate Professor, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

## Abstract

**Background:** In the last two decades, one of the critical factors of assisted reproductive technology (ART) has been identified as ovarian capacity. Consequently, ovarian capacity markers have evolved to be part of diagnostic testing conducted regularly before IVF. Several predictor factors, such as antral follicle count (AFC), is one of several factors to predict ovarian capacity. These diagnostic procedures and tests have the purpose of identifying women at high-risk patient who respond poorly toward controlled ovarian stimulation throughout the IVF cycle. The ovarian response toward ovulation stimulation within this research was assessed by peak estradiol (E2) serum levels and the total number of mature follicles. There is no clear relationship between AFC and levels of estradiol serum and the number of mature ovarian follicles to identify women who respond poorly. Methods: This study is an analytical study with a cross-sectional design. Samples were taken from secondary data of medical records using the total sampling method on women undergoing the IVF program at the Graha Amerta fertility clinic of General Hospital of Dr. Soetomo from January 1 to December 31 2018 according to the inclusion and exclusion criteria. The AFC variables, estradiol serum levels, and the number of mature ovarian follicles in the study were analyzed using Spearman's correlation. Conclusion: There is a significant relationship between antral follicle count (AFC) with the levels of estradiol serum and the number of mature follicles in women undergoing the IVF program.

**Keywords:** Antral follicle count (AFC), estradiol (E2) serum levels, the number of mature ovarian follicles

## Introduction

Infertility considered as one of the global public health problems. Nowadays, infertility is an event that must be paid attention to because, corresponding to the World Health Organization, infertility in women currently being the severe fifth-highest incident in the

global scope. Currently, the highest pregnancy success rate achieved in the world by the IVF program has only reached 20%.<sup>1,2</sup> The rapidly developing medical science and technology answers the issue of difficulties in having offspring through Assisted Reproductive Technology (ART), such as In Vitro Fertilization (IVF).<sup>3</sup> Over the last twenty years, one of the critical factors of assisted reproductive technology (ART) has been identified as ovarian capacity. Consequently, ovarian capacity markers have been considered to be included in the routine diagnostic testing performed before IVF.<sup>4</sup>

---

### Corresponding author:

**RenyI'tishom,**

Department of Medical Biology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

E-mail: ritishom@fk.unair.ac.id



Based on several studies, ovarian capacity can be predicted by several predictor factors, including ovarian volume, antral follicle count (AFC), ovarian stromal blood flow, and hormonal markers such as follicle-stimulating hormone (FSH), estradiol (E2), anti-Mullerian hormone (AMH), and inhibin B.<sup>4</sup> These diagnostic procedures and tests have the purpose of identifying women at high-risk patient who respond poorly toward controlled ovarian stimulation throughout the IVF cycle.<sup>5</sup> The ovarian response to ovulation stimulation can be assessed by the total number of mature follicles, the total number of oocytes obtained in the ovum pick-up procedure, the number of mature oocytes obtained in the ovum pick-up procedure, and the total number of gonadotropin doses needed in the ovulation stimulation cycle.<sup>4</sup>

According to an article, some recent studies have shown that AFC is a useful predictor of ovarian response to ovulation induction drugs.<sup>6</sup> Moreover, AFC is considered to be the best single predictor of ovarian response in the ARB cycle. There is no clear relationship between antral follicle count and levels of estradiol serum and the number of mature ovarian follicles.

Based on the description above, it is necessary to conduct a study on "the relationship between antral follicle count (AFC) and levels of estradiol serum and the number of mature ovarian follicles in women undergoing the IVF program" with the hope of providing scientific information for later use in the initial evaluation of infertile patients in the procedures of IVF program. In the future, the success rate of IVF for infertile couples can increase in Surabaya and at the national and global levels.

## Materials and Methods

### Experimental Design

This research has been approved by Health Research Ethics Committee RSUD Dr. Soetomo Surabaya with ethical clearance number 1521/KEPK/IX/2019. It was an analytical study with a cross-sectional design. Samples were taken from secondary data of medical records on women undergoing the IVF program at the Graha Amerta fertility clinic of General Hospital of Dr.

Soetomo from the period of January 1 to December 31 2018 according to the inclusion and exclusion criteria.

Data was collected using the total sampling method. The data taken included age, occupation, residence, levels of E2 serum on day 3, levels of FSH serum on day 3, levels of LH serum on day 3, AFC, E2 serum levels before OPU, number of mature follicles, and type of ovulation stimulation drugs. Exclusion criteria were female patients with ovulation disorder Polycystic Ovary Syndrome (PCOS), endometriosis, and history of ovarian surgery. Incomplete data were excluded from the study.

### Statistical Analysis

The data obtained were then analyzed using observational analysis methods, processed using the SPSS v 25.0 program. The Kolmogorov Smirnov normality test was performed on research variables and with Spearman's correlation analysis and presented descriptively. P values less than 0.05 were considered statistically significant.

### Result

A total of 209 couples participated in the IVF program in the 2018 period. The results of medical record data that met the inclusion criteria were 126 samples. Most of the respondents were women of productive age with a mean of  $34.63 \pm 4.714$  years, and the largest age group was 30 - 34 years as many as 48 (38.1%). Based on occupation, most private employees (31.4%) followed by housewives (18.6%). In the category of self-employed and other groups, respectively 17, 1%. Other groups work as lecturers, teachers, doctors, midwives, honorary, and the Indonesian Navy. In addition, the majority of patients (99; 78.6%) resided in East Java. The data result (Table 1) showing the lowest AFC of 2 and the highest of 20 with a mean of  $8.17 \pm 3.64$  calculated from both right and left ovaries. The lowest level of E2 serum before OPU in patients was 14.80 and the highest of that was  $>3000$  pg/mL with a mean of  $1838.53 \pm 876.28$  pg/mL. The lowest and the highest number of mature follicles were 0 and 19 respectively, with a mean of  $6.21 \pm 3.78$ .

**Table 1. Descriptive analysis of AFC, levels of E2, and the number of mature ovarian follicles in women undergoing IVF at the GrahaAmerta fertility clinic in 2018**

Description	Total		Descriptive	
	n	%	Min - Max	Mean (SD)
AFC	126	100	2 – 20	8.17 (3.64)
Levels of E2 Serum			14.80 – 3001	1838.53 (876.28)
The number of mature follicles			0 – 19	6.21 (3.78)

The analysis of AFC based on the respondents' age (Table 2) showed that the maximum number of AFC in the age category of 20-24 years was 12 with a mean of  $9.33 \pm 2.517$ . In the age category of 25-29 years, the lowest AFC was 5 and the highest was 15 with a mean of  $9.38 \pm 3.070$ . The ages of 30-34 years had the lowest AFC of 4 and the highest AFC of 20 with a mean of  $9.00 \pm 3.345$ . The lowest AFC in the age category of 35 - 39 years was 2 and the highest AFC in this category was 20 with a mean of  $8.57 \pm 3.788$ . In the age category of 40 - 45 years, the mean AFC was the lowest compared to other age categories, in which the lowest AFC was 2 and the highest AFC was 8 with a mean of  $4.35 \pm 1.872$

**Table 2. Age-based AFC analysis of women undergoing IVF at the Graha Amerta fertility clinic in 2018**

Age (years)	Total (n=126)	AFC	
		Min - Max	Mean (SD)
20 – 24	3	7 – 12	9.33 (2.517)
25 – 29	13	5 – 15	9.38 (3.070)
30 – 34	48	4 – 20	9.00 (3.345)
35 – 39	42	2 – 20	8.57 (3.788)
40 – 45	20	2 – 8	4.35 (1.872)

Data of AFC, levels of E2 serum before ovum pick-up (OPU), and the number of mature ovarian follicles were tested for normality of data with the Kolmogorov Smirnov test with  $p=0.002$ ;  $0.002$ ;  $0.000$  respectively, indicating that the data was not normally distributed. Therefore, Spearman's correlation method is used.

Based on the statistical tests results on 50 samples using the Spearman method, the r-value was 0.526, indicating a moderate correlation between AFC and

levels of E2 serum stimulated by FSH. It was found that the  $p=0.000$  value was smaller than 0.05. Therefore, it can be concluded that the moderate correlation was statistically significant. In the group that received a combination of FSH - LH of a total of 76 samples (Table 3), the results of the Spearman's correlation test showed a moderate and significant correlation between the variables of AFC and levels of E2 serum with combined FSH - LH ovulation stimulation ( $r=0.571$ ;  $p=0.000$ ).

**Table 3. Relationship between AFC and levels of E2 serum before OPU based on the stimulation of FSH and a combination of FSH & LH**

Variable	Type of Ovulation Stimulation Drugs	Total (n)	r-Value (correlation coefficient)	p-Value (sig)
AFC à Levels of E2 serum before OPU	FSH	50	0.526**	0.000
	Combination of FSH & LH	76	0.571**	0.000

The results of statistical tests on 50 samples using the Spearman method show that the r-value was 0.603, indicating a moderate correlation between AFC and the number of mature follicles stimulated by FSH. Besides, it was found that the  $p=0.000$  value was smaller than 0.05, so that it can be concluded that the correlation was statistically significant. Patients with combination stimulation of FSH - LH show an r-value of 0.480, indicating that AFC correlated with the number of mature follicles (Table 4). This relationship was statistically significant ( $p=0.000$ ).

**Table 4. Relationship between AFC and Maturity Follicle Count based on stimulation of FSH and a combination of FSH & LH**

Variable	Type of Ovulation Stimulation Drugs	Total (n)	r-Value (correlation coefficient)	p-Value (sig)
AFC à Number of mature follicles	FSH	50	0.603**	0.000
	Kombinasi FSH & LH	76	0.480**	0.000

## Discussion

The data from this study follows the Human Fertilization and Embryology Authority or HFEA report that stated the characteristics of patients who participated in the FIV program in 2016 in the UK were patients aged less than 35 with an average of 35.5 years.<sup>7</sup> This is also following previous research that was conducted at the Graha Tunjung IVF Clinic at Sanglah Hospital, the study found that most patients in the IVF program were in the age group less than 35 years old.<sup>8</sup>

According to a research on the characteristics of infertile pairs shows that the most respondents' occupation is private workers (47%).<sup>9</sup> Another study that discusses the factors that affect infertility in women stated that most infertile women are women who work in the type of work most found are private workers.<sup>10</sup> The

results of this study related to respondent's occupation are in line with another research about infertility, which states that working women (career women) have an increased risk of infertility 3.91 times greater than women with normal stress levels and statistically significant ( $p = 0.046$ ).<sup>11</sup>

Ovarian aging appears to play a major part in reproductive aging and is associated with decreased oocyte quality and slowly decrease in the number of primordial follicles, which will be reduced exponentially through childhood and adulthood, resulting in nearly follicle-free ovaries at menopause. Likewise, the number of follicles that has entered during the growth phase through the antral development stage falls with age. A decrease in ovarian volume and the number of antral follicles are characteristics of ovarian aging. According to a study, AFC is a significant parameter compared to

patient age and other endocrine markers in indicating an ovarian reaction towards stimulation that has been given in advance.<sup>12</sup>

Demographic and clinical studies show that female fertility begins to subside from the age of 30, and the ability to get pregnant has become nearly zero at an average age of 41 years.<sup>13</sup> The outcome of this study shows that in the age category of 40 - 45, the average AFC was the lowest compared to other age categories, in which the lowest AFC was 2, and the highest was 8 with a mean of  $4.35 \pm 1.872$ . The older the women undergoing the IVF program at the Graha Amerta fertility clinic, the lower the average AFC. Another research stated that AFC could help determine the stimulation protocol; based on some studies, it has been proven to be the most calculable determinant of oocyte retrieval per initial FSH dose.<sup>14</sup>

All research subjects in this study were IVF participants using the GnRH antagonist protocol. Ovarian stimulation was begun on the third day using recombinant FSH drugs, and then the injection was controlled following each of the participant's reactions towards the dose given. Gonadotropin-releasing hormone (GnRH) antagonist administered to deny premature LH surge begins at the time follicle attained 14 mm in diameter.<sup>15</sup> Another research stated that GnRH antagonists' use in ovarian stimulation protocols had provided a new option in the promotion of multifollicular growth.<sup>16</sup> Antagonists cause direct suppression in releasing gonadotropin, in the absence of the flare-up effect. FSH is considered to play a more critical role in multifollicular growth. However, it is possible to suggest that LH can be advised to be included in all ovulation induction protocols, since it is difficult to ascertain which individual will be advantaged from LH administration. Currently, no evidence has been made regarding the negative effects of LH to the ovarian stimulation outcome.

According to the results of the statistical tests on 50 samples using the Spearman method, the r-value was 0.526, indicating a moderate correlation between AFC and levels of E2 serum stimulated by FSH. Besides, it was found that the  $p=0.000$  value was smaller than 0.05, so it can be concluded that the moderate correlation was statistically significant. Meanwhile, in the group that received a combination of FSH - LH of 76 samples, Spearman's correlation test results show a moderate and significant correlation between the variables of AFC and

levels of E2 serum with combined ovarian stimulation of FSH - LH ( $r=0.571$ ;  $p=0.000$ ). The same results as this study were found in research that examined the relationship of AFC among various ovarian examination parameters, including levels of E2 serum.<sup>17</sup> The results show that AFC was significantly associated with levels of E2 serum before OPU. Another research about the AFC application in clinical settings to indicate whether AFC acquired from various observers is a predictor of the quality of the stimulation with the parameter of peak levels of E2 serum, scientifically showing a major positive correlation between AFC and peak levels of E2 serum.<sup>6</sup>

The statistical tests on 50 samples using the Spearman method show that the r-value was 0.603, indicating a moderate correlation between AFC and the number of mature follicles stimulated by FSH. Besides, it was found that the  $p=0.000$  value was smaller than 0.05, so it can be concluded that the correlation was statistically significant. Patients with combination stimulation of FSH - LH show an r-value of 0.480, indicating that the number of mature follicles corresponds to the AFC. This relationship was statistically significant ( $p=0.000$ ). It is in line with the results of another study, stating that there was a significant relationship between AFC and ovarian response to ovarian stimulation.<sup>4</sup> In this study, the ovarian response was examined by four parameters, such as the quantity of mature follicles. The higher the number of AFC, the higher the number of mature follicles.

Poor ovarian response was first described in 1983 as having a peak estradiol concentration  $<300$  pg/mL and inadequate follicular production resulting in fewer eggs being acquired, thus declines the amount of embryos transferred. In that sense, a poor ovarian response leads to drop of oocyte production, cycle cancellation, and overall is corresponded with a significantly reduced chance of pregnancy.<sup>18</sup> Data presented by another study proved that AFC provides more significant prediction regarding the incident of a poor response to hormone stimulation in IVF compared to the patients' age and other examination that is recently used, for example, basal FSH, E2, or inhibin B.<sup>19</sup> It can be stated that AFC becomes a good prognosis of poor ovarian response. It is an essential evaluation in an IVF program because a patient undergoing an IVF program due to a poor ovarian response is generally regarded to have a lower pregnancy rate than those with a normal ovarian response.

## Conclusion

1. There is a relationship between antral follicle count (AFC) and estradiol (E2) serum levels in women undergoing the IVF program at the Graha Amerta fertility clinic at General Hospital of Dr. Soetomo for the period 2018.

2. There is a relationship between antral follicle count (AFC) and the number of mature follicles in women undergoing the IVF program at the Graha Amerta fertility clinic at General Hospital of Dr. Soetomo Hospital for the period 2018.

**Conflict of Interest:** The authors declare that there is no conflict of interest.

**Ethical Clearance:** This research has been approved by Health Research Ethics Committee RSUD Dr. Soetomo Surabaya with ethical clearance number 1521/KEPK/IX/2019.

**Source of Funding:** Self-funding.

**Acknowledgments:** The author would like to express gratitude for the authority of the Faculty of Medicine, Airlangga University and RSUD. Dr. Soetomo who is involved in providing the facilities for this research.

## References

1. World Health Organization. Infertility Definitions And Terminology [Internet]. 2019 [cited 2019 May 3]. Available from: <https://www.who.int/reproductivehealth/topics/infertility/definitions/en/>
2. Silvia M, Sofia A, Nadhira A, Harzif A. Karakteristik pasien yang menjalani terapi Fertilisasi In Vitro (FIV) di klinik Yasmin Kencana RSUP Dr. Cipto Mangunkusumo Jakarta pada Januari 2011 - Desember 2015. *Indones J Obstet Gynecol.* 2016;4:130.
3. Kementerian Kesehatan Republik Indonesia. Peraturan Kementerian Kesehatan nomor 39 tentang Tertibkan Klinik Teknologi Reproduksi Berbantu di Rumah Sakit. Indonesia; 2016.
4. Adnyana IBP. Hubungan Jumlah Folikel Antral dengan Respons Ovarium terhadap Stimulasi Ovulasi. *Indones J Obstet Gynecol* [Internet]. 2008;32(1). Available from: <http://www.inajog.com/index.php/journal/article/view/146>
5. Christianson MS, Shoham G, Tobler KJ, Zhao Y, Cordeiro CN, Leong M. et al. Measurement of Antral Follicle Count in Patients Undergoing In Vitro Fertilization Treatment: Results of A Worldwide Web-Based Survey. *J Asisted Reprod Genet.* 2015;32(10):1435–40.
6. Hansen KR, Morris JL, Thyer AC, Soules MR. Reproductive Aging and Variability in The Ovarian Antral Follicle Count: Application in The Clinical Setting. *Fertil Steril.* 2003;80(3):577–83.
7. Human Fertilisation & Embryology Authority. Fertility Treatment 2014–2016 Trends And Figures. United States: Human Fertilisation & Embryology Authority; 2018.
8. Dharma NDW, Budiana ING. Perbedaan Gambaran Oosit Berdasarkan Kelompok Usia Pada Pasien In Vitro Fertilization di Klinik Bayi Tabung RSUP Sanglah Denpasar. *E-Jurnal Med Udayana.* 2019;8(5).
9. Sa'adah N, Purnomo W. Karakteristik dan Perilaku Berisiko Pasangan Infertil di Klinik Fertilitas dan Bayi Tabung Tiara Cita Rumah Sakit Putri Surabaya. *J Biometrika dan Kependud* [Internet]. 2016;5(1):61–9. Available from: *Jurnal Biometrika dan Kependudukan*
10. Oktarina A, Abadi A, Bachsin R. Faktor-Faktor Yang Memengaruhi Infertilitas pada Wanita di Klinik Fertilitas Endokrinologi Reproduksi. *Maj Kedokt Sriwij.* 2014;46(4):295–300.
11. Indarwati I, Hastuti URB, Dewi YLR. Analysis of Factors Influencing Female Infertility. *J Matern Child Heal.* 2017;2(2):150–61.
12. Hestiantoro A, Atadisasatra M, Andonotopo W, Sumapraja K, Wiweko B. Ultrasonografi Endokrinologi Reproduksi & Infertilitas. Jakarta: Sagung Seto; 2013.
13. Scheffer GJ, Broekmans FJ, Dorland M, Habbema JD, Looman CW, te Velde ER. Antral Follicle Counts By Transvaginal Ultrasonography are Related To Age In Women With Proven Natural Fertility. *Fertil Steril.* 1999;72(5):845–51.
14. Hsu A, Army M, Knee AB, Bell C, Cook E, Novak AL, et al. Antral Follicle Count in Clinical Practice: Analyzing Clinical Relevance. *Fertil Steril.* 2011;95(2):474–9.
15. Hendarto H. Stres Infertilitas Menghambat Maturasi Oosit dan Hasil Fertilisasi in Vitro. *Maj Obstet Ginekol.* 2015;23(1):17–21.

16. Levi-Setti PE, Cavagna M, Baggiani A, Zannoni E, Colombo G V, Liprandi V. FSH And LH Together in Ovarian Stimulation. *Eur J Obstet Gynecol Reprod Biol.* 2004;115:S34-9.
17. Ben-Haroush A, Farhi J, Zahalka Y, Sapir O, Meizner I, Fisch B. Small Antral Follicle Count (2-5 Mm) And Ovarian Volume For Prediction Of Pregnancy Inin Vitrofertilization Cycles. *Gynecol Endocrinol.* 2011;27(10):748-52.
18. Badawy A, Wageah A, Gharib M El, Osman EE. Prediction and Diagnosis of Poor Ovarian Response: The Dilemma. *J Reprod Infertil.* 2011;12(4):241-8.
19. Banesi LFJM., Broekmans FJ., Eijkemans MJ., Jong FH de, Habbema JDF, Velde ER te. Predictors of Poor Ovarian Response in in Vitro Fertilization: a Prospective Study Comparing Basal Markers of Ovarian Reserve. *Fertil Steril* [Internet]. 2002;77(2):328-36. Available from: [https://www.fertstert.org/article/S0015-0282\(01\)02983-1/fulltext](https://www.fertstert.org/article/S0015-0282(01)02983-1/fulltext)

# Source details

## Indian Journal of Forensic Medicine and Toxicology

Scopus coverage years: from 2008 to Present

Publisher: Institute of Medico-Legal Publications

SSN: 0973-9122 E-ISSN: 0973-9130

Subject area: Social Sciences: Law Medicine: Pathology and Forensic Medicine

Pharmacology, Toxicology and Pharmaceuticals: Toxicology Environmental Science: Health, Toxicology and Mutagenesis

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

CiteScore 2020

0.1

SJR 2020

0.115

SNIP 2020

0.243

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### Improved CiteScore methodology

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

### CiteScore 2020

0.1 =  $\frac{387 \text{ Citations } 2017 - 2020}{3,509 \text{ Documents } 2017 - 2020}$

Calculated on 05 May, 2021

### CiteScoreTracker 2021

0.1 =  $\frac{393 \text{ Citations to date}}{3,814 \text{ Documents to date}}$

Last updated on 04 June, 2021 • Updated monthly

### CiteScore rank 2020

Category	Rank	Percentile
Social Sciences		
— Law	#639/722	11th
Medicine		
— Pathology and Forensic Medicine	#183/191	4th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site &](#)

### About Scopus

- [What is Scopus](#)
- [Content coverage](#)
- [Scopus blog](#)
- [Scopus API](#)

### Language

- [日本語に切り替える](#)
- [切换到简体中文](#)
- [切换到繁體中文](#)
- [Русский язык](#)

### Customer Service

- [Help](#)
- [Contact us](#)



## Journal Research

ReviewerCredits - Certification and rewarding of peer review and conference talks

reviewercredits.html

OPEN

# Indian Journal of Forensic Medicine and Toxicology

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
India <small>Universities and research institutions in India</small>	Environmental Science Health, Toxicology and Mutagenesis  Medicine Pathology and Forensic Medicine  Pharmacology, Toxicology and Pharmaceutics Toxicology  Social Sciences Law	Indian Journal of Forensic Medicine and Toxicology	<b>20</b>
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	09739122, 09739130	2008-2020	Homepage How to publish in this journal editor.ijfmt@gmail.com

### Scopus Indexed Jo

Call for Papers August Iss

Fast Track Peer Reviewed Put

toijqi.net

OPEN

### SCOPE

"Indian Journal of Forensic Medicine & Toxicology " is a double-blind peer reviewed international journal. The frequency is quarterly. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine, environmental medicine, Forensic Pathology, legal medicine and public health laws.

Join the conversation about this journal



# Scopus Indexed Journal

Fast Track Peer Reviewed Publication.

tojqi.net

OPEN

Quartiles

# Scopus Indexed Journal


Fast Track Peer Reviewed Publication.


tojqi.net

OPEN

## FIND SIMILAR JOURNALS

Rank	Journal Name	Country	Similarity
1	Medico-Legal Update	IND	53%
2	International Journal of Medical Toxicology and Legal	IND	41%
3	Journal of Indian Academy of Forensic Medicine	IND	34%
4	Journal of Punjab Academy of Forensic Medicine and	IND	30%
5	Journal of Association of Forensic Medicine and Toxicology	IND	





Indian Journal of Forensic Medicine and Toxicology

Q4

Health, Toxicology and Mutagenesis

best quartile

SJR 2020 0.12

powered by scimagojr.com

Show this widget in your own website

Just copy the code below and paste within your html code

```
<a href="https://www.scimagojr.com" data-bbox="230 174 354 184">
```

## How to Claim a Review

All your peer reviews will be certified by journals and you earn credits to get benefits

[reviewercredits.com](https://www.scimagojr.com)

OPEN

Metrics based on Scopus® data as of April 2021

A **Aloto Ximenes Belo Amaral** 2 months ago

hello sir/madam,

indian journal of forensic medicine and toxicology 'Coverage 2008-2019' is the journal out of Scopus? or someother reason?

thank you and please let me know.

reply



**Melanie Ortiz** 2 months ago

SCImago Team

Dear Aloto,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was released on 11 June 2020. We suggest you consult the Scopus database directly to see the current index status as SJR is a static image of Scopus, which is changing every day.

Best Regards, SCImago Team

S **sekar anggung gp** 5 months ago

how can i see the impact factor of Indian Journal of Forensic Medicine

reply



**Melanie Ortiz** 5 months ago

SCImago Team

Dear Sekar, thank you very much for your comment. SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR (Check it on our website). We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team