

# Male Infertility An Epidemiological and Clinical Profile at the Andrology Unit of Dr. Soetomo Hospital, Surabaya, Indonesia

*by Antonius Wahyudi*

---

**Submission date:** 20-Jun-2022 02:28PM (UTC+0800)

**Submission ID:** 1859960341

**File name:** e\_Infertility\_An\_Epidemiological\_and\_Clinical\_Profile\_at\_the.pdf (4.76M)

**Word count:** 2552

**Character count:** 13939

## Male Infertility: An Epidemiological and Clinical Profile at the Andrology Unit of Dr. Soetomo Hospital, Surabaya, Indonesia

Antonius Wahyudi<sup>1\*</sup>, Petrus Supardi<sup>1</sup>, Pety Narulita<sup>1</sup>, Reny Ptishom<sup>2</sup>, Silvia W Lestari<sup>3</sup>

<sup>1</sup>Andrology Specialist Program, <sup>2</sup>Faculty of Medicine of Universitas Airlangga/ Dr. Soetomo Hospital Surabaya, Indonesia.

<sup>2</sup>Department of Medical Biology, Faculty of Medicine of Universitas Airlangga, Surabaya, Indonesia.

<sup>3</sup>Department of Medical Biology, Faculty of Medicine of Universitas Indonesia, Jakarta, Indonesia.

\*Corresponding Author: Antonius Wahyudi

### Abstract

Objective: To discover and to lay out epidemiological features and clinical characteristics of patients with infertility problems who seek treatment at the Andrology Unit of Dr. Soetomo Hospital. Methods: A cross sectional study was conducted to men with infertility who sought treatment in the Andrology Unit of Dr. Soetomo Hospital ranging from January to December 2017. Only patients who lived together with their spouses were included as research subjects. Results: The average age of the patients was 33.76 years (age ranged from 25 to 46 years) and the average age of their spouses was 30.4 years (age ranged from 19 to 42 years). The average duration of infertility is 4.25 years (between 1 year and 13 years). Average age of puberty was 14.5 years old. First consultations were late with average of 4.46 years. From overall infertile patients, 90% of them were categorized as primary infertility. History of smoking was found in 84 people (68.29%), history of wearing tight pants 27 people (21.95%). In terms of Body Mass Index(BMI), we found that underweighted patients comprise 7% (9 patients) from total liable subjects, while 55 people (55%) were in the area of ideal body mass, 46 people were overweight (35%), whereas obesity was found in 13 subjects (10%). The results of semen analysis showed that the most case were asthenoteratozoospermia 34 people (27.6%). A careful history, physical examination and semen analysis can determine the cause of male infertility. The cause of male infertility in Surabaya, Indonesia is very complex, which involves older age, delayed consultation and background as risk factors that cause abnormalities in semen analysis.

**Keywords:** Male infertility, epidemiology, andrology, Dr. Soetomo Hospital, Surabaya, Indonesia.

### Introduction

Infertility is defined as the inability to have offspring within one year with regular intercourse (2-3 times a week), without any contraception.[1] Male infertility comprises almost half of the total infertility cases.[2] Despite many advances in medical science and technology, especially in infertility management, the number of infertility cases is still unceasingly increasing.[3,4] Meanwhile, only a few researches on male infertility were conducted in Indonesia,

particularly epidemiological study. The aim of this study was to describe the epidemiological side of infertility cases in Indonesia, which consist etiological and clinical profile of male infertility, especially in Surabaya.

### Patients and Methods

A descriptive, cross-sectional design was implemented in this one year study, starting from 1<sup>st</sup> January to 31<sup>st</sup> December 2017.

Samples of this study were taken from the patients' medical records in Andrology Unit of Dr. Soetomo Hospital, Surabaya, Indonesia. Men who were diagnosed as infertile by the attending andrology specialists were included in the study. Semen analysis is a must in this study, for at least 2 times with 3 months interval. Semen was collected in glass container by masturbation, with the prior abstinence period of 2 to 7 days. Semen analysis was performed according to the WHO guidelines. In several patients, additional hormonal examinations

consist of FSH and testosterone level were done.

### Results

During the study period, 172 patients visited the Andrology Unit of Dr. Soetomo Hospital, Surabaya, Indonesia, and 123 of them were eligible to the study according to the inclusion criteria (71.5%). The average age of men was 33.76 years with the youngest was 25 years old and the oldest 46 years old. The highest percentage of patients were in the 30 -40 years old age group (Table 1).

**Table 1: Distribution of patient according age group**

Age group (year)	Number of Patients	Percentage (%)
20 - 29	34	27.64
30 - 40	60	48.78
40 - 50	17	13.28

The average age of the spouses was 30.4 years with the youngest was 20 years old and the oldest was 42 years old. The majority of the patients were employee in government or private office (40%), while other occupation consist of farmer and trader, respectively 30% and 15%, and the other 15% were unspecified. Eighty percent of patients reside

in urban areas. More than a third of the patients were referred by other doctor, in which 25% of them were referred by gynecologist, and the other 12% by general practitioner, whereas 63% of cases were without any reference. The average consultation delay was 4.46 years (ranged from 1 year to 8 years) (Table 2).

**Table 2: Distribution of patients according to first consultation delay**

Delay of First Consultation (years)	Number of Patients	Percentage (%)
1 - 2	84	68.29
3 - 4	27	21.95
5 - 6	84	68.29
7 - 8	27	21.95

The average duration of infertility was 4.25 years, ranged from 1 year and 13 years. Primary infertility was the majority of the case, consisting 90% of the total cases, while the other 10% were secondary. The average age of puberty was 14.5 years old, ranged from 12 to 20 years old. The average frequency of sexual intercourse was 2 to 3 times per week in 95 cases (78%). In terms of medical history, sexual transmitted diseases

were found in 2 cases (1.6%). History of smoking was found in 84 cases (68.29%) and wearing tight pants was in 27 cases (21.95%). In addition, from general examination, obesity with body mass index (BMI) more than 30% was observed in 13 cases (8.13 %). Furthermore, from andrological examination, varicocele was observed in 28 cases (22.76%). (Table 3).

**Table 3: Distribution of patients according background**

Background	Number	Percentage (%)
Sexual transmitted diseases	2	1.19
Smoking	84	50.0
Wearing tight pants	27	16.07
Obesity	13	7.73
Varicocele	28	16.67
Small testicle volume	12	7.14
Undescended Testis	2	1.19

Semen analysis was performed in all patients. The three most frequent results of semen analysis were asthenoteratozoospermia in 34 cases (34.3%), followed by teratozoospermia in 30 cases (24.3%) and oligoasthenoteratozoospermia in 21 cases (17.0%). In the sperm concentration parameter, the oligozoospermia was very dominant, and followed by severe

oligozoospermia and non-obstructive azoospermia. The complete semen analysis profile was showed in Table 4. In azoospermia cases (11 cases), additional hormonal analysis was performed. Elevated FSH levels were observed in five cases, while low testosterone levels were observed in three cases.

**Table 4: Seminal analysis result**

Semen analysis result	Number	Percentage (%)
Normozoospermia	4	3.2
Oligozoospermia	1	0.8
Asthenozoospermia	2	1.6
Teratozoospermia	30	24.3
Oligo-astheno-zoospermia	1	0.8
Oligo-terato-zoospermia	4	3.2
Astheno-terazoospermia	34	27.6
Oligo-astheno-teratozoospermia	21	17.0
Severe oligo-asthenozoospermia	10	8.1
Azoospermia	14	11.8
Aspermia	1	0.8
Retrograde ejaculation	1	0.8

**Discussions**

Actually, other than andrology, there are several other specialties that involved in the management of infertility, namely obstetrics gynecology and urology. Obstetrics gynecology is the leading specialist in the management of infertility, because most of the infertile couples came to them to have consultations in the first chance.

This condition is related to the belief of the society that infertility is only caused by female factors. Some other patients was referred to urology specialist for sperm abnormalities related diseases such as varicocele, undescended testis, ejaculatory duct obstruction and sperm retrieval for in vitro fertilization (IVF) – intracytoplasmic sperm injection (ICSI). Meanwhile, andrology was still unbeknownst to infertile patients, even to some doctors.

In contrast, in our clinic, if compared to other infertility centers, the number of male infertility patients were quite high. These patients comprised 50% of the total patients who visited the Andrology Unit. Most of the patients were not referred, but rather they came on their own volition. This is understandable because our andrology unit also served as an infertility centers in academic hospital in Surabaya, Indonesia.

Age is also considered as a regulating factor in couples of infertility. In women, the increase in age will be inversely affecting the quality of the egg. And, in line with that, the increase in age also decrease male fertility. Researches in seminal analysis showed there were several kind of damages that were inflicted by age.[5] This phenomenon was related to genetic material alterations of the sperm with age.[6] Nocturnal emission was considered as an accurate marker of male coming of age, which is the start of further development of the male reproductive system.

The age of first nocturnal emission, or commonly known as the onset of puberty is very closely related to adulthood semen quality. Late onset of puberty may resulting poor quality of semen in adulthood, which due to the differences in reproductive hormones level. In delayed puberty, testosterone peak level will unlikely be reached, or may be reached later than usual. Sometimes, even the normal level of testosterone might not reached, which ultimately disrupt sperm production in adulthood [7].

The harmful effects of smoking on male fertility have long been known. A few studies mentioned some mechanisms by which lead to infertility, such as the decrease in the quality of semen, dysfunction of the reproductive hormone, disruption of spermatogenesis, sperm maturation and sperm function.[8, 9, 10] The use of tight pants will generate excessive heat, in which induced heat stress that could be interfering with the spermatogenesis process. Heat stress is associated with decreased sperm motility, concentration and viability of sperm in humans [11]. Oxidative stress is believed to be the responsible mechanism in this case[12].

Varicocele was the most frequent physical examination finding in this study. The underlying mechanism of varicocele as a cause to male infertility is due to the increase of temperature in the dilated blood vessels, reflux of toxic metabolite and hypoxia of the testis caused by the stasis of blood vessels [13]. These mechanisms may contribute negatively to the sperm concentration, motility and morphology. Furthermore, aside of those parameters, varicocele may also interfere with sperm DNA fragmentation as well [13].

Obesity is associated with several medical conditions including fertility [14]. According to the guidelines provided by WHO, obesity is defined as body mass index (BMI) above 30 kg / m<sup>2</sup> [15]. Male fertility is influenced by obesity through the reduction of testosterone level which resulted in the interference to the sperm quality [14]. However, another review article which recently published, showed that there is no significant effect between BMI and sperm quality [16]. Semen analysis is a routine examination to evaluate male

fertility [17]. WHO had published a guideline for semen analysis, including the criteria for normal semen parameters [18]. Sperm abnormalities may cause inability to fertilize oocytes. The two most frequent semen analysis results in this study, were asthenoteratozoospermia and teratozoospermia, which in observation through electron microscopic may show immature sperm shape, morphological apoptotic patterns of sperm and cytoplasm with many sperm fragments. Oligozoospermia, severe oligozoospermia and non-obstructive azoospermia may be related to the small testicle volume.

Azoospermia were categorized into two, namely obstructive and non-obstructive azoospermia. There was no obstructive azoospermia reported in this study. This phenomenon was supported by the low prevalence of infection history. Meanwhile, the high prevalence of non-obstructive azoospermia was associated with elevated FSH levels. It means that the damage primarily occurred in the testis. In vitro fertilization (IVF) – intracytoplasmic sperm injection (ICSI) was chosen as the further management of this type of infertility, testicular biopsy should be done prior the sperm retrieval. In addition, we also reported one case of retrograde ejaculation and one case of aspermia.

### Conclusion

A careful history, physical examination and semen analysis can determine the cause of male infertility. The cause of male infertility in Surabaya, Indonesia is very complex, which involves older age, delayed consultation and background as risk factors that cause abnormalities in semen analysis.

### Reference

1. Anwar S, Anwar A (2016) Infertility: A Review on Causes, Treatment and Management, Women's Health & Gynecology, vol 2
2. Inhorn MC (2003) Global Infertility and the Globalization of New Reproductive Technologis; Illustration From Egypt, Social Science & Medicine, 56:1837-1851
3. Matzuk MM, Lamb, DJ (2008) The Biology of Infertility: Research Andvances and Clinical Challenges, Nature Medicine, 14:1197-1213
4. Mukhopadhyay D, Varghese AC, Pal M, Banerjee SK, Bhattacharyya AK, (2010) Semen Quality and Age-Spesific Changes: A Study between Two decades on 3,729 Male Partners of Couples with Normal Sperm Count and Attending an Andrology Laboratory for Infertility-Related Problems

- in an Indian City. *Fertility and Sterility*, 93:2247-2254.
5. Plas E, Berger P, Hermann M, Pfluger H (2000) Effect of Aging on Male Infertility? *Experimental Gerontology*, 35:543-551.
  6. Belloc S, Hazout A, Zini A, Merviel P, Cabry R, Chahine H (2014) How to Overcome Male Infertility After 40: Influence of Paternal Age on Fertility, *Maturitas*, 78:22-29.
  7. Lauridsen L, Arendt L, Stovring H, Olsen J, Ramlau-Hansen Caecilia (2014) Is age puberty associated with semen quality and reproductive hormones in young adult life?, *Asian Journal of Andrology*, 19:625-632
  8. Dai J, Wang Z, Qiao Z (2015) The Hazardous effects of tobacco smoking on male fertility, *Asian Journal of Andrology*, 17:954-960
  9. Harlev A, Agarwal A, Gunes S, Shetty A, Plessis S (2015) Smoking and Male Infertility: An Evidence-Based Review, 33(3):143-160
  10. Vine MF, Tse CK, Hu P, Truong KY (1996) Cigarette smoking and semen quality, *Fertil Steril*, 41:11-5
  11. P Thonneau L, Bujan L, Multigner, R, Micusset (1998) Occupational heat exposure and male infertility: a review," *Human Reproduction*, 13(8):2122-2125
  12. Hamilton T, Mendes C, Castro L, Assis P, Siquera A (2015) Evaluation of Lasting Effects of Heat Stress on Sperm Profile and Oxidative Status of Ram Semen and Epididymal Sperm, *Oxidative Medicine and Cellular Longevity*, 1-12
  13. Blumer CG, Restelli AE, Giudice PTD, Soller TB, Fraietta R, Nichi M, (2011) Effect of varicocele on sperm function and semen oxidative stress. *BJUI*, 109:259-65
  14. Hammoud AO, Wide N, Gibson M, Barks A, Carrell DT, Meikle AW (2008) Male obesity and alteration in sperm parameters, *Fertil Steril*, 90:2222-2225
  15. World Health Organization: Obesity and overweight. Fact sheet No. 311, updated March 2013
  16. Rufus O, James O, Michael A (2017) Male obesity and semen quality: Any association? *Int J Reprod Biomed*, 16(4):285-290
  17. Rowe PH, Combaire FH, Hargreave TB, Mellows HJ, (1993) WHO manual for standardized and diagnosis of the infertile couple. Cambridge, England: Cambridge University Press
  18. World Health Organization, WHO laboratory manual for the examination of human semen and semen-cervical interaction. Singapore: Press Concern, 1980.

# Male Infertility An Epidemiological and Clinical Profile at the Andrology Unit of Dr. Soetomo Hospital, Surabaya, Indonesia

## ORIGINALITY REPORT

7%

SIMILARITY INDEX

5%

INTERNET SOURCES

5%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.e-journal.unair.ac.id">www.e-journal.unair.ac.id</a> Internet Source	1%
2	<a href="http://e-journal3.unair.ac.id">e-journal3.unair.ac.id</a> Internet Source	1%
3	"Male Infertility", Springer Science and Business Media LLC, 2020 Publication	1%
4	Brahima Kirakoya, Zango Barnabé, Paré Abdoul Karim, Kaboré Fasnéwendé Aristide, Yaméogo Clotaire, Nikièma Amélie. "Epidemiological and Clinical Profile of Male Hypofertility in Consultation at the Urology-Andrology of Yalgado Ouedraogo Teaching Hospital (Burkina Faso)", Advances in Sexual Medicine, 2015 Publication	1%
5	<a href="http://dspace.nuph.edu.ua">dspace.nuph.edu.ua</a> Internet Source	1%
6	<a href="http://www.ijhsonline.com">www.ijhsonline.com</a> Internet Source	<1%

7	<a href="http://www.scirp.org">www.scirp.org</a> Internet Source	<1 %
8	<a href="http://www.actamedindones.org">www.actamedindones.org</a> Internet Source	<1 %
9	"OS18: Interstitial Lung Disease 1", Respirology, 2013 Publication	<1 %
10	<a href="http://cyberleninka.org">cyberleninka.org</a> Internet Source	<1 %
11	<a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a> Internet Source	<1 %
12	"1-A1: Lung Cancer 1 : Poster Sessions", Respirology, 2013. Publication	<1 %
13	Submitted to Universitas Airlangga Student Paper	<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On



# Male Infertility An Epidemiological and Clinical Profile at the Andrology Unit of Dr. Soetomo Hospital, Surabaya, Indonesia

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---