

# CARPATHIAN JOURNAL OF FOOD SCIENCE AND TECHNOLOGY

## SPECIAL ISSUE

### 5<sup>TH</sup> INTERNATIONAL SYMPOSIUM ON PROBIOTICS AND PREBIOTICS

*Vol. 11(5)*  
*2019*



*Technical University of Cluj Napoca*  
*U.T.Press Publishing House*



**Carpathian Journal of Food Science and Technology**

***Print : ISSN 2066-6845***  
***Online : ISSN 2344-5459***  
***ISSN-L 2066-6845***

***Vol. 11, Nr.(5) 2019***  
***Special issue***



***Technical University of Cluj Napoca***  
***U.T.Press Publishing House***



**Editor in Chief:**

Liviu Giurgiulescu -Technical University of Cluj Napoca, North University Center of Baia Mare, Chemistry-Biology Department, [giurgiulescul@yahoo.com](mailto:giurgiulescul@yahoo.com)

**Executive-editor:**

Anca Mihaly-Cozmuta- Technical University of Cluj Napoca, North University Center of Baia Mare, [ancamihalycozmuta@gmail.com](mailto:ancamihalycozmuta@gmail.com)

**Editors:**

Anca Peter- Technical University of Cluj Napoca, North University Center of Baia Mare, [peteranca@yahoo.com](mailto:peteranca@yahoo.com)

Camelia Nicula- Technical University of Cluj Napoca, North University Center of Baia Mare, [vargacamelia@yahoo.com](mailto:vargacamelia@yahoo.com)

Leonard Mihaly Cozmuta - Technical University of Cluj Napoca, North University Center of Baia Mare, [mihalyl@yahoo.com](mailto:mihalyl@yahoo.com)

**Editorial board:**

Prof. dr. Michael Eskin, University of Manitoba, Canada

Prof.dr. Vizireanu Camelia - University of Galați, Faculty of Food Science and Engineering, Romania

Prof.dr. Chifiriuc Mariana Carmen - University of Bucharest, Faculty of Biology, Romania

Prof.dr. Trașcă Teodor - USAMV of Banat, Timisoara, Romania

Dr. Qian Lu-College of Food, Agricultural and Natural Resources Sciences, University of Minnesota, USA

Prof.dr. Monye Felicia Nwanne- University of Nigeria, Faculty of Law, Nigeria

Prof. dr. Jan Bojkovski - Faculty of Veterinary Medicine – University of Belgrade, Serbia

Assoc. prof. PhD, NG EYK, School of Mechanical & Aerospace Engineering, Nanyang Technological University N3.2-02-70, 50 Nanyang Avenue, Singapore 639798

Prof.dr. Vagelas Ioannis -Technological Institute of Larissa, TEI, Department of Crop Protection and Plant Pathology, Greece

Prof. Dr. Claudio De Pasquale, Department Scienze Agrarie, Alimentari e Forestali, Università degli Studi di PALERMO, Italy

Prof.dr. Gerhard Schleining, Department of Food Sciences and Technology BOKU - University of Natural Resources and Life Sciences, Secretary General of the ISEKI-Food Association, Vienna, Austria

Technical University of Cluj Napoca, Romania

U.T. Press Publishing House



CONTENT

|  |              |
|--|--------------|
| <b>Athiyyah A.F., Giurgiulescu L., Editorial introduction special issue “Probiotic, prebiotic, and micronutrients in immune system”</b>  | <b>5</b>     |
| <b>Rahardja F., Shahib M.N., Tjahjani S., Prasetyo D., The inhibition of <i>Salmonella Typhi</i> growth by the cell free supernatans of <i>Lactobacillus acidophilus</i> cultures</b>  | <b>6-10</b>  |
| <b>Athiyyah A.F., Darma A., Ranuh R.G., Riawan W., Sudiana I.K., Surono I.S., Sudarmo S.M., The effect of <i>Lactobacillus plantarum</i> IS-10506 on accelerating the regeneration of the intestinal mucosa structure examined by the scanning electron microscope</b> | <b>11-18</b> |
| <b>Wibowo S., Pratiwi T.P., Proborini A., Susanto W.P., Correlation between mucosal and systemic adaptive immune response after probiotic administration in mouse model</b>  | <b>19-28</b> |
| <b>Darma A., Putra A.M., Fitri P., Athiyyah A.F., Ranuh R.G., Sudarmo M.S., Responses of dendritic and NK cells after multispecies probiotic administration in <i>Balb/c</i> mice</b>  | <b>29-36</b> |
| <b>Athiyyah A.F., Darma A. , Ranuh R.G., Setyawati A., Setyo boedi B., Endaryanto A., Arief S., Sudarmo S.M. Effect of probiotic on innate immune response in the liver of <i>mus musculus Balb/c</i></b>  | <b>37-42</b> |
| <b>Sudarmo S. M., Yamin A., Athiyyah A.F., Darma A., Ranuh R.G. The effect of probiotics, prebiotics and synbiotics in formula milk towards the incidence of acute respiratory infection on 1-5 years old healthy children in day care centres</b>                     | <b>43-49</b> |
| <b>Ranuh R.G., Athiyyah A.F., Darma A., Pramono B., Handini L.S., Shanty M.C., Sudarmo S.M. Effects of heat-killed probiotic complex on acute diarrhoea in children</b>  | <b>50-56</b> |
| <b>Darma A., Athiyyah A. F., Candra L. E., Ranuh R. G., Sudarmo S. M., Effect of preventative zinc supplementation on damage to intestinal integrity caused by <i>Escherichia coli</i> lipopolysaccharide administration: experimental study in an animal model</b>    | <b>57-66</b> |

|   |         |
|---|---------|
| Ranuh R. G., Athiyyah A. F., Darma A., Candra R., Setyoningrum R.A.,<br>Endaryanto A., Hidajat B., Sudarmo S.M. <i>Zinc supplementation altered bronchus<br/>mucosal immune status expressed by IFN-<math>\gamma</math>, IL-6, dendritic cells and sIgA</i> | 67-72   |
| Etika R., <b>Ranuh R. G.</b> , Athiyyah A. F., Darma A., Sudarmo S. M., Suwarno,<br><b><i>Analysis of immunoglobulin levels after exposure of cow's milk protein in mice</i></b>  | 73-77 ✓ |
| Hidayat T., Utamayasa I. K. A., Rahman M. A. , Ontoseno T. <i>Effects of pomegranate<br/>extracts on MMP-1, TIMP-1, MMP-1/TIMP-1 ratio, and type 1 collagen to inhibit<br/>right ventricular fibrosis in animal models</i>                                  | 78-82   |
| Prihaningtyas R. A., Widjaja N.A., Irawan R., Hanindita H.M., Hidajat B. <i>Dietary<br/>intakes and high sensitivity CRP (hsCRP) in adolescents with obesity</i>  | 83-88   |
| Widjaja Nur Aisyah , Irawan R., Hanindita H.M., Prihaningtyas R.A., Hidajat B.<br><i>Anthropometric measurements and inflammatory biomarkers in obese adolescents</i>   | 89-95   |
| Hanindita M. H., Widjaja N.A., Irawan R., Hidayat B. <i>Comparison between baby<br/>led weaning and traditional spoon-feeding on iron status and growth in breastfed<br/>infants</i>  | 96-100  |
| Limanto T. L., Sampurna M. T.A., Handayani K. D., Angelika D., Utomo M. T.,<br>Etika R., Harianto A. <i>The effect of early parenteral nutrition on return to birth<br/>weight and gain weight velocity of premature infants with low birth weight</i>      | 101-107 |
| Handayani K.D., Etika R., Dwi Wijayanti N.A.D., Harianto A., Utomo M.T., Dina<br>Angelika D., Sampurna M.T.A., <i>Gain velocity and IgA secretory<br/>fecal between preterm baby received human milk and human milk fortified</i>                           | 108-121 |



## ANALYSIS OF IMMUNOGLOBULIN LEVELS AFTER EXPOSURE OF COW'S MILK PROTEIN IN MICE

Risa Etika<sup>1</sup>, **Reza Gunadi Ranuh<sup>1</sup>**, Alpha Fardah Athiyyah<sup>1</sup>, Andy Darma<sup>1</sup>, Subijanto Marto Sudarmo<sup>1\*</sup>, Suwarno<sup>2</sup>

<sup>1</sup>Department of Child Health, Dr. Soetomo Hospital, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

<sup>2</sup>Virology and Immunology Laboratory, Faculty of Veterinary, Universitas Airlangga  
\*subijanto.sudarmo@gmail.com

<https://doi.org/10.34302/crpifst/2019.11.5.10>

### Article history:

Received:

9 March 2019

Accepted:

20 September 2019

### Keywords:

Allergen;

Pregnant Balb/c mice;

Neonates Balb/c mice;

Immunoglobulin levels.

### ABSTRACT

This research aimed to analyze immunoglobulin levels after exposure of cow's milk protein in mice. We conducted a true experiment with a posttest-only control group method. This experiment used female Balb/c mice and neonate mice (pregnant mice 10 weeks old n=15, neonatal mice 2 weeks old). The sample was divided into low dose group (1 mg CMP per gram weight of mice), high dose group (10 mg CMP per gram weight of mice) and control group that consisted 5 pregnant mice respectively. The cow's milk protein allergens were induced in pregnant mice. We took blood from the pregnant mice, and their fetuses were taken for subclass immunoglobulin examination (Ig A, IgM, IgE, Ig G) by using an ELISA (Enzyme-Linked Immunosorbent Assay) kit. In low dose group we found a significant increase ( $p<0,05$ ) in IgG-1, IgG-2a, and IgG-2b levels in pregnant mice and IgG-1, IgG-2a, IgG-2b, and IgG-3 in neonate mice which described in table 1. While for the high dose group described in table 2, similar results occurred in IgG-1, IgG-2a, and IgG-3 levels ( $p<0.05$ ) in pregnant mice and IgG-1, IgG-2a, IgG-2b, and IgG-3 in neonate mice. In high-dose group we also found an increase in IgA, IgM, and IgE ( $p<0.05$ ) in either the neonate or pregnant mice. However, in low dose group there was only a significant increase in IgM ( $p=0.004$ ) and IgE ( $p=0,000$ ) in neonate mice and only IgE ( $P=0,000$ ) in pregnant mice. In conclusion, in low dose group we found a significant increase in IgG-1, IgG-2a, IgG-2b in both pregnant and neonate mice, IgG-3 and IgE in pregnant mice, IgM and IgE in neonate mice. While for the high dose group similar results occurred in IgG-1, IgG-2a, IgG-3, IgA, IgM, and IgE levels in both pregnant and neonate mice and IgG-2b in neonate mice.

### 1.Introduction

Allergy is a health problem with a high prevalence. In the human population, the prevalence of allergy has reached 20%, with 5%–15% occurring in children. In Southeast Asia, the prevalence of allergy reached 3.3% in children (Platts-Mills et al., 2003). While the prevalence is getting higher, efforts to prevent allergy is yet to be optimized, and the immunology process of *in utero* prevention has

not been understood (Jones, Holloway, and Warner, 2000; Boyle, Robins-Browne, and Tang, 2006; Fusaro et al., 2009). Today, suspected allergic events are associated with the fetus's exposure to allergens, but this has yet to be explained. Some studies suggest that the child of a pregnant mother whose family has a history of dust mite allergy may become allergic to dust mites while exposure to cat hair

may induce tolerance to cat hair within the child (Prescott et al., 2003). It is predicted that different doses and types of allergen will induce different immune responses from the fetus. A study shows that fetal dendritic cells do not express the Th2 allergen, but they express the Th1 allergen (Platts-Mills et al., 2003). This leads to a discussion on how to treat allergy and whether mothers should expose allergens to their child to prevent allergies from developing or avoid them. Allergy is a chronic disease that can alter the growth and development of the child. Therefore, an experiment on pregnant Balb/C mice is conducted exposing them to low-dose and high-dose cow's milk protein allergen to analyze immunoglobulin levels after exposure of cow's milk protein in mice (Kunert and Lavitska, 2001; Prescott et al., 2003; Lara-Villoslada, Olivares, and Xaus, 2005).

## 2. Materials and methods

We conducted a true experiment with a posttest-only control group method to analyze immunoglobulin levels after exposure of cow's milk protein in mice.

### 2.1. Animals

The study was conducted in the Virology and Immunology Laboratory, Department of Microbiology, Airlangga University, Surabaya, East Java.

This experiment used female Balb/C mice and neonates (pregnant mice 10 weeks old  $n=15$ , neonatal mice 2 weeks old) acquired from the Farma Center of Veterinary, Surabaya. All the animals were inspected by a veterinary consultant to ensure pregnancy and the neonates' health condition. Animals were excluded if they were pregnant with fetuses with congenital disorders, have differences in meal behavior, or show signs of sickness such as decreased weight, breathing patterns, and diarrhea. Cow's milk protein allergens were induced in pregnant mice. After the pregnant mice gave birth, the neonates were kept for two weeks before having their blood taken, and the serum of both the female mice and their fetuses

were examined. The samples were homogeneous in gender, age, and weight. The sample was divided into low dose group (1 mg CMP per gram weight of mice), high dose group (10 mg CMP per gram weight of mice) and control group. Then the data were analyzed using inferential statistics to achieve the research objectives.

### 2.2. Cows Milk Protein Allergen

Materials used were cow's milk protein allergen (Indoor Biotechnologies, Natural Bos d5 (NA-BD5-1). The cow's milk protein allergens were administered intraperitoneal in pregnant mice 5mg/5ml: low dose 1 mg CMP per gram weight of mice (Jones, Holloway, and Warner, 2000; Kunert and Lavitska, 2001; Prescott et al., 2003; Lara-Villoslada, Olivares, and Xaus, 2005). The high dose used the same reagent with 10 times the low dose.

### 2.3. ELISA

Immunoglobulin was analyzed using an IgG, IgM, and IgA ELISA kit with the Sandwich method. The ELISA kit (Ig isotyping Mouse Instant: Thermo Fisher Scientific company, catalog number 88-50660-22) was used to examine the optical density (OD) of IgG1, IgG2a, IgG2b, IgG3, IgM, IgA and IgE.

The 80 ml serums were collected into a microplate tube and processed by duplo. A standard 20  $\mu$ l solution of IgE with a titer of 0, 50, 100, 200, 500, and 1000 was added into each tube, commencing to the A 1 and 2 until the G 1 and 2 columns. The microplate was incubated for an hour in room temperature and then washed three times with buffer washing. Then 100  $\mu$ l of HRP enzyme was added to each tube. The microplate was then incubated for an hour and washed with buffer washing. Next, 100  $\mu$ l of TMB was added and incubated for 30 minutes.

Absorbant value was analyzed using the 450 nm wavelength of the ELISA reader and then interpreted by linear regression analysis.

### 2.4. Statistics Analysis

The experiment was analyzed using ANOVA homogeneity test and Kolmogorov-

Smirnov and normal probability plot for normality test for normal distribution. For testing differences between each group, we used ANOVA for normal distributed data and Kruskal Wallis A, Brown-Forsythe, and Mann-Whitney for abnormal distributed data.

### 3. Results and discussions

#### 3.1. Results

In low dose group we found a significant increase in IgG-1, IgG-2a, and IgG-2b levels in pregnant mice ( $p < 0.05$ ) and IgG-1, IgG-2a, IgG-2b, and IgG-3 ( $p < 0.05$ ) in neonate mice which described in table 1. While for the high dose group described in table 2, similar results occurred in IgG-1, IgG-2a, and IgG-3 levels ( $p < 0.05$ ) in pregnant mice and IgG-1, IgG-2a,

IgG-2b, and IgG-3 in neonate mice ( $p < 0.05$ ). Between high-dose and low-dose group, we found significant differences in IgG-1, IgG-2a, IgG-2b, and IgG-3 in neonate mice ( $p(r-t) < 0.05$ ) and IgG-1, IgG-2a, and IgG-2b in pregnant mice ( $p(r-t) < 0.05$ ). In high-dose group we found an increase in IgA, IgM, and IgE ( $p < 0.05$ ) in either the neonate or pregnant mice. However, in low dose group there was only a significant increase in IgM ( $p = 0.004$ ) and IgE ( $p = 0.000$ ) in neonate mice and only IgE ( $P = 0.000$ ) in pregnant mice. In the high-dose group and the low-dose group, there was a significant increase in IgA, IgM, and IgE in either the neonate or pregnant mice ( $p(r-t) < 0.05$ ).

**Table 1.** Immunoglobulin level after being exposed to low dose cow's milk allergen

|        | Low Dose Group (p)          |                             | Control Group |              |
|--------|-----------------------------|-----------------------------|---------------|--------------|
|        | Pregnant mice               | Neonate mice                | Pregnant mice | Neonate mice |
| IgG-1  | 1.85± 0.07<br>(p = 0.000)   | 1.90± 0.08<br>(p = 0.004)   | 1.26± 0.07    | 0.18± 0.03   |
| IgG-2a | 2.66 ± 0.04<br>(p = 0.000)  | 2.88 ± 0.23<br>(p = 0.000)  | 1.41 ± 0.05   | 1.48 ± 0.18  |
| IgG-2b | 3.38± 0.08<br>(p = 0.000)   | 2.87± 0.42<br>(p = 0.001)   | 2.58± 0.06    | 1.95± 0.18   |
| IgG-3  | 0.47± 0.14<br>(p = 0.078)   | 2.09± 0.47<br>(p = 0.000)   | 0.36± 0.04    | 0.09± 0.01   |
| IgA    | 0.16 ± 0.05<br>(p = 0.749)  | 0.12 ± 0.03<br>(p = 0.192)  | 0.15 ± 0.09   | 0.10 ± 0.04  |
| IgM    | 1.01 ± 0.05<br>(p = 0.184)  | 0.73 ± 0.10<br>(p = 0.004)  | 1.16 ± 0.22   | 0.24 ± 0.08  |
| IgE    | 183.21± 5.91<br>(p = 0.000) | 109.04±10.61<br>(p = 0.000) | 21.89 ± 0.84  | 22.37 ± 3.01 |

**Table 2.** Immunoglobulin level after being exposed to high dose cow's milk allergen

|        | High Dose Group (p)         |                             | Control Group |              |
|--------|-----------------------------|-----------------------------|---------------|--------------|
|        | Pregnant mice               | Neonate mice                | Pregnant mice | Neonate mice |
| IgG-1  | 1.49± 0.04<br>(p = 0.000)   | 1.31± 0.09<br>(p = 0.004)   | 1.26± 0.07    | 0.18± 0.03   |
| IgG-2a | 1.33 ± 0.46<br>(p = 0.688)  | 0.76 ± 0.17<br>(p = 0.000)  | 1.41 ± 0.05   | 1.48 ± 0.18  |
| IgG-2b | 2.68± 0.05<br>(p = 0.009)   | 2.39± 0.12<br>(p = 0.001)   | 2.58± 0.06    | 1.95± 0.18   |
| IgG-3  | 0.45± 0.05<br>(p = 0.007)   | 0.20± 0.05<br>(p = 0.001)   | 0.36± 0.04    | 0.09± 0.01   |
| IgA    | 0.32 ± 0.05<br>(p = 0.002)  | 0.04 ± 0.03<br>(p = 0.020)  | 0.15 ± 0.09   | 0.10 ± 0.04  |
| IgM    | 1.55 ± 0.05<br>(p = 0.007)  | 0.52 ± 0.11<br>(p = 0.000)  | 1.16 ± 0.22   | 0.24 ± 0.08  |
| IgE    | 192.96± 7.32<br>(p = 0.000) | 124.26± 9.31<br>(p = 0.000) | 21.89 ± 0.84  | 22.37 ± 3.01 |



### 3.2. Discussion

After exposure to cow's milk allergen, there were different responses from IgG2b both in pregnant mice and neonate mice and with high doses and low doses whereas for IgG3, significant response was only in pregnant and neonate mice with high-dose exposure and neonate mice with low-dose exposure. It can be explained in general that exposure to high doses and low doses of both pregnant and neonate mice gives the same response, meaning that the cow's milk antigen can be transferred through the placenta to the neonate and responded to by neonate plasma cells even with a lower response ability in neonate mice. Neonatal B cells express low levels of coreceptors including CD28 and CD40 ligand on Th2 or follicular T helper cells with their corresponding binding partners HLA-peptide, CD80/86, and CD40 on antigen-specific B cells. This limits their capacity to respond to B cells from neonates and infants aged less than two months. Hence, the immune system is immature and markedly impaired in neonates and more so in fetuses (Mcmichael, Simon, and Hollander, 2015).

Response to the exposure to high-dose cow's milk allergen occurred in IgA and IgM whereas for low-dose cow's milk, there was no significant increase from the control. It can be explained that the immune system remains inferior to exposure whereas the IgE response shows a potent allergy parameter both in pregnant mice and neonates with allergy. In food allergy, it is believed that food-specific IgE antibodies bind to high-affinity FcεRI receptors on mast cells, basophils, macrophages, and dendritic cells, as well as to low-affinity FcεRII receptors on macrophages, monocytes, lymphocytes, eosinophils, and platelets. When food allergens penetrate mucosal barriers and contact IgE antibodies bound to mast cells or basophils, histamine and other mediators that induce symptoms of immediate hypersensitivity are released (Li et al., 2005)

We found significant differences in IgE levels between pregnant mice and neonate mice

exposed to low-dose and high-dose allergens. Efforts to induce allergy tolerance in mice exposed to high-dose allergens were unsuccessful. This is contrary to a study conducted by Adel-Patient (2005), where the dose of allergens exposed did not affect IgE regulation. There was an increase in IgE levels both after the exposure to low-dose or high-dose allergens (Adel-Patient, Ah-Leung, Creminon, Nouaille, Chatel, Langella, and Wal, 2005; Wavrin, 2015).

### 4. Conclusions

In low dose group we found a significant increase in IgG-1, IgG-2a, IgG-2b in both pregnant and neonate mice, IgG-3 and IgE in pregnant mice, IgM and IgE in neonate mice. While for the high dose group similar results occurred in IgG-1, IgG-2a, IgG-3, IgA, IgM, and IgE levels in both pregnant and neonate mice and IgG-2b in neonate mice.

### 5. References

- Adel-Patient, K. et al. (2005). Oral administration of recombinant *Lactococcus lactis* expressing bovine beta-lactoglobulin partially prevents mice from sensitization, *Clinical and Experimental Allergy: Journal of the British Society for Allergy and Clinical Immunology*, 35(4), pp. 539–46.
- Adel-Patient, K. et al. (2011). Immunological and metabolomic impacts of administration of Cry1Ab protein and MON 810 maize in mouse, *PLoS ONE*. Edited by M. Fessler, 6(1), p. e16346.
- Aichbhaumik, N. et al. (2008). Prenatal exposure to household pets influences fetal immunoglobulin E production, *Clinical & Experimental Allergy*.
- Boyle, R. J., Robins-Browne, R. M., and Tang, M. L. K. (2006). Probiotic use in clinical practice: what are the risks?, *American Journal of Clinical Nutrition*.
- Briner, T. J. et al. (1993). Peripheral T-cell tolerance induced in naive and primed mice by subcutaneous injection of peptides from the major cat allergen Fel d I., *Proceedings of the National Academy of Sciences of the*

- United States of America*, 90(16), pp. 7608–12.
- Bufford, J. D. et al. (2008). Effects of dog ownership in early childhood on immune development and atopic diseases, *Clinical & Experimental Allergy*, 38(10), pp. 1635–1643.
- Fusaro, A. E. et al. (2009). Balance between early life tolerance and sensitization in allergy: dependence on the timing and intensity of prenatal and postnatal allergen exposure of the mother, *Immunology*, 128(1 Suppl), pp. e541–50.
- Jones, C. A., Holloway, J. A., and Warner, J. O. (2000). Does atopic disease start in foetal life?, *Allergy*, 55(1), pp. 2–10.
- Kunert, H. W. and Lavitska, E. (2001). Stresses and strains in anisotropic cubic ultra-thin overlayers, *Crystal Research and Technology*.
- Lara-Villoslada, F., Olivares, M., and Xaus, J. (2005). The balance between caseins and whey proteins in cow's milk determines its allergenicity, *Journal of Dairy Science*.
- Li, X. et al. (2005). A murine model of IgE-mediated cow's milk hypersensitivity, *Journal of Allergy and Clinical Immunology*, 103(2), pp. 206–214. doi: 10.1016/s0091-6749(99)70492-6
- McMichael, A., Simon, A. K., and Hollander, G. A. (2015). Evolution of the immune system in humans from infancy to old age, *Proceedings of the Royal Society B, Biological Sciences*, 282: 20143, pp. 1–9. doi: 10.1098/rspb.2014.3085
- Platts-Mills, T. A. E. et al. (2003). The relevance of maternal immune responses to inhalant allergens to maternal symptoms, passive transfer to the infant, and development of antibodies in the first 2 years of life, *Journal of Allergy and Clinical Immunology*.
- Prescott, S. L. et al. (2003). The value of perinatal immune responses in predicting allergic disease at 6 years of age, *Allergy: European Journal of Allergy and Clinical Immunology*.

# Source details

## Carpathian Journal of Food Science and Technology

Open Access ⓘ

Scopus coverage years: from 2009 to Present

Publisher: North University of Baia Mare

ISSN: 2066-6845

Subject area: Agricultural and Biological Sciences: Food Science

[View all documents >](#) [Set document alert](#) [Save to source list](#)

CiteScore 2019  
**0.3** ⓘ  
[Add CiteScore to your site](#)

SJR 2019  
**0.135** ⓘ

SNIP 2019  
**0.172** ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

**Improved CiteScore methodology** ⓘ

CiteScore 2019 counts the citations received in 2016-2019 to articles, reviews, conference papers, book chapters and data papers published in 2016-2019, and divides this by the number of publications published in 2016-2019. [Learn more >](#)

**CiteScore 2019** ▾

**0.3** =  $\frac{100 \text{ Citations 2016 - 2019}}{290 \text{ Documents 2016 - 2019}}$

Calculated on 06 May, 2020

**CiteScoreTracker 2020** ⓘ

**0.4** =  $\frac{93 \text{ Citations to date}}{221 \text{ Documents to date}}$

Last updated on 10 June, 2020 • Updated monthly

### CiteScore rank 2019 ⓘ

| Category                             | Rank     | Percentile |
|--------------------------------------|----------|------------|
| Agricultural and Biological Sciences | #257/299 | 14th       |
| Food Science                         |          |            |

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [🔗](#)

#### About Scopus

- [What is Scopus](#)
- [Content coverage](#)
- [Scopus blog](#)
- [Scopus API](#)
- [Privacy matters](#)

#### Language

- [日本語に切り替える](#)
- [切换到简体中文](#)
- [切换到繁體中文](#)
- [Русский язык](#)

#### Customer Service

- [Help](#)
- [Contact us](#)



Ads by Google

Stop seeing this ad Why this ad?

# Carpathian Journal of Food Science and Technology

Country [Romania](#) - SIR Ranking of Romania

Subject Area and Category [Agricultural and Biological Sciences](#)  
[Food Science](#)

Publisher [North University of Baia Mare](#)

Publication type Journals

ISSN 20666845

Coverage 2009-2019

**Scope** The Carpathian Journal of Food Sciences and Technology publishes original scientific peer-reviewed double blinded papers that cover all aspects of food science from raw materials through processing and economical aspects up to final products and their nutritional characteristics. Both fundamental and applied research is accepted. The main topics are: - Food Chemistry - Food Processing Technology - Health, Nutrition and Food - Toxicology and Food Safety - Food Analysis - Food Policy - Food marketing - Consumer protection - Food packaging



[Homepage](#)

[How to publish in this journal](#)

[Contact](#)



[Join the conversation about this journal](#)


# 6


H Index

Ads by Google


Stop seeing this ad Why this ad?

## Quartiles










Citations per document



● Cites / Doc. (4 years)  
● Cites / Doc. (3 years)  
● Cites / Doc. (2 years)



Carpathian Journal of Food Science and Technology  
 Food Science  
**Q4**  
 SJR 2019 0.14  
 best quartile  
 powered by scimagojr.com

← Show this widget in your own website  
 Just copy the code below and paste within your html code:  

```
<a href="https://www.scimagojr.com" style="color: red; text-decoration: none;">

```

Ads by Google

[Stop seeing this ad](#)   [Why this ad?](#)