ISSN 2599 - 0993



J. Parasite Sci. Vol. 5 No. 1 Hal. 1 - 34 Surabaya, Maret 2021

PUBLISHER COLLABORATION

Journal of Parasite Science



Collaboration Division of Veterinary Parasitology Faculty of Veterinary Medicine, Universitas Airlangga and Indonesian Veterinary Parasitology Association



EDITOR'S ADDRESS

Editorial Office Journal of Parasite Science

Division of Veterinary Parasitology Faculty of Veterinary Medicine, Universitas Airlangga Kampus "C" Jl. Mulyorejo Surabaya 60115 Phone: (031) 5992785; 5993016; Fax. (031) 5993015 e-mail: jparasitol@gmail.com ; jps@fkh.unair.ac.id Homepage : https://e-journal.unair.ac.id/JOPS

EDITORIAL BOARD

Chief Editor

Dr. Kusnoto, DVM., M.Si., Universitas Airlangga, Indonesia

Chief's Assistant Editor

Prof. Muchammad Yunus, DVM., M.Kes., Ph.D., Universitas Airlangga, Indonesia

Editorial Board Member

Prof. Dr. Lucia Tri Suwanti, DVM., MP. Universitas Airlangga, Indonesia
 Prof. Dr. Makoto Matsubayashi, DVM., Ph.D., Osaka Perfecture University, Japan
 Assoc. Prof. Dr. Mahmut Kabalak, Hacettepe University, Turkey
 Dr. Attaur Rahman, Abdul Wali Khan University Mardan, Pakistan
 Dr. Muhammad Hassan Mushtaq, University of Veterinary and Animal Sciences Lahore, Pakistan
 Aondohemba Samuel Nege, B.Fish., M.Sc., National Taiwan University, Taiwan, Province of China
 Muhammad Thohawi Elziyad Purnama, DVM., M.Si. PSDKU Universitas Airlangga, Banyuwangi

Administation Staff

Dyah Ayu Kurniawati, DVM., M.Sc, Universitas Airlangga, Indonesia Moch Arifudin, Universitas Airlangga, Indonesia

REVIEWERS

Editors, Authors and Readers of Journal of Parasite Science give the highest appreciation and gratitude to the experts below, as reviewers who has reviewed all the articles, whether published or rejected, according to the recommendations submitted to the editor in Vol. 5 No. 1 March 2021

April Hari Wardhana, DVM, M.Sc, Ph.D., Balai Besar Penelitian Veteriner Bogor, Bogor, Indonesia

Prof. Dr. Sri Subekti, DVM., DEA., Perkumpulan Pemberantasan Penyakit Parasit Indonesia

Prof. Dr. Upiek Kesumawati Hadi, DVM., MS., Institut Pertanian Bogor, Bogor, Indonesia

Dr. Raden Wisnu Nurcahyo, DVM., Universitas Gadjah Mada, Yogyakarta, Indonesia

Dr. Dwi Priyowidodo, DVM., MP., Universitas Gadjah Mada, Yogyakarta, Indonesia

Dr. Nyoman Adi Suratma, DVM., MP., Universitas Udayana, Bali, Indonesia

Prof. Hiroshi Sato, DVM., Ph.D., Yamaguchi University, Jepang

Prof. Graeme Martin, B.Sc(Agric)., Ph.D W.Aust., University of Western Australia School of Agriculture and Environment, Australia

Prof. Dr. Muhammad Salisu Abubakar., Usmanu Danfodiyo University Sakoto, Nigeria Dr. Zainab Alwan Makawi., University of Baghdad, Iraq

VISION DAN MISSION

Journal of Parasite Science (JoPS) with registered number <u>P-ISSN (2599-0993)</u>; <u>E-ISSN (2656-5331)</u> published by the Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga.

Journal of Parasite Science (JoPS) presenting research articles, case reports, community development activities and literature studies of veterinary medicine. Published twice a year on March and September. Since 2002 the Faculty of Veterinary Medicine has had Veterinary Medicine Media, because considering that there is only one journal that must accommodate many articles written by lecturers and students, various journals have been developed according to the disciplines in each department, one of which is the JoPS. JoPS can publish articles from various faculties and institutions related to the parasitology. JoPS which has been published since 2017 is here for accommodate publication obligations for undergraduate and postgraduate students as well as other parties conducting research in the field of parasitology. Loading of articles in the JoPS through the Open Journal System (OJS). Complete information for article loading and article writing instructions are available on the website and every issue. Incoming articles will go through a selection process by editors and reviewers.

VISION

To become a leading and reputable journal at the national and international level in veterinary parasitology science.

MISSION

- 1. Provide scientific communication media of Parasitology to participate in advancing science and technology in related sector;
- 2. Organizing the publication of scientific journals of Parasitology with a high impact factor in the development of science and technology;
- 3. Making journals as terms for the advancement and intellectual development of the academic community in welcoming the Universitas Airlangga World Class University;
- 4. Organizing an accountable and quality journal management to increase the number of intellectual products in the form of scientific journals;
- 5. Become a leading reference for the academic community and researchers of veterinary parasitology and published as a scientific journal.

Journal Scope: Journal of Parasite Science (JoPS) publishes the results of original research in all aspects of basic and applied parasitology, and ranging from parasites biodiversity, parasites of all wildlife, invertebrate and vertebrate, as well as host-parasite relationships of intrinsic biological interest to those of social and economic importance predominately in veterinary, human medicine and agriculture aspect. Original research includes the development of novel and innovative concepts and ideas, as well as experimental and observational science that raises new theory.

Language:

Main: English

The articles accepted by the Journal of Parasite Science (JoPS) are:

- 1. Research articles;
- 2. Case reports;
- 3. Community development articles;
- 4. Literature study articles.

TABLE OF CONTENS

Page

1.	The Potency of Anting Anting (<i>Acalypha indica</i> L.) Leaf Extract as An Acaricide on <i>Boophilus microplus</i> in Larvae and Adult Stages <i>In Vitro</i> (Fitri Nurdiana, Rahmi	
	Sugiharuti, Bambang Sektiari Lukiswanto)	1 - 4
	Prevalence and Scabies Severity Rate on a Goat at Mempawah Hilir, Mempawah	
2.	District West Kalimantan Province (Arum Maula Hidayah, Wiwiek Tyasningsih,	- 10
	Social Social Derecitor Information on Stray Cat and Dat Cat at Plitar Decency Fast	5 - 10
	Lava Province (Egga Johan Prantanto, Bambang Purnomo Sunardi, Budiarto, Agus	
3.	Wijava)	11 – 14
	The Effectiveness of Bandotan (Ageratum convzoides L.) Leaf Extract on Mortality of	11 14
4	Boophilus microplus In Vitro (Dwi Retno Widyawati, Iwan Sahrial Hamid, Moh.	
4.	Sukmanadi, Sri Agus Sudjarwo, Muhammad Hambal)	15 - 18
	Identification of Ectoparasites and Endoparasites on Java Langurs (Trachypithecus	
_	sp.) and Silvery Gibbons (Hylobates moloch) in The Aspinall Foundation Indonesia	
5.	Program (Ghifari Lutfi Fauzi, Endang Suprihati, Poedji Hastutiek, Boedi Setiawan,	
	Retno Wulansari)	19 - 24
	Repellent Effectiveness of Permot Leaf Ethanol Extract (<i>Passiflora Foetida</i> Linn.)	
6.	against Aedes Aegypti Adult Mosquitoes (David Mohamad Qadafi, Poedji Hastutiek,	
	LIIIK Masiachan, Endang Suprinati, Munammad Hambal)	25 - 30
	Surabaya City Based on Fecal Examination (Nurin Ardhiani Endang Supribati Aditya	
7.	Yudhana Faisal Fikri)	21 - 24
	i danana, i albai i inti j	דכ יכ

AUTHOR INDEX

Aditya Yudhana,..... 31 Agus Wijaya,.... 11 Ahmed,..... 33, 34 Arum Maula Hidayah,..... 5 Astuti,.... 2, 3, 4 Bambang Purnomo Sunardi,.... 11 Bambang Sektiari Lukiswanto,..... 1 Boedi Setiawan,.... 19 Budiarto,.... 11 David Mohamad Qadafi,..... 25 Dinata,..... 3, 4 Dwi Retno Widyawati,..... 15 Egga Johar Praptanto,..... 11 Endang Suprihati,..... 19, 25, 31 Estuningsih,..... 11, 13 Faisal Fikri,..... 31 Farjana,..... 32, 34 Fitri Nurdiana,.... 1 Ghifari Lutfi Fauzi,.... 19 Gunandini,..... 15, 18 Hanafiah,..... 19, 23 Hartati,..... 8, 9 Hastutiek P,..... 26, 28, 29 Hidayat,..... 12, 13

Husna,.... 1, 4 Iwan Sahrial Hamid,..... 15 Lastuti,..... 5, 9 Lilik Maslachah,..... 25 Manurug,..... 15, 19 Melinda,..... 17, 19 Moh. Sukmanadi,..... 15 Mufasirin,..... 33, 36 Muhammad Hambal,..... 15, 25 Nurin Ardhiani,..... 31 Padmasari,..... 26, 29 Poedji Hastutiek,..... 19, 25 Rahmi Sugihartuti,.... 1 Retno Wulansari,.... 5, 19 Samsudin,..... 16, 17, 19 Sasmita,.... 5, 9 Soeharsono,..... 5 Sri Agus Sudjarwo,..... 15 Suprihati E,..... 23, 36 Susilowati,..... 26, 29 Wardhana,..... 2, 4 Werdiningsih,..... 28, 30 Wiwiek Tyasningsih, 5

SUBJECT INDEX

Acalypha indica,.... 1, 2, 4, 5 acaricide,..... 1, 3 Adult stage, 1 Aedes aegypti..... 25, 27, 28, 29, 30 Ageratum conyzoides..... 15, 16, 18, 19 Blitar regency,..... 11 Boophilus microplus.... 1, 2, 4, 5, 15, 16, 19 Chelonoidis carbonaria..... 31, 32, 34, 35 Ectoparasite..... 19, 24 Endoparasite 19, 31, 32, 34 ethanol ectract..... 25 fecal examination..... 31, 32 Gastrointestinal..... 11, 12, 31, 32, 34 goats..... 5, 6, 7, 8, 9 in vitro, 1, 2, 3 Java Langur.... 19, 20, 23 leaf extract..... 1, 2, 3, 15, 16, 17, 18 Mempawah Hilir..... 5, 6, 8, 9

mosquito..... 25, 26, 27, 28 neguvon.... 15, 16 Passiflora foetida..... 25, 26, 29, 30 Permot Leaf..... 25 pet cat..... 11, 12, 13 prevalence..... 5, 9, 11, 13, 19, 20, 23 primate..... 19, 23 probit analysis..... 15, 17 Repellent..... 25, 26, 27, 28, 29, 30 scabies..... 5, 6, 7, 8, 9 severity..... 5, 6, 7, 8, 9 Silvery Gibbon.... 19 skin disease..... 5 stray cat..... 11, 12, 13 Surabaya city, 31 tortoise..... 31, 32, 34, 35 Toxocara cati..... 11, 12, 13

SUBSCRIBED FORM

	Issue* (give sign √)			
AREA	1 year	2 year	3 year	
Surabaya	□ Rp. 200.000,-	□ Rp. 350.000,-	🗆 Rp. 450.000,-	
East Java	🗆 Rp. 250.000,-	□ Rp. 400.000,-	🗆 Rp. 500.000,-	
Other Java	□ Rp. 300.000,-	□ Rp. 450.000,-	🗆 Rp. 550.000	
Other Countries	🗆 USD 50	□ USD 75	□ USD 100	

*(published 2 times a year)

I propose to subscribe to Journal of Parasite Science
Name:
Institution:
Office address and Phone No:
Home address and Phone No:
Country:
Fax:
Email:
Postal code:
Date:
Signature:
I pay for this journal subscription using (give sign $$):
Western Union Cheque with MTCN No:
□ Bank transfer / online to:
Bank Mandiri
Account no 142 oo 1670625 o
Dr. Kusnoto, drh., M.Si
Send proof of payment with the subscription form to email jps@fkh.unair.ac.id

AUTHOR GUIDELINES

Journal of Parasite Science (JoPS) with registered number <u>P-ISSN (2599-0993)</u>; <u>E-ISSN</u> (2656-5331) published by the Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga.

Journal of Parasite Science (JoPS) presenting research articles, case reports, community development activities and literature studies of veterinary medicine. Published twice a year on March and September. Since 2002 the Faculty of Veterinary Medicine has had Veterinary Medicine Media, because considering that there is only one journal that must accommodate many articles written by lecturers and students, various journals have been developed according to the disciplines in each department, one of which is the JoPS. JoPS can publish articles from various faculties and institutions related to the parasitology. JoPS which has been published since 2017 is here for accommodate publication obligations for undergraduate and postgraduate students as well as other parties conducting research in the field of parasitology. Loading of articles in the JoPS through the Open Journal System (OJS). Complete information for article loading and article writing instructions are available on the website and every issue. Incoming articles will go through a selection process by editors and reviewers.Manuscripts should be typed in double spaced, except for titles, abstracts, titles of tables and tables, headings of figures, reference, and attachments, typed according to separate provisions. Paragraph formatting: new paragraph begins (first line) 0.3 inches. The standard font for writing is Times New Roman 12. Using A4 paper size (8.27 x 11.69)". Using English. Tables / Illustrations / Images must be of high contrast, also include separate scanning files (photos) with papers in JPG file format. Table captions, pictures or other explanations in the attachment should be typed 1 (one) space.

Title should be concise but informative, as far as possible in no more than 12 words.

Abstract written in English, do not exceed 250 words, containing elements of background, material and methods, results and conclusions.

Keywords maximum of 6 (six) words or phrases, written after the abstract in each languages, alphabetically ordered. As far as possible avoid using keywords from the title.

Introduction consists of the background, research objectives, problem formulation and benefits. The background includes four main ideas, namely: problem statement, problem scale, problem chronology and research approach. Contains a description of the facts and expectations for the completion of the research. Described based on concepts built from references in the literature. The research question and/or hypothesis (if any) in the introduction should not write in the form of points or should not be numbered.

Method in details so that other author could do the similar research with the written method in the article. Materials and methods are brief descriptions of the research design, how to conduct research and mention the materials/ objects and tools used.

Results and Discussion compared with other research that ever conducted or discuss based on the theoretical framework that is used in the research. Use up-to-date references such as books that published in the last ten years and journal articles that published in the last five years (except the reference that being recommended by the reviewer in this field).

Tables are made without vertical lines (use only lines at the top and bottom of the table as well as for separating heading from the main table), with table title placed before the table, numbered in Arabic numerals (Table X), and have to be referred in the text. The description of the table is placed after the table; it must be concise but clear enough so that the table separately can be understood without referring to the text. The table along with the title and description are placed after the References.

Figures presented are only those that support the findings of the study, and not restatement of data from tables in the form of figures. When resulted data in the form of figure is more informative, interesting or significant, presentation of data in table form is not required. Figure title is placed after the figure, numbered in Arabic numerals (Figure Y), and have to be referred in the text. The description of the figure is placed under the title of the figure; it must be concise but clear enough so that the figure separately can be understood without referring to the text. Image (in JPEG format) is sent in separate file. The title and description of the figure are placed after the References.

Conclusion in one until two paragraphs without using the bullet points or number.

Approval of Ethical Commission have to be stated (number and institution) if the manuscript is constructed based on a research using live animals.

Acknowledgements are delivered to the research funders (state the name, number and recipient name of the grant, if applicable), and to those who have helped carry out the research.

References should be written according to the APA guidelines, that uses the family name of the author at the beginning of the sequence. Example: Family Name, Abbreviated the First Name and the Middle Name. Do not use commas or point for the abbreviated name. For the year of the references give bracket without points. The reference from a journal article, anthology books or book chapter must include the page. References from the internet not journals are not recommended.

The example of the references as below: <u>Online journal article</u>

Johnson, K. D., Winkelman, C., Burant, C. J., Dolansky, M., & Totten, V. (2014). The factors that affect the frequency of vital sign monitoring in the emergency department. JEN: Journal of Emergency Nursing, 12(2), 27-35. https://doi.org/10.1016/j.jen.2012.07.023

Book (authored or entire book) *

Judd, D. M., Sitzman, K., & Davis, M. (2010). A history of American nursing: Trends and eras. Jones and Bartlett Publishers.

Chapter in an edited book *

Lynch, S. (2010). Pain management. In S. R. Hardin and R. Kaplow (Eds.), Cardiac surgery essentials for critical care nursing (3rd ed., pp. 287-309). Jones & Bartlett. PowerPoint slides (from classroom website)

Nicholson, A. (n.d.). RN to BSN program orientation [PowerPoint slides]. Canvas @UIowa: <u>https://uiowa.instructure.com/</u>

<u>Webpage</u>

Nelson, L. (2019). Nurses say violent assaults against healthcare workers are a silent epidemic. <u>https://nurse.org/articles/-</u> workplace-violence-in-nursing-and-<u>hospitals/</u>

<u>Report</u>

Institute of Medicine (2010). The future of nursing: Leading change, advancing health [report brief]. <u>https://www.nap.edu/resource/12956/Future-of-Nursing-2010-Report-Brief.pdf</u>

Review Article

Journal of Parasite Science received review articles on topics included in the scope of the journal which are of current interest. Reviews are invited by the editor. The review must be a comprehensive analysis and perspective on the state of the field and where it is heading. Reviews will be subject to the same peer review process as is applied for original papers. The manuscript is arranged in the same way as the original article with an unstructured abstract (maximum of 250 words). Manuscripts should be typed in double spaced, except for titles, abstracts, titles of tables and tables, headings of figures, reference, and attachments, typed according to separate provisions. Paragraph formatting: new paragraph begins (first line) 0.3 inches. The standard font for writing is Times New Roman 12. Using A4 paper size (8.27 x 11.69)". Using English. Tables / Illustrations / Images must be of high contrast, also include separate scanning files (photos) with papers in JPG file format. Table captions, pictures or other explanations in the attachment should be typed 1 (one) space.

Case reports:

Journal of Parasite Science receives quality and interesting case reports with topics included in the scope of this journal. All reports are peer reviewed as is applied for original articles. All instructions are the same as for the original articles.





Division of Veterinary Parasitology Faculty of Veterinary Medicine, Universitas Airlangga Kampus "C" Jl. Mulyorejo Surabaya 6015 Phone: (031) 5992785; 5993016; Fax. (031) 5993015 e-mail: jparasitol@gmail.com ; jps@fkh.unair.ac.id Website : https://e-journal.unair.ac.id/JoPS





Dimensions **PKP**|INDEX

Google Scrossref

JOURNAL OF PARASITE SCIENCE INDEXING BY :

OpenAIRE ROAD - C GARUDA A MENDELEY

HARVARD HOLLIS

Editorial Office Journal of Parasite Science Division of Veterinary Parasitology

Faculty of Veterinary Medicine, Universitas Airlangga

Kampus "C" Jl. Mulyorejo Surabaya 60115

Phone: (031) 5992785; 5993016; Fax. (031) 5993015 e-mail: jparasitol@gmail.com ; jps@fkh.unair.ac.id

00023950

View My Stats

000

International License.

₩





Pusat Pengembangan Jurnal dan Publikasi Ilmiah Universitas Airlangga copyright@2017 Template PPJPI

Repellent Effectiveness of Permot Leaf Ethanol Extract (*Passiflora Foetida* Linn.) against *Aedes Aegypti* Adult Mosquitoes

¹⁾David Mohamad Qadafi, ²⁾Poedji Hastutiek^(b), ³⁾Lilik Maslachah, ²⁾Endang Suprihati, ⁴⁾Muhammad Hambal

¹⁾Student, Faculty of Veterinary Medicine, Universitas Airlangga, davidqadafi@gmail.com
 ²⁾Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga
 ³⁾Division of Basic Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga
 ⁴⁾Department of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Syiah Kuala
 Corresponding author: poedjihastutiek@gmail.com

Abstract

Indonesia is one of the largest tropical countries in the world and various diseases can arise in the tropics which are caused by animals as vectors. An example of a vector that can carry diseases is a mosquito. Mosquitoes are insects that live side by side with humans buy act as vectors of disease. Mosquito *Aedes aegypti* is a type of mosquito that can carry the virus that causes Dengue Hemorrhagic Fever. This research was conducted to determine the effectiveness of Permot leaf ethanol extract (*Passiflora foetida* L.) as an alternative repellent against adult mosquitoes *Aedes aegypti*. This research was conducted from October to December 2020 and used Permot leaf ethanol extract consisting of 3 cream concentrations, namely 2.5%, 5%, 7.5%, negative control using cream without permot leaf ethanol extract and positive control using mosquito cream. The data of this study were tested using one way ANOVA to find the effectiveness rate and comparations of the each Permot leaf repellent extract. This study proven that the permot leaf ethanol extract is effective as a repellent for *Aedes aegypti* mosquitoes.

Keywords : Permot Leaf, *Passiflora foetida*, *Aedes aegypti*, Repellent, ethanol ectract, mosquito.

Introduction

The Aedes aegypti mosquito is a type of mosquito that can carry the virus that causes Dengue Hemorrhagic Fever (DHF) (Ishartadiati, 2010). This mosquito usually bites in the morning until noon and reproduces by laying eggs in ditches, as well as in other places.

According to Taslimah (2014) mosquitoes are insects that live side by side with humans but act as disease vectors. Various diseases have been spread by no less than 2,500 species of mosquitoes (Herawati, 2010). Containers such as plastic waste, used bottles, used tires, cans and so on during the rainy season can become a breeding ground for mosquito larvae (Romandani, 2019).

Infectious diseases from the mosquito vector *Ae. aegypti* which has recently been endemic in Indonesia is dengue fever, the infectious disease which spread by the mosquito vector *Ae. aegypti*. Dengue Hemorrhagic Fever is an infectious disease caused by the dengue virus which can be transmitted through the bite of *Ae*.

aegypti, this disease is still a health problem in Indonesia. Effective prevention efforts are through controlling the mosquito in the environment.

Mosquitoes can also transmit diseases to animals not only to humans. The impact of disease on animals can cause losses for farmers due to decreased production of livestock. The Ministry of Health of the Republic of Indonesia (Depkes RI, 2000) states that mosquitoes are the most dangerous animals on earth in terms of the number of diseases that cause death, and the economic loss caused by mosquitoes.

Efforts to control *Ae. aegypti* as a vector of dengue disease which is widely applied today using a fumigation system and the use of insecticide Abate SG (Temephos 1%). One of the practical vector control methods is to use mosquito repellent (Murrukmihadi and Dias, 2015). Repellent is a compound that works locally or at a certain distance, which has the ability to prevent mosquitoes from flying, landing or biting the surface of human skin



EY NO SA This work is licensed under a Creative Commons Attribution-Share Alike 4.0 International License.

(Djatmiko *et al.*, 2010). Repelent that is often used by the public is in the form of cream. Synthetic repellents in the form of creams on the market still contain N, N-diethyl-m-toluamide (DEET) which can cause toxic effects. DEET is a synthetic repellent which has a broad spectrum and has been widely used in Indonesia.

The active substance DEET in this synthetic repellent product is corrosive so it can harm the injured skin. The use of natural repellents from plant materials can reduce the harmful effects of using DEET. This has prompted many researchers to conduct research on natural repellents derived from plants. Safe repellents are those that come from natural compounds (Murrukmihadi and Dias, 2015). Apart from being safe, repellent from plant materials has a distinctive aroma that is liked by humans but is not liked by mosquitoes (Aini et al., 2016). Permot plant (Passiflora foetida Linn.) is a medicinal plant that grows wild and is thought to have potential as a natural repellent (Susilowati, 2016). The leaf part of the permot plant is the part that contains the active compound as a repellent. Permot leaf ethanol extract contains alkaloids, terpenoids and phenols. Terpenoids contain isophytol and phytol which have properties as insecticides against Ae. aegypti (Hastutiek and Sunarso, 2013). Concentrations of 2.5%, 5%, 7.5% were determined based on Faig's (2015) previous research, the amount of which had been adjusted. Based on the description above, it is necessary to conduct research to determine the effectiveness of the ethanol extract of permot leaves as a mosquito repellent for Ae. aegypti adult stage.

Materials and Methods

The sample used in this study waa Ae. aegypti as many as 650 adult females from the Institute of Tropical Disease (ITD) Campus C UNAIR, Surabaya. Permot leaves were taken intentionally from the Keputih area in Surabaya. The leaves are sorted and then cleaned of adhering dirt by washing with clean water, drained and spread on newspapers so that the water evaporates, weighed as much as 7.5 kg dry weight, aerated for \pm one week by placing in a place protected from sunlight then mashed to form a smooth simplicia prepared for extraction. Extraction was carried out by maceration method using ethanol 96% PA. Permot leaf simplicia used as much as 1000 grams was put in a maceration device and 96% PA ethanol solvent was added until everything was submerged. Then macerated for 3x24 hours at room temperature then filtered and the filtrate was accommodated. The filtrate was collected and evaporated at a temperature of 40°C using a rotary evaporator with decreasing pressure until a thick extract was obtained and then weighed. Evaporation was carried out to obtain a thick extract using an oven at 40°C to obtain a thick ethanolic extract of Permot leaves (Padmasari et al., 2013).

The basic cream preparation was carried out to obtain a suitable base cream formulation to be used as a repellent. The cream was made by mixing compounds (A) consisting of stearic acid, lanolin, and aquadest with compounds (B) such as glycerol, sodium hydroxide, triethanolamine, and aquadest in a ratio of 1:1 in a beaker over a bath at 80°C and stirred with a stirrer so that it is evenly distributed. The best formula is then used as the basis for making repellant creams. The percentage of materials used can be seen in Table 1.

 Table 1. Base Cream Formulation

Compounds	Α	В
Stearic acid	16	-
Lanolin	0.5	-
Sodium Hidroxide	-	0.6
Trietanolamine	0.5	1.0
Glycerol	-	17
Aquadest	ad 100%	ad 100%
C_{1}		

Source: Rahayu and Naimah, (2010)

After obtaining the best formula for making the base cream, the formula was used for the negative control and treatment. To make repellant cream, the active compound of permot leaf ethanol extract is added to the base cream, with the variables of permot leaf ethanol extract being 2.5%, 5%, and 7.5% (P1, P2, P3) and the mixing method can be seen in Table 2.

Table 2. Repellant Cream Formulation			
Compound	Base Cream	Permot Extract	
1	97.5	2.5	
2	95	5	
3	92.5	7.5	
K-	100	0	

Permot leaf ethanol extract was weighed as needed. Permot leaf ethanol extract used is a concentration of 2.5%, 5%, 7.5%. The determination of the concentration of the cream is based on previous research by Faiq (2015) whose concentration has been adjusted. The cream concentration of 2.5% was made by weighing the ethanol extract of Permot leaves as much as 2.5 grams then mixed with 97.5 grams of o/w cream then ground with a mortar until homogeneous, while the cream concentration of 5% was carried out by weighing the ethanol extract of Permot leaves. as much as 5 grams and then mixed with 95 grams of cream and making cream with a concentration of 7.5% by weighing 7.5 grams of ethanol extract of Permot leaves then mixed with 92.5 grams of cream. After being homogeneous, it was put into a medicine pot and labeled for each treatment and ready to be used.

Three medicine containers are provided, each of which contains cream of ethanol extract of permot leaves with different concentrations. The experiment was carried out in the morning because the peak activity of mosquitoes occurred in the morning from 09.00-11.00 A.M. This research requires a trial volunteer. Volunteers used repellant cream on the right and left arms to then be tested directly on exposure to mosquitoes. Volunteers have certain conditions that must be met, among others; 17-44 years old, healthy, not affected by malaria, filariasis, DHF, allergies, and have no injuries to the arms (Millati *et al.*, 2018).

The first test procedure is the finger skin surface area to the arm skin surface (test object) smeared with 3 grams of repellent. The repellants were allowed to dry at room temperature. Both arms were put in a cage containing 25 female mosquitoes and then tested through the hole in the repellent test cage and counted with a stopwatch for 10 minutes referring to the research method of Fradin and Day (2002) which modified the time of observation of their perch. Then every 10 minutes thereafter until 5 replications observed the number of mosquitoes perched. The number of mosquitoes that landed on the arm was counted as soon as the hand entered the cage. The same thing was done for all treatments with 5 replications. Arms were put back in cages after 5 minutes pause after treatment. Each treatment was added with repellent, using a new cage, and different mosquitoes.

Results and Discussion

The results of observations made on the repellency of *Ae. aegypti* adults based on various treatments can be seen in Table 3

Research on the effectiveness of permot leaf ethanol extract (*P. foetida* Linn.) against adult

stage of *Aedes aegypti* mosquitoes showed that permot leaf ethanol extract had potential as a repellent. In this study, several different treatments were used, namely the provision of negative control, positive control and ethanol extract of permot leaves with different concentrations. The test was carried out by smearing the finger to the wrist using cream without ethanol extract of permot leaf for negative control, positive control using mosquito repellent lotion (Autan)® containing N, N-diethyl-m-toluamide (DEET) 13% and ethanol extract of Permot leaf. with different concentrations. This observation was carried out for 10 minutes. Before the test, the air temperature in the room when carrying out the test ranged from 26.3°C-28.8°C which is a proper temperature for the life, development and activity of mosquitoes in biting or searching for food (Rizki *et al.*, 2015).

Table 3. Average and Standard Deviation of theNumber of Mosquitoes Perched on VariousTreatments

Treatment Mean±SD	
K-	$24.00^{e} \pm 0.70$
K+	$1.00^{a} \pm 0.00$
P1	17.00 ^d ± 0.70
P2	$11.00^{\circ} \pm 0.70$
P3	$3.00^{b} \pm 0.70$

Note: Different superscripts show significant differences (P<0.05).

Moreover, at a concentration of 7.5%, an effective protection potency of 88% was obtained. Repellent testing using ethanol extract of permot leaves has a high average protection potency at the beginning of the test. The results of this study indicate that the effective concentration of ethanolic extract of permot leaves is at a concentration of 7.5% because the mosquito repel potency at this concentration is 84% at the beginning of use, namely at 0-10 minutes. The effectiveness and duration of repellency of the repellent depends on the active compounds of the repellent formulation, application method, temperature, humidity, wind, and the attraction of mosquitoes to individuals (Suwasono and Blondine, 2009). This effectiveness can also be reduced due to evaporation, abrasion, absorption by skin and sweat, volunteer characteristics such as age, sex and biochemical attractiveness of mosquitoes in repellent test volunteers (Hidayah et al., 2018).

The phytochemical results that have been carried out by Mena et al., (2016) show that the ethanolic extract of Permot leaves contains alkaloids, saponins, and flavonoids. This compound can attack nerve organs which can cause nerve weakness, such as breathing and death. Flavonoids can work as respiratory inhibitors. Inhibitors are substances that inhibit or reduce the rate of chemical reactions, and flavonoids can also interfere with the energy mechanism in the mitochondria by inhibiting the electron transport system (Basundari et al., 2018). Alkaloids have the ability to be good contact or stomach poisons because of their ability to penetrate insect cuticles (Hastutiek et al., 2017). Alkaloids are effective against a wide range of insects, particularly soft-bodied insects. In the nervous system this active compound acts on the ganglia of the central nervous system. Alkaloid compounds as contact and stomach poisons can kill insects slowly followed by stopped eating activities (Hastutiek and Sunarso, 2015).

Mosquitoes get fluids from food to maintain water balance. Mosquitoes also like a warm atmosphere because body temperature will affect the number of mosquitoes that perch. Mosquito behavior when searching for food is influenced by stimuli released by food sources (humans and animals), namely visual stimuli, warmth, humidity, smell and the presence or absence of carbon dioxide. Mosquitoes like a warm atmosphere and hands that emit a little moisture (Sanjaya et al., 2014). The repellent enters the respiratory system in the form of gas or fine particles which are carried to the mosquito's body tissues. The smell of permot leaf ethanol extract contains active compounds that mix with gases in the air which causes mosquitoes to not be attracted to this smell and will avoid it. The smell of the repellent is detected by chemical receptors found on the mosquito antenna, then the mosquito will convert it into impulses and forwarded by nerve axons to the central nervous system, which causes integration of the motor nerves to the brain so that mosquitoes avoid (Werdiningsih and Amalia, 2018). In addition, physiology, hunger, or satiety, sex and sexual development and eating habits at different developmental stages in each species affect the olfactory organs (Sanjaya et al., 2014).

The higher the concentration, the active compounds content of permot leaf ethanol extract, the higher the effectiveness of the repellent against *Ae. aegypti*. The active

compounds of permot leaf ethanol extract as a repellent emits a pungent odor that will eventually keep mosquitoes away (Pettagading, 2018). This study proved that the ethanolic extract of permot leaf (*P. foetida* Linn.) has active compounds that can be used as a mosquito repellent predominately for *Ae. aegypti*.

Conclusion

Based on the results of the research that has been done, it can be concluded that the ethanol extract of permot leaf (P. foetida Linn.) has repellency against Ae. aegypti adult stage and ethanol extract with a concentration of 7.5% had a protective potency of 88% so that it could be said to be effective as a repellent against Ae. aegypti adult stage for 10 minutes according to the standards of the Ministry of Health of the Republic of Indonesia. Moreover, further research is needed such as using permot leaf extract (P. foetida Linn.) with non-polar solvents such as n-hexane or semipolar solvents such as ethyl acetate. Furthermore, conducting further research on permot leaf extract (P. foetida Linn.) with a higher concentration or more than 15% to reveal the effectiveness of the repellent against Ae. aegypti. In addition, it is also necessary to conduct further research on the effectiveness of the repellent ethanol extract of permot leaf (P. foetida Linn.) using the WHO standard with a protective potency above 90%.

References

- Aini, R., R. Widyastuti dan N.A. Nadhifa. 2016. Uji Efektifitas Formula Spray dari Minyak Atsiri Herba Kemangi (*Ocium sanctum*. L) sebagai *Repellent* Nyamuk *Aedes aegypti*. J. Ilmiah Manuntung. 2(2):189-197.
- Basundari, S. A., Tarwotjo, U., dan E. Kusdiyantini. 2018. Pengaruh Kandungan Ekstrak Daun Zodia *(Evodia suaveolens)* terhadap Mortalitas Larva Nyamuk *Aedes aeqypti*. Bioma. 20(1): 51-58.
- Departemen Kesehatan RI. 2000. Modul Epidemiologi Malaria. Ditjen P2M dan PLP. Jakarta. Hal 37.
- Djatmiko M, Y. Anas, dan S.M. Handayani. 2011. Uji aktivitas *repellent* fraksi n-heksan ekstrak etanolik daun mimba (*Azadirachta indicaa juss*) terhadap nyamuk *Aedes aegypti*. E-Publikasi Fakultas Farmasi. 3(1):24-30.

- Faiq, M. 2015. Efektifitas repelen ekstrak daun Permot (*Passiflora foetida* L.) terhadap nyamuk *culex fatigans* dewasa [Skripsi]. Fakultas Kedokteran Hewan. Universitas Airlangga.
- Fradin, M.S., and D.F. Day. 2002, Comparative Efficacy of Insect *Repellents* against Mosquito Bites, The New England Journal of Medicine, Chapel Hill Dermatology, 3(4): 13-18.
- Hastutiek P., dan A. Sunarso. 2013. Identifikasi Komponen Kimia Ekstrak daun Permot (*Passiflora foetida* Linn.) dengan TLC dan GC-MS sebagai Kandidat Bioinsektisida terhadap Nyamuk. Seminar Nasional Biologi/IPA dan Pembelajarannya di Universitas Negeri Malang 1-2 Nop 2014.
- Hastutiek, P., dan A. Sunarso. 2015. Gambaran Histopatologi Saluran Pencernaan Larva Instrar IV Nyamuk *Aedes aegypti* setelah Perendaman dengan Senyawa Aktif Ekstrak Daun Permot (*Passiflora foetida* Linn.) dan Potensinya sebagai Bioinsektisida. Veterinaria Medika. 8(2): 137-144.
- Hastutiek, P., Sunarso, A., and R.H. Prasetyo. 2017. Permot (*Passiflora Foetida* Linn.) Leaf Extracts as Bioinsecticide Against *Aedes aegypti* Larvae. The Southest Asian Journal of Tropical Medicine and Public Health. 48(6): 1169-1174.
- Herawati, R. 2010. Ekstrak Daun Sirih (Piper betle L.) sebagai Insektisida Nabati untuk Membasmi Larva Nyamuk Aedes aegypti L. [Skripsi]. Fakultas Teknobiologi. Universitas Atma Jaya Yogyakarta. Yogyakarta.
- Hidayah, N., Mustafa, H., Murni, dan I. Tolistiawaty. 2018. Efektivitas Repelen Losion Minyak Atsiri Kulit Jeruk Bali (*Citrus maxima* (Burm.) Merr.) terhadap *Aedes aegypti*. Balaba. 14(2):159-168.
- Ishartadiati, K. 2010. *Aedes aegypti* sebagai Vektor Demam Berdarah Dengue. Universitas Wijaya Kusuma Surabaya. Surabaya. Hal 31.
- Mena, P., Cirlini, M., Tassotti, M., A. Herrlinger, K., Dall' Asta, C., and D. Del Rio. 2016. Phytochemical Profiling of Flavonoids, Phenolic Acids, Terpenoids, and Volatile Fraction of a Rosemary (*Rosmarinus* officinalis L.) Extract. PubMed. 21(11): 1-15.

- Millati, F. F., Ferry, dan F. Sofian. 2018. Review Artikel: Kandungan Senyawa Minyak Atsiri pada Tanaman Pengusir Nyamuk. Jurnal Farmaka. 16(2):572-580.
- Murrukmihadi, M and D.A. Fajarini. 2015. Repelen Activity Test of Essential Oil of Basil Leaves (Ocimum basilicum L.) (f. Citratum Back) against Aedes aegypti Lotion and Physical Characteristics Test of The Lotion. Traditional Medicine J. 20(2): 91-97.
- Padmasari, P.D., K.W. Astuti, dan N.K. Warditiani. 2013. Skrining Fitokimia Ekstrak Etanol 70% Rimpang Bangle *(Zingiber purpureum Roxb.)*. Fakultas Matematika dan Ilmu Pengetahuan Alam. Universitas Udayana. Denpasar. Hal 20.
- Rahayu, S. P., dan S. Naimah. 2010. Pembuatan formulasi krim anti nyamuk dari fraksi minyak sereh. Jurnal Kimia dan Kemasan, 32(2): 53-61.
- Rizki, D. A., Budiyono, dan R. Hestiningsih. 2015. Daya Tolak *Repellent* Bentuk Lotion dengan Ekstrak Daun Alpukat (*Persea americana* Mill) Terhadap Nyamuk *Aedes aegypti* Linn. Jurnal Kesehatan Masyarakat. 3(3): 702-711.
- Romandoni, F. N. 2019. Hubungan Upaya Pencegahan terhadap Kejadian Penyakit DBD pada Masyarakat di Desa Gemaharjo Wilayah Kerja Puskesmas Gemaharjo Kabupaten Pacitan. [Skripsi] Stikes Bhakti Husada Mulia Madiun.
- Sanjaya Y., Adisenjaya, Yusuf H., dan L. Wijayanti. 2014. Efektivitas Daya Tolak Ekstrak *Geranium radula* C. Terhadap Nyamuk *Aedes aegypti* (Linn.). Bionatura. 16(2): 62-67.
- Susilowati, R.P. 2016. Uji Toksisitas Obat Nyamuk Bakar Herbal Berbahan Ekstrak Daun Permot (*Passiflora foetida*): Kajian Histopatologis Hati dan Ginjal Mencit. Prosiding SNPBS (Seminar Nasional Pendidikan Biologi dan Saintek) Ke-1 hal 492-497.
- Suwasono, H., dan Ch. P, B. 2009. Uji Efikasi Repelen "X" terhadap Nyamuk *Aedes aegypti, Culex quinquefasciatus* dan *Anopheles aconitus* di Laboratorium. Jurnal Vektor 1(2): 101-108.
- Taslimah. 2014. Uji Efikasi Ekstrak Biji Srikaya (Annona squamosal L.) sebagai

Bioinsektisida dalam Upaya *Integrated Vector Management* terhadap *Aedes aegypti* [Skripsi]. Fakultas Kedokteran dan Ilmu Kesehatan. UIN Syarief Hidayatullah Jakarta.

Werdiningsih, I., dan R. Amelia. 2018. Lotion Ekstrak Daun Ekstrak Daun Zodia *(Evodia sauveolens)* sebagai *Repellent* Nyamuk *Aedes* sp. Jurnal Sains dan Seni. 1(1): 1-4.



Kampus C Mulyorejo Surabaya 60115 Telp. (031) 5992785, 5993016 Fax (031) 5993015 Laman: http://www.fkh.unair.ac.id, e-mail: info@fkh.unair.ac.id

SURAT KETERANGAN Nomor : 2934/UN3.1.6/KP/2022

Yang bertanda tangan di bawah ini :

Nama	: Prof. Dr. Mirni Lamid, drh., MP
NIP	: 196201161992032001
Pangkat / Golongan	: Pembina Utama Madya / (Gol. IV/d)
Jabatan	: Dekan

Dengan ini menerangkan bahwa :

Nama	: Dr. Endang Suprihati, drh., MS
NIP	: 195810211983112001
Pangkat / Golongan	: Pembina TK I / (Gol. IV/b)
Jabatan	: Lektor Kepala

Telah melaksanakan penelitian dengan judul sebagai berikut :

No.	Judul Karya Ilmiah	Tahun pelaksanaan Penelitian
1.	Morphological Detection of The Intestinal Parasite Blastocystis sp.	2019
	(Mammalia: Petauridae) In Surabaya, Indonesia.	2018
2.	Identification of Active Compounds of Ethanol Extract of Citrus amblycarpa leaves by Analysis of Thin-layer Chromatography and Gas Chromatography-Mass Spectrometry as Bioinsecticide Candidates for Mosquitoes	2020
3.	Histopathological studies on <i>Leucocytozoon Caulleryi</i> infection on broiler in endemic area of Indonesia	2020
4.	Potential Extract Ethanol Citrus Amblycarpa as a Bioinsecticide Against Aedes Aegypti Larvae	2021
5.	Protein Profile of Sporozoite of Leucocytozoon sp. from Culicoides sp.	2010
6.	Deteksi Cryptosporidium canis pada Anjing di Kota Surabaya	2020
7.	Eksplorasi Protein Antigenik <i>Leucocytozoon caulleryi</i> sebagai Kit Diagnostik Leucocytozoonosis pada Ayam Broiler	2013

















UNIVERSITAS AIRLANGGA

FAKULTAS KEDOKTERAN HEWAN

Kampus C Mulyorejo Surabaya 60115 Telp. (031) 5992785, 5993016 Fax (031) 5993015 Laman: http://www.fkh.unair.ac.id, e-mail: info@fkh.unair.ac.id

8.	Uji reaktivitas protein 30 kDA bakteri <i>Aeromonas hydrophila</i> yang diisolasi dari ikan air tawar dengan teknik indirect ELISA.	2016	
9.	Penambahan Sari Air Laut (Nigarin) Dalam Pengencer Skim Kuning		
	Telur Terhadap Viabilitas Dan Motilitas Spermatozoa Sapi Limousin	2018	
	Post Thawing		
10.	The Effectiveness of Ethanol Extract of Red Betel Leaf (Piper	2020	
	crocatum) Againts Mortality of Boophilus microplus Larvae In Vitro	2020	
11.	Prevalence of Ectoparasites in Bean Goats on the Sub-District of	2020	
	Prambon, District of Nganjuk	2020	
12.	Repellent Effectiveness of Permot Leaf Ethanol Extract (Passiflora)	2021	
	Foetida Linn.) against Aedes Aegypti Adult Mosquitoes	2021	
13.	Detection of Goat Digestive Tract Protozoa Through Feces	2021	
	Examination in Kwanyar Sub-District, Bangkalan District	2021	
14.	Identification and Prevalence of Digestive Tract Endoparasites of	2021	
	Goats in Ujungpangkah, Gresik District	2021	
15.	Morphology of surface ultrastructure of Duthiersia expansa(Cestoda		
	Diphyllobothriidea) from water lizards (Varamus salvator) from	2014	
	Sidoarjo, Indonesia		
16.	Antigenic Protein of Leucocytozoon caulleryi schizont Inducing	2017	
	Cellular Immune Resonse: TLR-2 and CD4 as Marker	2017	

Adapun penelitian tersebut <u>tidak perlu</u> dilakukan *Uji Etical Clearence* karena tidak menggunakan hewan coba.

Demikian surat kerangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional <u>Guru Besar</u>

Surabaya, 8 Agustus 2022

Dekan, HRSITAS AIRL THE Dr. Mirni Lamid, drh., MP of 196201161992032001 NI















