

ISSN 1365-3113
CODEN SYRREV

Volume 17, Number 1
January 2005

Systematic Reviews

in Epidemiology

Journal of the International Society for Systematic Reviews

www.systematicreviews.com




Systematic Reviews in Epidemiology is a peer-reviewed journal that provides a forum for the publication of original research, reviews, and commentaries on the methodology and practice of systematic reviews. The journal is published quarterly by Wiley-Blackwell.


WILEY-BLACKWELL



Systematic Reviews in Pharmacy





- [Online First](#)
- [Archive](#)
- [Aims and Scope](#)
- [Abstracting & Indexing](#)
- [Most Accessed Articles](#)
- [Most Downloaded Articles](#)



[Google Scholar citation report](#)

Citations : 9175

[Systematic Reviews in Pharmacy received 9175 citations as per google scholar report](#)



[Online First](#)


[Archive](#)

[Aims and Scope](#)

[Abstracting & Indexing](#)

[Most Accessed Articles](#)

[Most Downloaded Articles](#)



[Google Scholar citation report](#)

Citations : 9175

Systematic Reviews in Pharmacy received 9175 citations as per google scholar report

Review Article**Collaborative Governance Framework in Health Care: A Qualitative Exploration of Hospital Pharmacy Management Reform at Hospital Setting in Indonesia**

N Hardi Warsana, Retna Harani, Malisa Dwi Putra.

SRP 2020; 11(4): 397-408

>> Abstract >> PDF [DOI: 10.31838/srp.2020.4.40](https://doi.org/10.31838/srp.2020.4.40)Review Article**Study of Carcinoma Embryonic Antigen (CEA) and its Relationship to TSH Polymorphisms in Patients with Thyroid Gland Dysfunction.**

S Syasari Hassan Kadhim.

SRP 2020; 11(4): 399-402

>> Abstract >> PDF [DOI: 10.31838/srp.2020.4.46](https://doi.org/10.31838/srp.2020.4.46)Review Article**Pharmacy Service Quality for Pharmacy Customer Satisfaction, Trust, and Loyalty**

N Trialy Chandra, Lajla Hitti, Geany Naitra

SRP 2020; 11(4): 403-407

>> Abstract >> PDF [DOI: 10.31838/srp.2020.4.50](https://doi.org/10.31838/srp.2020.4.50)Review Article**Globalization and Problems of Modern Regional Economy**

S Wawan Ropizalevina Warkanyan, Lidiana Kristina Wikauleana, Kamohatowa Silaterisa Waresna, Chetsona Vera Alexandrussa, Keshusidico Henna Anatalyeana

SRP 2020; 11(4): 408-412

>> Abstract >> PDF [DOI: 10.31838/srp.2020.4.51](https://doi.org/10.31838/srp.2020.4.51)Review Article**Interleukin-6 Associated with Insulin Resistance in Non-Diabetic Predialysis Chronic Kidney Disease Patients**

N Hermy Nevanadana Bramantya, Candra Arumadhi Mubandri, Suciagijjo Adi Soelastjo

SRP 2020; 11(4): 414-418

>> Abstract >> PDF [DOI: 10.31838/srp.2020.4.52](https://doi.org/10.31838/srp.2020.4.52)Review Article**Study the Laser Bleaching for Several Human Teeth and Their Structural Properties as a Model of Therapy**

S Elhazgham M. Abd. Ali AL- Haddad, Mithaq M. Mehdy AL- Soltani

SRP 2020; 11(4): 419-422

>> Abstract >> PDF [DOI: 10.31838/srp.2020.4.57](https://doi.org/10.31838/srp.2020.4.57)Review Article**Comparison of the 25(OH)D Levels Between Sarcopenia and Frailty in Elder Women: A Cross-Sectional Observation Analytic Study in Elderly Community in Surabaya**

N Wiranti Ayu Adyanita, Nurica Mubandari, Sony Wibisono

SRP 2020; 11(4): 423-428

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.53

Review Article

Visualization of Information in the Educational Process: Current Trends

Dr. Kabanov Oleg Vladimirovich, Starostina Iulia Evgenievna, Zhukan Vladimir Mikhailovich, Kurbanov Denis Anatolevich, Kazantseva Irina Anatolevna, Studinova Anna Vasilevna, Lova Raza Suleymanovna, Morozov Sergey Aleksandrovich

SRP. 2020; 11(4): 1 – 5

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.51

Review Article

Comparison between Levofloxacin Based Therapy and Clarithromycin Based Therapy through 14 Days Period for H-Pylori Eradication

Dr. Nasser Hassan Abbas Alsaadi, Marwan Hassan Abbas Alsaadi, Zoya Tahir Ali

SRP. 2020; 11(4): 583 – 588

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.50

Review Article

Hemoglobin Levels in Patients with Human Immunodeficiency Virus Naïve Therapy Containing Zidovudine in the First Three Months

Dr. Pradyumn Singh, Estu Pratiwi, Usman Hani, Muhammad Alimuzza Aziz

SRP. 2020; 11(4): 365 – 370

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.54

Review Article

A Case of Death as the Result of Firing from Self-Made Firearms

Dr. Yu.V. Zhurava, Dr. V. Bogdanov, V.A. Potvinov

SRP. 2020; 11(4): 6 – 8

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.52

Review Article

Differences in VO2 Max Based on Age, Gender, Hemoglobin Levels, and Leukocyte Counts in Hajj Prospective Pilgrims in Hulu Sungai Tengah Regency, South Kalimantan

Dr. Ulfahri, Harun Achmad, Anwar Asyraf, Aminuddin Praditama Putra, Bayu Indah Sukmana, Dwi Salsawa Adipatry, Idris Kasih

SRP. 2020; 11(4): 9 – 14

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.03

Review Article

Sequence-based Detection and Identification Biodiversity of Uncultivated Fungi in Soils.

Dr. Wijan Alimuzza Ali

SRP. 2020; 11(4): 529 – 603

>> Abstract >> PDF DOI: 10.31838/srp.2020.4.02

Review Article

Analysis of Dental Caries & Gingivitis with the Occurrence of Stunting in Children in Makassar City (Tamalanrea Subdistrict)

H. Harun Achmas, Hendriantuti Mandayanti, Marhamah E. Singsih, Shedy Hanae, Siti Ramadhany, Febeila Setiawati, Runita Felty Ramadhany

DOI: 10.55530/srp.2020.11.04

[Submit Article](#)

» [HUSUDU](#) » [FDR](#) » [www.sysrevpharm.org](#)

[Login](#)

[Register](#)

Most Viewed Articles

Dental Development between Assisted Reproductive Therapy (ART) and Natural Conceived Children: A Comparative Pilot Study Norzati Mohd Norzal, Maimah Hasmah Mohd Fadzil, Nurhayati Bohari, Ahmad Faisal Samad, Roszaman Ramli SRP. 2020; 11(1): 01-06 » doi: 10.55530/srp.2020.11.01

Psychometric properties of the World Health Organization Quality of Life-instrument, short form: Validity in the Vietnamese healthcare context Trung Quang GGT, Bao Tran Duy Tran, Ngan Duy Nguyen, Tran Thiuyen Nguyen, Thy Pham Chung Tran SRP. 2020; 11(1): 04-22 » doi: 10.55530/srp.2020.11.03

A Review of Pharmacoeconomics: The Key to "Healthcare for All" Hasmi AA, Pali SS, Shaik Imran, Nazirah K SRP. 2019; 10(1): s40-s42 » doi: 10.55530/srp.2019.10.21

Desamun (Spiced Watermelon) as an Treatment of Cancer: Antic-Squamous, Oxydewitaya, Hewan (Supernatant), Tumbuhan (Plantawa), Dan (Rumay)na, Dan (Ernakova) SRP. 2019; 10(1): 102-107 » doi: 10.55530/srp.2019.11.03

Most Downloaded

Dental Development between Assisted Reproductive Therapy (ART) and Natural Conceived Children: A Comparative Pilot Study Norzati Mohd Norzal, Maimah Hasmah Mohd Fadzil, Nurhayati Bohari, Ahmad Faisal Samad, Roszaman Ramli SRP. 2020; 11(1): 01-06 » doi: 10.55530/srp.2020.11.01

Misokarsapada (L.) Rujer (Tub) Pesi: A Phytochemical and Pharmacological Review Kati Pravin P, Shweta Shrikar C SRP. 2019; 10(1): 11-14 » doi: 10.55530/srp.2019.10.2

Pharmacognostic and Phytochemicals Overview on Bonina on the Parlay Warbika, Comarby, Bukal Gunung Tawar SRP. 2019; 10(1): 20-25 » doi: 10.55530/srp.2019.10.4

A Review of Pharmacoeconomics: The Key to "Healthcare for All" Hasmi AA, Pali SS, Shaik Imran, Nazirah K SRP. 2019; 10(1): s40-s42 » doi: 10.55530/srp.2019.10.21

A Prospective Review on Phyto-Pharmacological Aspects of Anthyraghis gentiana (Gawitaj) Mikadawon, Asumagan Wijay Asani, Palanisamy Sampathkumar, Pulkarnan Vinayaga Murugli, Saswanja Pruthi SRP. 2019; 10(1): 15-19 » doi: 10.55530/srp.2019.11.3

Copyright © 2022 Systematic Reviews in Pharmacy All Rights Reserved. Subject to change without notice from or liability to Systematic Reviews in Pharmacy.



Copyright © 2022 Systematic Reviews in Pharmacy All Rights Reserved. Subject to change without notice from or liability to Systematic Reviews in Pharmacy. For best results, please use Internet Explorer or Google Chrome

POLICIES & JOURNAL LINKS

- [Advertising Policy](#)
- [Author's Rights and Obligations](#)
- [Conflict of Interest Policy](#)
- [Digital Archiving & Preservation Policies](#)
- [Editorial Policies](#)
- [Peer Review Policy](#)
- [Editorial & Peer Review Process](#)
- [License Information](#)
- [Plagiarism Policy](#)
- [Privacy Policy](#)
- [Protection of Research Participants \(Statement On Human And Animal Rights\)](#)
- [Publishing Ethics](#)
- [Corrections, Retractions & Expressions of Concern](#)
- [Self-Archiving Policies](#)
- [Statement of Informed Consent](#)
- [Terms of Use](#)




Systematic Reviews Pharmacy

FOLLOW

GET MY OWN PROFILE

Editor
 Verified email at sysrevpharma.org - [Homepage](#)
 Pharmacy Pharmaceutical sciences pharmacology Biochemistry Biomedicine

Cited by	VIEW ALL	
	All	Since 2017
Citations	9175	8587
h-index	36	35
i10-index	225	219



TITLE	CITED BY	YEAR
Nanoemulsion: A pharmaceutical review. P Shah, D Bhalodia, P Shelat Systematic reviews in pharmacy 11 (1):	399	2010
Emulsion micro emulsion and nano emulsion: a review SN Kale, SL Deore Systematic Reviews in Pharmacy 8 (1): 39	279	2017
COVID-19 pandemic and home online learning system: Does it affect the quality of pharmacy school learning M Suryaman, Y Cahyono, D Mulliansyah, O Bustani, P Suryani, M Fahien Systematic Reviews in Pharmacy 11 (8): 524-530	158	2020
Optimization of MSMEs empowerment in facing competition in the global market during the COVID-19 pandemic time A Affandi, AS Sarwani, H Ertangga, AO Siagian, A Purwanto, AA Effendy Systematic Reviews in Pharmacy 11 (11): 1506-1515	107	2020
Biopharmaceutics classification system H Chavda, C Patel, I Anand Systematic reviews in pharmacy 1 (1): 62	106	2010
University students online learning system during Covid-19 pandemic: Advantages, constraints and solutions NA Fatonia, E Nurkhyati, E Nurdiaatid, GP Fidziah, S Adhag Systematic Reviews in Pharmacy 11 (7): 570-576	93	2020
Coumarins: the antimicrobial agents YK Al-Majedy, AAH Kadhum, AA Al-Ansary, AB Mohamad Systematic Reviews in Pharmacy 8 (1): 62	91	2017
Ophthalmic drug delivery system: challenges and approaches P Patel, D Shastri, P Shelat, A Shukla Systematic Reviews in Pharmacy 1 (2): 113	85	2010

Systematic Reviews in Pharmacy peer review process verified at publons

Indexed In

- > Genamics JournalSeek
- > JournalTOCs
- > China National Knowledge Infrastructure (CNKI)
- > Scimago
- > Ulrich's Periodicals Directory
- > EBSCO A-Z
- > Pollution Abstracts
- > OCLC- WorldCat
- > Proquest Summons
- > ROAD
- > CABI full text
- > SciLit - Scientific Literature
- > Publons
- > Google Scholar
- > J-Gate
- > Chemical Abstract
- > SHERPA ROMEO

Editorial Board

Editor-in-Chief

Tweets by @in_systematic
Dr. Ayad F. Alkaim

University of Babylon, College of Science for Women, Babylon, Scopus Author ID: 55255310600, Iraq

Systematic Reviews in Pharmacy

@in_systematic

Editorial Board

According to the new clinical data, 50-µg booster dose of mRNA-1273.214 had a potent neutralizing antibody response against the #OmicronVariant in all individuals, regardless of prior infection. [#Medscape](#), PhD

Professor, Department of Experimental and Translation Oncology

Croatian Institute for experimental and translation oncology, Zagreb, Croatia, Croatia

15h

Dr. Lucius, MBBS

Systematic Reviews in Pharmacy

General Practice
@in_systematic

Ludwig Maximilians University Munich, Germany

A phase 3 clinical trial of #brexpiprazole showed positive results in the treatment of agitation in patients with #Alzheimer dementia

Dr. Aygul Z. Ibatova

Department of Natural Sciences

Yuzhen Industrial University, Scopus Author ID: 57191110632 <http://orcid.org/0000-0003-0565-8533>, Russia

[View on Twitter](#)

Dr Ahmad Faisal Ismail

Kulliyah of Dentistry

International Islamic University Malaysia, Kuantan Campus, 25200 Kuantan, Pahang, Scopus Author ID: 35388596700, Malaysia

Dr. Huijiang ZHAO, Ph.D

Guizhou Minzu University, Huaxi District, Guiyang, China

Dr. Mohd Armi Abu Samah

International Islamic University Malaysia, (IIUM) 25200 Kuantan Pahang

Dr. Baded ramji

Sri Lanka

Dr. Chris randea

South Africa

Dr. Yingwen ZHAO

Researcher of Guizhou Rural Economic and Social Development Research Institute, China

Dr. Li Zihan, Ph.D

University of Glasgow, UK

Gabriela Cioca

Faculty of Medicine, Pharmacology Department

Lucian Blaga University of Sibiu, Romania, Lucian Blaga street, no 2A, Sibiu, Romania

Dariusz Nowak

Municipal Hospital, Mickiewicza street no 12, 42-200 Czestochowa, Poland

Aleksandra Zyska

Faculty of Medicine, Department of Physiology

Opole University, Oleska street no 48, 45-052 Opole, Poland

Katarzyna Sznajder

Faculty of Medicine, Clinical Department of Diagnostic Imaging

Opole University, Oleska street no 48, 45-052 Opole, Poland

Jacek Jankowski

Faculty of Medicine, Department of Family Medicine and Public Health

Opole University, Oleska street no 48, 45-052 Opole, Poland

Luciano Benedini

Universidad Nacional del Sur (National University of South-UNS), Bahía Blanca 8000, Argentina

Paula Messina

Departamento de Biología

Universidad Nacional del Sur (National University of South-UNS), Bioquímica y Farmacia, Bahía Blanca 8000, Argentina

Michael Walsh

Washington State University, College of Pharmacy and Pharmaceutical Sciences (CPPS), USA

Prof. Dr. Kittisak Jermstittiparsert

Henan University, China

Amel Dawod Kamel Gudia, PhD

Faculty of nursing
Cairo University, Egypt

Arif Nur Muhammad Ansori
Airlangga University, Scopus Author ID: 57195995342, <https://orcid.org/0000-0002-1279-3904>, Indonesia

Mohammed Nader Shalaby
Suez Canal University, Associate Professor of Biological Sciences and Sports Health, Egypt

Dr. Faten Abo-Aziza Mohamed, PhD
Associate Professor, Clinical Pathology and Stem Cell Research
National Research Centre, Manager of Veterinary Division Central Lab (605), 33 El-Beheos St, Dokki, Cairo, Egypt

Professor Asim Ahmed Elnour Ahmed
College of Pharmacy
Al-Ain University of Science and Technology, UAE

S. Parasuraman, M.Pharm., Ph.D
AIMST University, Malaysia

Ebenezer Wiafe, PhD
Pharmacy
University of Kwazulu-Natal, South Africa

[Submit Article](#)

[Login](#)

[Register](#)

Most Viewed Articles

Dental Development between Assisted Reproductive Therapy (Art) and Natural Conceived Children: A Comparative Pilot Study Norzaiti Mohd Kenali, Naimah Hasanah Mohd Fathil, Norbasyirah Bohari, Ahmad Faisal Ismail, Roszaman Ramli SRP. 2020; 11(1): 01-06 » doi: 10.5530/srp.2020.1.01

Psychometric properties of the World Health Organization Quality of life-instrument, short form: Validity in the Vietnamese healthcare context Trung Quang Vo*, Bao Tran Thuy Tran, Ngan Thuy Nguyen, Tram ThiHuyen Nguyen, Thuy Phan Chung Tran SRP. 2020; 11(1): 14-22 » doi: 10.5530/srp.2019.1.3

A Review of Pharmacoeconomics: the key to "Healthcare for All" Hasamnis AA, Patil SS, Shaik Imam, Narendiran K SRP. 2019; 10(1): s40-s42 » doi: 10.5530/srp.2019.1s.21

Deuterium Depleted Water as an Adjuvant in Treatment of Cancer Anton Syroeshkin, Olga Levitskaya, Elena Uspenskaya, Tatiana Pleteneva, Daria Romaykina, Daria Ermakova SRP. 2019; 10(1): 112-117 » doi: 10.5530/srp.2019.1.19

Most Downloaded

Dental Development between Assisted Reproductive Therapy (Art) and Natural Conceived Children: A Comparative Pilot Study Norzaiti Mohd Kenali, Naimah Hasanah Mohd Fathil, Norbasyirah Bohari, Ahmad Faisal Ismail, Roszaman Ramli SRP. 2020; 11(1): 01-06 » doi: 10.5530/srp.2020.1.01

Manilkara zapota (L.)-Roeys Fruit Peel: A Phytochemical and Pharmacological Review Karle Pravin P, Dhawale Shashikant C SRP. 2019; 10(1): 11-14 » doi: 10.5530/srp.2019.1.2

Pharmacognostic and Phytopharmacological Overview on Bombax ceiba Pankaj Haribhau Chaudhary, Mukund Ganeshrao Tawar SRP. 2019; 10(1): 20-25 » doi: 10.5530/srp.2019.1.4

A Review of Pharmacoeconomics: the key to "Healthcare for All" Hasamnis AA, Patil SS, Shaik Imam, Narendiran K SRP. 2019; 10(1): s40-s42 » doi: 10.5530/srp.2019.1s.21

A Prospective Review on Phyto-Pharmacological Aspects of Andrographis paniculata Govindraj Akilandeswari, Arumugam Vijaya Anand, Palanisamy Sampathkumar, Puthamohan Vinayaga Moorthi, Basavaraju Preethi SRP. 2019; 10(1): 15-19 » doi: 10.5530/srp.2019.1.3

Copyright © 2022, Systematic Reviews in Pharmacy. All Rights Reserved. Subject to change without notice from or liability to Systematic Reviews in Pharmacy. For best results, please use Internet Explorer or Google Chrome.

[Advertising Policy](#)
[Author's Rights and Obligations](#)
[Conflict of Interest Policy](#)
[Digital Archiving & Preservation Policies](#)
[Editorial Policies](#)
[Peer Review Policy](#)
[Editorial & Peer Review Process](#)
[License Information](#)
[Plagiarism Policy](#)
[Privacy Policy](#)
[Protection of Research Participants \(Statement On Human And Animal Rights\)](#)
[Publishing Ethics](#)

Comparison of the 25(OH)D Levels Between Sarcopenia and Frailty in Elder Women: A Cross-Sectional Observation Analytic Study in Elderly Community in Surabaya

Viranti Ayu Adyanita¹, Novira Widajanti^{2*}, Sony Wibisono³

¹Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

²Geriatric and Gerontology Division, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

³Endocrine and Metabolic Division, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya 60131, Indonesia

Article History:

Submitted: 11.01.2020

Revised: 15.03.2020

Accepted: 20.04.2020

ABSTRACT

The prevalence of vitamin D deficiency in the elderly, specifically in post-menopausal women, are very high. Deficiency of 25(OH)D has a direct impact on decreasing muscle strength and mass that can lead into sarcopenia and frailty. This study aimed to compare the difference levels of 25(OH)D between sarcopenia and frailty in elderly community.

Methods: This was a cross-sectional design study. The blood sample were collected to evaluate the levels of 25(OH)D using The Chemiluminescent Immune Assay (CLIA) method.

Results: Most of the subjects were at deficiency of 25(OH)D with mean of 16.1 (range level of 6.2-32.7). Subjects with deficiency level of 25(OH)D were 20 (71.4%) with sarcopenia and 23 (85.1%) with frailty. Subjects with an insufficiency level of 25(OH)D were 6 (21.4%) with sarcopenia and 23 (85.1%) with frailty. Subjects with sufficiency level of 25(OH)D were 2 (7.1%) with sarcopenia and 1

(3.7%) with frailty. The median 25(OH)D levels were 17.1 and 13.2 for sarcopenia and frailty, respectively. There was significant differences in levels of 25(OH)D in both groups ($p = 0.014$).

Conclusions: The levels of 25(OH)D of subjects with sarcopenia was significantly higher than those with frailty.

Keywords: 25(OH)D, vitamin D, sarcopenia, frailty, elder women.

Correspondence:

Novira Widajanti, MD

Geriatric Division, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya 60131, Indonesia
Jalan Mayjand Prof. Dr. Moestopo No. 6-8 Surabaya 60131, Indonesia.

E-mail: novirawidajanti@yahoo.com

DOI: 10.31838/srp.2020.4.53

©Advanced Scientific Research. All rights reserved

INTRODUCTION

Health problems in the elderly come from declining body cells, so the function and

body endurance decreases along with increased risk factors for diseases and infections.^{1,2} One of health problems is vitamin D deficiency. The prevalence of vitamin D deficiency in the elderly, specifically in post-menopausal women, is very high. The incidence rate reaches 90% in Asia and 41.1% in Europe.³ Vitamin D deficiency in the elder women in Jakarta reaches 35%.⁴ The elderly with obesity are at risk for suffering vitamin D deficiency.⁵ Vitamin D deficiency can lead to sarcopenia and frailty. Sarcopenia and frailty can influence the quality of life of the elderly, resulting in an adverse outcome, such as falls, fractures, hospitalization, and high comorbidity, disability, and mortality rates.

Technically, vitamin D is a prohormone that is produced by the photochemical process on the skin with the initial form of 7-dehydrocholesterol. The synthesis of vitamin D is mainly in the skin, and less than 10% is earned from the food source. Vitamin D deficiency can lead into the decrease in muscle mass and muscle strength. Besides, vitamin D deficiency affects to tooth loss that may be considered an early marker of decline and weakness in the elderly.⁶ It mainly occurs in the elderly because there is a low intake of food that contains vitamin D, lowest of sun exposure, intestinal malabsorption, and decreased vitamin D hydroxylase activity in the kidneys. The main synthesis of vitamin D is on the skin. Less than 10% comes from the diet. 7-dehydrocholesterol converts to pre-vitamin D3 on the skin with the aid of ultraviolet-B exposure from sunlight. Pre-vitamin D3 undergoes an isomerase, which depends on the temperature to form vitamin D3.

Enzymatically, vitamin D3 will be converted to 25-hydroxyvitamin D3 (25(OH)D) by the enzyme 25-hydroxylase in the liver and converted to 1,25-dihydroxy vitamin D3 (biologically active form) in the kidney by the enzyme 1-alpha-hydroxylase. The 25(OH)D is in the form of an inactive version of vitamin D and has a longer half time compare with another form of vitamin D.⁷⁻⁹

The person in advanced age is likely to experience a decrease in physical quality, such as a weakness in his legs.¹⁰ The aging process in bones and muscles is characterized by the decrease of muscle mass and strength, and if both of that process occurs in a person, it is called sarcopenia. If sarcopenia develops continuously, it will lead to frailty. Deficiency of 25(OH)D has a direct impact on decreasing muscle strength and mass. The muscle strength is correlated with physical performance.¹¹ Decreased strength and muscle mass can lead to the lowering of mechanical myokines, which can also lead to the lowering of mineral bone density.¹²

Sarcopenia is described as the loss of muscle mass and strength inherent to the aging process.¹³ Frailty is a condition that is prone to the exposure of a stressor related to an aging process with the disorder of neuromuscular, metabolic, and immune systems that are at risk of comorbidities, disability, and mortality.¹⁴ Previous study in America showed that in the elder women (age >65 y.o), the deficiency of 25(OH)D can be used as a predictor of the risk of sarcopenia and frailty.¹⁵ Other studies in South Korea showed that levels of 25(OH)D less than 10 ng/dL are associated with the incidence of sarcopenia.¹⁶ Sarcopenia is a condition of pre-frailty and also the precursor of frailty.¹⁷ The elderly people with sarcopenia are three times as likely to develop frailty.¹⁸

Generally, the pathogenesis of sarcopenia and frailty are broadly the same. It begins with the changes in size and length of muscle fibers, loss of type-2 muscle fibers, the resistance of anabolic signal, and low physical performance which lead to the decrease of ATP production by muscle, denervation of type-2 muscle fiber, and decrease of the neuromuscular junction. All the processes mentioned above can cause a decrease in muscle mass, muscle strength, and physical performance.

The involvement of vitamin D in physical performance is at the cellular level and the structure of muscle. Deficiency of 25(OH)D can cause a decrease/atrophy of type II muscle fibers and increase infiltration adipocyte between muscle fibers. Another study stated that in post-menopausal women who consumed an additional intake of vitamin D and calcium had better physical performance than if they only consumed calcium.^{19,20} This study aimed to compare the different levels of 25(OH)D between sarcopenia and frailty in the elderly community in Surabaya.

MATERIALS AND METHODS

This study used a cross-sectional design to compare the levels of 25(OH)D between the subjects with sarcopenia and frailty by using a ratio scale data. The subjects of this study were the elder women from five regions of public health center (Puskesmas) in Surabaya, Indonesia, they were: Puskesmas Menur from East region of Surabaya, Puskesmas Perak Timur from North Region of Surabaya, Puskesmas Tambak Rejo from Central Region of Surabaya, Puskesmas Semerni from West Region of Surabaya, and Puskesmas Putat Jaya from South Region of Surabaya. The subjects must undergo the interview, physical examinations, and undersign the inform consent and informed to consent.

The subjects were categorized into two groups; sarcopenia and frailty based on the examination before. The diagnose of sarcopenia was based on physical examination that found a decrease in muscle mass, muscle strength, and physical performance. The diagnose of frailty was based on the criteria of the Cardiovascular Health Study (CHS). Afterward, the subject must be screened and divided into inclusion and exclusion criteria. The inclusion criteria in this study were elder women >60 years old, able to do active communication, met the criteria of sarcopenia, met the criteria of frailty, and willing to be involved in this study by undersigning the inform consent an informed to consent. The exclusion criteria in this study were obesity (BMI >25kg/m²), history of kidney disease, history of liver disease, and subjects with supplementation of vitamin D. Subjects that were qualified into inclusion criteria must be examined the level of 25(OH)D serum by using The Chemiluminescent Immune Assay (CLIA) method. The levels of 25(OH)D were classified into three groups, they were; sufficient (levels of 25(OH)D>30ng/dL), insufficient (levels of 25(OH)D29–20ng/dL), and the deficiency (levels of 25(OH)D<20ng/dL).

Statistical Analysis

Baseline characteristics were presented descriptively, in frequency, and in percentage for all categorical data. All data of normality distribution were tested using the Kolmogorov Smirnov test because the subjects of this study were more

than 50 subjects. To compare the different levels of 25(OH)D between sarcopenia and frailty, we used the Mann-Whitney test. The Mann-Whitney test was utilized because the data did not have normal distribution. The statistical significance was achieved if p-value was <0.05. All statistical analyses were undertaken using the SPSS program version 25.0.

RESULTS

The general characteristics of subjects are shown in Table 1. The total subjects of this study were 55 samples; they were 28 subjects with sarcopenia and 27 subjects with frailty. The ranged age of the total subjects were 60–100 years with the median age of 66 years. Most of the subjects in this study were in sarcopenia (88%) and frailty (89%) and living at their own house (not joining their relatives). The marital status of the subjects was mostly marriage and widow, most of them were unemployed, some of them were retirement, and some of them were still working as an enterpriser. The financial support of the subjects was varying, most of them got their financial support from their child, some of them were from retirement fund, and some others got from their own. Physical activity in the daily exercise of the subjects found that subjects with sarcopenia were more active than those with frailty. Most of the subjects had sarcopenia of 85.7% and frailty of 92.6% and did not use walking aid. Mostly, from both groups, subjects did not have a history of falls for the last one year.

For the comorbid factor, this study found that subjects mostly got a musculoskeletal disorder, diabetes, and hypertension. Based on Table 2, comprehensive geriatric assessment for Barthel Activity Daily Living (ADL) showed that subjects with sarcopenia (71.4%) were more independent than those with frailty (66.7%). For Mini Nutritional Assessment (MNA), both groups of sarcopenia (50%) and frailty (51.8%) were at risk of malnutrition.

In this study, Table 3 shows most of the total subjects, as many as 43 (78.2%), were at deficiency 25(OH)D with range level of 6.2–32.7 and mean of 16.1. Subjects with deficiency level of 25(OH)D were 20 (71.4%) with sarcopenia and 23 (85.1%) with frailty. Subjects with an insufficiency level of 25(OH)D were 6 (21.4%) with sarcopenia and 23 (85.1%) with frailty. Subjects with sufficiency level of 25(OH)D were 2 (7.1%) with sarcopenia and 1 (3.7%) with frailty.

Physical examination of the subjects are shown in Table 4. The sarcopenia's subjects had a better physical condition than frailty in each component of the physical examination. While Table 5 shows the mean and median differences between two groups. The mean 25(OH)D levels of sarcopenia was 18.05 and frailty was 14.1, the median value was 17.1 for sarcopenia and 13.2 for frailty. Analysis for normality of distribution data concluded that this study data was not normal, and the analysis of the Mann-Whitney test showed that p-value = 0.014 so that sarcopenia subjects had higher levels of 25(OH)D compared to frailty subjects.

DISCUSSION

Most of the subjects in this study were at the deficiency of 25(OH)D. The comparison between two groups revealed there was a higher level of 25(OH)D in subjects with sarcopenia than those with frailty. The subjects with

sarcopenia will third times be at risk to fall into frailty, so it was concluded that sarcopenia could be an indicator of frailty.^{18,21}

This study only used woman subjects. The study in Malaysia showed that the prevalence of 25(OH)D deficiency was greater in elder women than men.²² Decreased growth hormone (GH) and Insulin-Like Growth Factor (IGF-1) in post-menopausal women causes a decrease in muscle protein synthesis. Levels of IGF and estrogen decrease in post-menopausal women.²³ On the contrary, an increase in proinflammatory cytokines such as IL-6 and TNF- α can increase the risk of sarcopenia and frailty by reducing muscle protein synthesis. Estrogen has an anabolic mechanism in the muscles; the actions of estrogen in the muscle are by stimulating the IGF-1 receptor. Estrogen receptors in post menopause are lower than at perimenopause, so it is prone to get sarcopenia and frailty.²⁴

Most of the subjects in this study were retirement and unemployment and got financial support from their child or retirement fund. It is similar to another study revealed that most of the subjects were widows with financial support from the retirement fund.²⁵ Most of the subjects were at risk of malnutrition based on the MNA scoring system. It is related to the circumstances of socio-economics of the subjects with their nutritional status. It was shown that the elderly are really close to the malnutrition.

In this study, socio-economic data had a strong relation to employment status. It had a statistically significant value. Subjects who were unemployed were more frequent in frailty than sarcopenia. The impact of socio-economic factors on low levels of 25(OH)D in the elderly had an adverse nutritional status.²⁵ Relationship between subject's income with low levels of 25(OH)D was the subjects with low income were unable to buy nutritious food sources containing vitamin D. It was in line with study in Poland that suggested that low-income subjects rarely consumed food sources of vitamin D.²⁶ As many as 90.4% in female subjects had levels of 25(OH)D <30 ng/dL. The main source of vitamin D in the body is through exposure to UV-B rays and vitamin D-rich food ingredients.²⁶

Based on US Endocrine society, the levels of 25(OH)D were classified into three classifications: sufficiency when the levels of 25(OH)D serum >30ng/mL; insufficiency when the levels of 25(OH)D serum 21-29 ng/mL; deficiency when the levels of 25(OH)D serum <20ng/mL. It is suitable according to the theory that explained the involvement of 25(OH)D levels with physical performance and muscle structure. This study found that at each component of the physical examination, the sarcopenia's subjects had a better physical condition than frailty. Vitamin D deficiency can lead to a decrease/atrophy of type II muscle fibers and increase fat cell infiltration between muscle fibers so that it can cause sarcopenia and frailty.⁷⁻⁹

This study has a several limitations. It is only a cross sectional research in a several times, so it cannot describe the process of the occurrence of deficiency 25(OH)D. The exclusion criteria of this study were only based on interviews and physical examinations of the subjects so that exclusion criteria were difficult to objectively assess. This study has several confounding factors that cannot be excluded, such as

a history of nutrient content of daily food, a history of sun exposure, and the magnitude/level of spectrum of UV-B ray exposure.

CONCLUSION

In the conclusion of this study, it is obvious that subjects with sarcopenia has better physical condition than frailty, so the results of this research showed that the levels of 25(OH)D of subjects with sarcopenia were significantly higher than those with frailty. Further research with retrospective design could be done, so the relationship between deficiency 25(OH)D with sarcopenia and frailty will be clear, and the research of vitamin D supplementation with a certain dose in some certain periods is needed to be done to find out the optimal dose to reach optimal levels of 25(OH)D.

ACKNOWLEDGMENTS

This study did not receive any funding nor grant.

CONFLICT OF INTEREST

The authors in this study declared that they do not have any conflict of interest concerning this manuscript.

REFERENCES

1. Amelia R, Harahep J, Wahyuni AS, Pratama A. Health status of elderly based on daily activities living, cholesterol and uric acid profile in Medan city. In: IOP Conference Series Earth and Environmental Science. Institute of Physics Publishing; 2018.
2. Miftahussurur M, Yamaoka Y. Helicobacter pylori virulence genes and host genetic polymorphisms as risk factors for peptic ulcer disease. *Expert Rev Gastroenterol Hepatol*. 2015;9(12):1535.
3. Capatina C, Carsote M, Caragheorghieopol A, Poiana C, Berteanu M. Vitamin d deficiency in postmenopausal women - biological correlates. *Maedica (Buchar)*. 2014 Dec;9(4):316-22.
4. Vera V, Setiati S, Roosheroe AG. Determinan diagnostik klinis defisiensi vitamin D pada wanita berusia lebih dari 50 tahun. *J Penyakit dalam Indones*. 2015;2(1):38-48.
5. Suryadinata RV, Lorensia A. Food Frequency, Knowledge about Vitamin D and Obesity among Elderly. *Amerita Nutr*. 2020;4(1):43-8.
6. Thalib B, Horas B, Machmud E, Asmawati, Hasyim R. Alteration of memory and depression in elderly with full overdenture - Pilot study. *J Int Dent Med Res*. 2018;11(1):62-5.
7. Arik G, Ulger Z. Vitamin D in sarcopenia: Understanding its role in pathogenesis, prevention and treatment. *Eur Geriatr Med*. 2016;7(3):207-13.
8. Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem Biol*. 2014 Mar;21(3):319-29.
9. Christakos S, Ajibade D V, Dhawan P, Fechner AJ, Mady LJ. Vitamin D: metabolism. *Endocrinol Metab Clin North Am*. 2010 Jun;39(2):243-53, table of contents.
10. Siregar B, Andayani U, Bahri RP, Seniman, Fahmi F.

- Real-time monitoring system for elderly people in detecting falling movement using accelerometer and gyroscope. In: *Journal of Physics: Conference Series* [Internet]. Faculty of Computer Science and Information Technology, Universitas Sumatera Utara, Medan, Indonesia: Institute of Physics Publishing; 2018. Available from: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85045135350&doi=10.1088%2F1742-6596%2F978%2F1%2F012110&partnerID=40&md5=296f8bd2d19e2a2baba95660d452efda>
11. Kristiana T, Widajanti N, Satyawati R. Association between Muscle Mass and Muscle Strength with Physical Performance in Elderly in Surabaya. *Surabaya Phys Med Rehabil J*. 2020;2(1):24–34.
 12. Fornelli G, Izain GC, D'amelio P. Ageing, muscle and bone. *J Gerontol Geriatr*. 2016;3:75–80.
 13. Ogawa S, Yakabe M, Akishita M. Age-related sarcopenia and its pathophysiological bases. *Inflamm Regen*. 2016;36: 17.
 14. Lappe JM, Binkley N. Vitamin D and Sarcopenia/Falls. *J Clin Densitom Off J Int Soc Clin Densitom*. 2015;19(4):473–82.
 15. Ensrud KE, Ewing SK, Fredman L, Hochberg MC, Cauley JA, Hillier TA, et al. Circulating 25-hydroxyvitamin D levels and frailty status in older women. *J Clin Endocrinol Metab*. 2010 Dec;95(12):5266–73.
 16. Park S, Ham J-O, Lee B-K. A positive association of vitamin D deficiency and sarcopenia in 50 year old women, but not men. *Clin Nutr*. 2014 Oct;33(5):900–5.
 17. Wilson D, Jackson T, Sapey E, Lord JM. Frailty and sarcopenia: The potential role of an aged immune system. *Ageing Res Rev*. 2017 Jul;36:1–10.
 18. Frisoli AJ, Chaves PH, Ingham SIM, Fried LP. Severe osteopenia and osteoporosis, sarcopenia, and frailty status in community-dwelling older women: results from the Women's Health and Aging Study (WHAS) II. *Bone*. 2011 Apr;48(4):952–7.
 19. Pfeifer M, Bergerow B, Minne HW, Abrams C, Nachtigall D, Hansen C. Effects of a short-term vitamin D and calcium supplementation on body sway and secondary hyperparathyroidism in elderly women. *J Bone Miner Res Off J Am Soc Bone Miner Res*. 2000 Jun;15(6):1113–8.
 20. Pfeifer M, Bergerow B, Minne HW, Suppan K, Fahrleitner-Pammer A, Dobnig H. Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community-dwelling older individuals. *Osteoporos Int a J Establ as result Coop between Eur Found Osteoporos Natl Osteoporos Found USA*. 2009 Feb;20(2):315–22.
 21. Viana JJ, Silve SLA, Torres JL, Dias JMD, Pereira LSM, Dias RC. Influence of sarcopenia and functionality indicators on the frailty profile of community-dwelling elderly subjects: a cross-sectional study. *Brazilian J Phys Ther*. 2013;17(4):373–81.
 22. Moy FM. Vitamin D status and its associated factors of free living Malay adults in a tropical country, Malaysia. *J Photochem Photobiol B*. 2011 Sep;104(3):444–8.
 23. Randah, Masni, Theha RIM. Relationship between hormonal contraception use with age of menopause among elderly at Posyandu Sidodadi village Wonorejo subdistrict Polewali Mandar district. In: *Proceedings of the International Conference on Healthcare Service Management 2018. Association for Computing Machinery*; 2018. p. 68–72.
 24. Maltais ML, Desroches J, Dionne I. Changes in muscle mass and strength after menopause. *J Musculoskelet Neuronal Interact*. 2009;9(4):186–97.
 25. Sari DK, Rasyid H A I, Lipoeto NI, Lubis Z. Occurrence of vitamin D deficiency among women in North Sumatera, Indonesia. *Malays J Nutr [Internet]*. 2014;20(1):63–70. Available from: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-84909608080&partnerID=40&md5=42b00cd544df499d82639bab6fcb2d2>
 26. Wyskida M, Owczarek A, Szybalska A, Brzozowska A, Szczerbowska I, Wieczorowska-Tobis K, et al. Socio-economic determinants of vitamin D deficiency in the older Polish population: results from the PolSenior study. *Public Health Nutr*. 2018 Aug;21(11):1995–2003.

Table 1: Baseline characteristics

Characteristics	Sarcopenia (n=28)		Frailty (n=27)		p-value
	n	%	n	%	
Residence					
• Private house	25	89.3	24	88.9	0.136
• Joint house (Family)	2	7.1	1	3.7	
• Rent house	1	3.6	1	3.7	
• etc.	0	0.0	1	3.7	
Status					
• Marriage	13	46.4	15	55.6	0.659
• Widow	13	46.4	11	40.7	
• Unmarried	2	7.1	1	3.7	
Job					
• Unemployment	17	60.7	23	85.2	0.039
• Enterpriser	6	21.4	2	7.4	
• Employee of a private company	1	3.6	1	3.7	
• Retirement	3	10.7	0	0.0	
• Etc.	1	3.6	1	3.7	
Financial Support					
• Independent cost	6	21.4	5	18.5	0.325
• Support from another family member and independent cost	4	14.3	2	7.4	
• Support from another family member	14	50.0	18	66.7	
• Retirement fund	4	14.3	2	7.4	
Gait style					
• No walking aid	24	85.7	25	92.6	0.104
• With walking aid	4	14.3	2	7.4	
Exercise					
• Yes	22	78.6	11	40.7	0.070
• No	6	21.4	16	59.3	
History of fall					
• Yes	2	7.1	4	14.8	0.070
• No	26	92.9	23	85.2	

Table 2: Comprehensive Geriatric Assessment

Comprehensive Geriatric Assessment	Sarcopenia(n=28)		Frailty(n=27)		p-value
	n	%	n	%	
Barthel Activity Daily Living (ADL)					
Independent	20	71.4	18	66.7	0.17
Mild dependency	8	28.6	8	29.6	
Moderate dependency	0	0.0	1	3.7	
MMSE					
Normal	26	92.9	22	81.5	0.095
Mild Cognitive disorder	1	3.6	4	14.8	
Severe Cognitive disorder	1	3.6	1	3.7	
Mini Nutritional Assessment (MNA)					
Malnutrition	6	21.4	7	25.9	0.19
Risk of Malnutrition	14	50.0	14	51.8	
Normal	8	28.5	6	22.2	
GDS					0.95

Comprehensive Geriatric Assessment	Sarcopenia(n=28)		Frailty(n=27)		p-value
	n	%	n	%	
Normal	27	96.4	26	96.3	
Possible Depression	1	3.6	1	3.7	

Table 3: 25(OH)D Levels in sarcopenia and frailty

25(OH)D Levels	Sarcopenia (n=28)		Frailty (n=27)	
	n	%	n	%
Deficiency	20	71.4	23	85.1
Insufficiency	6	21.4	3	11.1
Sufficiency	2	7.1	1	3.7

Table 4: Physical examination of the subjects

Physical examination	Sarcopenia		Frailty		P-value
	Median	Mean±SD	Median	Mean±SD	
Muscle strength:					
<18Kg	14.5	14.91±1.5	15	13.72±3.1	0.04
>18Kg	21	21.75±2.3	20	20.77±2.8	
Muscle mass					
<3Kg/m ²	2.54	2.6±0.39	2.63	2.56±0.2	0.037
>3Kg/m ²	-	-	3.36	3.50±0.4	
Gait speed test					
>0.8m/mnt	0.64	0.60±0.1	0.61	0.58±0.1	0.51
<0.8m/mnt	-	-	-	-	
PASE score					
<126.50Kkal/week	58.09	64.27±29.1	55.25	54.19±32.2	0.001
>126.51Kkal/week	180.39	187.46±42.2	-	-	

Table 5: Physical examination components

Variable	Groups	Median	Mean	p-value	Conclusion
25(OH)D	Sarcopenia	17.1ng/dL	18.05 ng/dL	0.014	Significantly different
	Frailty	13.2ng/dL	14.1 ng/dL		