

23. Effect of Acute Persadia Gymnastic on The Ability to Concentrate in Diabetic Patients

by Desi Rianti Rahmadhani

Submission date: 23-Aug-2022 02:25PM (UTC+0800)

Submission ID: 1885840184

File name: Gymnastic_on_The_Ability_to_Concentrate_in_Diabetic_Patients.pdf (239.61K)

Word count: 4220

Character count: 22042

Original Research

Effect of Acute Persadia Gymnastic on The Ability to Concentrate in Diabetic Patients

Desi Rianti Rahmadhani^{1*}, Lilik Herawati², Sony Wibisono Mudjanarko³, Kristanti Wanito Wigati², Laylatul Fitriah Mukarromah⁴, Winna Adelia Amru⁵


¹Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

²Department of Physiology, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

³Department of Internal Medicine, Dr. Soetomo General Hospital, Universitas Airlangga, Surabaya, East Java, Indonesia

⁴Faculty of Medicine, Universitas Jember, East Java, Indonesia

⁵Faculty of Medicine, Universitas Brawijaya, East Java, Indonesia

*Corresponding Author: 

Desi Riati Rahmadhani, Faculty of Medicine, Universitas Airlangga Jl. Mayjen Prof. Dr. Moestopo No.47, Surabaya, East Java, Indonesia

Email: desi.rianti.rahmadhani-2017@fk.unair.ac.id

Article info:

Received: November 29th, 2020;

Received in revised: November 1st, 2021;

Accepted: November 3rd, 2021

Published: February 25th, 2022



This is an open access article under the CC-BY license

(<http://creativecommons.org/licenses/by/4.0/>)



Cite this as:

Rahmadhani DR, Herawati L, Mudjanarko SW, Wigati KW, Mukarromah LF, Amru WA. Effect of acute persadia gymnastic on the ability to concentrate in diabetic patients. *SPMRJ*. 2022;4(1):7-14.

ABSTRACT

Background: Diabetes Mellitus (DM) can affect the peripheral and central nervous system. It can affect cognitive function, one of which can be seen from the decrease in the concentration ability. The concentration ability can be assessed by measuring the reaction time. Reaction time is the time interval between the onset of a stimulus and the initiations of a response. One of the important treatments for DM is physical exercises. Persadia gymnastic is one example of physical exercise that is specially designed for people with DM.

Aim: The purpose of this study was to identify the effect of acute Persadia gymnastic on the concentration ability of people with DM.

Material and Methods: This was an experimental study in type 2 DM patients. Fourteen subjects who met the inclusion criteria were randomly grouped into control (without performing Persadia gymnastic) and intervention group (performing Persadia gymnastic). The subjects' reaction times were measured before and after the intervention. The mean results of the reaction time of all subjects were then compared between the control and the intervention groups using the Independent-T test and Mann-Whitney tests.

Results: The analyzed data were only 7 subjects per group with a total of 14 subjects. The mean value of reaction time in the intervention group was decreased, both audio (-0.04 ± 0.85) and visual (-0.09 ± 0.56). However, there was no significant difference of the reaction time between control and intervention groups ($p > 0.05$).

Conclusion: Many factors can affect reaction time and some still cannot be controlled in this study due to a lot of limitations. Further research is still needed to figure out the effect of persadia gymnastics on the ability to concentrate in people with Diabetes Mellitus.

Keywords: *cognitive function, Diabetes Mellitus, Persadia, physical exercise, reaction time*

Introduction

Diabetes Mellitus (DM) is a metabolic disease characterized by an increase in blood glucose levels above normal or it is commonly referred to as hyperglycemia.¹ The Indonesian Basic Health Result 2018 showed the prevalence of DM based on blood tests in people aged ≥ 15 years old according to the 2015 Indonesian Society for Endocrinology Consensus criteria was 10.9% and according to the 2011 Indonesian Society for Endocrinology Consensus criteria was 8.5%. This number showed an increase in prevalence from 2013 which was only 6.9%.²

High blood sugar levels in people with type 2 DM often caused neurological negative effects that affect the peripheral and central nerves.³ People with DM are more likely to experience impaired cognitive function compared to those without DM.^{4,5} Cognitive function is a process that includes executive function, attention or reasoning, speed of processing, memory, and language.^{4,5,6}

The most common way to evaluate the speed of processing is to assess reaction times (RTs).⁷ Reaction time or response latency is defined as the time interval between the delivery of a stimulus and the initiation of a response.^{7,8} Reaction time is an index of processing speed in the central nervous system with the stimulation of sensory input modalities such as visual, auditory, pain, touch or temperature.³

In the management of DM, there are important pillars in controlling the disease which are education, medical nutritional therapy, physical exercise, and pharmacological therapy.⁹ Physical exercise can increase specific improvements in neuropathy symptoms, including an increase of nerve fibers branches.¹⁰ Other research showed that there was an increase in speed of reaction time after aerobic exercise, which was 4% 7% faster reaction than the initial reaction.¹¹

The criteria of physical exercise recommended for people with DM are aerobic low impact and rhythmic.^{12,13} In

Indonesia, there is a physical exercise that was specially designed for people with DM, namely Indonesian Diabetes Association (IDA) exercises or commonly known as Persadia (*Persatuan Diabetes Indonesia*) gymnastic.^{12,13,14} However, until now no one has discussed the effect of acute Persadia gymnastic on the concentration ability of people with DM. Based on this background, a study was conducted to identify the effect of acute Persadia gymnastic on the concentration ability of people with DM.

Material and Methods

This research was an experimental study with a pre-post evaluation design. The population in this study were people with type 2 DM. The sample size of this study was determined by using the Lemeshow formula, and based on research conducted by Ando et al.,¹⁵ we obtain the value of $n = 6$. To anticipate the possibility of the subject dropping out, the researchers set a correction factor of 30%, so that the result became 8 samples per study group. The sampling technique used in this study was purposive sampling. Samples were determined according to the inclusion and exclusion criteria determined by the researchers.

Inclusion criteria of this study were people with type 2 DM who were willing to participate in research by filling out the informed consent, age between 40-65 years old, having diabetes for at least 5 years, Random Blood Glucose (RBG) level 100 - 250 mg/dl, and Body Mass Index (BMI) 18.5 - 29.9 kg/m². Exclusion criteria of this study were people with impaired physical mobility, significant cardiovascular disorders (e.g. coronary heart disease, heart failure, and stroke), vision and hearing problems (related to the reaction time examination), end-stage renal disease (nephropathy), and uncontrolled hypertension (blood pressure $\geq 140/90$ mmHg). Drop out's criteria of this study were subjects who were not able to complete the research protocol and subjects who were suffered from illness or injury during the research.

The instruments used in this study were whole-body reaction measuring equipment, active speakers, scale and height measuring device, blood glucose kit, tensimeter, stethoscope, and pulse-oximetry. This research was conducted at the Department of Physiology Laboratory, Faculty of Medicine, Universitas Airlangga on 22 – 23 February 2020. This research has been declared to be ethically appropriate by Health Research Ethics Committee, Faculty of Medicine, Universitas Airlangga (No. 27/EC/KEPK/FKUA/2020). Signed informed consent was obtained from all participants.

This research was divided into 2 days. Each day was attended by 8 research subjects. Subjects were instructed to have breakfast 1 hour before doing the exercise. The screening was done before doing the exercise by measuring RBG levels, blood pressure (BP), heart rate (HR), oxygen saturation (SpO₂), height, and weight of all subjects. After screening, those 8 subjects were divided randomly into 2 groups (intervention and control group) using a lottery paper which also had the serial number written for the queue. According to the 2015 Indonesian Society for Endocrinology Consensus,⁹ if the subject has RBG level of more than 250 mg/dl, the subject is advised to postpone the physical exercise. Therefore, the researchers decided to immediately assign the subject to the control group if such a case was found.

The concentration ability test was assessed by measuring the speed of subject's auditory and visual reaction time using the whole-body reaction measuring equipment alternately, according to the serial number on the lottery paper (alternating between the intervention and control groups). Each subject was tested with audio stimulation for 3 times and visual stimulation for 3 times. Every time a stimulus was given, the subject must respond as quickly as possible by pressing the button that matches the stimulus.

After measuring the reaction time of all subjects, the intervention group immediately continued to perform Persadia gymnastic together within 35 minutes that consists of warm-up (5 minutes), core (15

minutes), transition (10 minutes), and cool-down (5 minutes). It was a single bout or one session of aerobic low-impact exercise, with moderate intensity (50-70% HR max). After exercising, the heart rate, oxygen saturation, blood pressure, and RBG level were measured again immediately. The control group who did not perform Persadia gymnastic at that time was asked to wait for 35 minutes according to the exercise time. Then, their heart rate, oxygen saturation, blood pressure, and RBG level were also immediately measured. After that, the concentration ability was assessed again alternately according to the lottery serial number. The mean value of reaction times changes before and after intervention were then compared between the control and the intervention groups. In this study, the normality test was done by using the Shapiro-Wilk test. The data that were not normally distributed ($p < 0.05$) were then continued by using Mann-Whitney test, while the data that were normally distributed were then continued by using the T-test.

Results

Of the total 16 subjects with type 2 DM who participated in this study, 2 subjects were dropped out. Therefore, the data analyzed were became 14 subjects with 7 subjects per group.

Table 1. General Characteristics of Research Subjects

	Control (n=7)			Intervention (n=7)		
	Min	Max	Mean ± SD	Min	Max	Mean ± SD
Age (years old)	49	67	58.57 ± 6.45	49	60	56.43 ± 3.70
BMI (kg/m ²)	20.77	29.77	26.78 ± 3.18	21.17	27.04	24.83 ± 1.91
HR-pre (bpm)	65	98	88.57 ± 11.33	73	91	85.00 ± 6.66
BPsys-pre (mmHg)	120	170	144.29 ± 17.18	110	150	128.57 ± 13.45
BPdya-pre (mmHg)	80	100	90.00 ± 8.17	80	90	85.71 ± 5.34
O ₂ Sat.-pre (%)	98	100	98.71 ± 0.76	95	99	97.57 ± 1.81
RBG-pre (mg/dl)	128	462	210.00 ± 116.30	142	250	192.86 ± 41.06
RTvisual-pre (sec)	0.414	2.922	1.32 ± 0.93	0.33	1.49	0.81 ± 0.45
RTaudio-pre (sec)	0.767	2.327	1.59 ± 0.64	0.41	1.95	1.33 ± 0.51

Table 2. Posttest Results

	Control (n=7)			Intervention (n=7)		
	Min	Max	Mean ± SD	Min	Max	Mean ± SD
HR-post (bpm)	67	105	87.71 ± 13.20	82	114	96.29 ± 10.44
BPsys-post (mmHg)	120	170	145.71 ± 15.12	120	160	132.86 ± 14.96
BPdya-post (mmHg)	80	110	88.57 ± 10.69	80	90	81.43 ± 3.78
O ₂ Sat-post (%)	98	99	98.43 ± 0.53	97	99	97.86 ± 0.69
RBG-post (mg/dl)	82	313	214.00 ± 85.77	83	252	163.14 ± 56.51
RTvisual-post (sec)	0.38	2.80	1.31 ± 1.05	0.33	1.67	0.72 ± 0.44
RTaudio-post (sec)	0.87	4.46	2.27 ± 1.19	0.59	2.03	1.29 ± 0.51

Table 3. P Value of Delta between The Control and Intervention Groups

Variable	Group	Mean ± SD	P value
Delta RT visual (sec)	Control	-0.01 ± 1.04	0.87*
	Intervention	-0.09 ± 0.56	0.87*
Delta RT audio (sec)	Control	0.68 ± 0.92	0.14**
	Intervention	-0.04 ± 0.85	0.16*

* = Independent T-test; ** = Mann-Whitney test

The mean value of the intervention group that performed Persadia gymnastic showed a decrease in the reaction time, both audio (-0.04 ± 0.85) and visual (-0.09 ± 0.56). This can be seen from the minus value in the delta which indicates that the value after the intervention is smaller than the value before the intervention. This decrease in value indicates that the average speed of the subject's reaction time was faster after being given the intervention.

From the results of the Independent T-test and the Mann-Whitney test, all the data in the intervention group and the control group obtained a $p > 0.05$ which means that there was no significant difference between the control group and the intervention group.

Discussion

Physical exercise could have acute or chronic effects. The effect will be chronic if it is done repetitively, and will be acute if it is only done once. A previous study used the ergocycle to analyze the effect of a single bout or acute moderate-intensity physical exercise on the level of concentration in teenagers. In this study significant differences of concentration level between pre- and post-exercise were found.¹⁶ Other types of physical exercise such as sprints were also improved response accuracy in another study.¹⁷ Several studies also showed that after moderate-intensity physical exercise of 50-70% HR max, the executive function, especially response accuracy was increased.^{18,19,20}

In this study, it was difficult to find subjects that really met the expected criteria and willingly took part in the research. The total number of subjects obtained were 16 people, and 2 were dropped out. Therefore, the total subjects analyzed were only 14 people and all of them were females. This research was divided into 2 days due to the limited instrument availability.

Diabetics are advised to do regular physical exercise for 3-5 days a week with moderate-intensity (50-70% maximum HR) for 30-45 minutes.⁹ In this research, the physical exercise taken was in moderate intensity and only a single bout or one session exercise. The intervention group

performed the gymnastic for once and the changes observed here are the acute responses from the result of exercise. Before and after the intervention, the visual and audio reaction time of each subject was measured in turn according to the queue serial number because there was only one instrument used to measure the reaction time. The instrument can measure the visual and audio reaction time.

Reaction time is an important physiological parameter to provide information on how fast a person reacts to a stimulus and is often used to evaluate the timing of the thought process and coordination between the sensory and motor systems.²¹ The process stages that determine the reaction time include the arrival of stimuli, the process of sensation, perception, which is then manifested in the form of a response to stimuli.²² If there was interference in one of these factors, then the person's reaction time will not be on optimum value.

The executive function of the brain, especially the function to concentrate, could be improved by acute physical exercise. Inhibition control which is one of the executive functions had an important role in concentration. When other stimuli were suppressed, it could lead the subjects to only focus on one stimuli.²³ Many factors can affect the speed of reaction time, including age, gender, arousal, right/left-handed dominance, diet, exercise, practice and errors, intelligence, fatigue, distraction, personality type (psychological), alcohol, stimulants (caffeine), brain injury, illness, type of stimulus (audio or visual), the sequence of stimulus given, and intensity of the stimulus.²⁴

In this study, only 3 of the total 14 subjects (21.43%) have routinely exercised for at least 3 times a week as recommended by American Diabetes Association and Indonesian Society for Endocrinology.^{1,9} For diet and nutritional intake, consumption of alcohol and stimulant drugs or caffeine were not observed in this study. The level of understanding and the ability to remember the type of stimulus varied between

subjects due to the differences of intelligence and education background. In this research, mental and psychological conditions of each subject were also not observed by the researchers.

Exercise can increase the Nerve Growth Factor (NGF)²⁵ and BDNF (brain-derived neurotrophic factor) release.²⁶ Exercise can also increase Dopamine (DA), noradrenaline (NE), and serotonin (5-HT) levels.²⁷ In addition, exercise can increase insulin sensitivity,²⁸ heart rate, cardiac output, and cerebral blood flow to blood oxygenation to the brain.²⁹ Exercise can also increase the amplitude of P300 and decrease the latency of P300, a component of ERP (event-related potential) that has a role in the decision-making function or the allocation of the body's response which is related to the attention function.³⁰

Indonesian Diabetes Association (2000), as quoted by Rahim et al,¹⁴ stated that diabetes exercise is a physical exercise designed based on age and physical status and is part of the treatment of DM. Previously, diabetes exercise had 6 series of movements in the form of rhythmic stretching and relaxation in muscles, joints, vascular, and nerves.^{12,13,31} According to *Senam Diabetes Persadia 1* (2017), as quoted by Asrizal,³² Persadia gymnastic series 1 is a continuation of the previous series diabetes exercises with a structured, rhythmic, aerobic low impact, moderate-intensity exercise dominated by eccentric movements consisting of warm-up, core, transition and cool-down sections. This exercise is in accordance with American Diabetes Association and Indonesian Society for Endocrinology recommendations as moderate-intensity aerobic exercise, but the main movement is eccentric movement, that previous research has shown to increase the activity of GLUT 4 and GLUT 1 in the function of blood glucose uptake so that it is expected to further improve blood glucose, glucose tolerance, and insulin resistance.^{1,9,12,13,33}

In this study, the reaction time was measured after the intervention and cannot be done simultaneously and immediately on all subjects because there was only 1 instrument available to assess all the

subjects. Because of these limitations, the acute effects caused after exercise cannot be detected immediately. For subjects who have waited for a long time, the levels of the neurotrophic factor, dopamine, epinephrine, and HR that initially increased, can fall back to normal again so that the increase cannot be detected. The differences that occurred when the measurement test was carried out between one subject to another may be the reason for the insignificant difference in reaction time.

Conclusion

There was no significant difference in both audio and visual reaction time ($p > 0.05$), between the control and intervention groups. Although the difference was not significant, there was a change in reaction time before and after the intervention.

Some factors that can affect the reaction time were still cannot be controlled in this study due to a lot of limitations. Although no significant difference was found, Persadia Gymnastic still can be recommended for people with Diabetes Mellitus in daily living activities to maximize their quality of life and can be used as a treatment and prevention because it can increase physical fitness, improve glucose tolerance and reduce HbA1c.^{12,13,14} Further research with a proper instrument and appropriate subject is still needed to figure out the effect of persadia gymnastics on the ability to concentrate in people with Diabetes Mellitus.

Acknowledgement

This research was supported by Department of Physiology, Faculty of Medicine, Universitas Airlangga for the reaction time measuring equipment and pulse-oximetry. For the process of finding subjects, this research was supported by Indonesian Diabetes Association - Surabaya Diabetes and Nutrition Center (SDNC) Dr. Soetomo General Hospital. There was no funding for this research.

References

1. Association AD. Diabetes basics: Common terms. Retrieved March. 2014;24:2017.
2. Kementerian Kesehatan RI. Riskendas 2018. Lap Nas Riskendas 2018. 2018;44(8):181–222.
3. Niruba R, Maruthy KN. Assessment of Auditory and Visual Reaction Time in Type 2 Diabetics – A Case Control Study. *Al Ameen J Med Sci.* 2011;4(3):274–9.
4. Xue M, Xu W, Ou YN, Cao XP, Tan MS, Tan L, et al. Diabetes mellitus and risks of cognitive impairment and dementia: A systematic review and meta-analysis of 144 prospective studies. *Ageing Res Rev.* 2019;55(August):100944.
5. Lyu F, Wu D, Wei C, Wu A. Vascular cognitive impairment and dementia in type 2 diabetes mellitus: An overview. *Life Sci.* 2020;254(February):117771.
6. Guo M, Kang K, Wang A, Jia J, Zhang J, Wang Y, et al. Association of diabetes status with cognitive impairment in two Chinese rural communities. *J Neurol Sci.* 2020;415(October 2019):116894.
7. Rigoli M, Facchin A, Cardile D, Beschin N, Luzzatti C. Open-source open-access reaction time test (OORTT): an easy tool to assess reaction times. *Neurol Sci.* 2021;42(6):2461–9.
8. Levakova M, Lansky P. Latency of inhibitory response. *BMC Neurosci.* 2013;14(S1):2202.
9. Soelistijo S, Novida H, Rudijanto A, Soewondo P, Suastika K, Manaf A, et al. *Konsesus Pengelolaan Dan Pencegahan Diabetes Melitus Tipe2 Di Indonesia 2015.* Perkeni. 2015. 82 p.
10. Kluding PM, Pasnoor M, Singh R, Jernigan S, Farmer K, Rucker J, et al. The effect of exercise on neuropathic symptoms, nerve function, and cutaneous innervation in people with diabetic peripheral neuropathy. *J Diabetes Complications.* 2012;26(5):424–9.
11. Morrison S, Colberg SR, Parson HK, Vinik AI. Exercise improves gait, reaction time and postural stability in older adults with type 2 diabetes and neuropathy. *J Diabetes Complications.* 2014;28(5):715–22.
12. Pranoto A. *Senam Persadia 1.* Indonesia: Persatuan Diabetes Indonesia; 2017.
13. Wibisono S. *Surabaya International Physiology Symposium.* Surabaya Univ Airlangga. 2017;
14. Rahim NK. Pengaruh Senam Diabetes Terhadap Penurunan Kadar Gula Darah Sewaktu Pada Pasien dengan DM tipe II di Wilayah Kerja Puskesmas Global Kecamatan Limboto Kabupaten Gorontalo. *Skripsi.* 2015;1(841411044).
15. Ando S, Kokubu M, Kimura T, Moritani T, Araki M. Effects of acute exercise on visual reaction time. *Int J Sports Med.* 2008;29(12):994–8.
16. Wijaya MA, Sulistiawaty NN, Wigati KW, Herawati L. A Single Bout of Moderate Intensity Exercise Improves Concentration Level on Teenagers. *Surabaya Phys Med Rehabil J.* 2020;2(1):1.
17. Cooper SB, Bandelow S, Nute ML, Dring KJ, Stannard RL, Morris JG, et al. Sprint-based exercise and cognitive function in adolescents. *Prev Med Reports.* 2016;4:155–61.
18. Lucas SJE, Ainslie PN, Murrell CJ, Thomas KN, Franz EA, Cotter JD. Effect of age on exercise-induced alterations in cognitive executive function: relationship to cerebral perfusion. *Exp Gerontol.* 2012;47(8):541–51.
19. Martins AQ, Kavussanu M, Willoughby A, Ring C. Moderate intensity exercise facilitates working memory. *Psychol Sport Exerc.* 2013;14(3):323–8.
20. Pesce C, Audiffren M. Does acute exercise switch off switch costs? A study with younger and older athletes. *J Sport Exerc Psychol.* 2011;33(5):609–26.
21. Gavkare AM, Surdi AD, Nanaware NL. Auditory Reaction Time, Visual Reaction Time and Whole Body Reaction Time in Athletes. 2013;
22. Syafitri AW, Supatmo Y, Indraswari DA. Perbedaan Waktu Reaksi Tanganantara Cabang Olahraga Permainan Dan Beladiri. *J Kedokt Diponegoro.* 2017;6(2):177–87.
23. Diamond A. Executive functions. *Annu Rev Psychol.* 2013;64:135–68.
24. Kosinski RJ. A Literature Review on Reaction Time. Retrieved [http://biaeclemsonedubpcbpLab110reactionhtm06042010.2010;10\(August\):2006](http://biaeclemsonedubpcbpLab110reactionhtm06042010.2010;10(August):2006).
25. Park SJ, Yong MS, Na SS. Effect of exercise on the expression of nerve growth factor in the spinal cord of rats with induced osteoarthritis. *J Phys Ther Sci.* 2015;27(8):2551–4.
26. De Azevedo KPM, De Oliveira Segundo VH, De Medeiros GCBS, De Sousa Mata AN, García DÁ, De Carvalho Leitão JCG, et al. Effects of exercise on the levels of BDNF and executive function in adolescents: A protocol for systematic review and meta-Analysis. *Med (United States).* 2019;98(28).
27. Lin TW, Kuo YM. Exercise benefits brain function: The monoamine connection. *Brain Sci.* 2013;3(1):39–53.
28. Vetrivel Venkatasamy V, Pericherla S, Manthuruthil S, Mishra S, Hanno R. Effect of physical activity on insulin resistance, inflammation and oxidative stress in diabetes mellitus. *J Clin Diagnostic Res.* 2013;7(8):1764–6.
29. Kirk-Sanchez NJ, McGough EL. Physical exercise and cognitive performance in the elderly: Current perspectives. *Clin Interv Aging.* 2013;9:51–62.
30. Kumar N, Sood S, Sakshi, Behera J, Singh M, Beena, et al. Effect of acute moderate exercise on cognitive P300 in persons having sedentary lifestyles. *Int J Appl Basic Med Res.* 2012;2(1):67.
31. Damayanti S. Diabetes mellitus dan penatalaksanaan keperawatan. *Yogyakarta Nuha Med.* 2015;
32. CYNTHIA WAHYU ASRIZAL N. PENGARUH SENAM PERSADIA 1

TERHADAP GLUKOSA DARAH PUASA,
GLUKOSA DARAH 2 JAM POSTPRANDIAL
DAN THE HOMEOSTASIS MODEL
ASSESSMENT OF INSULIN RESISTANCE
(HOMA IR). Universitas Airlangga; 2018.

33. Mice SDM, Adelia H, Bambang P, Arifa M.
The Effect of Eccentric Activity on Glucose
Transporter Type 4 in Gastrocnemius Muscle of
Streptozot. 3(3):39-43.

23. Effect of Acute Persadia Gymnastic on The Ability to Concentrate in Diabetic Patients

ORIGINALITY REPORT

16%

SIMILARITY INDEX

14%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	ners.unair.ac.id Internet Source	1%
2	oamjms.eu Internet Source	1%
3	www.dovepress.com Internet Source	1%
4	journalcra.com Internet Source	1%
5	Arifa Mustika, Nurmawati Fatimah, Gadis Meinar Sari. "The self-nanoemulsifying drug delivery system of <i>Petiveria alliacea</i> extract reduced the homeostatic model assessment-insulin resistance value, interleukin-6, and tumor necrosis factor- α level in diabetic rat models", <i>Veterinary World</i> , 2021 Publication	1%
6	Rika Wahyuni Arsianti, Dewy Haryanti Parman, Hendy Lesmana, Muhammad Taufiqqurohman. "Effect of Electrical	1%

Stimulation in Lower Extremity as Physical Exercise in Type 2 Diabetes Mellitus Patients", The Indonesian Biomedical Journal, 2018

Publication

7	kb.psu.ac.th:8080 Internet Source	1 %
8	www.researchgate.net Internet Source	1 %
9	jebmh.com Internet Source	1 %
10	repository.unair.ac.id Internet Source	1 %
11	hdl.handle.net Internet Source	1 %
12	www.atlantis-press.com Internet Source	1 %
13	medicopublication.com Internet Source	<1 %
14	www.journal-jams.org Internet Source	<1 %
15	www2.gre.ac.uk Internet Source	<1 %
16	www.semanticscholar.org Internet Source	<1 %

17	Sutrisno Sutrisno, Hardianti Aprina, Happy Marthalena Simanungkalit, Asti Andriyani et al. "Genistein modulates the estrogen receptor and suppresses angiogenesis and inflammation in the murine model of peritoneal endometriosis", Journal of Traditional and Complementary Medicine, 2017 Publication	<1 %
18	www.parasitol.kr Internet Source	<1 %
19	mdpi-res.com Internet Source	<1 %
20	www.jove.com Internet Source	<1 %
21	Walker, Jan, Almond, Palo. "EBOOK: Interpreting Statistical Findings: A Guide For Health Professionals And Students", EBOOK: Interpreting Statistical Findings: A Guide For Health Professionals And Students, 2010 Publication	<1 %
22	cyberleninka.org Internet Source	<1 %
23	persadia.or.id Internet Source	<1 %
24	www.science.gov	

<1 %

25

"1-B2: Respiratory Infections (Non-Tuberculosis) 2 : Poster Sessions",
Respirology, 2013.

Publication

<1 %

26

Ryan A. Williams, Simon B. Cooper, Karah J. Dring, Lorna Hatch, John G. Morris, Caroline Sunderland, Mary E. Nevill. "Effect of football activity and physical fitness on information processing, inhibitory control and working memory in adolescents", BMC Public Health, 2020

Publication

<1 %

27

ejobios.org

Internet Source

<1 %

28

ejournal.unsri.ac.id

Internet Source

<1 %

29

tsukuba.repo.nii.ac.jp

Internet Source

<1 %

30

Benjamin S. Boyd, Robert J. Nee, Betty Smoot. "Safety of lower extremity neurodynamic exercises in adults with diabetes mellitus: a feasibility study", Journal of Manual & Manipulative Therapy, 2016

Publication

<1 %

31

Isabelle Riedl, Megan E. Osler, Boubacar Benziane, Alexander V. Chibalin, Juleen R. Zierath. "Association of the ACTN3 R577X polymorphism with glucose tolerance and gene expression of sarcomeric proteins in human skeletal muscle", *Physiological Reports*, 2015

Publication

<1 %

32

Pritesh Hariprasad Gandhi, Pradnya A. Gokhale, H. B. Mehta, C. J. Shah. "A Comparative Study of Simple Auditory Reaction Time in Blind (Congenitally) and Sighted Subjects", *Indian Journal of Psychological Medicine*, 2020

Publication

<1 %

33

Stacy A Clemes, Daniel D Bingham, Natalie Pearson, Yu-Ling Chen et al. "Sit-stand desks to reduce sedentary behaviour in 9- to 10-year-olds: the Stand Out in Class pilot cluster RCT", *Public Health Research*, 2020

Publication

<1 %

34

acervodigital.ufpr.br

Internet Source

<1 %

35

asean-endocrinejournal.org

Internet Source

<1 %

36

journals.plos.org

Internet Source

<1 %

37

sjosm.org

Internet Source

<1 %

38

www.clinicaltrials.gov

Internet Source

<1 %

39

www.mdpi.com

Internet Source

<1 %

40

www.sid.ir

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

23. Effect of Acute Persadia Gymnastic on The Ability to Concentrate in Diabetic Patients

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
