

# Publish fast. Openly. Without restrictions.



F1000Research is an **Open Research** publishing platform for scientists, scholars and clinicians offering rapid publication of articles and other research outputs without editorial bias.

## HOW IT WORKS (/ABOUT)

SUBJECT AREAS | [Browse all →](#)

Natural Sciences

Medical and Health  
Sciences

Social Sciences

Engineering and  
Technology

Agricultural and  
Veterinary Sciences

Humanities and the Arts

RECENT ARTICLES | [Browse all →](#)

11 MARCH 2022

Synthesis and characterization of silica xerogel from corn husk waste as cationic ...

11 MARCH 2022

Interplay among positive and negative symptoms, neurocognition, social cogniti...

11 MARCH 2022

Cut-off points for low skeletal muscle mass in older adults: Colombia versus other po...

[Home](#) » [Advisory Board](#)

## Advisory Board

---

The Advisory Board of F1000Research comprises a large group of leading experts across biology and medicine. They do not act as Editors in the traditional sense (they do not handle manuscripts or make decisions to accept or reject a paper), but they provide strategic input on the direction we should take with F1000Research. They occasionally advise us on issues arising with specific articles, and many members of the board also review for us.



### B

- Ian Beales
- Nelson Bennett
- Avri Ben-Ze'ev
- Benedikt Berninger
- Eric Beyer
- Azra Bihorac
- Daniel Bikle
- Kevin J Black
- Chellakkan Selvanesan Blesson
- Erin Alello Bowles
- Bruce Brew

### C

- David Catcheside
- Andrew Chalmers
- Tak Mao Chan
- Karen Chapman
- Declan Chard
- Walter Chazin
- Jonathan Chernoff
- Cheng-Ming Chiang
- Ryan Chisholm
- Wei-Sheng Chong
- Sandra Citi
- Vitaly Citovsky
- Tim Clark
- James Coker
- Giuseppe Colloca
- William Colmers
- Jason Crawford
- David Criddle

# F1000Research

Eric Dannaoul  
Vinicio de Jesus Perez  
Sharon DeMorrow  
Gonzalo G de Polavieja  
Saskia de Wildt  
Harriet de Wit  
Eleftherios P Diamandis  
Phedias Diamandis  
Betty Diamond  
J Alan Diehl  
Petya Dimitrova  
Annette Dolphin  
Lucy Donaldson  
Sylvie Doublie  
Paschalis-Thoma Doulias  
Crislyn D'Souza-Schorey  
James Duffin  
Janice Du Mont

**E** Sharyn Endow  
Markus Engstler  
Sam Enna  
Erim Erdem

**F** Alastair Ferguson  
Gerardo Ferrara  
Richard Festenstein  
Thomas Finger  
Céline Fiset  
Heike Fölsch  
Steven Frank  
Bernd Fritsch

**G** Gus Gazzard  
Jozef Gécz  
Robert Gerlai  
Ivan Gerling  
Carole Goble  
Richard Gomer  
Andrew Goryachev  
John Greenspan  
Guy Griebel  
W Sue Griffin  
Elizabeth Grove  
Jaime Grutzendler  
Wei Guo

# F1000Research

Stephen Hoffman  
Stephen Holgate  
Thorsten Hoppe  
Wolfgang Huber  
Arthur Hurwitz

I  
Radu Iliescu  
Robert Insall  
Harry Ischiropoulos

J  
Jan Jakobsson  
Guilhem Janbon  
Michael Joannidis  
Norman Johnson  
Etienne Joly

K  
Dieter Kabelitz  
Wael Kafienah  
Chaya Kalcheim  
Lynn Kamerlin  
Mikhail Kazachkov  
Johannes S Kern  
Jean-Pierre Kinet  
Edward Kipreos  
Fenella Kirkham  
Gordon Klein  
Alisa Koch  
Amos Korczyn  
Benoit Kornmann  
Jan Kucera  
Anuj Kumar  
Saravana Kumar

L  
Eileen Lafer  
Hans Lassmann  
Mario Lebediker  
John Lee  
Laurel Lenz  
Simon Levin  
Stefan Linder  
Ton Lisman  
Creighton M Litton  
Hartmut Lode  
Theresa Lu  
Robyn Lucas  
Ben Lugtenberg  
Paul Lyons

Jocelyn McDonald

Robert McPeck

Anthony Means

Julien Mendlewicz

Arthur Mercurio

Ralph Mislberger

Ali Mobasherl

David Moher

Randall Moon

Carlos Morel

Dimitrios Morikis

Nicola Mulder

**N** | Corey Nislow

**O** | Chiadi Onyike

**P** | Leonid Padyukov  
Eleftherios Paschalis  
Graham Pawelec  
Ming Pei  
Giampaolo Perna  
Stephen Pinfield  
Michel Pohl  
Simon Portsmouth  
David Potter  
Chaim Putterman

**R** | Adam Ratner  
Ana Recober  
Victor Reus  
José Luis Riechmann  
Karin Romisch  
Vincent Rotello  
Barry Rouse  
Gloria Rudenko  
James Russell

Werner Scheithauer  
Tamar Schlick  
Thomas Schnider  
Alfons Schnitzler  
Irene Schulz  
Michael Sendtner  
Andrew D Sharrocks  
Nilabh Shastri  
Kazim Sheikh  
Andrew Shennan  
Xiao Shifu  
Chiara Simonelli  
Helmy Siragy  
Cassian Sitaru  
Richard Smith  
H Peter Soyer  
Pamela Stanley  
Christoph Stein  
Carly Stevens  
Charles Stevens  
Bruno Stieger

T  
Paul-Peter Tak  
Paul Terry  
Igor Tetko  
Jacques Thibodeau  
Jakub Tolar  
Peter Tonellato  
Francis Tsai  
Takeshi Tsubata  
Tom Tullius  
Burkhard Tümmler

U  
Hisashi Umemori  
Shiro Urayama  
Vladimir Uversky

V  
Hans van Beek  
Hans van Bokhoven  
Martin van den Berg  
Peter Van Endert  
Dirk van Helden  
Chandra Verma  
Jan Vermorken  
David Voehringer

# F1000Research

Stephen Waxman

Alan Wein

Tom Woodcock

Long-Jun Wu

Jeremy C Wyatt

Kevan Wylie

**X** | Yongblao Xue

**Y** | Michael B Yaffe

Kenneth Yamada

Helen Yap

Dorothy Yuan

**Z** | Yunde Zhao

Deyou Zheng

Guy Zimmerman

Christos Zouboulis

# F1000Research

An innovative open access publishing platform offering rapid publication and open peer review, whilst supporting data deposition and sharing.

[BROWSE](#)

[GATEWAYS](#)

[COLLECTIONS](#)

[HOW IT WORKS](#)

[BLOG](#)

[CONTACT](#)

[RSS](#)



[Home](#) » [Browse Articles](#)

# You searched for Seroconversion rates among different designs of COVID-19 vaccines: a network meta-analysis of randomized controlled trials

[ARTICLES](#) [FACULTY REVIEWS](#) [DOCUMENTS](#) [POSTERS](#) [SLIDES](#)

4

FILTERS ▾

1:1 of 1 ARTICLE

SYSTEMATIC REVIEW  metrics

AWAITING PEER REVIEW

## Seroconversion rates among different designs of COVID-19 vaccines: a network meta-analysis of randomized controlled trials [version 1; peer review: awaiting peer review]

Gatot Soegiarto, Jonny Fajar, Laksmi Wulandari, Muhammad Anshory, Muhammad Ilmawan, Anisa Asmiragani, Himma Illiyana, Azaria Adam, Sutini Lamadi, Umi Sa'adah, Tubagus Yuantoko, Esi Nanda, Farida Rachmawati, Nabila Rahmadani, Randy Talilah, Madyline Katipana, Sharon Susanto, Maria Hindom, Ufi Anjasari, Nur Hidayah, Nanda Fadilla, Vanela Lekatompessy, Uzi Phoenna, Fredo Tamara, Dessy Kartini, Aditya Mahendra, Andi Permana, Erwin Pasaribu, Kuldeep Dhama, Harapan Harapan

 PEER REVIEWERS *Invited*

PUBLISHED 10 Mar 2022

unclear. OBJECTIVE: To assess the seroconversion rates among different COVID-19 vaccines using ... the seroconversion rates among different COVID-19 vaccine designs. Eligibility criteria Articles were included in our ...





## SYSTEMATIC REVIEW

# Seroconversion rates among different designs of COVID-19 vaccines: a network meta-analysis of randomized controlled trials [version 1; peer review: awaiting peer review]

Gatot Soegiarto<sup>1</sup>, Jonny Fajar<sup>2</sup>, Laksmi Wulandari<sup>3</sup>, Muhammad Anshory<sup>4</sup>, Muhammad Ilmawan<sup>5</sup>, Anisa Asmiragani<sup>5</sup>, Himma Illiyana<sup>6</sup>, Azaria Adam<sup>7</sup>, Sutini Lamadi<sup>5</sup>, Umi Sa'adah<sup>8</sup>, Tubagus Yuantoko<sup>6</sup>, Esi Nanda<sup>6</sup>, Farida Rachmawati<sup>8</sup>, Nabila Rahmadani<sup>5</sup>, Randy Talilah<sup>9</sup>, Madyline Katipana<sup>10</sup>, Sharon Susanto<sup>11</sup>, Maria Hindom<sup>12</sup>, Ufi Anjasari<sup>11</sup>, Nur Hidayah<sup>13</sup>, Nanda Fadilla<sup>14</sup>, Vanela Lekatompessy<sup>8</sup>, Uzi Phoenna<sup>15</sup>, Fredo Tamara<sup>2</sup>, Dessy Kartini<sup>5</sup>, Aditya Mahendra<sup>16</sup>, Andi Permana<sup>16</sup>, Erwin Pasaribu<sup>5</sup>, Kuldeep Dhama<sup>16</sup>, Harapan Harapan<sup>17</sup>

<sup>1</sup>Division of Allergy & Immunology, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, 60286, Indonesia

<sup>2</sup>Brawijaya Internal Medicine Research Center, Department of Internal Medicine, Faculty of Medicine, Universitas Brawijaya, Malang, East Java, 65145, Indonesia

<sup>3</sup>Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, 60286, Indonesia

<sup>4</sup>Division of Allergy & Immunology, Department of Internal Medicine, Faculty of Medicine, Universitas Brawijaya, Malang, East Java, 65145, Indonesia

<sup>5</sup>Faculty of Medicine, Universitas Brawijaya, Malang, East Java, 65145, Indonesia

<sup>6</sup>Faculty of Public Health, Universitas Indonesia, Jakarta, Jakarta, 10430, Indonesia

<sup>7</sup>Department of Psychiatry, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, 60286, Indonesia

<sup>8</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Malang, East Java, 65145, Indonesia

<sup>9</sup>Faculty of Nursing, Universitas Indonesia, Jakarta, Jakarta, 10430, Indonesia

<sup>10</sup>Department of Anesthesiology and Intensive Care, Faculty of Medicine, Universitas Udayana, Denpasar, Bali, 80361, Indonesia

<sup>11</sup>School of Pharmacy, Institut Teknologi Bandung, Bandung, East Java, 40132, Indonesia

<sup>12</sup>Department of Child Health, Faculty of Medicine, Public Health and Nursing, University of Gadjah Mada, Yogyakarta, Yogyakarta, 55281, Indonesia

<sup>13</sup>Faculty of Animal Science, Universitas Brawijaya, Malang, East Java, 65145, Indonesia

<sup>14</sup>Faculty of Mathematics and Natural Sciences, Universitas Negeri Malang, Malang, East Java, 65145, Indonesia

<sup>15</sup>Department of Clinical Pathology, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Central Java, 57126, Indonesia

<sup>16</sup>Division of Pathology, Indian Veterinary Research Institute, Izatnagar, Uttar Pradesh, 243122, India

<sup>17</sup>Medical Research Unit, School of Medicine, Universitas Syiah Kuala, Banda Aceh, Aceh, 23111, Indonesia

**v1** First published: 10 Mar 2022, 11:299  
<https://doi.org/10.12688/f1000research.110281.1>

Latest published: 10 Mar 2022, 11:299  
<https://doi.org/10.12688/f1000research.110281.1>

## Abstract

BACKGROUND: The COVID-19 vaccination program, which uses

## Open Peer Review

**Approval Status** AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

various types of vaccines, has been applied since the beginning of 2021. However, the efficacy in the context of seroconversion rate remains unclear.

**OBJECTIVE:** To assess the seroconversion rates among different COVID-19 vaccines using a network meta-analysis approach.

**METHODS:** A network meta-analysis of randomized controlled trials (RCTs) was conducted during the study period. Data of interest, such as seroconversion rate and the type of COVID-19 vaccine, were extracted from each study. The analysis was performed using single-arm analysis by calculating the cumulative seroconversion rate. A network meta-analysis was conducted using the Bayesian method.

**RESULTS:** A total of 31 RCTs were included in our analysis. Our pooled calculation revealed that the seroconversion rates of inactivated messenger ribonucleic acid (mRNA), protein subunit, and vector COVID-19 vaccines during the follow-up periods were 93.2%, 93.9%, 65.3%, and 54.7%, respectively, at  $\leq 15$  days; 96.0%, 94.8%, 91.2%, and 89.7%, respectively, between days 16–30; and 98.5%, 98.6%, 98.5%, and 96.2%, respectively, between days 31–60. The indirect comparison revealed that in the follow-up periods of  $\leq 15$  and 16–30 days, the inactivated and mRNA COVID-19 vaccines had superior seroconversion rates compared with those of the protein subunit and vector vaccines. In the follow-up period of 31–60 days, the highest seroconversion rates were found in the inactivated, mRNA, and protein subunit COVID-19 vaccines.

**CONCLUSION:** This study provides valuable information regarding the comparison of seroconversion rates of COVID-19 vaccines.

#### **Keywords**

COVID-19; vaccine; seroconversion; efficacy; immunization.

**Corresponding authors:** Gatot Soegiarto ([gatot\\_soegiarto@fk.unair.ac.id](mailto:gatot_soegiarto@fk.unair.ac.id)), Jonny Fajar ([gembyok@gmail.com](mailto:gembyok@gmail.com))

**Author roles:** **Soegiarto G:** Conceptualization, Formal Analysis, Investigation, Methodology, Resources, Supervision, Validation, Writing – Review & Editing; **Fajar J:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Resources, Software, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Wulandari L:** Conceptualization, Formal Analysis, Investigation, Methodology, Supervision; **Anshory M:** Conceptualization, Formal Analysis, Investigation, Methodology, Supervision; **Ilmawan M:** Formal Analysis, Investigation, Methodology, Software; **Asmiragani A:** Formal Analysis, Investigation, Methodology, Software; **Iliyana H:** Project Administration, Resources; **Adam A:** Data Curation, Formal Analysis, Software; **Lamadi S:** Data Curation, Project Administration, Resources; **Sa'adah U:** Data Curation, Project Administration, Resources; **Yuantoko T:** Data Curation, Project Administration, Resources; **Nanda E:** Data Curation, Project Administration, Resources; **Rachmawati F:** Data Curation, Project Administration, Resources; **Rahmadani N:** Data Curation, Project Administration, Resources; **Talilah R:** Data Curation, Project Administration, Resources; **Katipana M:** Data Curation, Project Administration, Resources; **Susanto S:** Data Curation, Project Administration, Resources; **Hindom M:** Data Curation, Project Administration, Resources; **Anjasari U:** Data Curation, Project Administration, Resources; **Hidayah N:** Data Curation, Project Administration, Resources; **Fadilla N:** Data Curation, Project Administration, Resources; **Lekatompessy V:** Data Curation, Project Administration, Resources; **Phoenna U:** Data Curation, Project Administration, Software; **Tamara F:** Data Curation, Project Administration, Resources; **Kartini D:** Data Curation, Project Administration, Resources; **Mahendra A:** Data Curation, Project Administration, Resources; **Permana A:** Data Curation, Project Administration, Resources; **Pasaribu E:** Data Curation, Project Administration, Resources; **Dhama K:** Data Curation, Project Administration, Resources; **Harapan H:** Data Curation, Project Administration, Resources

**Competing interests:** No competing interests were disclosed.

**Grant information:** The author(s) declared that no grants were involved in supporting this work.

**Copyright:** © 2022 Soegiarto G *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Soegiarto G, Fajar J, Wulandari L *et al.* Seroconversion rates among different designs of COVID-19 vaccines: a network meta-analysis of randomized controlled trials [version 1; peer review: awaiting peer review] F1000Research 2022, 11:299 <https://doi.org/10.12688/f1000research.110281.1>

**First published:** 10 Mar 2022, 11:299 <https://doi.org/10.12688/f1000research.110281.1>

## Introduction

COVID-19 remains a global challenge.<sup>1</sup> While the incidence of COVID-19 was reported to have decreased in recent months, recent reports have suggested that there has been a rising trend in COVID-19 incidence.<sup>2</sup> This fluctuating incidence might be affected by the variants of concern,<sup>3</sup> in which COVID-19 management also remained challenging,<sup>4</sup> and the management of new variants of concern might differ from that of previous variants.<sup>5</sup> The guidelines for COVID-19 management have been periodically published and updated.<sup>6</sup> However, the efficacy of each treatment was inconclusive, particularly in the case of severe or critical illness.<sup>7-9</sup> Therefore, the proper treatment of COVID-19 remains under investigation. While the potential of new drugs has been explored,<sup>10</sup> the vaccination program appears to have the potential to end this pandemic.<sup>11</sup>

The COVID-19 vaccination program was introduced in early 2021 and implemented worldwide.<sup>12</sup> This vaccination program was initially targeted to health workers, a population with high risk of infection, and continued to the public.<sup>12,13</sup> To date, a wide variety of COVID-19 vaccines have been available, such as inactivated, messenger ribonucleic acid (mRNA), protein subunit, and vector vaccines.<sup>14</sup> However, efficacy differs between vaccines, and the results from each study have varied.<sup>15</sup> In this circumstance, conflict between pharmaceutical companies might occur. Therefore, the question of which vaccine has the best efficacy remains. In the context of vaccination, seroconversion was used to assess the early response of neutralizing antibody production.<sup>16</sup> However, the report of seroconversion of COVID-19 vaccines varied in each study, particularly in the special cases with comorbidity.<sup>17-19</sup> Moreover, to date, no study has directly compared the efficacy of COVID-19 vaccines. Therefore, the present study aimed to assess the indirect comparison of seroconversion rates among different COVID-19 vaccines using a network meta-analysis approach. Our present study provides preliminary evidence regarding potential COVID-19 vaccines.

## Methods

### Study design

A meta-analysis, following the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) protocols,<sup>20</sup> was conducted to compare the seroconversion rates of different COVID-19 vaccine designs. To formulate a comprehensive comparison, the relevant articles were collected from PubMed, Embase, and Scopus, and the information of interest was extracted to compare the seroconversion rates among different COVID-19 vaccine designs.

### Eligibility criteria

Articles were included in our analysis if they met the following inclusion criteria: (1) assessed the seroconversion rate of a COVID-19 vaccine; and (2) provided standardized data to determine the seroconversion rate of a COVID-19 vaccine. The following articles were excluded: reviews, commentaries, letters to the editor, non-randomized controlled trials (RCTs), and double publications.

### Search strategy and data extraction

As of January 10, 2022, we searched for potential articles in PubMed, Scopus, and Web of Science. We determined potential COVID-19 vaccine designs to be involved in our study prior to searching the primary outcomes (seroconversion rate). We used the keywords adapted from medical subject headings: "COVID-19 vaccine" or "inactivated COVID-19 vaccine" or "mRNA COVID-19 vaccine" or "protein subunit COVID-19 vaccine" or "vector COVID-19 vaccine" and "efficacy" or "seroconversion". The search was limited to RCTs and articles published in English. If a double publication was found, only articles with a larger sample size were included. We also browsed the reference list of relevant systematic reviews to obtain additional references. Subsequently, the following information of interest from the potential articles was extracted by two independent investigators: (1) first author name, (2) publication year, (3) study design, (4) age of patients, (5) sample size, (6) design of COVID-19 vaccine, (7) trade name of COVID-19 vaccine, (8) dosage of COVID-19 vaccine, (9) modified JADAD scale, and (10) seroconversion rate.

### Assessment of the methodological quality

Prior to inclusion in our analysis, articles were appraised for quality using the modified JADAD scale. The scores ranged from 0 to 7. Scores of 5-7, 3-4, and 0-2 indicated high-, moderate-, and low-quality papers, respectively.<sup>21</sup> Low-quality articles were excluded from the analysis. Using a pilot form, quality assessment was performed by two independent authors (JKF & MI). Discrepancies between the two authors were resolved by discussion.

### Outcome measure

The primary outcome was the seroconversion rate of the COVID-19 vaccine, defined as the level of geometric mean titer (GMT) of neutralizing antibodies of greater than or equal to four-fold from the baseline. The predictors were different COVID-19 vaccine designs. To identify the potential COVID-19 vaccine designs, an initial evaluation of the available data in PubMed, Scopus, and Web of Science was performed. Of those, inactivated, mRNA, protein subunit, and vector COVID-19 vaccines were available for the analysis.


## Statistical analysis

Before analyzing the data, the potential publication bias and heterogeneity across the studies was assessed. Publication bias was assessed using an Egger test, and a p-value < 0.05 indicated a publication bias. Heterogeneity among the studies was evaluated using the Q test. A p-value < 0.10 suggested that heterogeneity existed among studies, and that a random effect model should be applied for the data analysis. Otherwise, a fixed-effect model should be used. The cumulative seroconversion rate of COVID-19 vaccines was determined using a single arm model meta-analysis by calculating the pooled seroconversion events from the total sample size. The effect size was presented using the seroconversion percentage and 95% confidence interval (CI). The data were analyzed using R package software (R package, MA, US, RRID:SCR\_001905). The pooled seroconversion rates are summarized in a forest plot. The seroconversion rates between different designs of COVID-19 vaccines were compared by calculating the effect size of each COVID-19 vaccine design. The highest seroconversion rate was considered the highest efficacy, and the Confidence in Network Meta-Analysis software version 1.9.1 (Bern, Switzerland, RRID:SCR\_016488) was used to outline the network diagram of comparison among COVID-19 vaccines.

## Results

### Study selection

We collected 8,070 potential papers from the databases. Of these, 23 duplication papers were found, and 7,966 papers had irrelevant topics; therefore, those papers were excluded. Subsequently, 81 papers were included for further full-text reviews. Among these, 19 reviews and 32 papers with insufficient data and were excluded. Finally, a total of 31 papers were analyzed to compare the seroconversion rates among COVID-19 vaccines.<sup>22-52</sup> The process of article selection in our study is presented in [Figure 1](#), and the characteristics of the papers included in our analysis are summarized in [Table 1](#).



**Figure 1.** A PRISMA flow chart of study selection.

**Table 1. Baseline characteristics of study included in our meta-analysis.**

Author	Age ( $\pm$ SD)	Sample size	Type of vaccine	Merk vaccine	Dose of vaccine	Jadad Modified Scale
Al Kaabi et al 2021	36.2 (9.2)	40832	Inactivated	WIV04, HB02, alum-only	WIV04 = 5 $\mu$ g/dose; HB02 = 4 $\mu$ g/dose; alum-only = 0.5mg	6
Ali et al 2021	14.2 (1.6)	3732	mRNA	mRNA-1273	200 $\mu$ g	5
Baden et al 2021	51.3 (24.7)	30420	mRNA	mRNA-1273	200 $\mu$ g	4.5
Che et al 2020	41.4 (10.84)	750	Inactivated	Inactivated SARS-CoV-2 vaccine	50 EU, 100 EU, 150 EU	6
Ella et al 2021	32.5 (24)	375	Inactivated	BBV152	Algel-IMDG = 3 $\mu$ g; Algel-IMDG = 6 $\mu$ g with; Algel = 6 $\mu$ g	7
Fadiyana et al 2021 (1)	35.6 (11.3)	1620	Inactivated	Inactivated SARS-CoV-2 vaccine	3 $\mu$ g	6.5
Falsey et al 2021	64.8 (21.4)	32379	Vector vaccine	AZD1222 (ChAdOx1 nCoV-19)	$5 \times 10^{10}$ viral particles	7
Feng et al 2021	N/A	809	Inactivated	Inactivated SARS-CoV-2 vaccine	3 $\mu$ g	5.5
Formica et al 2021	38.9 (12.40)	1288	Protein subunit vaccine	NVXCoV2373	5 $\mu$ g, 25 $\mu$ g	6.5
Han et al 2021	8.4 (4.2)	550	Inactivated	CoronaVac	1.5 $\mu$ g, 3 $\mu$ g	6
Jackson et al 2020	36.7 (7.9)	45	mRNA	mRNA-1273	25 $\mu$ g, 100 $\mu$ g, 250 $\mu$ g	4.5
Kremsner et al 2020	38.6 (12.9)	248	mRNA	CvCoV	2-12 $\mu$ g	5.5
Li et al 2021	37.9 (9.6)	144	mRNA	BNT162b1	10 $\mu$ g, 30 $\mu$ g	7
Liu et al 2021	58.2 (4.81)	830	mRNA	ChAd, BNT	ChAd = 0.5 mL, BNT = 0.3 mL	6.5
Melo-González et al 2021	N/A	94	Inactivated	CoronaVac	3 $\mu$ g	5.5
Pan et al 2021	38.0 (9.5)	560	Inactivated	KCONVAC	5 $\mu$ g, 10 $\mu$ g	7
Polack et al 2020	N/A	43448	mRNA	BNT162b2	30 $\mu$ g	5.5
Pu et al 2021	18-59	96	Inactivated	NA	NA	5.5


Table 1. Continued

Author	Age ( $\pm$ SD)	Sample size	Type of vaccine	Merk vaccine	Dose of vaccine	Jadad Modified Scale
Richmond et al 2021	37.6 (11.9)	148	Protein subunit vaccine	SCB-2019	3 $\mu$ g, 9 $\mu$ g, 30 $\mu$ g	7
Sedoff et al 2021	36.1 (10.1)	805	Vector vaccine	Ad26_COV2.5	$5 \times 10^{10}$ , $1 \times 10^{11}$ viral particles	5.5
Shu et al 2021	43.9 (11.3)	880	Protein subunit vaccine	V-01	10 $\mu$ g, 25 $\mu$ g	7
Tanriover et al 2021	N/A	10218	Inactivated	CoronaVac	3 $\mu$ g	7
Thomas et al 2021	N/A	44060	mRNA	BNT162b2	30 $\mu$ g	5.5
Wu et al 2021	65.6 (4.3)	395	Inactivated	CoronaVac	1.5 $\mu$ g, 3 $\mu$ g, 6 $\mu$ g	7
Xia et al 2020 (a)	36.0 (8.5)	320	Inactivated	Inactivated SARS-CoV-2 vaccine	2.5 $\mu$ g, 5 $\mu$ g, 10 $\mu$ g	6
Xia et al 2020 (b)	42.7 (8.1)	640	Inactivated	BBIBP-CoV	2 $\mu$ g, 4 $\mu$ g, 8 $\mu$ g	7
Yang et al 2020	32.6 (9.41)	950	Protein subunit vaccine	ZF2001	25 $\mu$ g, 50 $\mu$ g	6
Zeng et al 2021	45.2 (9.1)	540	Inactivated	CoronaVac	1.5 $\mu$ g, 3 $\mu$ g, 6 $\mu$ g	7
Zhang et al 2020	41.8 (9.4)	744	Inactivated	CoronaVac	3 $\mu$ g, 6 $\mu$ g	6
Zhang et al 2021	40.0 (9.2)	180	Protein subunit vaccine	V-01	10 $\mu$ g, 25 $\mu$ g, 50 $\mu$ g	7
Zhu et al 2020	37.2 (10.7)	108	Vector vaccine	Ad5 vectored COVID-19 vaccine	$5 \times 10^{10}$ , $1 \times 10^{11}$ , $1.5 \times 10^{11}$ viral particles	4

Note: NA, not available; SD, standard deviation; mRNA, messenger ribonucleic acid.


### The seroconversion rates among different COVID-19 vaccines

In the follow-up period of  $\leq 15$  days, 24 papers assessing the seroconversion rate of COVID-19 vaccines were collected. In total, the seroconversion rate was 91.1% (Figure 2A). The seroconversion rates of inactivated (Figure 2B), mRNA (Figure 2C), mRNA




**Figure 2.** A forest plot of the seroconversion rate among different type of COVID-19 vaccines at day  $\leq 15$ . A). All vaccine types; B). Inactivated vaccine; C). mRNA vaccine; D). Protein subunit vaccine; and E). Vector vaccine.





**Figure 3. A forest plot of the seroconversion rate among different type of COVID-19 vaccines at day 16-30. A). All vaccine types; B). Inactivated vaccine; C). mRNA vaccine; D). Protein subunit vaccine; and E). Vector vaccine.**


(Figure 2C), protein subunit (Figure 2D), and vector COVID-19 vaccines (Figure 2E) were 93.2%, 93.9%, 65.3%, and 54.7%, respectively. Subsequently, in the follow-up period of 16–30 days, the cumulative seroconversion rate of COVID-19 vaccines was 94.1% (Figure 3A). The seroconversion rates of inactivated (Figure 3B), mRNA (Figure 3C), protein subunit (Figure 3D), and vector COVID-19 vaccines (Figure 3E) were 96.0%, 94.8%, 91.2%, and 89.7%, respectively. Alternatively, in the follow-up period of 31–60 days, the pooled seroconversion rate of COVID-19 vaccines was 98.0% (Figure 4A). The seroconversion rate of inactivated (Figure 4B), mRNA (Figure 4C), protein subunit (Figure 4D), and vector COVID-19 vaccines (Figure 4E) were 98.5%, 98.6%, 98.5%, and 96.2%, respectively.



**Figure 4.** A forest plot of the seroconversion rate among different type of COVID-19 vaccines at day 31-60. A). All vaccine types; B). Inactivated vaccine; C). mRNA vaccine; D). Protein subunit vaccine; and E). Vector vaccine.

**The indirect comparison of seroconversion rates among different COVID-19 vaccines**

In the follow-up period of  $\leq 15$  days (Figure 5A), the seroconversion rates of inactivated and mRNA COVID-19 vaccines were superior to those of the protein subunit and vector COVID-19 vaccines. In the follow-up period of 16–30 days (Figure 5B), the seroconversion rates of inactivated and mRNA COVID-19 vaccines were significantly higher than those of protein subunit and vector COVID-19 vaccines. In the follow-up period of 31–60 days (Figure 5C), the seroconversion rate of the vector COVID-19 vaccine was inferior to that of the inactivated, mRNA, and protein subunit COVID-19 vaccines. The network diagrams of seroconversion rates among different COVID-19 vaccine designs are presented in Figure 5D, 5E, and 5F for the follow-up periods of  $\leq 15$ , 16–30, and 31–60 days, respectively. The summary of indirect comparison of seroconversion rate among COVID-19 vaccines is outlined in Table 2.



**Figure 5. The indirect comparison between different types of COVID-19 vaccines.** A). Follow up period at day  $\leq 15$ ; B). Follow up period at day 16-30; C). Follow up period at day 31-60; D). The networking among studies at day  $\leq 15$ ; E). The networking among studies at day 16-30; and F). The networking among studies at day 31-60.

**Table 2.** The summary of analysis on the indirect comparison of seroconversion rate between different types of COVID-19 vaccines.

Indirect comparison	NS	Sample size	OR	95%CI	p
<b>Follow up at day ≤15</b>					
Inactivated vs. mRNA	16 vs. 5	15464 vs. 53454	4.06	3.58 – 4.60	<0.0001
Inactivated vs. protein subunit	16 vs. 1	15464 vs. 150	29.53	20.66 – 42.19	<0.0001
Inactivated vs. vector	16 vs. 2	15464 vs. 21623	30.95	27.37 – 34.99	<0.0001
mRNA vs. protein subunit	5 vs. 1	53454 vs. 150	7.27	5.19 – 10.20	<0.0001
mRNA vs. vector	5 vs. 2	53454 vs. 21623	7.63	7.30 – 7.97	<0.0001
Protein subunit vs. vector	1 vs. 2	150 vs. 21623	1.05	0.75 – 1.47	0.7850
<b>Follow up at day 16-30</b>					
Inactivated vs. mRNA	13 vs. 4	990 vs. 159	0.97	0.37 – 2.53	0.9540
Inactivated vs. protein subunit	13 vs. 6	990 vs. 453	1.97	1.17 – 3.32	0.0100
Inactivated vs. vector	13 vs. 2	990 vs. 194	2.51	1.33 – 4.73	0.0040
mRNA vs. protein subunit	4 vs. 6	159 vs. 453	2.03	0.77 – 5.35	0.1520
mRNA vs. vector	4 vs. 2	159 vs. 194	2.58	0.92 – 7.26	0.0730
Protein subunit vs. vector	6 vs. 2	453 vs. 194	1.27	0.66 – 2.44	0.4690
<b>Follow up at day 31-60</b>					
Inactivated vs. mRNA	1 vs. 5	32 vs. 499	0.59	0.03 – 11.20	0.7260
Inactivated vs. protein subunit	1 vs. 6	32 vs. 448	0.36	0.02 – 7.74	0.5170
Inactivated vs. vector	1 vs. 1	32 vs. 158	2.77	0.15 – 50.42	0.4910
mRNA vs. protein subunit	5 vs. 6	499 vs. 448	0.56	0.10 – 3.04	0.4980
mRNA vs. vector	5 vs. 1	499 vs. 158	4.89	1.36 – 17.54	0.0150
Protein subunit vs. vector	6 vs. 1	448 vs. 158	8.80	1.76 – 44.08	0.0080

Note: NS, number of studies; OR, odd ratio; CI, confidence interval; mRNA, messenger ribonucleic acid.

### Source of heterogeneity

In the follow-up period of ≤ 15 days, evidence of heterogeneity ( $p$  heterogeneity > 0.05) was observed in the models of analyses for all COVID-19 vaccines, inactivated COVID-19 vaccines, mRNA COVID-19 vaccines, and vector COVID-19 vaccines. Therefore, a random effect model was used. In the follow-up period of 16–30, evidence of heterogeneity was observed in all models of analyses; therefore, a random effect model was used. In the follow-up period of 31–60 days, all analyses were performed using a fixed effect model, because no evidence of heterogeneity was found (**Supplementary files**).<sup>20</sup>

### Discussion

Our study reported on the seroconversion rate among different COVID-19 vaccines in the follow-up periods of ≤ 15, 16–30, and 31–60 days. We revealed that, in the follow-up period of ≤ 15 days, the highest seroconversion rate was that of the inactivated vaccine. In the follow-up period of 16–30 days, the highest seroconversion rates were those of the inactivated and mRNA vaccines. Conversely, in the follow-up period of 31–60 days, the highest seroconversion rates were those of the inactivated, mRNA, and protein subunit vaccines. To date, our study is the first to report on the seroconversion rates among different COVID-19 vaccines; therefore, comparisons among meta-analyses were not discussed. However, similar meta-analyses have been performed to assess the efficacy of COVID-19 vaccines. In total, nine meta-analysis studies have been conducted.<sup>53–61</sup> While the majority of these studies focused on side effects of the vaccine, they also reported the efficacy of COVID-19 vaccines by assessing the reduced risk of infection, mortality, hospitalization, and admission to the ICU. The findings revealed that the mRNA vaccine had the highest efficacy in preventing COVID-19 infection.<sup>59</sup> Those previous meta-analyses support our findings that, besides the mRNA vaccine, the inactivated vaccine had a higher seroconversion rate than that of the protein subunit and vector COVID-19 vaccines.

The theory underlying the comparison of the efficacy of COVID-19 vaccines remains debatable. It has been widely proposed that various vaccines have various immunogens and may be engulfed, processed, and presented by antigen-presenting cells (APCs) along with MCH antigens to CD4+ T cells. Resultingly, cytokine synthesis may occur and may

activate humoral and cellular responses, including antibody production, CD8+ T cell activation, and macrophage stimulation. Subsequently, B lymphocytes may differentiate into plasma cells and produce specific antibodies to protect against the infection.<sup>62,63</sup> In the inactivated vaccine, the immunogenic property is a killed or modified virus containing the whole pathogen, virus fragments, or virus epitope.<sup>64</sup> In the mRNA vaccine, the mRNA will be taken up by APC and translated into protein *in situ*. The mRNA encodes the full-length, pre-fusion stabilized spike protein (S) of SARS-CoV-2.<sup>14</sup> In the protein subunit vaccine, the immunogenic property is specific isolated proteins from SARS-CoV2 virus (S glycoprotein), which is responsible for receptor binding to cellular ACE-2.<sup>65-67</sup> This type of vaccine is similar to a vector vaccine, in which the S protein is produced to confer protection against COVID-19.<sup>68</sup> Of those possible mechanisms, the inactivated vaccine may target a wide variety of epitopes,<sup>69</sup> and therefore, this type of vaccine may have a wide protection against COVID-19 variants of concern compared to other types of COVID-19 vaccines. However, vaccine specificity may differ, which requires evaluation by comparing the total GMT levels.

To the best of our knowledge, the present study is the first to report a comparison of seroconversion rates among different COVID-19 vaccine designs. Our study provided valuable evidence that inactivated and mRNA vaccines provided an early seroconversion rate, and the protein subunit vaccine achieved a similar seroconversion rate to that of inactivated and mRNA vaccines in the follow-up period of 31–60 days. The results of our research may be used as a basis for evaluating the development of COVID-19 vaccines in the future.<sup>13</sup> We hope that future vaccine development may take into account the results of seroconversion rates between different vaccine designs, as we reported in our study. Therefore, the expected protective effects of the vaccine could be achieved. However, our present study only assessed the seroconversion rate in a short follow-up period. Further studies assessing a long-term follow-up period are required.

Our current study had several important limitations. First, we did not include potential confounding factors in assessing the efficacy of COVID-19 vaccines, such as comorbidity, nutritional status, and transmission area. Therefore, the probability of the dependent effect remains open to discussion. Second, the different report formats among studies required manual calculation for the precise seroconversion rate information. Therefore, there is the possibility of human error in interpreting seroconversion rates. Third, the vaccine and booster dosages varied among the studies. Therefore, false-positive findings might exist. Fourth, we only focused on the seroconversion rate, in which this evaluation was only effective in assessing the sensitivity of vaccines. Further studies evaluating cumulative GMTs might be required. Fifth, in the current study, vaccine safety was not evaluated. Therefore, further studies assessing safety are needed.

## Conclusion

Our study revealed that the inactivated and mRNA COVID-19 vaccines provided the highest seroconversion rates at early follow-up. The protein subunit COVID-19 vaccine achieved a seroconversion rate similar to that of the inactivated and mRNA vaccines at the follow-up period of 31–60 days. Our study might contribute to better insight into the seroconversion rates of different COVID-19 vaccines.

## Data availability

All data underlying the results are available as part of the article and no additional sources of data are required.

## Reporting guidelines

Figshare. PRISMA checklist. DOI: <https://doi.org/10.6084/m9.figshare.19236714.v2><sup>20</sup>

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

## Competing interests

No competing interests were declared.

## Grant information

This study received no external funding.

## Author contribution

Idea/concept: GS, JKF. Design: GS, JKF, LW. Control/supervision: GS, LW, MA, KD, III. Data collection/processing: MI, AA, III, AAA, SL, US, TDY, EDN, FR, NR, RT, MVK, SS, MCH, UA, NII, NBF, VCL, UMP, FT, DAK, AIM, AP, EAP. Extraction/Analysis/interpretation: JKF, MI. Literature review: GS, JKF, MI. Writing the article: GS, JKF, MI. Critical review: GS, LW, MA, KD, III. All authors have critically reviewed and approved the final draft and are responsible for the content and similarity index of the manuscript.

## Acknowledgement

We thank Lembaga Pengelola Dana Pendidikan (LPDP) Republic of Indonesia for supporting this project.

## References

- Mutiawati E, Syahrul S, Fahriani M, et al.: **Global prevalence and pathogenesis of headache in COVID-19: A systematic review and meta-analysis.** *F1000Res.* 2020; **9**: 1316.  
[Publisher Full Text](#)
- Kibria HB, Jyoti O, Matin A: **Forecasting the spread of the third wave of COVID-19 pandemic using time series analysis in Bangladesh.** *Inform. Med. Unlocked.* 2022; **28**: 100815.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Thakur V, Bholra S, Thakur P, et al.: **Waves and variants of SARS-CoV-2: understanding the causes and effect of the COVID-19 catastrophe.** *Infection.* 2021.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Krishnan A, Hamilton JP, Alqahtani SA, et al.: **COVID-19: An overview and a clinical update.** *World J. Clin. Cases.* 2021; **9**: 8-23.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cascella M, Rajnik M, Aleem A, et al.: **Features, Evaluation, and Treatment of Coronavirus (COVID-19).** Treasure Island (FL): StatPearls; 2022.
- Agarwal A, Rochwerg B, Lamontagne F, et al.: **A living WHO guideline on drugs for covid-19.** *BMJ.* 2020; **370**: m3379.
- Ozluzen B, Kozan S, Akcan RE, et al.: **Effectiveness of favipiravir in COVID-19: a live systematic review.** *Eur. J. Clin. Microbiol. Infect. Dis.* 2021; **40**: 2575-2583.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wardhani SO, Fajar JK, Wulandari L, et al.: **Association between convalescent plasma and the risk of mortality among patients with COVID-19: a meta-analysis.** *F1000Res.* 2021; **10**: 64.  
[Publisher Full Text](#)
- Wardhani SO, Fajar JK, Soegiarto G, et al.: **The association between therapeutic plasma exchange and the risk of mortality among patients critically ill with COVID-19: a meta-analysis.** *F1000Res.* 2021; **10**: 1280.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Prasansuklab A, Theerasri A, Rangsinth P, et al.: **Anti-COVID-19 drug candidates: A review on potential biological activities of natural products in the management of new coronavirus infection.** *J. Tradit. Complement. Med.* 2021; **11**: 144-157.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- The LM: **COVID-19 vaccines: the pandemic will not end overnight.** *Lancet Microbe.* 2021; **2**: e1.  
[Publisher Full Text](#)
- Yu H, Yang J, Marziano V, et al.: **Can a COVID-19 vaccination program guarantee the return to a pre-pandemic lifestyle?** *Res. Sq.* 2021.
- Fajar JK, Harapan H: **Socioeconomic and attitudinal variables associated with acceptance and willingness to pay towards dengue vaccine: a systematic review.** *Arch. Clin. Infect. Dis.* 2017; **12**: e13914.  
[Publisher Full Text](#)
- Kaur SP, Gupta V: **COVID-19 Vaccine: A comprehensive status report.** *Virus Res.* 2020; **288**: 198114.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tentori K, Passerini A, Timberlake B, et al.: **The misunderstanding of vaccine efficacy.** *Soc. Sci. Med.* 2021; **289**: 114273.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Buttiron Webber T, Provinciani N, Musso M, et al.: **Predictors of poor seroconversion and adverse events to SARS-CoV-2 mRNA BNT162b2 vaccine in cancer patients on active treatment.** *Eur. J. Cancer.* 2021; **159**: 105-112.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Corti C, Antonarelli G, Scotte F, et al.: **Seroconversion rate after vaccination against COVID-19 in patients with cancer: a systematic review.** *Ann. Oncol.* 2022; **33**: 158-168.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Syahrul S, Maliga HA, Ilimawan M, et al.: **Hemorrhagic and ischemic stroke in patients with coronavirus disease 2019: incidence, risk factors, and pathogenesis - a systematic review and meta-analysis.** *F1000Res.* 2021; **10**: 34.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yusuf F, Fahriani M, Mamada SS, et al.: **Global prevalence of prolonged gastrointestinal symptoms in COVID-19 survivors and potential pathogenesis: A systematic review and meta-analysis.** *F1000Res.* 2021; **10**: 301.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fajar JK, Fajar, Jonny (2022): **Supplementary files: Seroconversion rates among different designs of COVID-19 vaccines: a network meta-analysis of randomized controlled trials.** *Figshare Dataset.* 2022; **1**: 1.  
[Publisher Full Text](#)
- Olivo SA, Macedo LG, Gadotti JC, et al.: **Scales to assess the quality of randomized controlled trials: a systematic review.** *Phys. Ther.* 2008; **88**: 156-175.  
[Publisher Full Text](#)
- Al Kaabi N, Zhang Y, Xia S, et al.: **Effect of 2 inactivated SARS-CoV-2 Vaccines on Symptomatic COVID-19 Infection in Adults: A Randomized Clinical Trial.** *JAMA.* 2021; **326**: 35-45.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ali K, Berman G, Zhou H, et al.: **Evaluation of mRNA-1273 SARS-CoV-2 Vaccine in Adolescents.** *N. Engl. J. Med.* 2021; **385**: 2241-2251.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Baden LR, El Sahly HM, Essink B, et al.: **Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine.** *N. Engl. J. Med.* 2021; **384**: 403-416.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Che Y, Liu X, Pu Y, et al.: **Randomized, Double-Blinded, Placebo-Controlled Phase 2 Trial of an Inactivated Severe Acute Respiratory Syndrome Coronavirus 2 Vaccine in Healthy Adults.** *Clin. Infect. Dis.* 2021; **73**: e3949-e3955.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ella R, Reddy S, Jogdand H, et al.: **Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBV152: interim results from a double-blind, randomised, multicentre, phase 2 trial, and 3-month follow-up of a double-blind, randomised phase 1 trial.** *Lancet Infect. Dis.* 2021; **21**: 950-961.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fadlyana E, Rusmil K, Tarigan R, et al.: **A phase III, observer-blind, randomized, placebo-controlled study of the efficacy, safety, and immunogenicity of SARS-CoV-2 inactivated vaccine in healthy adults aged 18-59 years: An interim analysis in Indonesia.** *Vaccine.* 2021; **39**: 6520-6528.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Falsey AR, Sobieszczyk ME, Hirsch L, et al.: **Phase 3 Safety and Efficacy of AZD1222 (ChAdOx1 nCoV-19) Covid-19 Vaccine.** *N. Engl. J. Med.* 2021; **385**: 2348-2360.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Feng Y, Chen J, Yao T, et al.: **Safety and immunogenicity of inactivated SARS-CoV-2 vaccine in high-risk occupational population: a randomized, parallel, controlled clinical trial.** *Infect. Dis. Poverty.* 2021; **10**: 138.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Formica N, Mallory R, Albert G, et al.: **Different dose regimens of a SARS-CoV-2 recombinant spike protein vaccine (NVX-CoV2373) in younger and older adults: A phase 2 randomized placebo-controlled trial.** *PLoS Med.* 2021; **18**: e1003769.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Han B, Song Y, Li C, et al.: **Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial.** *Lancet Infect. Dis.* 2021; **21**: 1645-1653.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jackson LA, Anderson EJ, Rouphael NG, et al.: **An mRNA Vaccine against SARS-CoV-2 - Preliminary Report.** *N. Engl. J. Med.* 2020; **383**: 1920-1931.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kremsner PG, Mann P, Kroidl A, et al.: **Safety and immunogenicity of an mRNA-lipid nanoparticle vaccine candidate against SARS-CoV-2: A phase 1 randomized clinical trial.** *Wien. Klin. Wochenschr.* 2021; **133**: 931-941.  
[PubMed Abstract](#) | [Publisher Full Text](#)

34. Li J, Hui A, Zhang X, et al.: **Safety and immunogenicity of the SARS-CoV-2 BNT162b1 mRNA vaccine in younger and older Chinese adults: a randomized, placebo-controlled, double-blind phase 1 study.** *Nat. Med.* 2021; **27**: 1062–1070. [PubMed Abstract](#) | [Publisher Full Text](#)
35. Liu X, Shaw RH, Stuart ASV, et al.: **Safety and immunogenicity of heterologous versus homologous prime-boost schedules with an adenoviral vectored and mRNA COVID-19 vaccine (Com-COV): a single-blind, randomised, non-inferiority trial.** *Lancet.* 2021; **398**: 856–869. [PubMed Abstract](#) | [Publisher Full Text](#)
36. Melo-Gonzalez F, Soto JA, Gonzalez LA, et al.: **Recognition of Variants of Concern by Antibodies and T Cells Induced by a SARS-CoV-2 Inactivated Vaccine.** *Front. Immunol.* 2021; **12**: 747830. [PubMed Abstract](#) | [Publisher Full Text](#)
37. Pan HX, Liu JK, Huang BY, et al.: **Immunogenicity and safety of a severe acute respiratory syndrome coronavirus 2 inactivated vaccine in healthy adults: randomized, double-blind, and placebo-controlled phase 1 and phase 2 clinical trials.** *Chin. Med. J.* 2021; **134**: 1289–1298. [PubMed Abstract](#) | [Publisher Full Text](#)
38. Polack FP, Thomas SJ, Kitchin N, et al.: **Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine.** *N. Engl. J. Med.* 2020; **383**: 2603–2615. [PubMed Abstract](#) | [Publisher Full Text](#)
39. Pu J, Yu Q, Yin Z, et al.: **The safety and immunogenicity of an inactivated SARS-CoV-2 vaccine in Chinese adults aged 18–59 years: A phase I randomized, double-blinded, controlled trial.** *Vaccine.* 2021; **39**: 2746–2754. [Publisher Full Text](#)
40. Richmond P, Hatchuel L, Dong M, et al.: **Safety and immunogenicity of S-Trimer (SCB-2019), a protein subunit vaccine candidate for COVID-19 in healthy adults: a phase 1, randomised, double-blind, placebo-controlled trial.** *Lancet.* 2021; **397**: 682–694. [PubMed Abstract](#) | [Publisher Full Text](#)
41. Sadoff J, Le Gars M, Shukarev G, et al.: **Interim Results of a Phase 1-2a Trial of Ad26.COV2.S Covid-19 Vaccine.** *N. Engl. J. Med.* 2021; **384**: 1824–1835. [PubMed Abstract](#) | [Publisher Full Text](#)
42. Shu YJ, He JF, Pei RJ, et al.: **Immunogenicity and safety of a recombinant fusion protein vaccine (V-01) against coronavirus disease 2019 in healthy adults: a randomized, double-blind, placebo-controlled, phase II trial.** *Chin. Med. J.* 2021; **134**: 1967–1976. [PubMed Abstract](#) | [Publisher Full Text](#)
43. Tannirver MD, Doganay HL, Akova M, et al.: **Efficacy and safety of an inactivated whole-virion SARS-CoV-2 vaccine (CoronaVac): interim results of a double-blind, randomised, placebo-controlled, phase 3 trial in Turkey.** *Lancet.* 2021; **398**: 213–222. [PubMed Abstract](#) | [Publisher Full Text](#)
44. Thomas SJ, Moreira ED Jr, Kitchin N, et al.: **Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine through 6 Months.** *N. Engl. J. Med.* 2021; **385**: 1761–1773. [PubMed Abstract](#) | [Publisher Full Text](#)
45. Wu Z, Hu Y, Xu M, et al.: **Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy adults aged 60 years and older: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial.** *Lancet Infect. Dis.* 2021; **21**: 803–812. [PubMed Abstract](#) | [Publisher Full Text](#)
46. Xia S, Zhang Y, Wang Y, et al.: **Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial.** *Lancet Infect. Dis.* 2021; **21**: 39–51. [PubMed Abstract](#) | [Publisher Full Text](#)
47. Xia S, Duan K, Zhang Y, et al.: **Effect of an Inactivated Vaccine Against SARS-CoV-2 on Safety and Immunogenicity Outcomes: Interim Analysis of 2 Randomized Clinical Trials.** *JAMA.* 2020; **324**: 951–960. [PubMed Abstract](#) | [Publisher Full Text](#)
48. Yang S, Li Y, Dai L, et al.: **Safety and immunogenicity of a recombinant tandem-repeat dimeric RBD-based protein subunit vaccine (ZF2001) against COVID-19 in adults: two randomised, double-blind, placebo-controlled, phase 1 and 2 trials.** *Lancet Infect. Dis.* 2021; **21**: 1107–1119. [PubMed Abstract](#) | [Publisher Full Text](#)
49. Zeng G, Wu Q, Pan H, et al.: **Immunogenicity and safety of a third dose of CoronaVac, and immune persistence of a two-dose schedule, in healthy adults: interim results from two single-centre, double-blind, randomised, placebo-controlled phase 2 clinical trials.** *Lancet Infect. Dis.* 2021. [PubMed Abstract](#) | [Publisher Full Text](#)
50. Zhang Y, Zeng G, Pan H, et al.: **Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18–59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial.** *Lancet Infect. Dis.* 2021; **21**: 181–192. [PubMed Abstract](#) | [Publisher Full Text](#)
51. Zhang J, Hu Z, He J, et al.: **Safety and immunogenicity of a recombinant interferon-armed RBD dimer vaccine (V-01) for COVID-19 in healthy adults: a randomized, double-blind, placebo-controlled, Phase I trial.** *Emerg. Microbes Infect.* 2021; **10**: 1589–1597. [PubMed Abstract](#) | [Publisher Full Text](#)
52. Zhu FC, Li YH, Guan XL, et al.: **Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial.** *Lancet.* 2020; **395**: 1845–1854. [Publisher Full Text](#)
53. Zheng C, Shao W, Chen X, et al.: **Real-world effectiveness of COVID-19 vaccines: a literature review and meta-analysis.** *Int. J. Infect. Dis.* 2022; **114**: 252–260. [PubMed Abstract](#) | [Publisher Full Text](#)
54. Chen M, Yuan Y, Zhou Y, et al.: **Safety of SARS-CoV-2 vaccines: a systematic review and meta-analysis of randomized controlled trials.** *Infect. Dis. Poverty.* 2021; **10**: 94. [PubMed Abstract](#) | [Publisher Full Text](#)
55. Harder T, Kulper Schiek W, Reda S, et al.: **Effectiveness of COVID-19 vaccines against SARS-CoV-2 infection with the Delta (B.1.617.2) variant: second interim results of a living systematic review and meta-analysis, 1 January to 25 August 2021.** *Euro Surveill.* 2021; **26**. [Publisher Full Text](#)
56. Ling Y, Zhong J, Luo J: **Safety and effectiveness of SARS-CoV-2 vaccines: A systematic review and meta-analysis.** *J. Med. Virol.* 2021; **93**: 6486–6495. [PubMed Abstract](#) | [Publisher Full Text](#)
57. Liu Q, Qin C, Liu M, et al.: **Effectiveness and safety of SARS-CoV-2 vaccine in real-world studies: a systematic review and meta-analysis.** *Infect. Dis. Poverty.* 2021; **10**: 132. [PubMed Abstract](#) | [Publisher Full Text](#)
58. Kalajj AGI, Dirjayanto VJ, Yusuf SM, et al.: **Immunogenicity and safety of adenovirus-based vector vaccines for COVID-19: a systematic review and meta-analysis.** *Med. J. Indones.* 2021; **30**: 264–278. [Publisher Full Text](#)
59. Pormohammad A, Zarei M, Ghorbani S, et al.: **Efficacy and Safety of COVID-19 Vaccines: A Systematic Review and Meta-Analysis of Randomized Clinical Trials.** *Vaccines (Basel).* 2021; **9**. [Publisher Full Text](#)
60. McDonald I, Murray SM, Reynolds CJ, et al.: **Comparative systematic review and meta-analysis of reactogenicity, immunogenicity and efficacy of vaccines against SARS-CoV-2.** *NPJ Vaccines.* 2021; **6**: 74. [PubMed Abstract](#) | [Publisher Full Text](#)
61. Sharif N, Alzahrani KJ, Ahmed SN, et al.: **Efficacy, Immunogenicity and Safety of COVID-19 Vaccines: A Systematic Review and Meta-Analysis.** *Front. Immunol.* 2021; **12**: 714170. [PubMed Abstract](#) | [Publisher Full Text](#)
62. Pulendran B, Ahmed R: **Immunological mechanisms of vaccination.** *Nat. Immunol.* 2011; **12**: 509–517. [PubMed Abstract](#) | [Publisher Full Text](#)
63. Fajar JK, Mahendra AI, Tamara F, et al.: **The association between complete blood count and the risk of coronary heart disease.** *Turkiye Klinikleri J. Med. Sci.* 2019; **39**: 56–64. [Publisher Full Text](#)
64. Kyriakidis NC, Lopez-Cortes A, Gonzalez EV, et al.: **SARS-CoV-2 vaccines strategies: a comprehensive review of phase 3 candidates.** *NPJ Vaccines.* 2021; **6**: 28. [PubMed Abstract](#) | [Publisher Full Text](#)
65. Wu Y, Huang X, Yuan L, et al.: **A recombinant spike protein subunit vaccine confers protective immunity against SARS-CoV-2 infection and transmission in hamsters.** *Sci. Transl. Med.* 2021; **13**. [PubMed Abstract](#) | [Publisher Full Text](#)
66. Fajar JK, Pikir BS, Sidarta EP, et al.: **The Gene Polymorphism of Angiotensin-Converting Enzyme Intron Deletion and Angiotensin-Converting Enzyme G2350A in Patients With Left Ventricular Hypertrophy: A Meta-analysis.** *Indian Heart J.* 2019; **71**: 199–206. [PubMed Abstract](#) | [Publisher Full Text](#)
67. Rohman MS, Fajar JK, Kuncahyo BH, et al.: **Angiotensin-converting enzyme (ACE) I/D and bradykinin B2 receptor T/C genes polymorphism in patients with ACE inhibitors-related cough.** *Egypt. J. Med. Hum. Genet.* 2018; **19**: 307–313. [Publisher Full Text](#)

68. Doerfler W: **Adenoviral Vector DNA- and SARS-CoV-2 mRNA-Based Covid-19 Vaccines: Possible Integration into the Human Genome - Are Adenoviral Genes Expressed in Vector-based Vaccines?** *Virus Res.* 2021; **302**: 198466. [PubMed Abstract](#) | [Publisher Full Text](#)
69. Ella R, Vadrevu KM, Jogdand H, et al.: **Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBV152: a double-blind, randomised, phase 1 trial.** *Lancet Infect. Dis.* 2021; **21**: 637-646. [PubMed Abstract](#) | [Publisher Full Text](#)



The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact [research@f1000.com](mailto:research@f1000.com)

**F1000Research**



# Source details

## F1000Research

Open Access ⓘ

Scopus coverage years: from 2012 to Present

Publisher: Taylor & Francis

E-ISSN: 2046-1402

Subject area: [Pharmacology, Toxicology and Pharmaceutics: General Pharmacology, Toxicology and Pharmaceutics](#)

[Biochemistry, Genetics and Molecular Biology: General Biochemistry, Genetics and Molecular Biology](#)

View all ▾

Source type: Journal

CiteScore 2020

4.4



SJR 2020

1.099



SNIP 2020

0.921



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### Improved CiteScore methodology



CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020 ▾

$$4.4 = \frac{13.426 \text{ Citations 2017 - 2020}}{3.027 \text{ Documents 2017 - 2020}}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021 ⓘ

$$4.9 = \frac{14.958 \text{ Citations to date}}{3.026 \text{ Documents to date}}$$

Last updated on 06 March, 2022 - Updated monthly

### CiteScore rank 2020 ⓘ

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#13/67	81st
General Pharmacology, Toxicology and Pharmaceutics		
Biochemistry, Genetics and	#60/204	70th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)



Ads by Google

[Stop seeing this ad](#)
[Why this ad?](#) 

## F1000Research

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
United Kingdom 	Biochemistry, Genetics and Molecular Biology <ul style="list-style-type: none"> <li>Biochemistry, Genetics and Molecular Biology (miscellaneous)</li> </ul> Immunology and Microbiology <ul style="list-style-type: none"> <li>Immunology and Microbiology (miscellaneous)</li> </ul> Medicine <ul style="list-style-type: none"> <li>Medicine (miscellaneous)</li> </ul> Pharmacology, Toxicology and Pharmaceutics <ul style="list-style-type: none"> <li>Pharmacology, Toxicology and Pharmaceutics (miscellaneous)</li> </ul>		<b>60</b>
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	20461402	2012-2020	<a href="#">Homepage</a> <a href="#">How to publish in this journal</a> <a href="mailto:research@f1000.com">research@f1000.com</a>

## SCOPE

F1000Research publishes articles and other research outputs reporting basic scientific, scholarly, translational and clinical research across the physical and life sciences, engineering, medicine, social sciences and humanities. F1000Research is a scholarly publication platform set up for the scientific, scholarly and medical research community; each article has at least one author who is a qualified researcher, scholar or clinician actively working in their speciality and who has made a key contribution to the article. Articles must be original (not duplications). All research is suitable irrespective of the perceived level of interest or novelty; we welcome confirmatory and negative results, as well as null studies. F1000Research publishes different type of research, including clinical trials, systematic reviews, software tools, method articles, and many others. Reviews and Opinion articles providing a balanced and comprehensive overview of the latest discoveries in a particular field, or presenting a personal perspective on recent developments, are also welcome. See the full list of article types we accept for more information.

Join the conversation about this journal

Quartiles

Ads by Google

[Stop seeing this ad](#)
[Why this ad?](#)

Advances in Experimental  
Medicine and Biology  
USA

EBioMedicine  
NLD

Science Translational  
Medicine  
USA

Frontiers of Medicine  
CHN

Journal of Biomedical  
Science  
GBR


79%  
similarity

78%  
similarity

73%  
similarity

73%  
similarity

71%  
similarity



F1000Research



Show this widget in your own website

Just copy the code below and paste within your html code

<https://www.scimagojr.com>

SCImago Graphica

Explore, visually communicate and make sense of data with our new free tool.

Get it



Metrics based on Scopus® data as of April 2021



Henry Akool Najj 8 months ago

I have been waiting about one year for my article to publish or review but till now nothing happened

reply



Melanie Ortiz 8 months ago

SCImago Team

Dear Henry,  
Thank you for contacting us.  
Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff, so they could inform you more deeply.  
Best Regards, SCImago Team



alenu 12 months ago

Background: Nowadays, people die due to metabolic aging than chronological age worldwide.

Ads by Google

Stop seeing this ad Why this ad?